



**HAL**  
open science

# Superposition with subunitary powers in Sobolev spaces

Petru Mironescu

► **To cite this version:**

| Petru Mironescu. Superposition with subunitary powers in Sobolev spaces. 2014. hal-01069611v1

**HAL Id: hal-01069611**

**<https://hal.science/hal-01069611v1>**

Preprint submitted on 6 Oct 2014 (v1), last revised 10 Mar 2015 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Superposition with subunitary powers in Sobolev spaces

Petru Mironescu \*

29 septembre 2014

**Abstract.** Let  $0 < a < 1$  and set  $\Phi(t) = |t|^a, \forall t \in \mathbb{R}$ . Let  $1 < p < \infty$  and  $n \geq 1$ . We prove that the superposition operator  $u \mapsto \Phi(u)$  maps the Sobolev space  $W^{1,p}(\mathbb{R}^n)$  into the fractional Sobolev space  $W^{a,p/a}(\mathbb{R}^n)$ . We also investigate the case of more general nonlinearities.

**Résumé.** Superposition avec des puissances sousunitaires dans les espaces de Sobolev. Pour  $0 < a < 1$ , soit  $\Phi(t) = |t|^a, \forall t \in \mathbb{R}$ . Soient  $1 < p < \infty$  et  $n \geq 1$ . Nous montrons que l'opérateur de superposition  $u \mapsto \Phi(u)$  envoie l'espace de Sobolev  $W^{1,p}(\mathbb{R}^n)$  dans l'espace de Sobolev fractionnaire  $W^{a,p/a}(\mathbb{R}^n)$ . Nous examinons aussi la superposition avec des nonlinéarités plus générales.

## 1 Introduction

Let  $0 < a < 1$  and set

$$\Phi(t) = |t|^a, \forall t \in \mathbb{R}. \tag{1}$$

We consider the superposition operator

$$u \mapsto \Phi(u), \forall u \in W^{1,p}(\mathbb{R}^n),$$

and investigate its mapping properties. This question, natural in the context of the analysis of singularities of supercritical viscous Hamilton-Jacobi equations [3], was asked to me by L. Véron.

Our main result is the following

**Theorem 1.1** *Let  $1 < p < \infty$  and  $n \geq 1$ . The superposition operator  $u \mapsto \Phi(u)$  maps  $W^{1,p}(\mathbb{R}^n)$  into  $W^{a,p/a}(\mathbb{R}^n)$ .*

As we will see, the above theorem does not hold when  $p = 1$ .

An immediate consequence of Theorem 1.1 is the answer to the original question in [3].

**Corollary 1.2** *Let  $0 < a < 1$ ,  $1 < p < \infty$  and  $n \geq 1$ . Let  $\tilde{\Phi}(t) = |t|^a \operatorname{sgn} t, \forall t \in \mathbb{R}$ . The superposition operator  $u \mapsto \tilde{\Phi}(u)$  maps  $W^{1,p}(\mathbb{R}^n)$  into  $W^{a,p/a}(\mathbb{R}^n)$ .*

---

\*Université de Lyon, Université Lyon 1, CNRS UMR 5208, Institut Camille Jordan, 43 blvd. du 11 novembre 1918, F-69622 Villeurbanne cedex, France. Email address: mironescu@math.univ-lyon1.fr

Keywords. Sobolev spaces, functional calculus

2002 AMS classification subject. 46E35

The heart of the proof of Theorem 1.1 consists in establishing the estimate

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\Phi(u(y)) - \Phi(u(x))|^{p/a}}{|y-x|^{n+p}} dx dy \lesssim \int_{\mathbb{R}^n} |\nabla u(x)|^p dx, \quad \forall u \in W^{1,p}(\mathbb{R}^n). \quad (2)$$

Estimate (2) is a special case of a family of inequalities involving more general nonlinearities  $\Phi$ . Assume that  $\Phi$  is even and that we may write, on  $[0, \infty)$ ,

$$\begin{aligned} \Phi &= \zeta - g, \text{ with } \zeta \text{ continuous concave non decreasing} \\ &\text{such that } \zeta(0) = 0 \text{ and with } g \text{ non increasing.} \end{aligned} \quad (3)$$

This assumption is satisfied e.g. when  $\Phi$  is either continuous concave nondecreasing or Lipschitz on  $[0, \infty)$ . If (3) holds, then it is always possible to choose in (3) a function  $\zeta$  such that in addition

$$\zeta : [0, \infty) \rightarrow [0, \infty) \text{ is a concave homeomorphism} \quad (4)$$

and

$$\text{the restriction to } (0, \infty) \text{ of the inverse map } \Psi = \zeta^{-1} \text{ is } C^2. \quad (5)$$

[Indeed, every concave function  $\zeta$  as in (3) is dominated by a concave continuous function  $\tilde{\zeta} : [0, \infty) \rightarrow [0, \infty)$  such that  $\tilde{\zeta}(0) = 0$ ,  $\tilde{\zeta} \in C^\infty((0, \infty))$ ,  $\tilde{\zeta}'(t) > 0, \forall t > 0$  and  $\lim_{t \rightarrow \infty} \tilde{\zeta}(t) = \infty$ . Any such  $\tilde{\zeta}$  satisfies (4) and (5). In addition, if we define  $\tilde{g} = g + \zeta - \tilde{\zeta}$ , then  $u = \tilde{\zeta} + \tilde{g}$  and  $\tilde{g}$  is non decreasing.]

Note that when  $\Phi$  is given by (1) we may take  $g = 0$  and  $\zeta$  to be the restriction of  $\Phi$  to  $[0, \infty)$ . The same holds when  $\Phi$  is even and the restriction  $\zeta : [0, \infty) \rightarrow [0, \infty)$  of  $\Phi$  is a concave homeomorphism whose inverse satisfies (5).

Define

$$F(t) = \int_0^t \left( \int_0^s [\Psi'(\tau)]^{1-1/p} [\Psi''(\tau)]^{1/p} d\tau \right)^p ds, \quad \forall t \geq 0. \quad (6)$$

We have the following

**Theorem 1.3** *Let  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  be even and satisfy (3)-(5). Let  $F$  be given by (6). We have*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{F(|\Phi(u(y)) - \Phi(u(x))|)}{|y-x|^{n+p}} dx dy \lesssim \int_{\mathbb{R}^n} |\nabla u(x)|^p dx, \quad \forall u \in W^{1,p}(\mathbb{R}^n). \quad (7)$$

The reader may check that, when  $\Phi$  is as in (1) and  $\zeta$  is the restriction of  $\Phi$  to  $[0, \infty)$ , we have  $F(t) = C_{p,a} t^{p/a}$ , and thus (7) includes (2) as a special case.

## 2 Proofs

The key estimate is the following special case of (7):

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{F(|\Phi(v(y)) - \Phi(v(x))|)}{|y-x|^{1+p}} dx dy \lesssim \int_{\mathbb{R}} |v'(x)|^p dx, \quad \forall v \in C_c^\infty(\mathbb{R}), \quad \forall p \in (1, \infty). \quad (8)$$

Assuming (8) established for the moment, we proceed to the **Proof of Theorems 1.1 and 1.3, and of Corollary 1.2.**

1. We start from the following calculation, valid for every measurable function  $f : \mathbb{R}^n \times \mathbb{R}^n \rightarrow [0, \infty]$ .

$$\begin{aligned} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x, y) \, dx dy &= \frac{1}{2} \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}} |r|^{n-1} f(x, x + r\omega) \, dr ds_\omega dx \\ &= \frac{1}{2} \int_{\mathbb{S}^{n-1}} \int_{\omega^\perp} \int_{\mathbb{R}} \int_{\mathbb{R}} |r|^{n-1} f(x' + s\omega, x' + (r+s)\omega) \, dr ds dx' ds_\omega \\ &= \frac{1}{2} \int_{\mathbb{S}^{n-1}} \int_{\omega^\perp} \int_{\mathbb{R}} \int_{\mathbb{R}} |r-s|^{n-1} f(x' + s\omega, x' + r\omega) \, dr ds dx' ds_\omega. \end{aligned} \quad (9)$$

Let  $I$  denote the left-hand side of (7) and let  $u \in C_c^\infty(\mathbb{R}^n)$ . If we take, in (9),

$$f(x, y) = \frac{F(|\Phi(u(y)) - \Phi(u(x))|)}{|y - x|^{n+p}}$$

and apply (8) with  $v(r) = u(x' + r\omega)$ ,  $\forall r \in \mathbb{R}$ , we obtain that

$$\begin{aligned} I &\lesssim \int_{\mathbb{S}^{n-1}} \int_{\omega^\perp} \int_{\mathbb{R}} |\nabla u(x' + r\omega) \cdot \omega|^p \, dr dx' ds_\omega \leq \int_{\mathbb{S}^{n-1}} \int_{\omega^\perp} \int_{\mathbb{R}} |\nabla u(x' + r\omega)|^p \, dr dx' ds_\omega \\ &= |\mathbb{S}^{n-1}| \int_{\mathbb{R}^n} |\nabla u(x)|^p \, dx, \end{aligned}$$

i.e., (7) holds for  $u \in C_c^\infty(\mathbb{R}^n)$ .

2. By a standard argument, (7) holds for  $u \in W^{1,p}(\mathbb{R}^n)$ . In particular, (2) holds.

3. Next, Theorem 1.1 is a consequence of (2) and of the identity  $\|\Phi(u)\|_{L^{p/a}(\mathbb{R}^n)}^{p/a} = \|u\|_{L^p(\mathbb{R}^n)}^p$  (with  $\Phi$  as in (1)).

4. Corollary 1.2 follows from Theorem 1.1 combined with the identity  $\tilde{\Phi}(u) = \Phi(u^+) - \Phi(u^-)$  and with the fact that  $u^\pm \in W^{1,p}(\mathbb{R}^n)$  whenever  $u \in W^{1,p}(\mathbb{R}^n)$ .  $\square$

We next proceed to the

**Proof of (8).**

1. Starting from the inequality

$$0 \leq \int_0^s [\Psi'(\tau)]^{1-1/p} [\Psi''(\tau)]^{1/p} \, d\tau \leq \int_0^s [\Psi'(\tau) + \Psi''(\tau)] \, d\tau = \Psi(s) + \Psi'(s), \quad \forall s > 0,$$

we find that  $F \in C^1([0, \infty))$  and that

$$F'(t) = \left( \int_0^t [\Psi'(s)]^{1-1/p} [\Psi''(s)]^{1/p} \, ds \right)^p > 0, \quad \forall t > 0, \quad \text{while } F'(0) = 0. \quad (10)$$

The identity

$$[F'(t)]^{1-p} [F''(t)]^p = C_p [\Psi'(t)]^{p-1} \Psi''(t), \quad \forall t > 0, \quad (11)$$

is a consequence of (10) combined with the formula of  $F''(t)$  obtained by differentiating (10).

Integrating (11), we obtain

$$\int_0^t [F'(s)]^{1-p} [F''(s)]^p \, ds = C_p' [\Psi'(t)]^p, \quad \forall t > 0. \quad (12)$$

If we let, in (12),  $t = \zeta(\tau)$ , and take into account the fact that  $\zeta = \Psi^{-1}$  on  $(0, \infty)$ , we find that

$$\int_0^{\zeta(\tau)} [F'(s)]^{1-p} [F''(s)]^p \, ds [\zeta'(\tau)]^p = C_p', \quad \forall \tau > 0. \quad (13)$$

2. We next note the following Hardy type inequality. If  $U \subset \mathbb{R}$  is a bounded open set and if  $w \in C(\bar{U}) \cap C^1(U)$  satisfies  $w = 0$  on  $\partial U$ , then

$$\int_{\mathbb{R} \setminus U} \left( \int_U \frac{|w(x)|^p}{|y-x|^{1+p}} dx \right) dy \leq C_p \int_U |w'(x)|^p dx, \quad \forall p \in (1, \infty). \quad (14)$$

Indeed, by the subadditivity of (14) with respect to  $U$ , it suffices to prove (14) when  $U = (a, b)$ , and in that case integration in  $y$  and Hardy's inequality in  $x$  lead to

$$\int_{\mathbb{R} \setminus U} \left( \int_U \frac{|w(x)|^p}{|y-x|^{1+p}} dx \right) dy \leq C'_p \int_U \frac{|w(x)|^p}{[\text{dist}(x, \partial U)]^p} dx \leq C_p \int_U |w'(x)|^p dx.$$

[The idea of using Hardy type inequalities in the study of superposition operators originates in [1].]

3.  $\Phi$  being even, it suffices to establish (8) when  $v$  is replaced by  $|v|$ . By a standard argument, it then suffices to establish (8) when  $v \in C_c^\infty(\mathbb{R})$  and  $v \geq 0$ . Moreover, by (3) and the fact that  $F$  is non decreasing, we have

$$F(|\Phi(t) - \Phi(s)|) \leq F(|\zeta(t) - \zeta(s)|), \quad \forall s, t \in \mathbb{R}^+,$$

and therefore it suffices to establish (8) when  $\Phi$  is replaced by  $\zeta$ . In conclusion, it suffices to establish the estimate

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{F(|\zeta(v(y)) - \zeta(v(x))|)}{|y-x|^{1+p}} dx dy \lesssim \int_{\mathbb{R}} |v'(x)|^p dx, \quad \forall v \in C_c^\infty(\mathbb{R}, \mathbb{R}^+), \quad \forall p \in (1, \infty), \quad (15)$$

with  $\zeta$  satisfying (4) and (5).

4. With  $v \in C_c^\infty(\mathbb{R}, \mathbb{R}^+)$  and  $x, y \in \mathbb{R}$  such that  $v(y) < v(x)$ , we have

$$\begin{aligned} F(|\zeta(v(y)) - \zeta(v(x))|) &= F(\zeta(v(x)) - \zeta(v(y))) \\ &= -F(\zeta(v(x)) - \zeta(t)) \Big|_{t=v(y)}^{t=v(x)} = \int_{v(y)}^{v(x)} [g_t(x)]^p dt, \end{aligned} \quad (16)$$

where

$$\begin{aligned} g_t(x) &= \left[ -\frac{d}{dt} [F(\zeta(v(x)) - \zeta(t))] \right]^{1/p} \\ &= [\zeta'(t)]^{1/p} [F'(\zeta(v(x)) - \zeta(t))]^{1/p}, \quad \forall x \in \mathbb{R}, \quad \forall t \in (0, v(x)). \end{aligned} \quad (17)$$

5. Integrating (16), we find that the left-hand side  $J$  of (15) is given by

$$J = 2 \int_0^\infty \left( \int_{v(y) < t} \left( \int_{v(x) > t} \frac{[g_t(x)]^p}{|y-x|^{1+p}} dx \right) dy \right) dt. \quad (18)$$

6. Let (with  $t > 0$  fixed)  $U = \{x; v(x) > t\}$ . Note that  $U$  is bounded. By (17) and the last assertion in (10), we have  $g_t = 0$  on  $\partial U$ , and thus we may apply (14) to  $x \mapsto g_t(x)$ ,  $\forall x \in U$ .

7. Combining (14) (applied with  $U = \{x; v(x) > t\}$  and with  $w = g_t$ ) with (18) and using the change of variables  $\zeta(v(x)) - \zeta(t) = s$ , we find that

$$\begin{aligned} J &\lesssim \int_{\mathbb{R}} \left( \int_0^{v(x)} \zeta'(t) [F'(\zeta(v(x)) - \zeta(t))]^{1-p} [F''(\zeta(v(x)) - \zeta(t))]^p dt \right) [\zeta'(v(x))]^p |v'(x)|^p dx \\ &= \int_{\mathbb{R}} \left( \int_0^{\zeta(v(x))} [F'(s)]^{1-p} [F''(s)]^p ds \right) [\zeta'(v(x))]^p |v'(x)|^p dx. \end{aligned} \quad (19)$$

Next (13) (applied with  $\tau = \zeta(v(x))$ ) combined with (19) leads to (15). The proof of (8) is complete.  $\square$

We conclude this note by proving that Theorem 1.1 does not hold when  $p = 1$ . The key observation is that  $\Phi$  given by (1) does not satisfy the estimate

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|\Phi(v(y)) - \Phi(v(x))|^{1/a}}{|y-x|^2} dx dy \lesssim \int_{\mathbb{R}} |v'(x)| dx, \quad \forall v \in C_c^\infty(\mathbb{R}). \quad (20)$$

Indeed, argue by contradiction and assume that (20) holds. By a standard limiting argument, (20) still holds for  $v \in \text{BV}(\mathbb{R})$ . However, if we take  $v = \mathbb{1}_{(0,1)}$ , then  $v \in \text{BV}(\mathbb{R})$  and the reader may check that the left-hand side of (20) is infinite. Thus (20) does not hold, as claimed.

Using the fact that (20) does not hold and the invariance of the estimate (20) by scaling and translations, we may find a sequence  $(v_j)_{j \geq 1} \subset C_c^\infty(\mathbb{R})$  such that

$$\text{supp } v_j \subset (j, j+1), \quad \int_{\mathbb{R}} |v'_j(x)| dx = \frac{1}{j^2}, \quad (21)$$

and

$$\int_j^{j+1} \int_j^{j+1} \frac{|\Phi(v_j(y)) - \Phi(v_j(x))|^{1/a}}{|y-x|^2} dx dy \geq \frac{1}{j}. \quad (22)$$

Note that

$$\int_{\mathbb{R}} |v_j(x)| dx \leq \int_{\mathbb{R}} |v'_j(x)| dx = \frac{1}{j^2}. \quad (23)$$

Set  $v := \sum_{j \geq 1} v_j$ . Using (21), (22) and (23), we find that  $v \in W^{1,1}(\mathbb{R})$  and, in addition,

$$\begin{aligned} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|\Phi(v(y)) - \Phi(v(x))|^{1/a}}{|y-x|^2} dx dy &\geq \sum_{j \geq 1} \int_j^{j+1} \int_j^{j+1} \frac{|\Phi(v(y)) - \Phi(v(x))|^{1/a}}{|y-x|^2} dx dy \\ &= \sum_{j \geq 1} \int_j^{j+1} \int_j^{j+1} \frac{|\Phi(v_j(y)) - \Phi(v_j(x))|^{1/a}}{|y-x|^2} dx dy = \infty. \end{aligned}$$

Thus  $v \notin W^{a,1/a}(\mathbb{R})$ , i.e., Theorem 1.1 does not hold when  $p = 1$  and  $n = 1$ .

The case where  $p = 1$  and  $n \geq 2$  is straightforward. Let  $v$  be as above and let  $\zeta \in C_c^\infty(\mathbb{R}^{n-1})$ ,  $\zeta \not\equiv 0$ . Set  $u(x', x_n) = \zeta(x')v(x_n)$ ,  $\forall x' \in \mathbb{R}^{n-1}$ ,  $\forall x_n \in \mathbb{R}$ . Then clearly  $u \in W^{1,1}(\mathbb{R}^n)$ . However, we claim that  $u \notin W^{a,1/a}(\mathbb{R}^n)$ . Indeed, argue by contradiction and assume that  $u \in W^{a,1/a}(\mathbb{R}^n)$ . By the Fubini property of the space  $W^{a,1/a}(\mathbb{R}^n)$  [2, Section 2.5.13, Theorem p. 115], for a.e.  $x' \in \mathbb{R}^{n-1}$  we have  $u(x', \cdot) \in W^{a,1/a}(\mathbb{R})$ . In particular, we have  $v \in W^{a,1/a}(\mathbb{R})$ , which is a contradiction. Therefore, when  $p = 1$  and  $n \geq 1$  Theorem 1.1 does not hold.

## Acknowledgements

The author was supported by the ANR project « Harmonic Analysis at its Boundaries », ANR-12-BS01-0013-03.

## References

- [1] G. Bourdaud and Y. Meyer. Fonctions qui opèrent sur les espaces de Sobolev. *J. Funct. Anal.*, 97(2):351–360, 1991.
- [2] H. Triebel. *Theory of function spaces*, volume 78 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1983.
- [3] L. Véron, personal communication.