



HAL
open science

Boundary singularities of solutions of semilinear elliptic equations with critical Hardy potentials

Konstantinos Gkikas, Laurent Véron

► **To cite this version:**

Konstantinos Gkikas, Laurent Véron. Boundary singularities of solutions of semilinear elliptic equations with critical Hardy potentials. 2014. hal-01071467v1

HAL Id: hal-01071467

<https://hal.science/hal-01071467v1>

Preprint submitted on 8 Oct 2014 (v1), last revised 26 Feb 2015 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Boundary singularities of solutions of semilinear elliptic equations with critical Hardy potentials

Konstantinos T. Gkikas

Centro de Modelamiento Matemático
 Universidad de Chile, Santiago de Chile, Chile
email: kugkikas@gmail.com

Laurent Véron

Laboratoire de Mathématiques et Physique Théorique
 Université François Rabelais, Tours, FRANCE
email: veronl@univ-tours.fr

Abstract

We study the boundary behaviour of the of (E) $-\Delta u - \frac{\kappa}{d^2(x)}u + g(u) = 0$, where $0 < \kappa < \frac{1}{4}$ and g is a continuous nondecreasing function in a bounded convex domain of \mathbb{R}^N . We first construct the Martin kernel associated to the the linear operator $\mathcal{L}_\kappa = -\Delta - \frac{\kappa}{d^2(x)}$ and give a general condition for solving equation (E) with any Radon measure μ for boundary data. When $g(u) = |u|^{q-1}u$ we show the existence of a critical exponent $q_c = q_c(N, \kappa) > 1$: when $0 < q < q_c$ any measure is eligible for solving (E) with μ for boundary data; if $q \geq q_c$, a necessary and sufficient condition is expressed in terms of the absolute continuity of μ with respect to some Besov capacity. The same capacity characterizes the removable compact boundary sets. At end any positive solution (F) $-\Delta u - \frac{\kappa}{d^2(x)}u + |u|^{q-1}u = 0$ with $q > 1$ admits a boundary trace $Tr_{\partial\Omega}(u)$ which is a positive outer regular Borel measure $\bar{\mu} \approx (\mathcal{S}, \mu)$, with compact singular set $\mathcal{S} \subset \partial\Omega$, and regular part μ which is a positive Radon measure on $\mathcal{R} := \partial\Omega \setminus \mathcal{S}$. When $1 < q < q_c$ to any outer regular positive Borel (F, μ) , it is possible to construct a positive solution u of (F) such that $Tr_{\partial\Omega}(u) = \bar{\mu}$.

Contents

1	Introduction	2
2	The linear operator $\mathcal{L}_\kappa = -\Delta - \frac{\kappa}{d^2(x)}$	7
2.1	The eigenvalue problem	7
2.2	Preliminaries	10
2.3	\mathcal{L}_κ -harmonic measure	17
2.4	The Poisson kernel of \mathcal{L}_κ	22

3	The nonlinear problem with measures data	25
3.1	The linear boundary value problem with L^1 data	25
3.2	General nonlinearities	28
3.3	The power case	32
4	Isolated boundary singularities	42
4.1	The spherical \mathcal{L}_κ -harmonic problem	43
4.2	The nonlinear eigenvalue problem	45
4.3	Isolated boundary singularities	47
5	The boundary trace of positive solutions	49
5.1	Construction of the boundary trace	50
5.2	Subcritical case	58
6	Appendix I: barriers and a priori estimates	66
6.1	Barriers	66
6.2	A priori estimates	70
7	Appendix II: The associated semilinear equation with singular potential	73

Key words: Semilinear elliptic equation; Hardy potentials; Harmonic measure; Singular integrals; Besov capacities; Boundary singularities; Boundary trace.

MSC2010: Primary 35J66, 35J10. Secondary 31A15, 35H25, 28A12.

1 Introduction

Let Ω be a bounded smooth domain in \mathbb{R}^N and $d(x) = \text{dist}(x, \Omega^c)$. In this article we study several aspects of the nonlinear boundary value associated to the equation

$$-\Delta u - \frac{\kappa}{d^2(x)}u + |u|^{p-1}u = 0 \quad \text{in } \Omega \quad (1.1)$$

where $p > 1$. The study of the boundary trace of solutions of (1.1) is a natural framework for a general study of several nonlinear problems where the nonlinearity, the geometric properties of the domain and the coefficient κ interact. On this point of view, the case $\kappa = 0$ has been thoroughly treated by Marcus and Véron [18], [19], [21], [20], for example and the synthesis presented in [22]. The associated linear Schrödinger operator

$$u \mapsto \mathcal{L}_\kappa u := -\Delta u - \frac{\kappa}{d^2(x)}u \quad (1.2)$$

plays an important role in functional analysis because of the particular singularity of its potential $V(x) := -\frac{\kappa}{d^2(x)}$. The case $\kappa < 0$ and more generally of nonnegative potential has been studied by Ancona [2] who has shown the existence of a Martin kernel which allows a general representation formula of nonnegative solutions of

$$\mathcal{L}_\kappa u = 0 \quad \text{in } \Omega, \quad (1.3)$$

This representation turned out to be the key ingredient of the full classification of positive solutions of

$$-\Delta u + u^q = 0 \quad \text{in } \Omega \quad (1.4)$$

which was obtained by Marcus [17]. In a more general setting, Véron and Yarur [27] constructed a capacity theory associated to the linear equation

$$\mathcal{L}_V u := -\Delta u + V(x)u = 0 \quad \text{in } \Omega, \quad (1.5)$$

where the potential V is nonnegative and singular near $\partial\Omega$. When $V(x) := -\frac{\kappa}{d^2(x)}$ with $\kappa > 0$, V is called a Hardy potential. There is a critical value $\kappa = \frac{1}{4}$. If $\kappa > \frac{1}{4}$, no positive solution of (1.3) exists. When $0 < \kappa \leq \frac{1}{4}$, there exist positive solutions and the geometry of the domain plays a fundamental role in the study of the mere linear equation (1.3). We define the constant c_Ω by

$$c_\Omega = \inf_{v \in H_0^1(\Omega) \setminus \{0\}} \frac{\int_\Omega |\nabla v|^2 dx}{\int_\Omega \frac{v^2}{d^2(x)} dx}. \quad (1.6)$$

It is known that c_Ω belongs to $(0, \frac{1}{4}]$. If Ω is convex or if the distance function d is super harmonic in the sense of distributions, then $c_\Omega = \frac{1}{4}$. Furthermore there holds $c_\Omega = \frac{1}{4}$ if and only if problem (1.6) has no minimizer. (see [16]). When $0 < \kappa \leq \frac{1}{4}$, which is always assumed in the sequel and there exists no minimizer to problem (1.6), it is possible to define the first eigenvalue λ_κ of the operator \mathcal{L}_κ . If we define the two fundamental exponents α_+ and α_- by

$$\alpha_+ = 1 + \sqrt{1 - 4\kappa} \quad \text{and} \quad \alpha_- = 1 - \sqrt{1 - 4\kappa} \quad (1.7)$$

then the first eigenvalue is achieved by an eigenfunction ϕ_κ which satisfies $\phi_\kappa(x) \approx d^{\frac{\alpha_+}{2}}(x)$ as $d(x) \rightarrow 0$. Similarly, the Green kernel $G_{\mathcal{L}_\kappa}$ associated to \mathcal{L}_κ inherits this type of boundary behaviour since there holds

$$\frac{1}{C_\kappa} \min \left\{ \frac{1}{|x-y|^{N-2}}, \frac{d^{\frac{\alpha_+}{2}}(x)d^{\frac{\alpha_+}{2}}(y)}{|x-y|^{N+\alpha_+-2}} \right\} \leq G_{\mathcal{L}_\kappa}(x, y) \leq C_\kappa \min \left\{ \frac{1}{|x-y|^{N-2}}, \frac{d^{\frac{\alpha_+}{2}}(x)d^{\frac{\alpha_+}{2}}(y)}{|x-y|^{N+\alpha_+-2}} \right\} \quad (1.8)$$

We show that \mathcal{L}_κ satisfies the maximum principle in the sense that if $u \in H_{loc}^1 \cap C(\Omega)$ is a subsolution i.e. $\mathcal{L}_\kappa u \leq 0$ such that

$$\begin{aligned} (i) \quad & \limsup_{x \rightarrow y} \frac{u(x)}{d^{\alpha_-}(x)} \leq 0 \quad \text{if } 0 < \kappa < \frac{1}{4} \\ (ii) \quad & \limsup_{x \rightarrow y} \frac{u(x)}{\sqrt{d(x)} |\ln d(x)|} \leq 0 \quad \text{if } \kappa = \frac{1}{4} \end{aligned} \quad (1.9)$$

for all $y \in \partial\Omega$, then $u \leq 0$. If $\xi \in \partial\Omega$ and $r > 0$, we set $\Delta_r(\xi) = \partial\Omega \cap B_r(\xi)$. We prove that a positive solution of $\mathcal{L}_\kappa u = 0$ which vanishes on a part of the boundary in the sense that

$$\begin{aligned} (i) \quad & \lim_{x \rightarrow y} \frac{u(x)}{d^{\alpha_-}(x)} = 0 \quad \forall y \in \Delta_r(\xi) \quad \text{if } 0 < \kappa < \frac{1}{4} \\ (ii) \quad & \lim_{x \rightarrow y} \frac{u(x)}{\sqrt{d(x)} |\ln d(x)|} = 0 \quad \forall y \in \Delta_r(\xi) \quad \text{if } \kappa = \frac{1}{4}, \end{aligned} \quad (1.10)$$

satisfies

$$\frac{u(x)}{\phi_\kappa(x)} \leq C_1 \frac{u(y)}{\phi_\kappa(y)} \quad \forall x, y \in \Delta_{\frac{r}{2}}(\xi), \quad (1.11)$$

for some $C_1 = C_1(\Omega, \kappa) > 0$.

For any $h \in C(\partial\Omega)$ we construct the unique solution $v := v_h$ of the Dirichlet problem

$$\begin{aligned} \mathcal{L}_\kappa v &= 0 & \text{in } \Omega \\ v &= h & \text{on } \partial\Omega \end{aligned} \quad (1.12)$$

Using this construction and estimates (1.9) we show the existence of the \mathcal{L}_κ -measure, which is a Borel measure ω^x with the property that for any $h \in C(\partial\Omega)$, the above function v_h satisfies

$$v_h(x) = \int_{\partial\Omega} h(y) d\omega^x(y). \quad (1.13)$$

Because of Harnack inequality, the measures ω^x and ω^z are mutually absolutely continuous for $x, z \in \Omega$ and for any $x \in \Omega$ we can define the Radon-Nikodym derivative

$$K(x, y) := \frac{d\omega^x}{d\omega^{x_0}}(y) \quad \text{for } \omega^{x_0}\text{-almost } y \in \partial\Omega. \quad (1.14)$$

There exists $r_0 := r_0(\Omega)$ such that for any $x \in \Omega$ such that $d(x) \leq r_0$, there exists a unique $\xi = \xi_x \in \partial\Omega$ such that $d(x) = |x - \xi_x|$. If we denote by Ω'_{r_0} the set of $x \in \Omega$ such that $0 < d(x) < r_0$, the mapping Π from $\overline{\Omega}'_{r_0}$ to $[0, r_0] \times \partial\Omega$ defined by $\Pi(x) = (d(x), \xi_x)$ is a C^1 diffeomorphism. If $\xi \in \partial\Omega$ and $0 \leq r \leq r_0$, we set $x_r(\xi) = \Pi^{-1}(r, \xi)$. Let W be defined in Ω by

$$W(x) = \begin{cases} d^{\frac{\alpha-}{2}}(x) & \text{if } \kappa < \frac{1}{4} \\ \sqrt{d(x)} |\ln d(x)| & \text{if } \kappa = \frac{1}{4}, \end{cases} \quad (1.15)$$

we prove that the \mathcal{L}_κ -harmonic measure can be equivalently defined by

$$\omega^x(E) = \inf \left\{ \psi : \psi \in C_+(\Omega), \mathcal{L}_\kappa\text{-superharmonic in } \Omega \text{ and s.t. } \liminf_{x \rightarrow E} \frac{\psi(x)}{W(x)} \geq 1 \right\} \quad (1.16)$$

for any compact set $E \subset \partial\Omega$ and then extended classically to Borel subsets of $\partial\Omega$.

The \mathcal{L}_κ -harmonic measure is connected to the Green kernel of \mathcal{L}_κ by the following estimates

Theorem A *There exists $C_3 := C_3(\Omega) > 0$ such that for any $r \in (0, r_0]$ and $\xi \in \partial\Omega$, there holds*

$$\frac{1}{C_3} r^{N+\alpha-2} G_{\mathcal{L}_\kappa}(x_r(\xi), x) \leq \omega^x(\Delta_r(\xi)) \leq C_3 r^{N+\alpha-2} G_{\mathcal{L}_\kappa}(x_r(\xi), x) \quad \forall x \in \Omega \setminus B_{4r}(\xi) \quad (1.17)$$

if $0 < \kappa < \frac{1}{4}$, and

$$\frac{1}{C_3} r^{N-1} |\ln d(x)| G_{\mathcal{L}_{\frac{1}{4}}}(x_r(\xi), x) \leq \omega^x(\Delta_r(\xi)) \leq C_3 r^{N-1} |\ln d(x)| G_{\mathcal{L}_{\frac{1}{4}}}(x_r(\xi), x) \quad \forall x \in \Omega \setminus B_{4r}(\xi). \quad (1.18)$$

As a consequence ω^x has the doubling property. The previous estimates allow to construct a kernel function of \mathcal{L}_κ in Ω , prove its uniqueness up to an homothety. When normalized, the kernel function denoted by $K_{\mathcal{L}_\kappa}$ is the Martin kernel, defined by

$$K_{\mathcal{L}_\kappa}(x, \xi) = \lim_{x \rightarrow \xi} \frac{G_{\mathcal{L}_\kappa}(x, y)}{G_{\mathcal{L}_\kappa}(x, x_0)} \quad \forall \xi \in \partial\Omega. \quad (1.19)$$

for some $x_0 \in \Omega$. An important property of the Martin kernel is that it allows to represent a positive \mathcal{L}_κ -harmonic function u by mean of a Poisson type formula which endows the form

$$u(x) = \int_{\partial\Omega} K_{\mathcal{L}_\kappa}(x, \xi) d\mu(\xi) \quad \text{for } \omega^{x_0}\text{-almost } x \in \Omega. \quad (1.20)$$

for some positive Radon measure μ on $\partial\Omega$. The measure μ is called the boundary trace of u . Furthermore $K_{\mathcal{L}_\kappa}$ satisfies the following two-side estimates

Theorem B *There exists $C_3 := C_3(\Omega, \kappa) > 0$ such that for any $(x, \xi) \in \Omega \times \partial\Omega$ there holds*

$$\frac{1}{C_3} \frac{d^{\frac{\alpha_+}{2}}}{|x - \xi|^{N+\alpha_+-2}} \leq K_{\mathcal{L}_\kappa}(x, \xi) \leq C_3 \frac{d^{\frac{\alpha_+}{2}}}{|x - \xi|^{N+\alpha_+-2}}. \quad (1.21)$$

Thanks to these estimates we can adapt the approach developed in [13] to prove the existence of weak solutions to the nonlinear boundary value problem

$$\begin{aligned} -\Delta u - \frac{\kappa}{d^2(x)} u + g(u) &= \nu & \text{in } \Omega \\ u &= \mu & \text{in } \partial\Omega, \end{aligned} \quad (1.22)$$

where g is a continuous nondecreasing function such that $g(0) \geq 0$ and ν and μ are Radon measures on Ω and $\partial\Omega$ respectively. We define the class $\mathbf{X}_\kappa(\Omega)$ of test functions by

$$\mathbf{X}_\kappa(\Omega) = \left\{ \eta \in L^2(\Omega) \text{ s.t. } \nabla(d^{-\frac{\alpha_+}{2}} \eta) \in L^2_{d^{\alpha_+}}(\Omega) \text{ and } \phi_\kappa^{-1} \mathcal{L}_\kappa \eta \in L^\infty(\Omega) \right\} \quad (1.23)$$

and we prove

Theorem C *Assume g satisfies*

$$\int_1^\infty (g(s) + |g(-s)|) s^{-2\frac{N-1+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} ds < \infty. \quad (1.24)$$

Then for any Radon measures ν on Ω and such that $\int_\Omega \phi_\kappa d|\mu| < \infty$ and μ on $\partial\Omega$ there exists a unique $u \in L^1_{\phi_\kappa}$ such that $g(u) \in L^1_{\phi_\kappa}$ which satisfies

$$\int_\Omega (u \mathcal{L}_\kappa \eta + g(u) \eta) dx = \int_\Omega (\eta d\nu + \mathbb{K}_{\mathcal{L}_\kappa}[\mu] \mathcal{L}_\kappa \eta dx) \quad \forall \eta \in \mathbf{X}_\kappa(\Omega). \quad (1.25)$$

When $g(r) = |r|^{q-1}r$ the critical value is $q_c = \frac{N+\frac{\alpha_+}{2}}{N+\frac{\alpha_+}{2}-2}$ and (1.24) is satisfied for $0 \leq q < q_c$. When $q \geq q_c$ not all the Radon measures are eligible for solving

$$\begin{aligned} -\Delta u - \frac{\kappa}{d^2(x)} u + |u|^{q-1} u &= 0 & \text{in } \Omega \\ u &= \mu & \text{in } \partial\Omega, \end{aligned} \quad (1.26)$$

We prove the following result in which statement $C_{2-\frac{2+\alpha_+}{2q'}, q'}^{\mathbb{R}^{N-1}}$ denotes the Besov capacity associated to the Besov space $B^{2-\frac{2+\alpha_+}{2q'}, q'}(\mathbb{R}^{N-1})$.

Theorem D *Assume $q \geq q_c$ and μ is a positive Radon measure on $\partial\Omega$. Then problem (1.26) admits a weak solution if and only if μ vanishes on Borel sets $E \subset \partial\Omega$ such that $C_{2-\frac{2+\alpha_+}{2q'}, q'}^{\mathbb{R}^{N-1}}(E) = 0$.*

This capacity plays a fundamental for characterizing the removable compact boundary sets which exist only in the *supercritical range* $q \geq q_c$.

Theorem E Assume $q \geq q_c$ and $K \subset \partial\Omega$ is compact. Then any function $u \in C(\overline{\Omega} \setminus K)$ which satisfies

$$\begin{aligned} -\Delta u - \frac{\kappa}{d^2(x)}u + |u|^{q-1}u &= 0 & \text{in } \Omega \\ u &= 0 & \text{in } \partial\Omega \setminus K, \end{aligned} \quad (1.27)$$

is identically zero if and only if $C_{2-\frac{2+\alpha_+}{2q'}, q'}^{\mathbb{R}^{N-1}}(K) = 0$.

We show that any positive solution u of (1.1) admits a boundary trace, and more precisely we prove that the following dichotomy holds: et

Theorem F Let u be a positive solution of (1.1) in Ω and $a \in \partial\Omega$. Then

(i) either for any $\epsilon > 0$

$$\lim_{\delta \rightarrow 0} \int_{\Sigma_\delta \cap B_\epsilon(a)} u d\omega_{\Omega'_\delta}^{x_0} = \infty, \quad (1.28)$$

where $\Omega'_\delta = \{x \in \Omega : d(x) > \delta\}$, $\Sigma_\delta = \partial\Omega'_\delta$ and $\omega_{\Omega'_\delta}^{x_0}$ is the harmonic measure in Ω'_δ ,

(ii) or there exists $\epsilon_0 > 0$ and a positive Radon measure λ on $\partial\Omega \cap B_{\epsilon_0}(a)$ such that for any $Z \in C(\overline{\Omega})$ with support in $\Omega \cup (\partial\Omega \cap B_{\epsilon_0}(a))$, there holds

$$\lim_{\delta \rightarrow 0} \int_{\Sigma_\delta \cap B_{\epsilon_0}(a)} Z u d\omega_{\Omega'_\delta}^{x_0} = \int_{\partial\Omega \cap B_{\epsilon_0}(a)} Z d\lambda. \quad (1.29)$$

The set of points $a \in \partial\Omega$ such that (i) (resp. (ii)) holds is closed (resp. relatively open) and denoted by \mathcal{S}_u (resp. \mathcal{R}_u). There exists a unique radon measure μ_u on \mathcal{R}_u such that, for any $Z \in C(\overline{\Omega})$ with support in $\Omega \cup \mathcal{R}_u$ there holds

$$\lim_{\delta \rightarrow 0} \int_{\Sigma_\delta} Z u d\omega_{\Omega'_\delta}^{x_0} = \int_{\mathcal{R}_u} Z d\mu_u. \quad (1.30)$$

The couple (\mathcal{S}_u, μ_u) is called *the boundary trace of u* and denoted by $Tr_{\partial\Omega}(u)$.

If $1 < q < q_c$ we denote by $u_{k\delta_a}$ positive solution of (1.1) with $\mu = k\delta_a$ for some $a \in \partial\Omega$ and $k \geq 0$. There exists $\lim_{k \rightarrow \infty} u_{k\delta_a} = u_{\infty, a}$. We prove the following

Theorem G Assume $1 < q < q_c$ and $a \in \partial\Omega$. Then If u is a positive solution of (1.1) such that $a \in \mathcal{S}_u$, then $u \geq u_{\infty, a}$.

In order to go further in the study of boundary singularities, we construct separable solutions of (1.1) in $\mathbb{R}_+^N = \{x = (x', x_N) : x_N > 0\} = \{(r, \sigma) \in \mathbb{R}_+ \times S_+^{N-1}\}$ which vanish on $\partial\mathbb{R}_+^N \setminus \{0\}$ under the form $u(r, \sigma) = r^{-\frac{2}{q-1}}\omega(\sigma)$, where $r > 0$, $\sigma \in S_+^{N-1}$. They are solutions of

$$\begin{aligned} -\Delta_{S^{N-1}}\omega - \ell_{q, N}\omega - \frac{\kappa}{e_{N, \sigma}}\omega + |\omega|^{q-1}\omega &= 0 & \text{in } S_+^{N-1} \\ \omega &= 0 & \text{in } \partial S_+^{N-1} \end{aligned} \quad (1.31)$$

where $\Delta_{S^{N-1}}$ is the Laplace-Beltrami operator, e_N the unit vector pointing toward the North pole and $\ell_{q, N}$ is a positive constant. We prove that if $1 < q < q_c$ problem (1.31) admits a unique positive solution

ω_κ while no such solution exists if $q \geq q_c$. To this phenomenon is associated a result of classification of positive solutions of (1.1) in Ω which vanishes of $\partial\Omega \setminus \{0\}$ (here we assume that $0 \in \partial\Omega$ and that the tangent plane to $\partial\Omega$ at 0 is $\{x : x \cdot e_N = 0\}$ and that $\Omega \subset \{x : x \cdot e_N \geq 0\}$)

Theorem H Assume $1 < q < q_c$ and let $u \in C(\bar{\Omega} \setminus \{a\})$ be a solution of (1.1) in Ω which vanishes of $\partial\Omega \setminus \{a\}$. Then

(i) Either $u = u_{\infty, a}$ and

$$\lim_{r \rightarrow 0} r^{\frac{2}{q-1}} u(r, \cdot) = \omega_\kappa \quad (1.32)$$

locally uniformly in S_+^{N-1} .

(ii) Or there exists $k \geq 0$ such that $u = u_{k\delta_a}$ and

$$u(x) = kK_{\mathcal{L}_\kappa}(x, a)(1 + o(1)) \quad \text{as } x \rightarrow 0 \quad (1.33)$$

If $1 < q < q_c$ we prove that to any couple (F, μ) where F is a closed subset of $\partial\Omega$ and μ a positive Radon measure on $R = \partial\Omega \setminus F$ we can associate a positive solution u of (1.1) in Ω with $Tr_{\partial\Omega}(u) = (F, \mu)$.

2 The linear operator $\mathcal{L}_\kappa = -\Delta - \frac{\kappa}{d^2(x)}$

Throughout this article c_j ($j=1,2,\dots$) denote positive constants the value of which may change from one occurrence to another. The notation κ is reserved to the value of the coefficient of the Hardy potential

2.1 The eigenvalue problem

We recall some known results concerning the eigenvalue problem (see [9], [12]).

1- Since Ω is convex, $c_\Omega = \frac{1}{4}$ and for any $\kappa \in (0, \frac{1}{4}]$ there exists

$$\lambda_\kappa = \inf_{u \in H_0^1(\Omega)} \frac{\int_\Omega (|\nabla u|^2 - \frac{\kappa}{d^2} u^2) dx}{\int_\Omega u^2 dx}.$$

2- If $0 < \kappa < \frac{1}{4}$ the minimizer ϕ_κ belongs $H_0^1(\Omega)$ and it satisfies

$$\phi_\kappa \approx d^{\frac{\alpha_+}{2}}(x), \quad (2.1)$$

where α_+ (as well as α_-) are defined by (1.7).

3- If $\kappa = \frac{1}{4}$, there exists a non-negative minimizer $\phi_{\frac{1}{4}} \in H_{loc}^1(\Omega)$ such that

$$\phi_{\frac{1}{4}} \approx d^{\frac{1}{2}}(x). \quad (2.2)$$

Furthermore, the function $\psi_{\frac{1}{4}} = d^{-\frac{1}{2}}$ belongs to $H_0^1(\Omega; d(x)dx)$

4- Let $H_0^1(\Omega, d^\alpha(x)dx)$ denote the closure of $C_0^\infty(\Omega)$ functions under the norm

$$\|u\|_{H_0^1(\Omega, d^\alpha(x)dx)}^2 = \int_\Omega |\nabla u|^2 d^\alpha(x)dx + \int_\Omega |u|^2 d^\alpha(x)dx. \quad (2.3)$$

If $\alpha \geq 1$ there holds [12, Th. 2.11]

$$H_0^1(\Omega, d^\alpha(x)dx) = H^1(\Omega, d^\alpha(x)dx) \quad \forall \alpha \geq 1. \quad (2.4)$$

5- Let $0 < \kappa \leq \frac{1}{4}$. Let $\mathbf{H}_\kappa(\Omega)$ be the subset of functions of $H_{loc}^1(\Omega)$ satisfying

$$\int_{\Omega} \left(|\nabla \phi|^2 - \frac{\kappa}{d^2} \phi^2 \right) dx < \infty. \quad (2.5)$$

Then the mapping

$$\phi \mapsto \left(\int_{\Omega} \left(|\nabla \phi|^2 - \frac{\kappa}{d^2} \phi^2 \right) dx \right)^{\frac{1}{2}} \quad (2.6)$$

is a norm on $\mathbf{H}_\kappa(\Omega)$. The closure $\mathbf{W}_\kappa(\Omega)$ of $C_0^\infty(\Omega)$ into $\mathbf{H}_\kappa(\Omega)$ satisfies

$$\mathbf{W}_\kappa(\Omega) = H_0^1(\Omega) \quad \forall 0 < \kappa < \frac{1}{4} \text{ and } \mathbf{W}_{\frac{1}{4}}(\Omega) \subset W_0^{1,q}(\Omega) \quad \forall 1 \leq q < 2, \quad (2.7)$$

see [6, Th B]. As a consequence $\mathbf{W}_\kappa(\Omega)$ is compactly imbedded into $L^r(\Omega)$ for any $r \in [1, 2^*)$.

6- Let $\alpha > 0$ and $\Omega \subset \mathbb{R}^N$ be a bounded domain. There exists $c^* > 0$ depending on $\text{diam}(\Omega)$, N and α such that for any $v \in C_0^\infty(\Omega)$

$$\left(\int_{\Omega} |v|^{\frac{2(N+\alpha)}{N+\alpha-2}} d^\alpha dx \right)^{\frac{N+\alpha-2}{N+\alpha}} \leq c^* \int_{\Omega} |\nabla v|^2 d^\alpha dx. \quad (2.8)$$

For a proof see [12, Th. 2.9].

The boundary behaviour of the first eigenfunction yield two-side similar estimates of the Green kernel for Schrödinger operators with a general Hardy type potentials [12, Corollary 1.9].

Proposition 2.1. *Consider the operator $E := -\Delta - V$, in Ω where $V = V_1 + V_2$, with*

$$|V_1| \leq \frac{1}{4d^2(x)} \quad \text{and } V_2 \in L^p(\Omega), \quad p > \frac{N}{2}.$$

We also assume that

$$0 < \lambda_1 := \inf_{u \in H_0^1(\Omega)} \frac{\int_{\Omega} (|\nabla u|^2 dx - V u^2) dx}{\int_{\Omega} u^2 dx},$$

and that to λ_1 is associated a positive eigenfunction ϕ_1 . If, for some $\alpha \geq 1$ and $C_1, C_2 > 0$, there holds

$$c_1 d^{\frac{\alpha}{2}}(x) \leq \phi_1(x) \leq c_2 d^{\frac{\alpha}{2}}(x) \quad \forall x \in \Omega,$$

then the Green kernel G_E^Ω associated to E in Ω satisfies

$$G_E^\Omega(x, y) \approx c_3 \min \left(\frac{1}{|x-y|^{N-2}}, \frac{d^{\frac{\alpha}{2}}(x) d^{\frac{\alpha}{2}}(y)}{|x-y|^{N+\alpha-2}} \right). \quad (2.9)$$

We set

$$\Omega_\delta = \{x \in \Omega : d(x) < \delta\}, \quad \Omega'_\delta = \{x \in \Omega : d(x) > \delta\} \text{ and } \Sigma_\delta = \{x \in \Omega : d(x) = \delta\}. \quad (2.10)$$

Definition 2.2. Let $G \subset \Omega$ and let $H_c^1(G) \subset H^1(G)$ denote the subspace of functions with compact support. A function $h \in W_{loc}^{1,1}(G)$ is \mathcal{L}_κ -harmonic in G if

$$\int_G \nabla h \cdot \nabla \psi dx - \kappa \int_\Omega \frac{1}{d^2(x)} h \psi dx = 0 \quad \forall \psi \in H_c^1(G).$$

A function $\underline{h} \in H_{loc}^1(G) \cap C(G)$ is \mathcal{L}_κ -subharmonic in G if

$$\int_G \nabla \underline{h} \cdot \nabla \psi dx - \kappa \int_\Omega \frac{1}{d^2(x)} \underline{h} \psi dx \leq 0 \quad \forall \psi \in H_c^1(G), \psi \geq 0.$$

We say that \underline{h} is a local \mathcal{L}_κ -subharmonic function if there exists $\delta > 0$ such that $\underline{h} \in H_{loc}^1(\Omega_\delta) \cap C(\Omega_\delta)$ is \mathcal{L}_κ -subharmonic in Ω_δ . Similarly, (local) \mathcal{L}_κ -superharmonics \bar{h} are defined with " \geq " in the above inequality.

Note that \mathcal{L}_κ -harmonic functions are C^∞ in G by standard elliptic equations regularity theory. The Phragmen-Lindelöf principle yields the following alternative.

Proposition 2.3. Let $\kappa \leq \frac{1}{4}$. If \underline{h} is a local \mathcal{L}_κ -subharmonic function, then the following alternative holds:

(i) either for every local positive \mathcal{L}_κ -superharmonic function \bar{h}

$$\limsup_{d(x) \rightarrow 0} \frac{\underline{h}(x)}{\bar{h}(x)} > 0, \quad (2.11)$$

(ii) or for every local positive \mathcal{L}_κ -superharmonic function \bar{h}

$$\limsup_{d(x) \rightarrow 0} \frac{\underline{h}(x)}{\bar{h}(x)} < \infty. \quad (2.12)$$

Definition 2.4. If a local \mathcal{L}_κ -subharmonic function \underline{h} satisfies (i) (resp. (ii)) it is called a large \mathcal{L}_κ -subharmonic ((resp. a small \mathcal{L}_κ -subharmonic)).

The next statement is [3, Theorem 2.9].

Proposition 2.5. Let \underline{h} be a small local \mathcal{L}_κ -subharmonic of \mathcal{L}_κ .

(i) If $\kappa < \frac{1}{4}$ then the following alternative holds:

$$\text{either } \limsup_{x \rightarrow \partial\Omega} \frac{\underline{h}}{d^{\frac{\alpha_-}{2}}} > 0 \quad \text{or} \quad \limsup_{x \rightarrow \partial\Omega} \frac{\underline{h}}{d^{\frac{\alpha_+}{2}}} < \infty.$$

(ii) If $\kappa = \frac{1}{4}$ then the following alternative holds:

$$\text{either } \limsup_{x \rightarrow \partial\Omega} \frac{\underline{h}}{d^{\frac{1}{2}} \log(\frac{1}{d})} > 0 \quad \text{or} \quad \limsup_{x \rightarrow \partial\Omega} \frac{\underline{h}}{d^{\frac{1}{2}}} < \infty.$$

Definition 2.6. Let $f_0 \in L_{loc}^2(\Omega)$. We say that a function $u \in H_{loc}^1(\Omega)$ is a solution of

$$\mathcal{L}_\kappa u = f_0 \quad \text{in } \Omega \quad (2.13)$$

if there holds

$$\int_\Omega \nabla u \cdot \nabla \psi dx - \kappa \int_\Omega \frac{1}{d^2(x)} u \psi dx = \int_\Omega f_0 \psi dx \quad \forall \psi \in C_0^\infty(\Omega). \quad (2.14)$$

2.2 Preliminaries

In this part we study some regularity properties of solutions of linear equations involving \mathcal{L}_κ .

Lemma 2.7. (i) If $\alpha > 1$ and $d^{-\frac{\alpha}{2}}u \in H^1(\Omega, d^\alpha(x)dx)$, then $u \in H_0^1(\Omega)$.

(ii) If $\alpha = 1$ and $d^{-\frac{1}{2}}u \in H^1(\Omega, d(x)dx)$, then $u \in W_0^{1,p}(\Omega)$, $\forall p < 2$.

Proof. There exists $\beta_0 > 0$ such that $d \in C^2(\overline{\Omega_{\beta_0}})$ and set $u = d^{\frac{\alpha}{2}}v$. In the two cases (i)-(ii), our assumptions imply

$$u \in L^2(\Omega) \quad \text{and} \quad \nabla u - \frac{\alpha}{2}ud^{-1}\nabla d \in L^2(\Omega). \quad (2.15)$$

(i) Let $0 < \varepsilon \leq \frac{\beta}{2} \leq \frac{\beta_0}{2}$ and ψ_ε be a cut of function such that $\psi_\varepsilon = 0$ in $\Omega_{\frac{\varepsilon}{2}} \cup \Omega_\beta$ and $\psi_\varepsilon = 1$ in $\Omega_{\frac{\beta}{2}} \setminus \overline{\Omega_\varepsilon}$. Then $u = d^{\frac{\alpha}{2}}(\psi_\varepsilon v + (1 - \psi_\varepsilon)v)$. Thus it is enough to prove that $u_\varepsilon = d^{\frac{\alpha}{2}}\psi_\varepsilon v$ remains bounded in $H^1(\Omega)$ independently of ε . Set $v_\varepsilon = \psi_\varepsilon v$, then

$$\int_\Omega |\nabla u_\varepsilon|^2 dx = \int_{\Omega_\beta} |\nabla u_\varepsilon|^2 dx \leq c_4 \left(\int_{\Omega_\beta} d^\alpha |\nabla v_\varepsilon|^2 dx + \int_{\Omega_\beta} d^{\alpha-2} v_\varepsilon^2 dx \right).$$

Note that $\alpha - 2 > -1$. Now

$$\int_{\Omega_\beta} d^{\alpha-2} v_\varepsilon^2 dx = \frac{1}{\alpha-1} \int_{\Omega_\beta} v_\varepsilon^2 \operatorname{div}(d^{\alpha-1} \nabla d) dx - \frac{1}{\alpha-1} \int_{\Omega_\beta} (d^{\alpha-1} (\Delta d)) v_\varepsilon^2 dx.$$

Now since $|\Delta d(x)| < c_5$, $\forall x \in \Omega_{\beta_0}$, we have

$$\left| \frac{1}{\alpha-1} \int_{\Omega_\beta} d^{\alpha-1} (\Delta d) v_\varepsilon^2 dx \right| \leq \frac{c_5 \beta_0^{\alpha-1}}{\alpha-1} \int_{\Omega_\beta} v_\varepsilon^2 dx.$$

Also

$$\begin{aligned} \left| \int_{\Omega_\beta} v_\varepsilon^2 \operatorname{div}(d^{\alpha-1} \nabla d) dx \right| &= 2 \left| \int_{\Omega_\beta} v_\varepsilon d^{\frac{\alpha}{2}} d^{\frac{\alpha}{2}-1} \nabla d \cdot \nabla v_\varepsilon dx \right| \\ &\leq c_6 \int_{\Omega_\beta} d^\alpha |\nabla v_\varepsilon|^2 dx + \delta \int_{\Omega_\beta} d^{\alpha-2} v_\varepsilon^2 dx. \end{aligned}$$

where $c_6 = c_6(\delta) > 0$, and the result follows in this case, if we choose β and δ small enough

(ii) By the same calculations we have

$$\int_\Omega d^{-\frac{p}{2}} |v_\varepsilon|^p dx \leq c_7 \int_{\Omega_\beta} d^{\frac{p}{2}} |\nabla v_\varepsilon|^p dx \leq c_7 \left(\int_\Omega d(x) dx \right)^{\frac{p}{2}} \int_{\Omega_\beta} d |\nabla v_\varepsilon|^2 dx.$$

□

In the following statement we prove regularity up to the boundary for the function $\frac{u}{\phi_\kappa}$.

Proposition 2.8. Let $f_0 \in L^2(\Omega)$. Then there exists a unique $u \in H_{loc}^1(\Omega)$ such that $\phi_\kappa^{-1}u \in H^1(\Omega, d^{\alpha+}(x)dx)$, satisfying (2.13). Furthermore, if $f_0 \in L^q(\Omega)$, $q > \frac{N+\alpha}{2}$, then there exists $0 < \beta < 1$ such that

$$\sup_{x,y \in \Omega, x \neq y} |x-y|^{-\beta} \left| \frac{u(x)}{\phi_\kappa(x)} - \frac{u(y)}{\phi_\kappa(y)} \right| < c_8 \|f_0\|_{L^q}. \quad (2.16)$$

Proof. If there exists a solution u , then $\psi = \frac{u}{\phi_\kappa}$ satisfies

$$-\phi_\kappa^{-2} \operatorname{div}(\phi_\kappa^2 \nabla \psi) + \lambda_\kappa \psi = \phi_\kappa^{-1} f_0. \quad (2.17)$$

and we recall that $\phi_\kappa(x) \approx d^{\frac{\alpha_+}{2}}(x)$. We endow the space $H^1(\Omega, \phi_\kappa^2 dx)$ with the inner product

$$\langle a, b \rangle = \int_\Omega (\nabla a \cdot \nabla b + \lambda_\kappa ab) \phi_\kappa^2 dx.$$

By a solution ψ of (2.17) we mean that $\psi \in H_0^1(\Omega, \phi_\kappa^2 dx)$ satisfies

$$\langle \nabla \psi, \nabla \zeta \rangle = \int_\Omega \nabla \psi \cdot \nabla \zeta \phi_\kappa^2 dx + \lambda_\kappa \int_\Omega \psi \zeta \phi_\kappa^2 dx = \int_\Omega f_0 \zeta \phi_\kappa dx \quad \forall \zeta \in H_0^1(\Omega, \phi_\kappa^2 dx). \quad (2.18)$$

By Riesz's representation theorem we derive the existence and uniqueness of the solution in this space. Since $H^1(\Omega, \phi_\kappa^2 dx) = H_0^1(\Omega, \phi_\kappa^2 dx)$ by [12, Th 2.11], any weak solution u of (2.13) such that $\phi_\kappa^{-1} u \in H^1(\Omega, \phi_\kappa^2 dx)$ is obtained by the above method.

Finally if $f_0 \in L^q(\Omega)$, where $q > \frac{N+\alpha}{2}$, we can apply Moser's approach thanks to (2.8) and prove first the estimate

$$\|\psi\|_{L^\infty(\Omega)} \leq c_8 \|f_0\|_{L^q(\Omega)} \quad (2.19)$$

where $c_8 = c_8(\Omega, \kappa, q)$, and then to derive the Hölder regularity up to the boundary (see e.g. [12]). \square

In the next results we make more precise the rate of convergence of a solution of (2.13) to its boundary value.

Proposition 2.9. *Assume $\kappa < \frac{1}{4}$. If $f_0 \in L^2(\Omega)$ and $h \in H^1(\Omega)$ there exists a unique weak solution u of (2.13) in $H_{loc}^1(\Omega)$ and such that $d^{-\frac{\alpha_+}{2}}(u - d^{\frac{\alpha_-}{2}} h) \in H^1(\Omega, d^{\alpha_+}(x) dx)$. Furthermore, if $f_0 \in L^q(\Omega)$, $q > \frac{N+\alpha}{2}$ and $h \in C^2(\bar{\Omega})$, then there exists $0 < \beta < 1$ such that*

$$\lim_{x \in \Omega, x \rightarrow y \in \partial\Omega} \frac{u(x)}{(d(x))^{\frac{\alpha_-}{2}}} = h(y) \quad \forall y \in \partial\Omega,$$

uniformly with respect to y ,

$$\left\| \frac{u}{d^{\frac{\alpha_-}{2}}} \right\|_{L^\infty(\Omega)} \leq c_9 \left(\|h\|_{C^2(\bar{\Omega})} + \|f_0\|_{L^q(\Omega)} \right),$$

and

$$\sup_{x, y \in \Omega, x \neq y} |x - y|^{-\beta} \left| \frac{u(x)}{(d(x))^{\frac{\alpha_-}{2}}} - \frac{u(y)}{(d(y))^{\frac{\alpha_-}{2}}} \right| < c_{10}. \quad (2.20)$$

with c_9 and c_{10} depending on Ω, N, q , and κ .

Remark. By Lemma 2.7 we already know that $u - d^{\frac{\alpha_-}{2}} h \in H_0^1(\Omega)$.

Proof. Let $\beta \leq \beta_0$ and $\eta \in C^2(\Omega)$ be a function such that $\eta = d^{\frac{\alpha_-}{2}}(x)$ in Ω_β and $\eta(x) > c > 0$, if $x \in \Omega'_\beta$. We set $u = \phi_\kappa v + \eta h$. Then v is a weak solution of

$$-\frac{\operatorname{div}(\phi_\kappa^2 \nabla v)}{\phi_\kappa^2} + \lambda_\kappa v = \frac{1}{\phi_\kappa} \left(f_0 + (\Delta \eta + \kappa \frac{\eta}{d^2}) h + 2 \nabla \eta \cdot \nabla h + \eta \Delta h \right), \quad (2.21)$$

in the sense that

$$\begin{aligned} \int_{\Omega} \nabla v \cdot \nabla \psi \phi_{\kappa}^2 dx + \lambda_{\kappa} \int_{\Omega} v \psi \phi_{\kappa}^2 dx &= \int_{\Omega} \left(f_0 + (\Delta \eta + \kappa \frac{\eta}{d^2}) h + 2 \nabla \eta \cdot \nabla h \right) \psi \phi_{\kappa} dx \\ &\quad - \int_{\Omega} \nabla h \cdot \nabla (\eta \psi \phi_{\kappa}) dx \quad \forall \psi \in C_0^{\infty}(\Omega). \end{aligned} \quad (2.22)$$

Let $\psi \in C_0^{\infty}(\Omega_{\beta})$. By an argument similar to the one in the proof of Lemma 2.7 we have

$$\int_{\Omega} \psi^2 dx = \int_{\Omega_{\beta}} \psi^2 dx = \int_{\Omega_{\beta}} \operatorname{div}(d \nabla d) |\psi|^2 dx - \int_{\Omega_{\beta}} (d \Delta d) |\psi|^2 dx,$$

which implies

$$\int_{\Omega_{\beta}} \psi^2 dx \leq c'_{10} \int_{\Omega_{\beta}} d^2 |\nabla \psi|^2 dx \leq c_{11} \int_{\Omega_{\beta}} d^{\alpha+} |\nabla \psi|^2 dx. \quad (2.23)$$

Now

$$\left| \int_{\Omega_{\beta}} \left((\Delta \eta + \kappa \frac{\eta}{d^2}) h + 2 \nabla \eta \cdot \nabla h \right) \psi \phi_{\kappa} dx \right| \leq c_{12} \int_{\Omega_{\beta}} \psi^2 dx,$$

and

$$\left| \int_{\Omega_{\beta}} \nabla h \cdot \nabla (\eta \psi \phi_{\kappa}) dx \right| \leq c_{13} \left(\int_{\Omega_{\beta}} |\nabla h|^2 dx + \int \int_{\Omega_{\beta}} d^{\alpha+} |\nabla \psi|^2 dx + \int \int_{\Omega_{\beta}} \psi^2 dx \right).$$

By (2.23) we can take $\psi \in H^1(\Omega, d^{\alpha+}(x) dx)$ for test function. Thus we can easily obtain that there exists a weak solution $v \in H^1(\Omega, d^{\alpha+}(x) dx)$ of (2.22).

To prove (2.20) we first obtain that if $\psi \in C_0^{\infty}(\Omega_{\varepsilon})$

$$\int_{\Omega} \psi dx = - \int_{\Omega_{\varepsilon}} d \nabla d \cdot \nabla \psi dx - \int_{\Omega_{\varepsilon}} d \Delta d \psi dx,$$

and since

$$\begin{aligned} \left| \int_{\Omega} \left((\Delta \eta + \kappa \frac{\eta}{d^2}) h + 2 \nabla \eta \cdot \nabla h + \eta \Delta h \right) \psi \phi_{\kappa} dx \right| &\leq c_{14} \|h\|_{C^2(\bar{\Omega})} \int_{\Omega} |\psi| dx \\ &\leq \frac{1}{2} \int_{\Omega_{\varepsilon}} d |\nabla \psi|^2 dx + c_{15}(\Omega) \|h\|_{C^2(\bar{\Omega})}. \end{aligned}$$

Using again (2.8) and Moser's iterative scheme as in Proposition 2.8, we obtain

$$\|v\|_{L^{\infty}(\Omega)} \leq c_9 \left(\|h\|_{C^2(\bar{\Omega})} + \|f_0\|_{L^q(\Omega)} \right),$$

where $c_9 = c_9(\Omega, q, \kappa) > 0$, from which it follows again that v is Hölder continuous up to the boundary and the uniform convergence holds. \square

Proposition 2.10. *Assume $\kappa = \frac{1}{4}$. If $f_0 \in L^2(\Omega)$ and $h \in H^1(\Omega)$, there exists a unique function u in $H_{loc}^1(\Omega)$ weak solution of*

$$\mathcal{L}_{\frac{1}{4}} u = f_0$$

verifying $d^{-\frac{1}{2}}(u - d^{\frac{1}{2}}|\log d|h) \in H^1(\Omega, d(x)dx)$. Furthermore, if $f_0 \in L^q(\Omega)$, $q > \frac{n+1}{2}$ and $h \in C^2(\bar{\Omega})$, then there exists $0 < \beta < 1$ such that

$$\lim_{x \in \Omega, x \rightarrow y \in \partial\Omega} \frac{u}{d^{\frac{1}{2}}|\log d|}(x) = h(y) \quad \forall y \in \partial\Omega,$$

uniformly with respect to y ,

$$\left\| \frac{u}{\sqrt{d}|\log \frac{d}{D_0}|} \right\|_{L^\infty(\Omega)} \leq c_{16} \left(\|h\|_{C^2(\bar{\Omega})} + \|f_0\|_{L^q(\Omega)} \right)$$

where $D_0 = 2 \sup_{x \in \Omega} d(x)$. Finally there holds

$$\sup_{x, y \in \Omega, x \neq y} |x - y|^{-\beta} \left| \frac{u(x)}{\sqrt{d(x)}|\log \frac{d(x)}{D_0}|} - \frac{u(y)}{\sqrt{d(y)}|\log \frac{d(y)}{D_0}|} \right| < c_{17} \left(\|h\|_{C^2(\bar{\Omega})} + \|f_0\|_{L^q(\Omega)} \right). \quad (2.24)$$

Proof. Using again Lemma 2.7, we know that $u - d^{\frac{1}{2}}|\log d|h \in W_0^{1,p}(\Omega)$, $\forall p < 2$. The proof is very similar to the proof of Proposition 2.9. The only differences are we impose $\eta = d^{\frac{1}{2}}|\log d|$ in Ω_β and we use the fact that $|\log d| \in L^p(\Omega)$, $\forall p \geq 1$. \square

In the next result we prove that the boundary Harnack inequality holds, provided the vanishing property of a solution is understood in an appropriate way.

Proposition 2.11. *Let $\delta > 0$ be small enough, $\xi \in \partial\Omega$ and $u \in H_{loc}^1(B_\delta(\xi) \cap \Omega) \cap C(B_\delta(\xi) \cap \bar{\Omega})$ be a positive $\mathcal{L}_{\frac{1}{4}}$ -harmonic function in $B_\delta(\xi) \cap \Omega$ vanishing on $\partial\Omega \cap B_\delta(\xi)$ in the sense that*

$$\lim_{\text{dist}(x, K) \rightarrow 0} \frac{u(x)}{d^{\frac{1}{2}}(x)|\log d(x)|} = 0 \quad \forall K \subset \partial\Omega \cap B_\delta(\xi), K \text{ compact}. \quad (2.25)$$

Then there exists a constant $c_{18} = c_{18}(N, \Omega) > 0$ such that

$$\frac{u(x)}{\phi_{\frac{1}{4}}(x)} \leq c_{18} \frac{u(y)}{\phi_{\frac{1}{4}}(y)} \quad \forall x, y \in \Omega \cap B_{\frac{\delta}{2}}(\xi).$$

Proof. We already know that $u \in C^2(\Omega)$. Let $\delta \leq \beta_0$ such that $B_\delta(\xi) \cap \Omega \subset \Omega_\delta \subset \Omega_{\beta_0}$.

By [3, Lemma 2.8] there exists a positive supersolution $\zeta \in C^2(\Omega_\delta)$ of (1.3) in Ω_δ with the following behaviour

$$\zeta(x) \approx d^{\frac{1}{2}}(x) \log \frac{1}{c_{19}d(x)} \left(1 + \left(\log \frac{1}{C_0(\Omega)d(x)} \right)^{-\beta} \right),$$

for some $\beta \in (0, 1)$ and $c_{19} = c_{19}(\Omega) > 0$. Set $v = \zeta^{-1}u$, then it satisfies

$$-\zeta^{-2} \text{div}(\zeta^2 \nabla v) \leq 0 \quad \text{in } B_\delta(\xi) \cap \Omega. \quad (2.26)$$

Let $\eta \in C_0^\infty(B_\delta(\xi))$ such that $0 \leq \eta \leq 1$ and $\eta = 1$ in $B_{\frac{3\delta}{4}}(\xi)$. We set $v_s = \eta^2(v - s)_+$. Since by assumption v_s has compact support in $B_\delta(\xi) \cap \Omega$, we can use it as a test function in (2.26) and we get

$$\int_{B_\delta(\xi) \cap \Omega} \zeta^2 \nabla v \cdot \nabla v_s dx = \int_{B_\delta(\xi) \cap \Omega} \zeta^2 \nabla(v - s)_+ \cdot \nabla v_s dx \leq 0, \quad (2.27)$$

which yields

$$\int_{B_\delta(\xi) \cap \Omega} |\nabla(v-s)_+|^2 \zeta^2 \eta^2 dx \leq 4 \int_{B_\delta(\xi) \cap \Omega} |\nabla \eta|^2 (v-s)_+^2 \zeta^2 dx.$$

Letting $s \rightarrow 0$ we derive

$$\int_{B_\delta(\xi) \cap \Omega} |\nabla v|^2 \zeta^2 \eta^2 dx \leq 4 \int_{B_\delta(\xi) \cap \Omega} |\nabla \eta|^2 v^2 \zeta^2 dx.$$

Since

$$|\nabla(v-s)_+|^2 \zeta^2 \eta^2 \uparrow |\nabla v|^2 \zeta^2 \eta^2 \quad \text{as } s \rightarrow 0,$$

and convergence of $\nabla(v-s)_+$ to ∇v holds a.e. in Ω , it follows by the monotone convergence theorem

$$\lim_{s \rightarrow 0} \int_{B_\delta(\xi) \cap \Omega} |\nabla(v - (v-s)_+)|^2 \zeta^2 \eta^2 dx = 0, \quad (2.28)$$

and finally $\zeta v_s \rightarrow \eta^2 \zeta v$ in $H^1(B_\delta(\xi) \cap \Omega)$, which yields in particular $\eta^2 u = \eta^2 \zeta v \in H_0^1(B_\delta(\xi) \cap \Omega)$.

Step 2. By [3, Lemma 2.8] there exists a positive subsolution $h \in C^2(\Omega_\delta)$ of (1.3) in Ω_δ with the following behaviour

$$h(x) \approx d^{\frac{1}{2}}(x) \log \frac{1}{c_{20}d(x)} \left(1 - \left(\log \frac{1}{c_{20}d(x)} \right)^{-\beta} \right),$$

where $\beta \in (0, 1)$ and $c_{20} = c_{20}(\Omega) > 0$. Set $w = h^{-1}u$ and $w_s = \eta^2(w-s)_+$. Then $w_s \rightarrow \eta^2 w$ in $H^1(B_\delta(\xi) \cap \Omega)$ by Step 1. Put $u_s = hw_s$, thus, for $0 < s, s'$, we have

$$\begin{aligned} \int_{B_\delta(\xi) \cap \Omega} |\nabla(u_s - u_{s'})|^2 dx - \frac{1}{4} \int_{B_\delta(\xi)} \frac{|u_s - u_{s'}|^2}{d^2(x)} dx &= \int_{B_\delta(\xi) \cap \Omega} h^2 |\nabla(w_s - w_{s'})|^2 dx \\ + \int_{B_\delta(\xi) \cap \Omega} |\nabla h|^2 |w_s - w_{s'}|^2 dx + \int_{B_\delta(\xi) \cap \Omega} h \nabla h \cdot \nabla(u_s - u_{s'})^2 dx - \frac{1}{4} \int_{B_\delta(\xi) \cap \Omega} \frac{h^2 |w_s - w_{s'}|^2}{d^2(x)} dx \\ &\leq \int_{B_\delta(\xi) \cap \Omega} h^2 |\nabla(w_s - w_{s'})|^2 dx, \end{aligned} \quad (2.29)$$

where, in the last inequality, we have performed by parts integration and then used the fact that h is a subsolution. Thus we have by (2.28) that

$$\lim_{s, s' \rightarrow 0} \int_{B_\delta(\xi)} |\nabla(u_s - u_{s'})|^2 dx - \frac{1}{4} \int_{B_\delta(\xi)} \frac{|u_s - u_{s'}|^2}{d^2(x)} dx = 0. \quad (2.30)$$

Let $\mathbf{W}(\Omega)$ denote the closure of $C_0^\infty(\Omega)$ in the space of functions ϕ satisfying

$$\|\phi\|_H^2 := \int_{\Omega} |\nabla \Phi|^2 dx - \frac{1}{4} \int_{\Omega} \frac{|\Phi|^2}{d^2(x)} dx < \infty.$$

Thus $\eta^2 u \in \mathbf{W}(\Omega)$, which implies

$$\frac{\eta u}{\phi_{\frac{1}{4}}} \in H_0^1(\Omega, d(x)dx).$$

Next we set $\tilde{v} = \phi_{\frac{1}{4}}^{-1} u$; then $\tilde{v} \in H^1(B_{\frac{3\delta}{4}}(\xi), d(x)dx)$ and it satisfies

$$-\phi_{\frac{1}{4}}^{-2} \operatorname{div}(\phi_{\frac{1}{4}}^2 \nabla \tilde{v}) + \lambda_{\frac{1}{4}} \tilde{v} = 0.$$

By the same approach based on Moser' iterative scheme applied to degenerate elliptic operators as the in [12, Theorem 1.5], we see that v satisfies a Harnack inequality up to the boundary of Ω . More precisely there exists a constant $c_{18} = c_{18}(\Omega) > 0$ such that

$$v(x) \leq c_{18}v(y) \quad \forall x, y \in B_{\frac{\delta}{2}}(\xi).$$

And the result follows. \square

In the case $\kappa < \frac{1}{4}$, the boundary Harnack inequality is the following,

Proposition 2.12. *Let $\delta > 0$ be small enough $\xi \in \Omega$, $\kappa < \frac{1}{4}$ and $u \in H_{loc}^1(B_\delta(\xi) \cap \Omega) \cap C(B_\delta(\xi) \cap \bar{\Omega})$ be a nonnegative \mathcal{L}_κ -harmonic in $B_\delta(\xi)$ vanishing on $\partial\Omega \cap B_\delta(\xi)$ in the sense that*

$$\lim_{\text{dist}(x,K) \rightarrow 0} \frac{u(x)}{(d(x))^{\frac{\alpha-}{2}}} = 0 \quad \forall K \subset \partial\Omega \cap B_\delta(\xi), K \text{ compact.} \quad (2.31)$$

Then there exists $c_{21} = c_{21}(\Omega, \kappa) > 0$ such that

$$\frac{u(x)}{\phi_\kappa(x)} \leq c_{21} \frac{u(y)}{\phi_\kappa(y)} \quad \forall x, y \in \bar{\Omega} \cap B_{\frac{\delta}{2}}(\xi).$$

Proof. The only difference with the preceding proof is that we take as subsolution and supersolution (see [3, Lemma 2.8]) $C^2(\Omega)$ the functions h and ζ respectively with the boundary behaviour

$$h \approx d^{\alpha-} (1 - c_{22}d^\beta) \quad \zeta \approx d^{\alpha-} (1 + c_{23}d^\beta),$$

where $\beta \in (0, \sqrt{1 - 4\kappa})$. \square

Proposition 2.13. *Let $u \in H_{loc}^1(\Omega) \cap C(\Omega)$ be a $\mathcal{L}_{\frac{1}{4}}$ -subharmonic function such that*

$$\limsup_{d(x) \rightarrow 0} \frac{u(x)}{d^{\frac{1}{2}}(x) |\log d(x)|} \leq 0.$$

Then $u \leq 0$.

Proof. We set $v = \max(u, 0)$ and we proceed as in Step 1 of the proof of Proposition 2.11 with $\eta = 1$. The result follows by letting $s \rightarrow 0$. \square

Similarly we have

Proposition 2.14. *Let $u \in H_{loc}^1(\Omega) \cap C(\Omega)$ be a \mathcal{L}_κ -subharmonic function such that*

$$\limsup_{d(x) \rightarrow 0} \frac{u(x)}{(d(x))^{\frac{\alpha-}{2}}} \leq 0.$$

then $u \leq 0$.

The two next statements shows that comparison holds provided comparable boundary data are achieved in way which takes into account the specific form of the \mathcal{L}_κ -harmonic functions

Proposition 2.15. Assume $\kappa < \frac{1}{4}$ and $h_i \in H^1(\Omega)$ ($i=1,2$). Let $u_i \in H_{loc}^1(\Omega)$ be two \mathcal{L}_κ -harmonic functions such that $d^{-\frac{\alpha_+}{2}} \left(u_i - d^{\frac{\alpha_-}{2}} h_i \right) \in H^1(\Omega, d^{\alpha_+}(x)dx)$. Then

If $h_1 \leq h_2$ a.e. in Ω , there holds

$$u_1(x) \leq u_2(x) \quad \forall x \in \Omega.$$

If $h_1 - h_2 \in H_0^1(\Omega)$, there holds

$$u_1(x) = u_2(x) \quad \forall x \in \Omega.$$

Proof. Set $w = \phi_\kappa^{-1}(u_1 - u_2)$, then $w \in H^1(\Omega, \phi_\kappa^2 dx)$ and

$$-\operatorname{div}(\phi_\kappa^2 \nabla w) + \lambda_\kappa \phi_\kappa^2 w = 0$$

Since $H^1(\Omega, \phi_\kappa^2 dx) = H_0^1(\Omega, \phi_\kappa^2 dx)$ by (2.4) we derive that w and w belongs to $H_0^1(\Omega, \phi_\kappa^2 dx)$ and, integrating by part, we derive $w_+ = 0$. The proof of the second statement is similar. \square

In the same way we have in the case $\kappa = \frac{1}{4}$.

Proposition 2.16. Assume $\kappa = \frac{1}{4}$. Let $h_i \in H^1(\Omega)$ ($i=1,2$) and let $u_i \in H_{loc}^1(\Omega)$ be two $\mathcal{L}_{\frac{1}{4}}$ -harmonic functions such that $d^{-\frac{1}{2}}(u_i - d^{\frac{1}{2}} |\log d| h_i) \in H^1(\Omega, d(x)dx)$.

(i) If $h_1 \leq h_2$ a.e. in Ω , then

$$u_1(x) \leq u_2(x) \quad \forall x \in \Omega.$$

(ii) If $h_1 - h_2 \in H_0^1(\Omega)$, then

$$u_1(x) = u_2(x) \quad \forall x \in \Omega.$$

We end with existence and uniqueness results for solving the Dirichlet problem associated to \mathcal{L}_κ .

Proposition 2.17. Assume $\kappa = \frac{1}{4}$. For any $h \in C(\partial\Omega)$ there exists a unique $\mathcal{L}_{\frac{1}{4}}$ -harmonic function u belonging to $H_{loc}^1(\Omega)$ satisfying

$$\lim_{x \in \Omega, x \rightarrow y \in \partial\Omega} \frac{u(x)}{d^{\frac{1}{2}}(x) |\log \frac{d(x)}{D_0}|} = h(y) \quad \text{uniformly for } y \in \partial\Omega.$$

Furthermore there exists a constant $c_{16} = c_{16}(\Omega) > 0 > 0$

$$\left\| \frac{u}{d^{\frac{1}{2}} |\log \frac{d}{D_0}|} \right\|_{L^\infty(\Omega)} \leq c_{24} \|h\|_{C(\partial\Omega)},$$

where $D_0 = 2 \sup_{x \in \Omega} d(x)$.

Proof. Uniqueness is a consequence of Proposition 2.13. For existence let $m \in \mathbb{N}$ and h_m be smooth functions such that $h_m \rightarrow h$ in $L^\infty(\partial\Omega)$. Then we can find a function $H_m \in C^2(\bar{\Omega})$ with trace h_m on $\partial\Omega$, and $\|H_m\|_{L^\infty(\Omega)} \leq \|h_m\|_{L^\infty(\partial\Omega)}$. By Lemma 2.10 there exists a unique weak solution u_m of $\mathcal{L}_{\frac{1}{4}} u = 0$ satisfying

$$\lim_{x \in \Omega, x \rightarrow y \in \partial\Omega} \frac{u_m}{d^{\frac{1}{2}} |\log d|}(x) = h_m(y) \quad \text{uniformly for } y \in \partial\Omega.$$

By Proposition 2.10 we have

$$\left\| \frac{u_m - u_n}{d^{\frac{1}{2}} |\log \frac{d}{D_0}|} \right\|_{L^\infty(\Omega)} \leq c_{16} \|h_m - h_n\|_{C(\partial\Omega)}.$$

Thus there exists u such that

$$\lim_{m \rightarrow \infty} \left\| \frac{u_m - u}{d^{\frac{1}{2}} |\log \frac{d}{D_0}|} \right\|_{L^\infty(\Omega)} = 0$$

and u is a solution of $L_{\frac{1}{4}} u = 0$.

Let $x \in \Omega$, with $d(x) < \frac{1}{2}$ and $y \in \partial\Omega$

$$\begin{aligned} \left| \frac{u}{d^{\frac{1}{2}} |\log d|}(x) - h(y) \right| &\leq \left| \frac{u}{d^{\frac{1}{2}} |\log d|}(x) - \frac{u_m}{d^{\frac{1}{2}} |\log d|}(x) \right| + \left| \frac{u_m}{d^{\frac{1}{2}} |\log d|}(x) - h_m(y) \right| \\ &\quad + |h(y) - h_m(y)|. \end{aligned}$$

The result follows by letting successively $x \rightarrow y$ and $m \rightarrow \infty$. \square

Similarly we have

Proposition 2.18. *Assume $\kappa < \frac{1}{4}$. Then for any $h \in C(\partial\Omega)$ there exists a unique \mathcal{L}_κ -harmonic function $u \in H_{loc}^1(\Omega)$ satisfying*

$$\lim_{x \in \Omega, x \rightarrow y \in \partial\Omega} \frac{u}{d^{\frac{\alpha-}{2}}}(x) = h(y) \quad \text{uniformly for } y \in \partial\Omega.$$

Furthermore there exists a constant $c_9 = c_9(\Omega, \alpha) > 0$ such that

$$\left\| \frac{u}{d^{\alpha-}} \right\|_{L^\infty(\Omega)} \leq c_9 \|h\|_{C(\partial\Omega)}.$$

A useful consequence of [3, Lemma 2.8] and Propositions 2.9 and 2.10 is the following local existence result.

Proposition 2.19. *There exists a positive \mathcal{L}_κ -harmonic function $Z_\kappa \in C(\overline{\Omega_{\beta_0}}) \cap C^2(\Omega_{\beta_0})$ satisfying*

$$\lim_{d(x) \rightarrow 0} \frac{Z_{\frac{1}{4}}(x)}{\sqrt{d(x)} |\ln d(x)|} = 0 \tag{2.32}$$

if $\kappa = \frac{1}{4}$, and

$$\lim_{d(x) \rightarrow 0} \frac{Z_\kappa(x)}{(d(x))^{\frac{\alpha-}{2}}} = 0 \tag{2.33}$$

if $0 < \kappa < \frac{1}{4}$.

2.3 \mathcal{L}_κ -harmonic measure

Let $x_0 \in \Omega$, $h \in C(\partial\Omega)$ and denote $L_{\kappa, x}(h) := v_h(x_0)$ where v_h is the solution of the Dirichlet problem (see Lemmas 2.17 and 2.18)

$$\begin{aligned} \mathcal{L}_\kappa v &= 0 & \text{in } \Omega \\ v &= h & \text{in } \partial\Omega \end{aligned}$$

where v take the boundary data in the sense of Lemmas 2.17 and 2.18. By Lemma's 2.14 and 2.13, the mapping $h \mapsto L_{\kappa, x_0}(h)$ is a linear positive functional on $C(\partial\Omega)$. Thus there exists a unique Borel measure on $\partial\Omega$, called \mathcal{L}_κ -harmonic measure in Ω , denoted by ω^{x_0} , such that

$$v_h(x_0) = \int_{\partial\Omega} h(y) d\omega^{x_0}(y).$$

Because of Harnack inequality the measures ω^x and ω^{x_0} , $x_0, x \in \Omega$ are mutually absolutely continuous. For every fixed x we denote the Radon-Nikodym derivative by

$$K(x, y) := \frac{d\omega^x}{d\omega^{x_0}}(y) \quad \text{for } \omega^{x_0}\text{-almost all } y \in \partial\Omega.$$

It is wellknown that the following formula is an equivalent definition of the \mathcal{L}_κ -harmonic measure: for any closed set $E \subset \partial\Omega$

$$\omega^{x_0}(E) = \inf \left\{ \psi : \psi \in C_+(\Omega), \mathcal{L}_\kappa\text{-superharmonic in } \Omega \text{ s.t. } \liminf_{x \rightarrow E} \frac{\psi(x)}{W(x)} \geq 1 \right\},$$

where

$$W(x) = \begin{cases} d^{\frac{\alpha_-}{2}}(x) & \text{if } \kappa < \frac{1}{4}, \\ d^{\frac{1}{2}}(x) |\log d(x)| & \text{if } \kappa = \frac{1}{4}. \end{cases}$$

The extension to open sets is standard. Let $\xi \in \partial\Omega$. We set $\Delta_r(\xi) = \partial\Omega \cap B_r(\xi)$ and $x_r = x_r(\xi) \in \Omega$, such that $d(x_r) = |x_r - \xi| = r$. Also $x_r(\xi) = \xi - r\mathbf{n}_\xi$ where \mathbf{n}_ξ is the unit outward normal vector to $\partial\Omega$ at ξ . We recall that $\beta_0 = \beta_0(\Omega) > 0$ has been defined in Lemma 2.20.

Lemma 2.20. *There exists a constant $c_{25} > 0$ which depends only on Ω and α such that if $0 < r \leq \beta_0$ and $\xi \in \partial\Omega$, there holds*

$$\frac{\omega^x(\Delta_r(\xi))}{W(x)} \geq c_{25} \quad \forall x \in \Omega \cap B_{\frac{r}{2}}(\xi). \quad (2.34)$$

Proof. Let $h \in C(\partial\Omega)$ be a function with compact support in $\Delta_r(\xi)$, $0 \leq h \leq 1$ and $h = 1$ on $\overline{\Delta_{\frac{3r}{4}}(\xi)}$. And let v_h, v_1 the corresponding \mathcal{L}_κ -harmonic functions with h and 1 as boundary data respectively (in the sense of Lemmas 2.17 and 2.18). Then $v_1(x) \geq v_h(x) \geq 0$ and

$$\lim_{x \in \Omega, x \rightarrow x_0} \frac{v_1(x) - v_h(x)}{W(x)} = 0 \quad \forall x_0 \in \Omega \cap B_{\frac{3r}{4}}(\xi).$$

By Lemmas 2.12 and 2.11, and $\phi_\kappa \approx d^{\frac{\alpha_+}{2}}$, there exists $c_{26} = c_{26}(\Omega) > 0$ such that

$$\frac{v_1(x) - v_h(x)}{d^{\frac{\alpha_+}{2}}(x)} \leq c_{26} \frac{v_1(y) - v_h(y)}{d^{\frac{\alpha_+}{2}}(y)}, \quad \forall x, y \in \Omega \cap \overline{B_{\frac{r}{2}}(\xi)}.$$

We consider first the case $\kappa = \frac{1}{4}$. By Lemma 2.17, we have

$$0 \leq \frac{v_1(y) - v_h(y)}{d^{\frac{1}{2}}(y)} \leq \frac{v_1(y)}{d^{\frac{1}{2}}(y)} \leq c_{24} |\log d(y)|.$$

Thus, combining all above we have that

$$\frac{v_1(x)}{d^{\frac{1}{2}}(x) |\log d(x)|} - c_{27} \frac{|\log d(y)|}{|\log d(x)|} \leq \frac{v_h(x)}{d^{\frac{1}{2}}(x) |\log d(x)|}.$$

Now by Lemma 2.10, there exists a $\varepsilon_0 > 0$ such that

$$\frac{v_1(x)}{d^{\frac{1}{2}}(x)|\log d(x)|} > \frac{1}{2} \quad \forall x \in \Omega_{\varepsilon_0}.$$

Thus if we choose y such that $d(y) = \frac{r}{4}$, there exists a constant $c_{27} = c_{27}(\Omega) > 0$ such that

$$c_{27} \frac{|\log d(y)|}{|\log d(x)|} = c_{27} \frac{|\log \frac{r}{4}|}{|\log d(x)|} \leq c_{27} \frac{|\log \frac{r}{4}|}{|\log \frac{r}{D_0}|} \leq \frac{1}{4} \quad \forall x \in \Omega_{\frac{r}{D_0}},$$

thus

$$\frac{v_h(x)}{d^{\frac{1}{2}}(x)|\log d(x)|} \geq \frac{1}{4} \quad \forall x \in \overline{B_{\frac{r}{2}}(\xi)} \cap \Omega_{\frac{r}{D_0}}. \quad (2.35)$$

In particular

$$\frac{v_h(x_{a^*r}(\xi))}{\sqrt{a^*r}|\log(a^*r)|} \geq \frac{1}{4}. \quad (2.36)$$

where $a^* = (\max\{2, D_0\})^{-1}$. If $D_0 \leq 2$ we obtain the claim. If not, set $k^* = \mathbb{E}[\frac{D_0}{2}] + 1$ (recall that $\mathbb{E}[x]$ denotes the largest integer less or equal to x). If $x \in \overline{B_{\frac{r}{2}}(\xi)} \cap \Omega$ there exists a chain of at most $4k^*$ points $\{z_j\}_{j=0}^{j_0}$ such that $z_j \in \overline{B_{\frac{r}{2}}(\xi)} \cap \Omega$, $d(z_j) \geq a^*r$, $z_0 = x_{a^*r}(\xi)$, $z_{j_0} = x$ and $|z_j - z_{j+1}| \leq \frac{a^*r}{4}$. By Harnack inequality (applied j_0 -times)

$$v_h(x_{a^*r}(\xi)) \leq c_{28}v_h(x). \quad (2.37)$$

Since

$$W(x_{a^*r}(\xi)) \geq \left(\frac{D_0}{2}\right)^{\frac{1}{2}} W(x),$$

we obtain finally

$$\frac{1}{4} \leq \frac{\omega^{x_{a^*r}(\xi)}(\Delta_r(\xi))}{\sqrt{a^*r}|\log(a^*r)|} \leq c_{28} \left(\frac{a}{a^*}\right)^{\frac{1}{2}} \frac{\omega^x(\Delta_r(\xi))}{W(x)} \quad \forall x \in \Omega \cap B_{\frac{r}{2}}(\xi). \quad (2.38)$$

In the case $\kappa < \frac{1}{4}$, the proof is simpler since no log term appears and we omit it. \square

The next result is a Carleson type estimate valid for positive \mathcal{L}_κ -harmonic functions.

Lemma 2.21. *There exists a constant c_{29} which depends only on Ω such that for any $\xi \in \partial\Omega$ and $0 < r \leq s \leq \beta_0$,*

$$\frac{\omega^x(\Delta_r(\xi))}{W(x)} \leq c_{29} \frac{\omega^{x_s(\xi)}(\Delta_r(\xi))}{W(x_s(\xi))} \quad \forall x \in \Omega \setminus B_s(\xi). \quad (2.39)$$

Proof. By Propositions 2.17 and 2.18 there exists a constant $c_{30} > 0$ such that for $0 < r < \beta_0$,

$$\frac{\omega^x(\Delta_r(\xi))}{W(x)} \leq \frac{\omega^x(\partial\Omega)}{W(x)} \leq c_{30} \quad \forall x \in \Omega. \quad (2.40)$$

By Propositions 2.17 and 2.18, there holds

$$\lim_{d(x) \rightarrow 0} \frac{v_1(x)}{W(x)} = 1, \quad (2.41)$$

thus we can replace W by v_1 in (2.39). Since $w_h = \frac{v_h(x)}{v_1(x)}$ is Hölder continuous in $\overline{\Omega}$ and satisfies

$$\begin{aligned} -\operatorname{div}(v_1^2 \nabla w_h) &= 0 && \text{in } \Omega \setminus \overline{B_s}(\xi) \\ 0 \leq w_h &\leq 1 && \text{in } \Omega \setminus \overline{B_s}(\xi) \\ w_h &= 0 && \text{in } \partial\Omega \setminus \overline{B_s}(\xi) \end{aligned} \quad (2.42)$$

the maximum of w_h is achieved on $\Omega \cap \partial B_s(\xi)$, therefore it is sufficient to prove the Carleson estimate

$$w_h(x) \leq c_{29} w_h(x_s(\xi)) \quad \forall x \in \Omega \cap \partial B_s(\xi). \quad (2.43)$$

If x such that $|x - \xi| = s$ is "far" from $\partial\Omega$, $w_h(x)$ is "controlled" by $w_h(x_s(\xi))$ thanks to Harnack inequality, while if it is close to $\partial\Omega$, $w_h(x)$ is "controlled" by the fact that it vanishes on $\partial\Omega \cap \partial B_s(\xi)$.

We also note that (2.34) can be written under the form

$$w_h(x) \geq c_{25} \quad \forall x \in \cap B_{\frac{r}{2}}(\xi). \quad (2.44)$$

Step 1. : $r \leq s \leq 4r$. By Lemma 2.20 and the above inequality we have that

$$w_h(x_{\frac{r}{2}}(\xi)) \geq \frac{c_{25}}{c_{30}} w_h(x) \quad \forall x \in \Omega.$$

Applying Harnack inequality to w_h in the balls $B_{\frac{(2+j)r}{4}}(x_{\frac{(2+j)r}{4}}(\xi))$ for $j = 0, \dots, j_0 \leq 14$ we obtain

$$w_h(x_{\frac{(2+j)r}{4}}(\xi)) \geq c_{31}^j w_h(x_{\frac{r}{2}}(\xi)) \quad \text{for } j = 1, \dots, j_0.$$

This implies

$$w_h(x_s(\xi)) \geq c_{32} w_h(x) \quad \forall x \in \Omega. \quad (2.45)$$

Step 2: $\beta_0 \geq s > 4r$. We apply Propositions 2.11, 2.12 to w_h in $B_{\frac{s}{2}}(\xi_1) \cap \Omega$ where $\xi_1 \in \partial\Omega$ is such that $|\xi - \xi_1| = s$ and we get

$$w_h(x) \leq c_{18} w_h(x_{\frac{s}{4}}(\xi_1)) \quad \forall x \in B_{\frac{s}{4}}(\xi_1) \cap \Omega \quad (2.46)$$

Then we apply six times Harnack inequality to w_h between $x_{\frac{s}{4}}(\xi_1)$ and $x_s(\xi)$ and obtain

$$w_h(x_{\frac{s}{4}}(\xi_1)) \leq c_{33} w_h(x_s(\xi)). \quad (2.47)$$

Combining (2.46) and (2.47) we derive (2.43).

Step 3. For $\epsilon > 0$, set $z_h = w_h - c_{33} w_h(x_s(\xi)) - \epsilon$. Then z_h^+ has compact support in $\Omega \setminus B_s(\xi)$ and thus belongs to $H_0^1(\Omega \setminus B_s(\xi))$. Integration by parts in (2.42) leads to

$$\int_{\Omega \setminus B_s(\xi)} v_1^2 |\nabla z_h^+|^2 dx = 0. \quad (2.48)$$

Then $z_h^+ = 0$ by letting $\epsilon \rightarrow 0$. Combining with (2.45) and $h \uparrow \chi_{\Delta_r(\xi)}$ implies (2.39). \square

Theorem 2.22. *There exists a constant c_{34} which depends only on Ω such that, for any $0 < r \leq \beta_0$ and $\xi \in \partial\Omega$, there holds*

$$\frac{1}{c_{34}} r^{N-\frac{1}{2}} |\log r| G_{\mathcal{L}_{\frac{1}{4}}}(x_r(\xi), x) \leq \omega^x(\Delta_r(\xi)) \leq c_{34} r^{N-\frac{1}{2}} |\log r| G_{\mathcal{L}_{\frac{1}{4}}}(x_r(\xi), x) \quad \forall x \in \Omega \setminus B_{4r}(\xi). \quad (2.49)$$

Proof. Let $\eta \in C_0^\infty(B_{2r}(\xi))$ such that $0 \leq \eta \leq 1$ and $\eta = 1$ in $B_r(\xi)$. We set

$$u = \eta(-\ln d)\sqrt{d} := \eta\psi,$$

(we assume that $4r < 1$), in order to have

$$\lim_{x \rightarrow x_0} \frac{u(x)}{\psi(x)} = \eta|_{\partial\Omega}(x_0) = \zeta(x_0) \quad \forall x_0 \in \partial\Omega,$$

uniformly with respect to x_0 . Since

$$-\Delta\psi - \frac{1}{4} \frac{\psi}{d^2(x)} = \frac{2 + \ln d}{2\sqrt{d}} \Delta d = -(N-1) \frac{2 + \ln d}{2\sqrt{d}} K$$

where K is the mean curvature of $\partial\Omega$, then u satisfies

$$\begin{aligned} -\Delta u - \frac{1}{4} \frac{u}{d^2(x)} &= -\psi\Delta\eta + \frac{2 + \ln d}{2\sqrt{d}} (2\nabla d \cdot \nabla\eta - (N-1)K\eta) := f && \text{in } \Omega \\ u &= \zeta && \text{on } \partial\Omega. \end{aligned}$$

Then $|f| \leq c_{35}(-\frac{\ln d}{\sqrt{d}})\chi_{\Omega \cap B_{2r}(\xi)}$ since η vanishes outside $B_{2r}(\xi)$. We have by the representation formula [12]

$$0 = u(x) = \int_{\Omega} G_{\mathcal{L}_{\frac{1}{4}}}(x, y) f dy + \int_{\partial\Omega} h(y) d\omega^x(y) \quad \forall x \in \Omega \setminus B_{2r}(\xi). \quad (2.50)$$

By Lemma 2.1, we have that for any $x \in \Omega \setminus B_{4r}(\xi)$ and $y \in B_{2r}(\xi)$

$$G_{\mathcal{L}_{\frac{1}{4}}}(x, y) \leq c_{36} G_{\mathcal{L}_{\frac{1}{4}}}(x, x_r(\xi)),$$

thus

$$\begin{aligned} \omega^x(\Delta_r(\xi)) &\leq \int_{\Omega \cap B_{2r}(\xi)} G_{\mathcal{L}_{\frac{1}{4}}}(x, y) |f(y)| dy \\ &\leq c_{37} G_{\mathcal{L}_{\frac{1}{4}}}(x, x_r(\xi)) \int_{\Omega \cap B_{2r}(\xi)} \frac{|\ln d(y)|}{\sqrt{d(y)}} dy \\ &\leq c_{38} G_{\mathcal{L}_{\frac{1}{4}}}(x, x_r(\xi)) r^{N-\frac{1}{2}} |\ln r|, \end{aligned} \quad (2.51)$$

since

$$\int_{\Omega \cap B_{2r}(\xi)} \frac{|\ln d(y)|}{\sqrt{d(y)}} dy \leq c_{39} r^{N-1} \int_0^{2r} \frac{|\ln t| dt}{\sqrt{t}} \leq 2c_{39} r^{N-\frac{1}{2}} |\ln r|.$$

This implies the right-hand side part of (2.49). For the opposite inequality we observe that if $x \in \partial B_{4r}(\xi)$, there holds by (2.34)

$$\begin{aligned} r^{N-\frac{1}{2}} |\log r| G_{\mathcal{L}_{\frac{1}{4}}}(x_r(\xi), x) &\leq c_{40} r^{N-\frac{1}{2}} |\log r| \min \left\{ \frac{1}{|x - x_r(\xi)|^{N-2}}, \frac{\sqrt{d(x)}\sqrt{d(x_r(\xi))}}{|x - x_r(\xi)|^{N-1}} \right\} \\ &\leq c_{41} \sqrt{d(x)} |\log r| \\ &\leq c_{42} W\left(\frac{x}{8}\right) \\ &\leq \frac{c_{42}}{c_{25}} \omega^{x_r(\xi)}(\Delta_r(\xi)). \end{aligned}$$

We end the proof by Harnack inequality between $\omega^{x_r(\xi)}(\Delta_r(\xi))$ and $\omega^{x_r(\xi)}(\Delta_r(\xi))$. \square

Replacing, in the last proof, the function $\psi = \sqrt{d}(-\ln d)$ by $\tilde{\psi} = d^{\frac{\alpha-}{2}}$, we obtain similarly.

Theorem 2.23. *Assume $\kappa < \frac{1}{4}$. There exists a constant c_{42} which depends only on Ω and κ such that, for any $0 < r \leq \beta_0$ and $\xi \in \partial\Omega$, there holds*

$$\frac{1}{c_{42}} r^{N-1+\frac{\alpha-}{2}} G_{\mathcal{L}_\kappa}(x_r(\xi), x) \leq \omega^x(\Delta_r(\xi)) \leq c_{42} r^{N-1+\frac{\alpha-}{2}} G_{\mathcal{L}_\kappa}(x_r(\xi), x) \quad \forall x \in \Omega \setminus B_{4r}(\xi).$$

As a consequence of Theorems 2.22 and 2.23 and the Harnack inequality, the harmonic measure for \mathcal{L}_κ possesses the doubling property.

Theorem 2.24. *Let $0 < \kappa \leq \frac{1}{4}$. There exists a constant c_{42} which depends only on Ω, κ such that for any $0 < r \leq \beta_0$, there holds*

$$\omega^x(\Delta_{2r}(\xi)) \leq c_{42} \omega^x(\Delta_r(\xi)) \quad \forall x \in \Omega \setminus B_{4r}(\xi).$$

Lemma 2.25. *Let $0 < r \leq \beta_0$ and u be a positive \mathcal{L}_κ -harmonic function such that*

(i) $u \in C(\overline{\Omega} \setminus B_r(\xi))$,

(ii)

$$\lim_{x \rightarrow x_0} \frac{u(x)}{W(x)} = 0 \quad \forall x_0 \in \Omega \setminus \overline{B_r(\xi)},$$

uniformly with respect to x_0 .

Then

$$c_{42}^{-1} u(x_r(\xi)) W^x(\Delta_r(\xi)) \leq u(x) \leq c_{42} u(x_r(\xi)) W^x(\Delta_r(\xi)) \quad \forall x \in \Omega \setminus \overline{B_{2r}(\xi)},$$

with c_{42} depends only on κ and Ω .

Proof. The result follows by Theorems 2.22 and 2.23 and the Harnack inequality like in the proof of Theorem 2.21. \square

2.4 The Poisson kernel of \mathcal{L}_κ

In this section we state some properties of the Poisson kernel associated to \mathcal{L}_κ .

Definition 2.26. *Fix $\xi \in \partial\Omega$. A function K defined in Ω is called a kernel function at ξ with pole at $x_0 \in \Omega$ if*

(i) $K(\cdot, \xi)$ is \mathcal{L}_κ -harmonic in Ω ,

(ii) $K(\cdot, \xi) \in C(\overline{\Omega} \setminus \{\xi\})$ and for any $\eta \in \partial\Omega \setminus \{\xi\}$

$$\lim_{x \rightarrow \eta} \frac{K(x, \xi)}{W(x)} = 0,$$

(iii) $K(x, \xi) > 0$ for each $x \in \Omega$ and $K(x, \xi) = 1$.

Proposition 2.27. *There exists one and only one kernel function for \mathcal{L}_κ at ξ with pole at x_0 .*

Proof. The proof is the same as the one of Theorem 3.1 in [7]. \square

Proposition 2.28. *The kernel function $K_{\mathcal{L}_\kappa}(x, \xi)$, is continuous in ξ on the boundary of Ω .*

Proof. The proof is the same as the one of Corollary 3.2 in [7]. \square

We have proved the uniqueness of Poisson kernel. By (1.14), Theorems 2.22, 2.23 and Proposition 2.1 we have the following result.

Theorem 2.29. *Assume $0 < \kappa \leq \frac{1}{4}$. There exists a positive constant c_{43} such that*

$$\frac{1}{c_{43}} \frac{d^{\frac{\alpha_+}{2}}(y)}{|\xi - y|^{N+\alpha_+-2}} \leq K_{\mathcal{L}_\kappa}(y, \xi) \leq c_{43} \frac{d^{\frac{\alpha_+}{2}}(y)}{|\xi - y|^{N+\alpha_+-2}}. \quad (2.52)$$

Remark 2.30. *As in [15, Remark 3.9], the Martin kernel which is classical defined by*

$$\tilde{K}_{\mathcal{L}_\kappa}(x, \xi) = \lim_{x \rightarrow \xi} \frac{G_{\mathcal{L}_\kappa}(x, y)}{G_{\mathcal{L}_\kappa}(x, x_0)} \quad \forall \xi \in \partial\Omega,$$

coincides with the Poisson kernel $K_{\mathcal{L}_\kappa}$.

Theorem 2.31. *Let u be a positive \mathcal{L}_κ -harmonic in the domain Ω . Then $u \in L^1_{\phi_\kappa}(\Omega)$ and there exists a unique Radon measure μ on $\partial\Omega$ such that*

$$u(x) = \int_{\partial\Omega} K_{\mathcal{L}_\kappa}(x, \xi) d\mu(\xi).$$

Proof. The proof is the same as the one of Theorem 4.3 in [15]. \square

Actually the measure μ is the boundary trace of u . This boundary trace can be achieved in a *dynamic way* as in [?, Sect 2]. We consider a increasing sequence of bounded smooth domains $\{\Omega_n\}$ such that $\overline{\Omega_n} \subset \Omega_{n+1}$, $\cup_n \Omega_n = \Omega$ and $\mathcal{H}^{N-1}(\Omega_n) \rightarrow \mathcal{H}^{N-1}(\Omega)$. such a sequence is a *smooth exhaustion* of Ω . For each n the operator $\mathcal{L}_\kappa^{\Omega_n}$ defined by

$$\mathcal{L}_\kappa^{\Omega_n} u = -\Delta u - \frac{\kappa}{d^2(x)} u \quad (2.53)$$

is uniformly elliptic and coercive in $H_0^1(\Omega_n)$ and its first eigenvalue $\lambda_\kappa^{\Omega_n}$ is larger than λ_κ . If $h \in C(\partial\Omega_n)$ the following problem

$$\begin{aligned} \mathcal{L}_\kappa^{\Omega_n} v &= 0 & \text{in } \Omega_n \\ v &= h & \text{on } \partial\Omega_n \end{aligned} \quad (2.54)$$

admits a unique solution which allows to define the $\mathcal{L}_\kappa^{\Omega_n}$ -harmonic measure on $\partial\Omega_n$ by

$$v(x_0) = \int_{\partial G_{w_n}} h(y) d\omega_{\Omega_n}^{x_0}(y). \quad (2.55)$$

Thus the Poisson kernel of $\mathcal{L}_\kappa^{\Omega_n}$ is

$$K_{\mathcal{L}_\kappa^{\Omega_n}}(x, y) = \frac{d\omega_{\Omega_n}^x(y)}{d\omega_{\Omega_n}^{x_0}(y)} \quad \forall y \in \partial\Omega_n. \quad (2.56)$$

Proposition 2.32. *Assume $0 < \kappa \leq \frac{1}{4}$ and let $x_0 \in \Omega_1$. Then for every $Z \in C(\overline{\Omega})$,*

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} Z(x) W(x) d\omega_{\Omega_n}^{x_0}(x) = \int_{\partial\Omega} Z(x) d\omega^{x_0}(x). \quad (2.57)$$

Proof. We recall that $d \in C^2(\overline{\Omega}_\varepsilon)$ for any $0 < \varepsilon \leq \beta_0$ and let $n_0 \in \mathbb{N}$ be such that

$$\text{dist}(\partial\Omega_n, \partial\Omega) < \frac{\beta_0}{2}, \quad \forall n \geq n_0.$$

For $n \geq n_0$ let w_n be the solution of

$$\begin{aligned} \mathcal{L}_\kappa^{\Omega_n} w_n &= 0 && \text{in } \Omega_n \\ w_n &= W && \text{on } \partial\Omega_n \end{aligned} \quad (2.58)$$

It is straightforward to see that the proof of Propositions 2.17 and 2.18 it is inferred that there exists a positive constant $c_{44} = c_{44}(\Omega, \kappa)$ such that

$$\|w_n\|_{L^\infty(\Omega_n)} \leq c_{44}, \quad \forall n \geq n_0.$$

Furthermore

$$w_n(x_0) = \int_{\partial\Omega_n} W(x) d\omega_{\Omega_n}^{x_0}(x) < c_{45}. \quad (2.59)$$

We extend $\omega_{\Omega_n}^{x_0}$ as a Borel measure on $\overline{\Omega}$ by setting $\omega_{\Omega_n}^{x_0}(\overline{\Omega} \setminus \Omega_n) = 0$, and keep the notation $\omega_{\Omega_n}^{x_0}$ for the extension. Because of (2.59) the sequence $\{W\omega_{\Omega_n}^{x_0}\}$ is bounded in the space $\mathfrak{M}_b(\overline{\Omega})$ of bounded Borel measures in $\overline{\Omega}$. Thus there exists a subsequence (still denoted by $\{W(x)\omega_{\Omega_n}^{x_0}\}$ which converges narrowly to some positive measure, say \tilde{w} which is clearly supported by $\partial\Omega$ and satisfies $\|\tilde{w}\|_{\mathfrak{M}_b} \leq c_{45}$ as in (2.59). For every $Z \in C(\overline{\Omega})$ there holds

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} Z(x) W d\omega_{\Omega_n}^{x_0} = \int_{\partial\Omega} Z d\tilde{w}.$$

Let $\zeta := Z|_{\partial\Omega}$ and $z := \mathbb{K}_{\mathcal{L}_\kappa}[\zeta]$. Then

$$\lim_{d(x) \rightarrow 0} \frac{z(x)}{W(x)} = \zeta \quad \text{and} \quad z(x_0) = \int_{\partial\Omega} \zeta d\omega^{x_0}.$$

By Propositions 2.17 and 2.18, $\frac{z}{W} \in C(\overline{\Omega})$. Since $\frac{z}{W}|_{\partial\Omega_n}$ converges uniformly to ζ , there holds

$$z(x_0) = \int_{\partial\Omega_n} z|_{\partial\Omega_n} d\omega_{\Omega_n}^{x_0} = \int_{\partial\Omega_n} W \frac{z|_{\partial\Omega_n}}{W} d\omega_{\Omega_n}^{x_0} \rightarrow \int_{\partial\Omega} \zeta d\tilde{w} \quad \text{as } n \rightarrow \infty.$$

It follows

$$\int_{\partial\Omega} \zeta d\tilde{w} = \int_{\partial\Omega} \zeta d\omega^{x_0}, \quad \forall \zeta \in C(\partial\Omega).$$

Consequently $\tilde{w} = d\omega^{x_0}$. Because the limit does not depend on the subsequence it follows that the whole sequence $W(x)d\omega_{\Omega_n}^{x_0}$ converges weakly to w . This implies (2.57). \square

In the same way we have

Proposition 2.33. *Let $x_0 \in \Omega_1$ and $\mu \in \mathfrak{M}(\partial\Omega)$. Put $v := \mathbb{K}_{\mathcal{L}_\kappa}[\mu]$, then for every $Z \in C(\overline{\Omega})$,*

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} Z(x) v d\omega_{\Omega_n}^{x_0} = \int_{\partial\Omega} Z(x) d\mu. \quad (2.60)$$

Proof. The proof is same as the proof of Lemma 2.2 in [?] and we omit it. \square

The next result is an analogous of Green formula for positive \mathcal{L}_κ -harmonic functions.

Proposition 2.34. *Let v be a positive \mathcal{L}_κ -harmonic function in Ω with boundary trace μ . Let $Z \in C^2(\bar{\Omega})$ and $\tilde{G} \in C(\Omega)$ which coincides with $G_{\mathcal{L}_\kappa}(x_0, \cdot)$ in Ω_δ for some $0 < \delta < \beta_0$ and some $x_0 \notin \bar{\Omega}_{\beta_0}$. Assume*

$$|\nabla \tilde{G} \cdot \nabla Z| \leq c'_{45} \phi_\kappa. \quad (2.61)$$

Then, if we set $\zeta = Z\tilde{G}$, there holds

$$\int_{\Omega} v \mathcal{L}_\kappa \zeta dx = \int_{\partial\Omega} Z d\mu. \quad (2.62)$$

Proof. Let $\{\Omega_j\}$ be a smooth exhaustion of Ω with Green kernel $G_{\mathcal{L}_\kappa}^{\Omega_j}$ and Poisson kernel $P_{\mathcal{L}_\kappa}^{\Omega_j} = -\partial_{\mathbf{n}} G_{\mathcal{L}_\kappa}^{\Omega_j}$. We assume that $j \geq j_0$ where $\bar{\Omega}'_\delta \subset \Omega_j$. Set $\zeta_j = Z\tilde{G}_j$, where the functions \tilde{G}_j are C^∞ in Ω_j , coincide with $G_{\mathcal{L}_\kappa}^{\Omega_j}(x_0, \cdot)$ in $\Omega_j \cap \bar{\Omega}'_\delta$ and satisfy $\tilde{G}_j \rightarrow \tilde{G}$ in $C^2(\Omega)$ -loc and such that $|\nabla \tilde{G}_j \cdot \nabla Z| \leq c'_{45} \phi_\kappa$.

$$\int_{\Omega_j} v \mathcal{L}_\kappa \zeta_j dx = - \int_{\partial\Omega_j} v \frac{\partial \zeta_j}{\partial \mathbf{n}} dS = - \int_{\partial\Omega_j} v Z \frac{\partial \tilde{G}_j}{\partial \mathbf{n}} dS = \int_{\partial\Omega_n} v Z P_{\mathcal{L}_\kappa}^{\Omega_j}(x_0, \cdot) dS = \int_{\partial\Omega_j} v Z d\omega_{\Omega_j}^{x_0}.$$

By (2.60)

$$\int_{\partial\Omega_j} v Z d\omega_{\Omega_j}^{x_0} \rightarrow \int_{\partial\Omega} Z(x) d\mu \quad \text{as } j \rightarrow \infty.$$

Next

$$\mathcal{L}_\kappa \zeta_j = Z \mathcal{L}_\kappa \tilde{G}_j + \tilde{G}_j \Delta Z + 2 \nabla \tilde{G}_j \cdot \nabla Z.$$

Since $v \in L^1_{\phi_\kappa}(\Omega)$, the proof follows. \square

Remark. In the statement of Proposition 2.34, it can be easier in some cases to replace \tilde{G} and \tilde{G}_j by ϕ_κ and $\phi_{\mathcal{L}_\kappa}^{\Omega_j}$. The assumption (2.61) has to be replaced by

$$|\nabla \phi_\kappa \cdot \nabla Z| \leq c'_{45} \phi_\kappa, \quad (2.63)$$

and the modification of the proof is straightforward.

3 The nonlinear problem with measures data

3.1 The linear boundary value problem with L^1 data

In the sequel we denote by $\omega = \omega^{x_0}$ the \mathcal{L}_κ -harmonic measure in Ω , for some fixed $x_0 \in \Omega$ and by $\mathfrak{M}_{\phi_\kappa}(\Omega)$ be the space of Radon measures ν in Ω such that $\phi_\kappa d|\nu|$ is a bounded measure. We also denote by $\mathfrak{M}(\partial\Omega)$ the space of Radon measures on $\partial\Omega$ with respective norms $\|\nu\|_{\mathfrak{M}_{\phi_\kappa}(\Omega)}$ and $\|\mu\|_{\mathfrak{M}(\partial\Omega)}$. Their respective positive cones are denoted by $\mathfrak{M}_{\phi_\kappa}^+(\Omega)$ and $\mathfrak{M}^+(\partial\Omega)$. By Fubini's theorem and (2.9), for any $\nu \in \mathfrak{M}_{\phi_\kappa}(\Omega)$ we can define

$$\mathbb{G}_{\mathcal{L}_\kappa}[\nu](x) = \int_{\Omega} G_{\mathcal{L}_\kappa}(x, y) d\nu(y),$$

and we have

$$\|\mathbb{G}_{\mathcal{L}_\kappa}[\nu]\|_{L^1_{\phi_\kappa}(\Omega)} \leq c_{46} \|\nu\|_{\mathfrak{M}_{\phi_\kappa}(\Omega)}. \quad (3.1)$$

If $\mu \in \mathfrak{M}(\partial\Omega)$, we set

$$\begin{aligned}\mathbb{K}_{\mathcal{L}_\kappa}[\mu](x) &= \int_{\partial\Omega} K_{\mathcal{L}_\kappa}(x, y) d\mu(y), \\ \|\mathbb{K}_{\mathcal{L}_\kappa}[\mu]\|_{L^1_{\phi_\kappa}(\Omega)} &\leq c_{47} \|\mu\|_{\mathfrak{M}(\partial\Omega)}.\end{aligned}\quad (3.2)$$

In the above inequalities c_{46} and c_{47} are positive constants depending on Ω and κ .

For $0 < \kappa \leq \frac{1}{4}$, we define the space of test functions $\mathbf{X}(\Omega)$ by

$$\mathbf{X}(\Omega) = \left\{ \eta \in H^1_{loc}(\Omega) : \frac{\eta}{d^{\frac{\alpha+1}{2}}} \in H^1(\Omega, d^{\alpha+1} dx), (\phi_\kappa)^{-1} \mathcal{L}_\kappa \eta \in L^\infty(\Omega) \right\}. \quad (3.3)$$

The next statement follows immediately from Propositions (2.9) and (2.10).

Lemma 3.1. *Let $0 < \kappa \leq \frac{1}{4}$. If $m \in L^\infty(\Omega)$, the solution η_m of*

$$\begin{aligned}\mathcal{L}_\kappa \eta_m &= m \phi_\kappa && \text{in } \Omega \\ \eta_m &= 0 && \text{on } \partial\Omega\end{aligned}\quad (3.4)$$

obtained by Propositions (2.9) and (2.10) with $f_0 = m$ and $h = 0$ belongs to $\mathbf{X}(\Omega)$. Furthermore

$$-\frac{\|m_-\|_{L^\infty(\Omega)}}{\lambda_\kappa} \phi_\kappa \leq -\eta_{m_-} \leq \eta_m \leq \eta_{m_+} \leq \frac{\|m_+\|_{L^\infty(\Omega)}}{\lambda_\kappa} \phi_\kappa. \quad (3.5)$$

In the next Proposition we give some key estimates for the weak solutions of

$$\begin{aligned}\mathcal{L}_\kappa u &= f && \text{in } \Omega \\ u &= h && \text{on } \partial\Omega\end{aligned}\quad (3.6)$$

Proposition 3.2. *For any $(f, h) \in L^1_{\phi_\kappa}(\Omega) \times L^1(\partial\Omega, d\omega)$ there exists a unique $u := u_{f,h} \in L^1_{\phi_\kappa}(\Omega)$ such that*

$$\int_{\Omega} u \mathcal{L}_\kappa \eta dx = \int_{\Omega} f \eta dx + \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[h\omega] \mathcal{L}_\kappa \eta dx \quad \forall \eta \in \mathbf{X}(\Omega). \quad (3.7)$$

There holds

$$u = \mathbb{G}_{\mathcal{L}_\kappa}[f] + \mathbb{K}_{\mathcal{L}_\kappa}[h\omega] \quad (3.8)$$

and

$$\|u\|_{L^1_{\phi_\kappa}(\Omega)} \leq c_{46} \|f\|_{L^1_{\phi_\kappa}(\Omega)} + c_{47} \|h\|_{L^1(\partial\Omega, d\omega)}. \quad (3.9)$$

Furthermore, for any $\eta \in \mathbf{X}(\Omega)$, $\eta \geq 0$, we have

$$\int_{\Omega} |u| \mathcal{L}_\kappa \eta dx \leq \int_{\Omega} f \eta \operatorname{sgn}(u) dx + \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[|h|\omega] \mathcal{L}_\kappa \eta dx, \quad (3.10)$$

and

$$\int_{\Omega} u_+ \mathcal{L}_\kappa \eta dx \leq \int_{\Omega} f \eta \operatorname{sgn}_+(u) dx + \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[h_+\omega] \mathcal{L}_\kappa \eta dx, \quad (3.11)$$

Proof. Step 1: proof of estimate (3.9). Assume u satisfies (3.7). If $\eta = \eta_{\operatorname{sgn}(u)}$, we have

$$\int_{\Omega} |u| \phi_\kappa dx = \int_{\Omega} u \mathcal{L}_\kappa \eta dx = \int_{\Omega} f \eta dx + \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[h\omega] \operatorname{sgn}(u) \phi_\kappa dx.$$

By (3.1), (3.2)

$$\int_{\Omega} f \eta dx \leq \frac{1}{\lambda_{\kappa}} \int_{\Omega} |f| \phi_{\kappa} dx,$$

$$\int_{\Omega} \mathbb{K}_{\mathcal{L}_{\kappa}} [h\omega] \operatorname{sgn}(u) \phi_{\kappa} dx \leq c_{47} \int_{\partial\Omega} |h| \omega \operatorname{sgn}(u) \phi_{\kappa} dx,$$

which implies (3.9) and uniqueness.

Step 2: proof of existence. If f and h are bounded, existence follows from Propositions 2.9, 2.10. In the general case let $\{(f_n, h_n)\}$ be a sequence of bounded measurable functions in Ω and $\partial\Omega$ which converges to $\{(f, h)\}$ in $L^1_{\phi_{\kappa}}(\Omega) \times L^1(\partial\Omega, d\omega)$. Let $\{u_n\} = \{u_{f_n, h_n}\}$ be the sequence weak solutions of (3.6). By estimate (3.9) it is a Cauchy sequence in $L^1_{\phi_{\kappa}}(\Omega)$ which converges to u . Letting $n \rightarrow \infty$ in identity

$$\int_{\Omega} u_n \mathcal{L}_{\kappa} \eta dx = \int_{\Omega} f_n \eta dx + \int_{\Omega} \mathbb{K}_{\mathcal{L}_{\kappa}} [h_n \omega] \mathcal{L}_{\kappa} \eta dx \quad (3.12)$$

where $\eta \in \mathbf{X}(\Omega)$ implies that $u = u_{f, h}$.

Step 3: proof of estimates (3.10), (3.11). We first assume that f is bounded and h is C^2 . For $\epsilon > 0$ we set $\mathcal{L}_{\kappa, \epsilon} = -\Delta - \frac{\kappa}{d^2 + \epsilon}$. Let $\lambda_{\kappa, \epsilon}$ be the first eigenvalue of $\mathcal{L}_{\kappa, \epsilon}$ in $H^1_0(\Omega)$. Then $\lambda_{\kappa, \epsilon} > \lambda_{\kappa}$, the mapping $\epsilon \mapsto \lambda_{\kappa, \epsilon}$ is increasing, and by Section 2.1-5 it converges to λ_{κ} when $\epsilon \rightarrow 0$. Furthermore the corresponding eigenfunction $\phi_{\kappa, \epsilon}$ normalized by $\sup \phi_{\kappa, \epsilon} = 1$ converges to ϕ_{κ} (with the same normalization) in $C^{\gamma}(\bar{\Omega})$, for some $0 < \gamma < \beta$. We denote by u_{ϵ} the solution of

$$\begin{aligned} \mathcal{L}_{\kappa, \epsilon} u_{\epsilon} &= f & \text{in } \Omega \\ u_{\epsilon} &= h & \text{on } \partial\Omega. \end{aligned} \quad (3.13)$$

We write $f = f^+ - f^-$, $h = h^+ - h^-$ and $u_{\epsilon} = u_{\epsilon, +} - u_{\epsilon, -}$ where

$$\begin{aligned} \mathcal{L}_{\kappa, \epsilon} u_{\epsilon, \pm} &= f^{\pm} & \text{in } \Omega \\ u_{\epsilon, \pm} &= h^{\pm} & \text{on } \partial\Omega. \end{aligned} \quad (3.14)$$

Then $u_{\epsilon} = u_{\epsilon, +} - u_{\epsilon, -}$, $u_{\epsilon, +} \uparrow u_+$ and $u_{\epsilon, -} \uparrow u_-$ as $\epsilon \rightarrow 0$ in $C^{\beta}(\bar{\Omega})$, where

$$\begin{aligned} \mathcal{L}_{\kappa} u_{\pm} &= f^{\pm} & \text{in } \Omega \\ u_{\pm} &= h^{\pm} & \text{on } \partial\Omega. \end{aligned} \quad (3.15)$$

Moreover $d^{\frac{\alpha+}{2}} u_{\epsilon, \pm} \rightharpoonup d^{\frac{\alpha+}{2}} u_{\pm}$ in $H^1(\Omega, d^{\alpha+} dx)$ weakly. Thus $u_{\epsilon, +} - u_{\epsilon, -} \rightharpoonup u_+ - u_-$ in $C^{\beta}(\bar{\Omega})$ and $d^{\frac{\alpha+}{2}} (u_{\epsilon, +} - u_{\epsilon, -}) \rightharpoonup d^{\frac{\alpha+}{2}} (u_+ - u_-)$ weakly in $H^1(\Omega, d^{\alpha+} dx)$ which yields $u_+ - u_- = u_{f, h}$.

Let $\eta \in \mathbf{X}(\Omega)$, $\eta \geq 0$. Then $\mathcal{L}_{\kappa} \eta = m_{\eta} \phi_{\kappa}$ with $m_{\eta} \in L^{\infty}$. By replacing m by $m + \delta$ ($\delta > 0$), we can always suppose that $\eta \geq \delta' \phi_{\kappa}$ for some $\delta' > 0$. There exists $\eta_{\epsilon} \in C^1_0(\bar{\Omega}) \cap W^{2,p}(\Omega)$ ($1 < p < \infty$) solution of

$$\begin{aligned} \mathcal{L}_{\kappa, \epsilon} \eta_{\epsilon} &= m_{\eta} \phi_{\kappa} & \text{in } \Omega \\ \eta_{\epsilon} &= 0 & \text{on } \partial\Omega. \end{aligned} \quad (3.16)$$

Since $\eta - \eta_{\epsilon}$ satisfies

$$\mathcal{L}_{\kappa, \epsilon} (\eta - \eta_{\epsilon}) = \kappa \left(\frac{1}{d^2} - \frac{1}{d^2 + \epsilon} \right) \eta \geq 0$$

there holds $\eta > \eta_{\epsilon}$. By Propositions 2.9, 2.10, $\eta_{\epsilon} \uparrow \eta$ in $C^{\gamma}(\bar{\Omega})$. We first assume that f and h are bounded functions. Assuming $\eta_{\epsilon} \geq 0$, and since $\mathcal{L}_{\kappa} \eta = \mathcal{L}_{\kappa, \epsilon} \eta_{\epsilon}$ we have classically [26, Th 2.4]

$$\int_{\Omega} |u_{\epsilon}| \mathcal{L}_{\kappa} \eta dx \leq \int_{\Omega} f \eta_{\epsilon} \operatorname{sgn}(u_{\epsilon}) dx + \int_{\Omega} \mathbb{K}_{\mathcal{L}_{\kappa, \epsilon}} [|h| \omega] \mathcal{L}_{\kappa} \eta dx, \quad (3.17)$$

Since $\mathbb{K}_{\mathcal{L}_{\kappa,\varepsilon}}[|h|\omega] \rightarrow \mathbb{K}_{\mathcal{L}_{\kappa}}[|h|\omega]$ uniformly in $\bar{\Omega}$ by Propositions 2.9 and 2.10, we derive (3.10) when the data f and h are regular enough. In the general case we consider a sequence $(f_n, h_n) \subset L^\infty(\Omega) \times C^2(\partial\Omega)$ which converges to (f, h) in $L^1(\Omega) \times L^1(\partial\Omega, d\omega)$. Since u_{f_n, h_n} converges to $u_{f, h}$ in $L^1_{\phi_\kappa}(\Omega)$ we obtain (3.10) from the inequality verified by any $\eta \in \mathbf{X}(\Omega)$

$$\int_{\Omega} |u_{f_n, h_n}| \mathcal{L}_{\kappa} \eta dx \leq \int_{\Omega} f_n \eta \operatorname{sgn}(u) dx + \int_{\Omega} \mathbb{K}_{\mathcal{L}_{\kappa}}[|h_n|\omega] \mathcal{L}_{\kappa} \eta dx.$$

The proof of (3.11) is similar. \square

3.2 General nonlinearities

Throughout this section Ω is a smooth bounded domain and κ a real number in $(0, \frac{1}{4}]$. Let $g : \mathbb{R} \mapsto \mathbb{R}$ be a nondecreasing continuous function, vanishing at 0 for simplicity. The problem under consideration is the following

$$\begin{aligned} -\Delta u - \frac{\kappa}{d^2} u + g(u) &= \nu & \text{in } \Omega \\ u &= \mu & \text{in } \partial\Omega \end{aligned} \quad (3.18)$$

where ν and μ are Radon measures respectively in Ω and $\partial\Omega$.

Definition. Let $\nu \in \mathfrak{M}_{\phi_\kappa}(\Omega)$ and $\mu \in \mathfrak{M}(\partial\Omega)$. We say that u is a solution of (3.18) if $u \in L^1_{\phi_\kappa}(\Omega)$, $g(u) \in L^1_{\phi_\kappa}(\Omega)$ and for any $\eta \in \mathbf{X}(\Omega)$ there holds

$$\int_{\Omega} (u \mathcal{L}_{\kappa} \eta + g(u) \eta) dx = \int_{\Omega} (\eta d\nu + \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu] \mathcal{L}_{\kappa} \eta) dx \quad (3.19)$$

Our main existence result for *subcritical nonlinearities* is the following.

Theorem 3.3. *Assume g satisfies*

$$\int_1^\infty (g(s) - g(-s)) s^{-2 \frac{N-1+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} ds < \infty. \quad (3.20)$$

Then for any $(\nu, \mu) \in \mathfrak{M}_{\phi_\kappa}(\Omega) \times \mathfrak{M}(\partial\Omega)$ problem (3.18) admits a unique solution $u = u_{\nu, \mu}$. Furthermore the mapping $(\nu, \mu) \mapsto u_{\nu, \mu}$ is increasing and stable in the sense that if $\{(\nu_n, \mu_n)\}$ converge to (ν, μ) in the weak sense of measures, $\{u_{\nu_n, \mu_n}\}$ converges to $u_{\nu, \mu}$ in $L^1_{\phi_\kappa}(\Omega)$.

The proof is based upon estimates of $\mathbb{M}_{\mathcal{L}_{\kappa}}$ and $\mathbb{K}_{\mathcal{L}_{\kappa}}$ into Marcinkiewicz spaces.

Lemma 3.4. *Let $\nu \in \mathfrak{M}_{\phi_\kappa}^+(\Omega)$, $\mu \in \mathfrak{M}^+(\partial\Omega)$ and for $s > 0$, $E_s(\nu) = \{x \in \Omega : \mathbb{G}_{\mathcal{L}_{\kappa}}[\nu](x) > s\}$ and $F_s(\mu) = \{x \in \Omega : \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu](x) > s\}$. If we denote*

$$\mathcal{E}_s(\nu) = \int_{E_s(\nu)} \phi_\kappa dx \quad \text{and} \quad \mathcal{F}_s(\mu) = \int_{F_s(\mu)} \phi_\kappa dx,$$

there holds

$$\mathcal{E}_s(\nu) + \mathcal{F}_s(\mu) \leq c_{47} \left(\frac{\|\nu\|_{\mathfrak{M}_{\phi_\kappa}(\Omega)} + \|\mu\|_{\mathfrak{M}(\partial\Omega)}}{s} \right)^{\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}}. \quad (3.21)$$

Proof. Step 1: estimate of $\mathcal{F}_s(\nu)$. By estimate (2.52), for any $\xi \in \partial\Omega$,

$$F_s(\delta_\xi) \subset \tilde{F}_s(\delta_\xi) := \left\{ x \in \Omega : \frac{d^{\frac{\alpha_+}{2}}(x)}{|x - \xi|^{N+\alpha_+-2}} \geq \frac{s}{c_{43}} \right\} \subset B_{(\frac{c_{43}}{s})^\theta}(\xi),$$

with $\theta = \frac{1}{N-2+\frac{\alpha_+}{2}}$. From (2.1), (2.2)

$$\mathcal{F}_s(\delta_\xi) \leq \int_{B_{(\frac{c_{43}}{s})^\theta}(\xi)} \phi_\kappa dx \leq c_{49} \int_{B_{(\frac{c_{43}}{s})^\theta}(\xi)} |x - \xi|^{\frac{\alpha_+}{2}} dx = c_{50} s^{-\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}}.$$

Therefore, for any $s_0 > 0$ and any Borel set $G \subset \Omega$

$$\begin{aligned} \int_G K_{\mathcal{L}_\kappa}(x, \xi) \phi_\kappa dx &\leq s_0 \int_G \phi_\kappa dx + \int_{F_{s_0}(\delta_\xi)} K_{\mathcal{L}_\kappa}(x, \xi) \phi_\kappa dx \\ &\leq s_0 \int_G \phi_\kappa dx - \int_{s_0}^\infty s d\mathcal{F}_s(\delta_\xi) \\ &\leq s_0 \int_G \phi_\kappa dx + c_{50} \int_{s_0}^\infty s^{-\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} ds \\ &\leq s_0 \int_G \phi_\kappa dx + c_{51} s_0^{-\frac{2}{N-2+\frac{\alpha_+}{2}}} \end{aligned}$$

Next we choose s_0 so that the two terms in the right part of the last inequality are equal and we get

$$\int_G K_{\mathcal{L}_\kappa}(x, \xi) \phi_\kappa dx \leq c_{52} \left(\int_G \phi_\kappa dx \right)^{\frac{2}{N+\frac{\alpha_+}{2}}}. \quad (3.22)$$

Henceforth, for any $\mu \in \mathfrak{M}(\partial\Omega)$, there holds by Fubini's theorem,

$$\int_G \mathbb{K}_{\mathcal{L}_\kappa}[|\mu|] \phi_\kappa dx = \int_\Omega \int_G K_{\mathcal{L}_\kappa}(x, \xi) \phi_\kappa(x) dx d|\mu|(\xi) \leq c_{52} \|\mu\|_{\mathfrak{M}(\partial\Omega)} \left(\int_G \phi_\kappa dx \right)^{\frac{2}{N+\frac{\alpha_+}{2}}}. \quad (3.23)$$

If we take in particular $G = F_s(|\mu|)$, we derive

$$s\mathcal{F}_s(|\mu|) \leq c_{52} \|\mu\|_{\mathfrak{M}(\partial\Omega)} (\mathcal{F}_s(|\mu|))^{\frac{2}{N+\frac{\alpha_+}{2}}},$$

which yields to (3.21) with $\nu = 0$.

Step 2: estimate of $\mathcal{E}_s(\nu)$. By estimate (2.9), for any $y \in \Omega$,

$$E_s(\delta_y) \subset \tilde{E}_s(\delta_y) := \left\{ x \in \Omega : \frac{d^{\frac{\alpha_+}{2}}(y) d^{\frac{\alpha_+}{2}}(x)}{|x - y|^{N+\alpha_+-2}} \geq \frac{s}{c_3} \right\} \cap \left\{ x \in \Omega : \frac{1}{|x - y|^{N-2}} \geq \frac{s}{c_3} \right\},$$

A simple geometric verification shows that there exists an open domain $\mathcal{O} \subset \bar{\mathcal{O}} \subset \Omega$ such that $y \in \mathcal{O}$, $\text{dist}(y, \mathcal{O}^c) > \lambda_1 d(y)$, $\mathcal{O} \subset B_{\lambda_2 d(y)}(y)$ for some $0 < \lambda_1 < \lambda_2 < 1$ independent of y with the following properties

$$\begin{aligned} x \in \mathcal{O} &\implies \frac{d^{\frac{\alpha_+}{2}}(y) d^{\frac{\alpha_+}{2}}(x)}{|x - y|^{N+\alpha_+-2}} \geq \frac{1}{|x - y|^{N-2}} \\ x \in \mathcal{O}^c &\implies \frac{d^{\frac{\alpha_+}{2}}(y) d^{\frac{\alpha_+}{2}}(x)}{|x - y|^{N+\alpha_+-2}} \leq \frac{1}{|x - y|^{N-2}}. \end{aligned}$$

Notice that if $\Omega = \mathbb{R}_+^N$ then $\mathcal{O} = B_{\frac{\sqrt{s}}{2}}(\tilde{y})$ where $d(\tilde{y}) = \frac{3}{2}d(y)$. Set

$$\tilde{E}_s^1(\delta_y) = \left\{ x \in \Omega : \frac{1}{|x-y|^{N-2}} \geq \frac{s}{c_3} \right\} \cap \mathcal{O}$$

and

$$\tilde{E}_s^2(\delta_y) = \left\{ x \in \Omega \setminus \mathcal{O} : \frac{d^{\frac{\alpha_+}{2}}(y)d^{\frac{\alpha_+}{2}}(x)}{|x-y|^{N+\alpha_+-2}} \geq \frac{s}{c_3} \right\}$$

In Appendix 2, we prove

$$\mathcal{E}_s(\delta_y) = \int_{E_s(\delta_y)} \phi_\kappa dx \leq \int_{\tilde{E}_s(\delta_y)} \phi_\kappa dx \leq c_{53} s^{-\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} (d(y))^{\frac{\alpha_+(N+\frac{\alpha_+}{2})}{2N-4+\alpha_+}}.$$

As in step 1, for any Borel subset $\Theta \subset \Omega$, we write

$$\begin{aligned} \int_{\Theta} G_{\mathcal{L}_\kappa}(x, y) \phi_\kappa dx &\leq s_0 \int_{\Theta} \phi_\kappa dx + \int_{E_{s_0}(\delta_y)} G_{\mathcal{L}_\kappa}(x, y) \phi_\kappa dx \\ &\leq s_0 \int_{\Theta} \phi_\kappa dx - \int_{s_0}^{\infty} s d\mathcal{E}_s(\delta_y) \\ &\leq s_0 \int_{\Theta} \phi_\kappa dx + c_{53} (d(y))^{\frac{\alpha_+(N+\frac{\alpha_+}{2})}{2N-4+\alpha_+}} \int_{s_0}^{\infty} s^{-\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} ds \\ &\leq s_0 \int_{\Theta} \phi_\kappa dx + c_{54} (d(y))^{\frac{\alpha_+(N+\frac{\alpha_+}{2})}{2N-4+\alpha_+}} s_0^{-\frac{2}{N-2+\frac{\alpha_+}{2}}} \\ \int_{\Theta} G_{\mathcal{L}_\kappa}(x, y) \phi_\kappa dx &\leq c_{55} (d(y))^{\frac{\alpha_+}{2}} \left(\int_G \phi_\kappa dx \right)^{\frac{2}{N+\frac{\alpha_+}{2}}} \leq c_{56} \phi_\kappa(y) \left(\int_G \phi_\kappa dx \right)^{\frac{2}{N+\frac{\alpha_+}{2}}}. \end{aligned} \quad (3.24)$$

Thus, for any $\nu \in \mathfrak{M}_{\phi_\kappa}(\Omega)$, we have

$$\int_{\Theta} \mathbb{G}_{\mathcal{L}_\kappa}[|\nu|] \phi_\kappa dx = \int_{\Omega} \int_{\Theta} G_{\mathcal{L}_\kappa}(x, y) \phi_\kappa(x) dx d|\nu|(y) \leq c_{55} \|\nu\|_{\mathfrak{M}_{\phi_\kappa}(\Omega)} \left(\int_{\Theta} \phi_\kappa dx \right)^{\frac{2}{N+\frac{\alpha_+}{2}}}. \quad (3.25)$$

Thus (3.21) holds. \square

Proof of Theorem 3.3. Step 1: existence and uniqueness. Let $\{(\nu_n, \mu_n)\} \subset C(\bar{\Omega}) \times C^1(\partial\Omega)$ which converges to (ν, μ) in the weak sense of measures in $\mathfrak{M}_{\phi_\kappa}(\Omega) \times \mathfrak{M}(\partial\Omega)$ and J_κ be the functional defined in $L^2(\Omega)$ by the expression

$$\mathcal{J}_\kappa(w) = \frac{1}{2} \int_{\Omega} \left(|\nabla w|^2 - \frac{\kappa}{d^2} w^2 + 2J(w) \right) dx - \int_{\Omega} \nu_n w \phi_\kappa dx - \int_{\partial\Omega} \mu_n w d\omega \quad (3.26)$$

where $J(w) = \int 0wg(t)dt$ with domain

$$D(\mathcal{J}_\kappa) = \{w \in \mathbf{H}_\kappa(\Omega) : J(w) \in L^1(\Omega)\},$$

(see definition in 2.1-5). By (2.7), \mathcal{J}_κ is a convex lower semicontinuous and coercive functional over $L^2(\Omega)$. Let $u_n = u_{\nu_n, \mu_n}$ be its minimum, it is the solution of

$$\begin{aligned} \mathcal{L}_\kappa u_n + g(u_n) &= \nu_n && \text{in } \Omega \\ u_n &= \mu_n && \text{in } \partial\Omega, \end{aligned} \quad (3.27)$$

and for any $\eta \in \mathbf{X}(\Omega)$, there holds

$$\int_{\Omega} (u_n \mathcal{L}_{\kappa} \eta + g(u_n) \eta) dx = \int_{\Omega} (\nu_n \eta + \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n \omega] \mathcal{L}_{\kappa} \eta) dx. \quad (3.28)$$

By Proposition 3.2 (3.10), there holds, with $\eta = \phi_{\kappa}$,

$$\begin{aligned} \int_{\Omega} (\lambda_{\kappa} |u_n| + |g(u_n)|) \phi_{\kappa} dx &\leq \int_{\Omega} (|\nu_n| + \mathbb{K}_{\mathcal{L}_{\kappa}}[|\mu_n| \omega]) \phi_{\kappa} dx \\ &\leq c_{46} \|\nu_n\|_{\mathfrak{M}_{\phi_{\kappa}}(\Omega)} + c_{47} \|\mu_n\|_{\mathfrak{M}(\partial\Omega)} \\ &\leq c_{57}. \end{aligned} \quad (3.29)$$

Moreover

$$-\mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^-] - \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^- \omega] \leq u_n \leq \mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^+] + \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^+ \omega]. \quad (3.30)$$

By using the local L^1 regularity theory for elliptic equations we obtain that the sequence $\{u_n\}$ is relatively compact in the L^1 -local topology in Ω and that there exist a subsequence still denoted by $\{u_n\}$ and a function $u \in L^1_{\phi_{\kappa}}(\Omega)$ such that $u_n \rightarrow u$ a.e. in Ω . By (3.30)

$$|g(u_n)| \leq g(\mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^+] + \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^+ \omega]) - g(-\mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^-] - \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^- \omega]). \quad (3.31)$$

We prove the convergence of $\{g(u_n)\}$ to $g(u)$ in $L^1_{\phi_{\kappa}}(\Omega)$ by the uniform integrability in the following way: let $G \subset \Omega$ be a Borel subset. Then for any $s_0 > 0$

$$\begin{aligned} \int_G |g(u_n)| \phi_{\kappa} dx &\leq \int_G (g(\mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^+] + \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^+ \omega]) - g(-\mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^-] - \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^- \omega])) \phi_{\kappa} dx \\ &\leq s_0 \int_G \phi_{\kappa} dx + \int_{E_s(\nu^+)} g(\mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^+]) \phi_{\kappa} dx + \int_{F_s(\mu^+)} g(\mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^+]) \phi_{\kappa} dx \\ &\quad - \int_{E_s(\nu^-)} g(-\mathbb{G}_{\mathcal{L}_{\kappa}}[\nu_n^-]) \phi_{\kappa} dx - \int_{F_s(\mu^-)} g(-\mathbb{K}_{\mathcal{L}_{\kappa}}[\mu_n^-]) \phi_{\kappa} dx \\ &\leq s_0 \int_G \phi_{\kappa} dx - \int_{s_0}^{\infty} g(s)(d\mathcal{E}_s(\nu_n^+) + d\mathcal{F}_s(\mu_n^+)) + \int_{s_0}^{\infty} g(-s)(d\mathcal{E}_s(\nu_n^-) + d\mathcal{F}_s(\mu_n^-)). \end{aligned}$$

But,

$$\begin{aligned} -\int_{s_0}^{\infty} g(s) d\mathcal{E}_s(\nu_n^+) &= g(s_0) \mathcal{E}_{s_0}(\nu_n^+) + \int_{s_0}^{\infty} \mathcal{E}_s(\nu_n^+) dg(s) \\ &\leq g(s_0) \mathcal{E}_{s_0}(\nu_n^+) + c_{47} (\|\nu_n^+\|_{\mathfrak{M}_{\phi_{\kappa}}})^{\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} \int_{s_0}^{\infty} s^{-\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} dg(s) \\ &\leq \frac{2N+\alpha_+}{2N-4+\alpha_+} c_{47} (\|\nu_n^+\|_{\mathfrak{M}_{\phi_{\kappa}}(\Omega)})^{\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} \int_{s_0}^{\infty} s^{-2\frac{N-1+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} g(s) ds. \end{aligned}$$

All the other terms yields similar estimates which finally yields to

$$\begin{aligned} \int_G |g(u_n)| \phi_{\kappa} dx &\leq s_0 \int_G \phi_{\kappa} dx \\ &\quad + c_{58} (\|\nu_n\|_{\mathfrak{M}_{\phi_{\kappa}}(\Omega)} + \|\mu_n\|_{\mathfrak{M}(\partial\Omega)})^{\frac{N+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} \int_{s_0}^{\infty} s^{-2\frac{N-1+\frac{\alpha_+}{2}}{N-2+\frac{\alpha_+}{2}}} (g(s) - g(-s)) ds \end{aligned} \quad (3.32)$$

Since $\|\nu_n\|_{\mathfrak{M}_{\phi_\kappa}(\Omega)} + \|\mu_n\|_{\mathfrak{M}(\partial\Omega)}$ is bounded independently of n , we obtain easily, using (3.20) and fixing s_0 first, that for any $\epsilon > 0$, there exists $\delta > 0$ such that

$$\int_G \phi_\kappa dx \leq \delta \implies \int_G |g(u_n)| \phi_\kappa dx \leq \epsilon. \quad (3.33)$$

Since

$$|u_n| \leq \mathbb{G}_{\mathcal{L}_\kappa}[\nu_n] + \mathbb{K}_{\mathcal{L}_\kappa}[\mu_n|\omega],$$

we have by (3.25), (3.25)

$$\int_G |u_n| \phi_\kappa dx \leq (c_{52}\|\mu_n\|_{\mathfrak{M}(\partial\Omega)} + c_{55}\|\nu_n\|_{\mathfrak{M}_{\phi_\kappa}(\Omega)}) \left(\int_G \phi_\kappa dx \right)^{\frac{2}{N + \frac{\alpha_\pm}{2}}}. \quad (3.34)$$

This implies the uniform integrability of the sequence $\{u_n\}$. Letting $n \rightarrow \infty$ in identity (3.28), we conclude that (3.19) holds. Uniqueness, as well as the monotonicity of the mapping $(\nu, \mu) \mapsto u_{\nu, \mu}$, is an immediate consequence of (3.10), (3.11) and the monotonicity of g .

Step 2: stability. The stability is a direct consequence of inequalities (3.32) and (3.34) which show the uniform integrability of the sequence $(u_n, g(u_n))$ in $L^1_{\phi_\kappa}(\Omega) \times L^1_{\phi_\kappa}(\Omega)$. \square

The proof of the following result is similar as the one of [23, Lemma 3.2, Def. 3.3].

Proposition 3.5. *Let $(\nu, \mu) \in \mathfrak{M}_{\phi_\kappa}(\Omega) \times \mathfrak{M}(\partial\Omega)$ such that problem (3.18) admits a solution $u_{\mu, \nu}$. Then*

$$u_{\mu, \nu} = \mathbb{G}_{\mathcal{L}_\kappa}[g(u_{\mu, \nu})] + \mathbb{K}_{\mathcal{L}_\kappa}[\mu]. \quad (3.35)$$

Conversely, if $u \in L^1_{\phi_\kappa}(\Omega)$ such that $g(u) \in L^1(\Omega)_{\phi_\kappa}$ satisfies (3.35), it coincides with the solution $u_{\mu, \nu}$ of problem (3.18).

3.3 The power case

In this section we study in particular the following boundary value problem with $\mu \in \mathfrak{M}(\partial\Omega)$

$$\begin{aligned} \mathcal{L}_\kappa u + |u|^{q-1}u &= 0 & \text{in } \Omega \\ u &= \mu & \text{in } \partial\Omega \end{aligned} \quad (3.36)$$

A Radon measure for which this problem has a solution (always unique) is called a *good measure*. The solution, whenever it exists, is unique and denoted by u_μ . For such a nonlinearity, the condition (3.20) is fulfilled if and only if

$$0 < q < q_c := \frac{N + \frac{\alpha_\pm}{2}}{N - 2 + \frac{\alpha_\pm}{2}}. \quad (3.37)$$

On the contrary, in the *supercritical case* i.e. if $q \geq q_c$, a continuity condition with respect to some Besov capacity is needed in order a measure be good. We recall some notations concerning Besov space. For $\sigma > 0$, $1 \leq p < \infty$, we denote by $W^{\sigma, p}(\mathbb{R}^d)$ the Sobolev space over \mathbb{R}^d . If σ is not an integer the Besov space $B^{\sigma, p}(\mathbb{R}^d)$ coincides with $W^{\sigma, p}(\mathbb{R}^d)$. When σ is an integer we denote $\Delta_{x, y} f = f(x + y) + f(x - y) - 2f(x)$ and

$$B^{1, p}(\mathbb{R}^d) = \left\{ f \in L^p(\mathbb{R}^d) : \frac{\Delta_{x, y} f}{|y|^{1 + \frac{d}{p}}} \in L^p(\mathbb{R}^d \times \mathbb{R}^d) \right\}$$

with norm

$$\|f\|_{B^{1,p}} = \left(\|f\|_{L^p}^p + \int \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|\Delta_{x,y} f|^p}{|y|^{p+d}} dx dy \right)^{\frac{1}{p}}.$$

Then

$$B^{m,p}(\mathbb{R}^d) = \{f \in W^{m-1,p}(\mathbb{R}^d) : D_x^\alpha f \in B^{1,p}(\mathbb{R}^d) \forall \alpha \in \mathbb{N}^d, |\alpha| = m-1\}$$

with norm

$$\|f\|_{B^{m,p}} = \left(\|f\|_{W^{m-1,p}}^p + \sum_{|\alpha|=m-1} \int \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|D_x^\alpha \Delta_{x,y} f|^p}{|y|^{p+d}} dx dy \right)^{\frac{1}{p}}.$$

These spaces are fundamental because they are stable under the real interpolation method as it was developed by Lions and Petree. For $\alpha \in \mathbb{R}$ we defined the Bessel kernel of order α by $G_\alpha(\xi) = \mathcal{F}^{-1}(1 + |\cdot|^2)^{-\frac{\alpha}{2}} \mathcal{F}(\xi)$, where \mathcal{F} is the Fourier transform of moderate distributions in \mathbb{R}^d . The Bessel space $L_{\alpha,p}(\mathbb{R}^d)$ is defined by

$$L_{\alpha,p}(\mathbb{R}^d) = \{f = G_\alpha * g : g \in L^p(\mathbb{R}^d)\},$$

with norm

$$\|f\|_{L_{\alpha,p}} = \|g\|_{L^p} = \|G_{-\alpha} * f\|_{L^p}.$$

It is known that if $1 < p < \infty$ and $\alpha > 0$, $L_{\alpha,p}(\mathbb{R}^d) = W^{\alpha,p}(\mathbb{R}^d)$ if $\alpha \in \mathbb{N}$ and $L_{\alpha,p}(\mathbb{R}^d) = B^{\alpha,p}(\mathbb{R}^d)$ if $\alpha \notin \mathbb{N}$, always with equivalent norms. The Bessel capacity is defined for compact subset $K \subset \mathbb{R}^d$ by

$$C_{\alpha,p}^{\mathbb{R}^d} = \inf\{\|f\|_{L_{\alpha,p}}^p, f \in \mathcal{S}'(\mathbb{R}^d), f \geq \chi_K\}.$$

It is extended to open set and then any set by the fact that it is an outer measure. Our main result is the following

Theorem 3.6. *Assume $0 < \kappa \leq \frac{1}{4}$. Then $\mu \in \mathfrak{M}^+(\partial\Omega)$ is a good measure if and only if it is absolutely continuous with respect to the Bessel capacity $C_{2-\frac{2+\alpha_+}{2q'}, q'}$ where $q' = \frac{q}{q-1}$, that is*

$$\forall E \subset \partial\Omega, E \text{ Borel}, C_{2-\frac{2+\alpha_+}{2q'}, q'}^{\mathbb{R}^{N-1}}(E) = 0 \implies \mu(E) = 0. \quad (3.38)$$

The striking aspect of the proof is that it is based upon potential estimates which have been developed by Marcus and Véron in the study of the supercritical boundary trace problem in polyhedral domains [23]. Before proving this result we need a key potential estimate.

Theorem 3.7. *Assume $0 < \kappa \leq \frac{1}{4}$ and $q \geq q_c$. There exists a constant $c_{59} > 1$ depending on Ω , q , and κ such that for any $\mu \in \mathfrak{M}^+(\partial\Omega)$ there holds*

$$\frac{1}{c_{59}} \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q'}, q}}^q \leq \int_{\Omega} (\mathbb{K}_{\mathcal{L}_\kappa}[\mu])^q \phi_\kappa dx \leq c_{59} \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q'}, q}}^q \quad (3.39)$$

Proof. Step 1: local estimates. Denote by $\xi = (\xi_1, \xi')$ the coordinates in \mathbb{R}_+^N , $\xi_1 > 0$, $\xi' \in \mathbb{R}^{N-1}$. The ball of radius $R > 0$ and center a in \mathbb{R}^{N-1} is denoted by $B'_R(a)$ (by B_R if $a = 0$). Let $R > 0$, $\nu \in \mathfrak{M}^+(\mathbb{R}_+^{N-1})$ with support in $B_{\frac{R}{2}}$ and

$$\mathbf{K}[\nu](\xi) = \int_{B'_{\frac{R}{2}}(a)} \frac{d\nu(\zeta')}{(\xi_1^2 + |\xi' - \zeta'|^2)^{\frac{N-2+\alpha_+}{2}}} \quad (3.40)$$

Then, by [23, Th 3.1],

$$\begin{aligned} \frac{1}{c_{60}} \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q^+},q}}^q &\leq \int_0^R \int_{B'_R} \xi_1^{(q+1)\frac{\alpha_+}{2}} \left(\int_{B'_{\frac{R}{2}}} \frac{d\nu(\zeta')}{(\xi_1^2 + |\xi' - \zeta'|^2)^{\frac{N-2+\alpha_+}{2}}} \right)^q d\xi' d\xi_1 \\ &\leq c_{60} \left(1 + R^{(q+1)\frac{\alpha_+}{2}}\right) \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q^+},q}}^q. \end{aligned} \quad (3.41)$$

There exists $R > 0$ such that for any $y_0 \in \partial\Omega$, there exists a C^2 diffeomorphism $\Theta := \Theta_{y_0}$ from $B_R(y_0)$ into \mathbb{R}^N such that $\Theta(y_0) = 0$, $\Theta_{y_0}(B_R(y_0)) = B_R$ and

$$\Theta(\Omega \cap B_R(y_0)) = B_R^+ := B_R \cap \mathbb{R}_+^N, \quad \Theta(\partial\Omega \cap B_{\frac{R}{2}}(y_0)) = B'_{\frac{R}{2}}, \quad \Theta(\partial\Omega \cap B_R(y_0)) = B'_R.$$

Moreover, Θ has bounded distortion, in the sense that since

$$\phi_\kappa(x) \int_{\partial\Omega \cap B_R(y_0)} \frac{d\mu(z)}{|x-z|^{N-2+\alpha_+}} = \phi_\kappa \circ \Theta^{-1}(\xi) \int_{B'_R} \frac{d(\mu \circ \Theta^{-1})(\zeta)}{|\Theta^{-1}(\xi) - \Theta^{-1}(\zeta)|^{N-2+\alpha_+}},$$

there holds

$$\begin{aligned} \frac{\xi_1^{\frac{\alpha_+}{2}}}{c_{61}} \int_{B'_{\frac{R}{2}}} \frac{d(\mu \circ \Theta^{-1})(\zeta)}{(\xi_1^2 + |\xi' - \zeta'|^2)^{\frac{N-2+\alpha_+}{2}}} \\ \leq \phi_\kappa \circ \Theta^{-1}(\xi) \int_{B'_{\frac{R}{2}}} \frac{d(\mu \circ \Theta^{-1})(\zeta)}{|\Theta^{-1}(\xi) - \Theta^{-1}(\zeta)|^{N-2+\alpha_+}} \\ \leq c_{61} \xi_1^{\frac{\alpha_+}{2}} \int_{B'_{\frac{R}{2}}} \frac{d(\mu \circ \Theta^{-1})(\zeta)}{(\xi_1^2 + |\xi' - \zeta'|^2)^{\frac{N-2+\alpha_+}{2}}} \end{aligned}$$

Since $\mu \mapsto \mu \circ \Theta^{-1}$ is a C^2 diffeomorphism between $\mathfrak{M}^+(\partial\Omega \cap B_{\frac{R}{2}}(y_0)) \cap B^{-2+\frac{2+\alpha_+}{2q^+},q}(\partial\Omega \cap B_{\frac{R}{2}}(y_0))$ and $\mathfrak{M}^+(B'_{\frac{R}{2}}) \cap B^{-2+\frac{2+\alpha_+}{2q^+},q}(B'_{\frac{R}{2}})$, we derive, using (2.52) and (3.41)

$$\frac{1}{c_{62}} \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q^+},q}}^q \leq \int_{\Omega \cap B_R(y_0)} (\mathbb{K}_{\mathcal{L}_\kappa}[\mu])^q \phi_\kappa dx \leq c_{62} \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q^+},q}}^q \quad (3.42)$$

Clearly the left-hand side inequality (3.39) follows. Combining Harnack inequality and boundary Harnack inequality we obtain

$$\int_{\Omega} (\mathbb{K}_{\mathcal{L}_\kappa}[\mu])^q \phi_\kappa dx \leq c_{63} \int_{\Omega \cap B_R(y_0)} (\mathbb{K}_{\mathcal{L}_\kappa}[\mu])^q \phi_\kappa dx \quad (3.43)$$

which implies the left-hand side inequality (3.39) when μ has its support in a ball $B_{\frac{R}{2}}$.

Step 2: global estimates. We write $\mu = \sum_{j=1}^{j_0} \mu_j$ where the μ_j are positive measures on $\partial\Omega$ with support in some ball $B_{\frac{R}{2}}(y_j)$ with $y_j \in \partial\Omega$ and such that

$$\frac{1}{c_{64}} \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q^+},q}} \leq \|\mu_j\|_{B^{-2+\frac{2+\alpha_+}{2q^+},q}} \leq c_{64} \|\mu\|_{B^{-2+\frac{2+\alpha_+}{2q^+},q}}.$$

Then

$$\|\mathbb{K}_{\mathcal{L}_\kappa}[\mu]\|_{L^q_{\phi_\kappa}} \leq \sum_{j=1}^{j_0} \|\mathbb{K}_{\mathcal{L}_\kappa}[\mu_j]\|_{L^q_{\phi_\kappa}} \leq c_{59}^{\frac{1}{q}} \sum_{j=1}^{j_0} \|\mu_j\|_{B^{-2+\frac{2+\alpha_\pm}{2q'}, q}}^q \leq j_0 c_{64} c_{59}^{\frac{1}{q}} \|\mu\|_{B^{-2+\frac{2+\alpha_\pm}{2q'}, q}}.$$

On the opposite side

$$\begin{aligned} \|\mathbb{K}_{\mathcal{L}_\kappa}[\mu]\|_{L^q_{\phi_\kappa}} &\geq \max_{1 \leq j \leq j_0} \|\mathbb{K}_{\mathcal{L}_\kappa}[\mu_j]\|_{L^q_{\phi_\kappa}} \\ &\geq \frac{1}{c_{59}^{\frac{1}{q}}} \max_{1 \leq j \leq j_0} \|\mu_j\|_{B^{-2+\frac{2+\alpha_\pm}{2q'}, q}} \\ &\geq \frac{1}{j_0 c_{59}^{\frac{1}{q}}} \sum_{j=1}^{j_0} \|\mu_j\|_{B^{-2+\frac{2+\alpha_\pm}{2q'}, q}} \\ &\geq \frac{1}{c_{64} c_{59}^{\frac{1}{q}}} \|\mu\|_{B^{-2+\frac{2+\alpha_\pm}{2q'}, q}}, \end{aligned}$$

which ends the proof. \square

Proof of Theorem 3.6: The condition is sufficient. Let μ be a boundary measure such that $\mathbb{K}_{\mathcal{L}_\kappa}[\mu] \in L^q_{\partial\Omega}$. For $k > 0$ set $g_k(u) = \text{sgn}(u) \min\{|u|^q, k^q\}$ and let u_k be the solution of

$$\begin{aligned} \mathcal{L}_\kappa u_k + g_k(u_k) &= 0 && \text{in } \Omega \\ u_k &= \mu && \text{in } \partial\Omega, \end{aligned} \quad (3.44)$$

which exists and is unique by Theorem 3.3. Furthermore $k \mapsto u_k$ is decreasing,

$$0 \leq u_k \leq \mathbb{K}_{\mathcal{L}_\kappa}[\mu]$$

and

$$0 \leq g_k(u_k) \leq g_k(\mathbb{K}_{\mathcal{L}_\kappa}[\mu]) \leq (\mathbb{K}_{\mathcal{L}_\kappa}[\mu])^q,$$

and the first terms on the right of the two previous inequalities are integrable for the measure $\phi_\kappa dx$ by Theorem 3.7. Finally for any $\eta \in \mathbf{X}_\kappa(\Omega)$, there holds

$$\int_{\Omega} (u_k \mathcal{L}_\kappa \eta + g_k(u_k) \eta) dx = \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[\mu] \mathcal{L}_\kappa \eta dx.$$

Since u_k and $g_k(u_k)$ converge respectively to u and $g(u)$ a.e. and in $L^1_{\phi_\kappa}(\Omega)$; we conclude that

$$\int_{\Omega} (u \mathcal{L}_\kappa \eta + g(u) \eta) dx = \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[\mu] \mathcal{L}_\kappa \eta dx.$$

If μ is a positive measure which vanishes on Borel sets $E \subset \partial\Omega$ with $C^{\mathbb{R}^{N-1}}_{2-\frac{2+\alpha_\pm}{2q'}, q}$ -capacity zero, there

exists an increasing sequence of positive measures in $B^{-2+\frac{2+\alpha_\pm}{2q'}, q}(\partial\Omega)$ $\{\mu_n\}$ which converges to μ (see [8], [11]). Let u_{μ_n} be the solution of (3.36) with boundary data μ_n . The sequence $\{u_{\mu_n}\}$ is increasing with limit u . Since, by taking ϕ_κ as test function, we obtain

$$\int_{\Omega} (\lambda_\kappa u_{\mu_n} + g(u_{\mu_n})) \phi_\kappa dx = \lambda_\kappa \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[\mu_n] \phi_\kappa dx,$$

it follows that $u, g(u) \in L^1_{\phi_\kappa}$. Thus

$$\int_{\Omega} (u \mathcal{L}_\kappa \eta + g(u) \eta) dx = \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[\mu] \mathcal{L}_\kappa \eta dx \quad \forall \eta \in \mathbf{X}_\kappa(\Omega),$$

and therefore $u = u_\mu$.

Definition A smooth lifting is a continuous linear operator $R[\cdot]$ from $C_0^2(\partial\Omega)$ to $C_0^2(\overline{\Omega})$ satisfying

$$\begin{aligned} (i) \quad & 0 \leq \eta \leq 1 \implies 0 \leq R[\eta] \leq 1, \quad R[\eta]|_{\partial\Omega} = \eta \\ (ii) \quad & |\nabla\phi_\kappa \cdot \nabla R[\eta]| \leq c_{65}\phi_\kappa \end{aligned} \quad (3.45)$$

where c_{65} depends on the C^1 -norm of η .

Our proof are based upon modification of an argument developed by Marcus and Véron in [18].

Lemma 3.8. *Assume there exists a solution u_μ of (3.36) with $\mu \geq 0$. For $\eta \in C^2(\Omega)$, $0 \leq \eta \leq 1$ set $\zeta = \phi_\kappa(R[\eta])^{q'}$ where R is a smooth lifting. Then*

$$\left(\int_{\partial\Omega} \eta d\mu \right)^{q'} \leq c_{67} \int_{\Omega} u^q \zeta dx + c_{67} \left(\int_{\Omega} u^q \zeta dx \right)^{\frac{1}{q}} \left(\left(\int_{\Omega} \phi_\kappa dx \right)^{\frac{1}{q'}} + q' \left(\int_{\Omega} (L[\eta])^{q'} dx \right)^{\frac{1}{q'}} \right) \quad (3.46)$$

where

$$L[\eta] = (R[\eta])^{q'-1} \left(2\phi_\kappa^{-\frac{1}{q}} |\nabla\phi_\kappa \cdot \nabla R[\eta]| + \phi_\kappa^{\frac{1}{q'}} |\Delta R[\eta]| \right) \quad (3.47)$$

and c_{67} depends on $\|\mu\|_{\mathfrak{M}(\partial\Omega)}$.

Proof. There holds

$$\mathcal{L}_\kappa \zeta = \lambda_\kappa (R[\eta])^{q'} \phi_\kappa - 2q' (R[\eta])^{q'-1} \nabla\phi_\kappa \cdot \nabla R[\eta] - q' (R[\eta])^{q'-2} \phi_\kappa (R[\eta] \Delta R[\eta] - (q' - 1) |\nabla R[\eta]|^2).$$

Then $\zeta \in \mathbf{X}_\kappa(\Omega)$ because of (3.45)-(ii) and

$$c_{66} \int_{\partial\Omega} \eta^{q'} d\mu \leq \int_{\Omega} (u \mathcal{L}_\kappa \zeta + u^q \zeta) dx.$$

Since

$$u \mathcal{L}_\kappa \zeta \leq u \left(\lambda_\kappa (R[\eta])^{q'} \phi_\kappa + 2q' (R[\eta])^{q'-1} |\nabla\phi_\kappa \cdot \nabla R[\eta]| + q' (R[\eta])^{q'-1} \phi_\kappa |\Delta R[\eta]| \right)$$

we obtain

$$\int_{\Omega} u \mathcal{L}_\kappa \zeta dx \leq \left(\int_{\Omega} u^q \phi_\kappa dx \right)^{\frac{1}{q}} \left(\left(\int_{\Omega} \phi_\kappa dx \right)^{\frac{1}{q'}} + q' \left(\int_{\Omega} (L[\eta])^{q'} dx \right)^{\frac{1}{q'}} \right),$$

where $L[\eta]$ is defined by (3.47). □

Lemma 3.9. *There exist a smooth lifting R such that $\eta \mapsto L[\eta]$ is continuous from $B^{2-\frac{2+\alpha_+}{2q'}, q'}(\partial\Omega)$ into $L^{q'}(\Omega)$. Furthermore,*

$$\|L[\eta]\|_{L^{q'}(\Omega)} \leq c'_{66} \|\eta\|_{L^\infty(\partial\Omega)}^{q'-1} \|\eta\|_{B^{2-\frac{2+\alpha_+}{2q'}, q'}(\partial\Omega)}. \quad (3.48)$$

Proof. The construction of the lifting is originated into [21, Sect 1]. For $0 < \delta \leq \beta_0$, we set $\Sigma_\delta = \{x \in \Omega : d(x) = \delta\}$ and we identify $\partial\Omega$ with $\Sigma := \Sigma_0$. The set $\{\Sigma_\delta\}_{0 < \delta \leq \beta_0}$ is a smooth foliation of $\partial\Omega$. For each $\delta \in (0, \beta_0]$ there exists a unique $\sigma(x) \in \Sigma_\delta$ such that $d(x) = \delta$ and $|x - \sigma(x)| = \delta$. The set of couples (δ, σ) defines a system of coordinates in Ω_{β_0} called the flow coordinates. The Laplacian obtain the following expression in this system

$$\Delta = \frac{\partial^2}{\partial \delta^2} + b_0 \frac{\partial}{\partial \delta} + \Lambda_\Sigma \quad (3.49)$$

where Λ_Σ is a linear second-order elliptic operator on Σ with C^1 coefficients. Furthermore $b_0 \rightarrow K$ and $\Lambda_\Sigma \rightarrow \Delta_\Sigma$, where K is the mean curvature of Σ and Δ_Σ the Laplace-Beltrami operator on Σ . If $\eta \in B^{-2+\frac{2+\alpha_+}{2q}, q}(\partial\Omega)$, we denote by $H := H[\eta]$ the solution of

$$\begin{aligned} \frac{\partial H}{\partial s} + \Delta_\Sigma H &= 0 & \text{in } (0, \infty) \times \Sigma \\ H(0, \cdot) &= \eta & \text{in } \Sigma \end{aligned} \quad (3.50)$$

Let $h \in C^\infty(\mathbb{R}_+)$ such that $0 \leq h \leq 1$, $h' \leq 0$, $h \equiv 1$ on $[0, \frac{\beta_0}{2}]$, $h \equiv 0$ on $[\beta_0, \infty]$. The lifting we consider is expressed by

$$R[\eta](x) = \begin{cases} H[\eta](\delta^2, \sigma(x))h(\delta) & \text{if } x \in \bar{\Omega}_{\beta_0} \\ 0 & \text{if } x \in \Omega'_{\beta_0}, \end{cases} \quad (3.51)$$

with $x \approx (\delta, \sigma) := (d(x), \sigma(x))$. Mutatis mutandis, we perform the same computation as the one in [18, Lemma 1.2], using local coordinates $\{\sigma_j\}$ on Σ and obtain

$$\nabla R[\eta] = 2\delta h(\delta) \frac{\partial H}{\partial \delta}(\delta^2, \sigma) \nabla \delta + \sum_{j=1}^{N-1} h(\delta) \frac{\partial H}{\partial \sigma_j}(\delta^2, \sigma) \nabla \sigma_j + h'(\delta) H(\delta^2, \sigma) \nabla \delta$$

In $\Omega_{\frac{\beta_0}{2}}$ there holds

$$\nabla R[\eta] \cdot \nabla \phi_\kappa = 2\delta h(\delta) \frac{\partial H}{\partial \delta}(\delta^2, \sigma) \nabla \phi_\kappa \cdot \nabla \delta + \sum_{j=1}^{N-1} h(\delta) \frac{\partial H}{\partial \sigma_j}(\delta^2, \sigma) \nabla \sigma_j \cdot \nabla \delta \quad (3.52)$$

Moreover $\phi_\kappa(x) \leq c_2(d(x))^{\frac{\alpha_+}{2}} = c_2\delta^{\frac{\alpha_+}{2}}$ and $|\nabla \phi_\kappa(x)| \leq c'_2(d(x))^{\frac{\alpha_+}{2}-1} = c'_2\delta^{\frac{\alpha_+}{2}-1}$. Similarly as in [18, (1.13)]

$$\nabla \phi_\kappa = \frac{\partial \phi_\kappa}{\partial \delta} \mathbf{n} + \sum_{j=1}^{N-1} \frac{\partial \phi_\kappa}{\partial \sigma_j}(\delta^2, \sigma) \nabla \sigma_j,$$

thus

$$\begin{aligned} |\nabla \phi_\kappa \cdot \nabla \sigma_j| &\leq c_{68} \delta^{\frac{\alpha_+}{2}}, \\ \phi_\kappa^{-\frac{1}{q}} |\nabla R[\eta] \cdot \nabla \phi_\kappa| &\leq c_{69} \delta^{\frac{\alpha_+}{2q}} \left(\left| \frac{\partial H}{\partial \delta}(\delta^2, \sigma) \right| + \sum_{j=1}^{N-1} \left| \frac{\partial H}{\partial \sigma_j}(\delta^2, \sigma) \right| - h'(\delta) H(\delta^2, \sigma) \right). \end{aligned}$$

Thus

$$\begin{aligned} \int_{\Omega} \phi_{\kappa}^{-\frac{q'}{q}} |\nabla R[\eta] \cdot \nabla \phi_{\kappa}|^{q'} dx &\leq c_{70} \int_{\Omega_{\beta_0}} \delta^{\frac{\alpha_{\pm}}{2}} \left| \frac{\partial H}{\partial \delta}(\delta^2, \sigma) \right|^{q'} dx \\ &+ c_{70} \sum_{j=1}^{N-1} \int_{\Omega_{\beta_0}} \delta^{\frac{\alpha_{\pm}}{2}} \left| \frac{\partial H}{\partial \sigma_j}(\delta^2, \sigma) \right|^{q'} dx \\ &+ c_{70} \int_{\Omega_{\beta_0} \setminus \Omega_{\frac{\beta_0}{2}}} \delta^{\frac{\alpha_{\pm}}{2}} H^{q'}(\delta^2, \sigma) dx \end{aligned}$$

Then

$$\begin{aligned} \int_{\Omega} \phi_{\kappa}^{-\frac{q'}{q}} |\nabla R[\eta] \cdot \nabla \phi_{\kappa}|^{q'} dx &\leq c_{71} \int_0^{\beta_0} \delta^{\frac{\alpha_{\pm}}{2}} \int_{\Sigma} \left| \frac{\partial H}{\partial t}(\delta^2, \sigma) \right|^{q'} dS d\delta \\ &\leq c_{71} \int_0^{\beta_0} \int_{\Sigma} \left(t^{\frac{2+\alpha_{\pm}}{4q'}} \left\| \frac{\partial H}{\partial t}(t, \cdot) \right\|_{L^{q'}(\Sigma)} \right)^{q'} \frac{dt}{t} \quad (3.53) \\ &\leq c_{72} \|\eta\|_{B^{2-\frac{2+\alpha_{\pm}}{2q'}, q'}(\Sigma)}^{q'} \end{aligned}$$

by using the classical real interpolation identity

$$\left[W^{2, q'}(\Sigma), L^{q'}(\Sigma) \right]_{1-\frac{2+\alpha_{\pm}}{4q'}, q'} = B^{2-\frac{2+\alpha_{\pm}}{2q'}, q'}(\Sigma). \quad (3.54)$$

Similarly (see [18, (1.17), (1.19)])

$$\sum_{j=1}^{N-1} \int_{\Omega_{\beta_0}} \delta^{\frac{\alpha_{\pm}}{2}} \left| \frac{\partial H}{\partial \sigma_j}(\delta^2, \sigma) \right|^{q'} dx + \int_{\Omega_{\beta_0} \setminus \Omega_{\frac{\beta_0}{2}}} \delta^{\frac{\alpha_{\pm}}{2}} H^{q'}(\delta^2, \sigma) dx \leq c_{72} \|\eta\|_{W^{2-\frac{2+\alpha_{\pm}}{2q'}, q'}(\Sigma)}^{q'}. \quad (3.55)$$

Next we consider the second term. Adapting in a straightforward manner the computation in [18, p. 886-887] we obtain the following instead of [18, (1.21)]

$$\begin{aligned} \int_{\Omega} \phi_{\kappa}^{\frac{q'}{q}} |\Delta R[\eta]|^{q'} dx &\leq c_{72} \int_0^{\beta_0} \int_{\Sigma} \left| \delta^{2+\frac{\alpha_{\pm}}{2q'}} \frac{\partial^2 H[\eta]}{\partial \delta^2} \right|^{q'}(\delta^2, \sigma) d\sigma d\delta \\ &+ c_{72} \int_0^{\beta_0} \int_{\Sigma} \delta^{\frac{\alpha_{\pm}}{2}} \left(\left| \frac{\partial H[\eta]}{\partial \delta} \right|^{q'} + |H|^{q'} + |\Lambda_{\Delta} - \Lambda_{\Sigma}|^{q'} \right) (\delta^2, \sigma) dx \quad (3.56) \end{aligned}$$

Then

$$\begin{aligned} \int_0^{\beta_0} \int_{\Sigma} \left| \delta^{2+\frac{\alpha_{\pm}}{2q'}} \frac{\partial^2 H[\eta]}{\partial \delta^2} \right|^{q'}(\delta^2, \sigma) d\sigma d\delta &= \int_0^{\beta_0} \int_{\Sigma} \left| t^{2\left(1-\frac{4q'-\alpha_{\pm}-2}{8q'}\right)} \frac{\partial^2 H[\eta]}{\partial t^2} \right|^{q'} d\sigma \frac{dt}{t} \\ &\leq c_{73} \|\eta\|_{B^{2-\frac{2+\alpha_{\pm}}{2q'}, q'}(\Sigma)}^{q'}, \quad (3.57) \end{aligned}$$

by using the real interpolation identity

$$\left[W^{4, q'}(\Sigma), L^{q'}(\Sigma) \right]_{\frac{4q'-\alpha_{\pm}-2}{8q'}, q'} = B^{2-\frac{2+\alpha_{\pm}}{2q'}, q'}(\Sigma). \quad (3.58)$$

The other term in the right-hand side of (3.56) yields to the same inequality as in (3.55). \square

Proof of Theorem 3.6: The condition is necessary. Let $K \subset \partial\Omega$ be a compact set and $\eta \in C_0^2(\partial\Omega)$ such that $0 \leq \eta \leq 1$ and $\eta = 1$ on K . Then, by (3.46)

$$\begin{aligned} (\mu(K))^{q'} &\leq c_{67} \int_{\Omega} u^q (R[\eta])^{q'} \phi_{\kappa} dx + \\ &c_{67} \left(\int_{\Omega} u^q (R[\eta])^{q'} \phi_{\kappa} dx \right)^{\frac{1}{q}} \left(\left(\int_{\Omega} \phi_{\kappa} dx \right)^{\frac{1}{q'}} + c'_{66} q' \|\eta\|_{B^{2-\frac{2+\alpha_{\pm}}{2q'}, q'}(\partial\Omega)} \right). \end{aligned} \quad (3.59)$$

From this inequality, we obtain classically the result since if $C_{2-\frac{2+\alpha_{\pm}}{2q'}, q'}^{\mathbb{R}^{N-1}}(K) = 0$ there exists a sequence $\{\eta_n\}$ in $C_0^2(\partial\Omega)$ with the following properties:

$$0 \leq \eta_n \leq 1, \quad \eta_n = 1 \text{ in a neighborhood of } K \text{ and } \eta_n \rightarrow 0 \text{ in } B^{2-\frac{2+\alpha_{\pm}}{2q'}, q'}(\partial\Omega) \text{ as } n \rightarrow \infty. \quad (3.60)$$

This implies that $u^q (R[\eta_n])^{q'} \rightarrow 0$ in $L^1_{\phi_{\kappa}}(\Omega)$. Therefore the right-hand side of (3.59) tends to 0 if we substitute η_n to η and thus $\mu(K) = 0$ for any K compact with zero capacity and this relation holds for any Borel subset. \square

Definition. We say that a compact set $K \subset \partial\Omega$ is *removable* if any positive solution $u \in C(\overline{\Omega} \setminus K)$ of

$$\mathcal{L}_{\kappa} u + |u|^{q-1} u = 0 \quad \text{in } \Omega \quad (3.61)$$

such that

$$\int_{\Omega} (u \mathcal{L}_{\kappa} \eta + |u|^{q-1} u \eta) dx = 0 \quad \forall \eta \in \mathbf{X}_{\kappa}^K(\Omega) \quad (3.62)$$

where $\mathbf{X}_{\kappa}^K(\Omega) = \{\eta \in \mathbf{X}_{\kappa}(\Omega) : \eta = 0 \text{ in a neighborhood of } K\}$, is identically zero.

Theorem 3.10. *Assume $0 < \kappa \leq \frac{1}{4}$ and $q \geq 1$. A compact set $K \subset \partial\Omega$ is removable if and only if $C_{2-\frac{2+\alpha_{\pm}}{2q'}, q}^{\mathbb{R}^{N-1}}(K) = 0$.*

Proof. The condition is clearly necessary since, if a compact boundary set K has positive capacity, there exists a capacitary measure $\mu_K \in \mathfrak{M}_+(\partial\Omega) \cap B^{-2+\frac{2+\alpha_{\pm}}{2q'}, q}(\partial\Omega)$ with support in K (see e.g. [1]). For such a measure there exists a solution u_{μ_K} of (3.36) with $\mu = \mu_K$ by Theorem 3.6. Next we assume that $C_{2-\frac{2+\alpha_{\pm}}{2q'}, q}^{\mathbb{R}^{N-1}}(K) = 0$. Then there exists a sequence $\{\eta_n\}$ in $C_0^2(\partial\Omega)$ satisfying (3.60). In particular, there exists a decreasing sequence $\{\mathcal{O}_n\}$ of relatively open subsets of $\partial\Omega$, containing K such that $\eta_n = 1$ on \mathcal{O}_n and thus $\eta_n = 1$ on $K_n := \overline{\mathcal{O}_n}$. We set $\tilde{\eta}_n = 1 - \eta_n$ and $\tilde{\zeta}_n = \phi_{\kappa} (R[\tilde{\eta}_n])^{2q'}$ where R is defined by (3.51). Then $0 \leq \tilde{\eta}_n \leq 1$ and $\tilde{\eta}_n = 0$ on K_n . Therefore

$$\tilde{\zeta}_n(x) \leq \phi_{\kappa} \min \left\{ 1, c_{74} (d(x))^{1-N} e^{-(4d(x))^{-2} (\text{dist}(x, K_n^c))^2} \right\} \quad (3.63)$$

Furthermore

$$\begin{aligned} (i) \quad &|\nabla R[\tilde{\eta}_n]| \leq c_{75} \min \left\{ 1, (d(x))^{-2-N} e^{-(4d(x))^{-2} (\text{dist}(x, K_n^c))^2} \right\} \\ (ii) \quad &|\Delta R[\tilde{\eta}_n]| \leq c_{75} \min \left\{ 1, (d(x))^{-4-N} e^{-(4d(x))^{-2} (\text{dist}(x, K_n^c))^2} \right\} \end{aligned} \quad (3.64)$$

Step 1. We claim that

$$\int_{\Omega} \left(u \mathcal{L}_{\kappa} \tilde{g} z_n + u^q \tilde{\zeta}_n \right) dx = 0. \quad (3.65)$$

By Proposition ?? there exists $c_{74} > 0$ such that

$$\begin{aligned} (i) \quad & u(x) \leq c_{76} (d(x))^{\frac{\alpha_+}{2}} (\text{dist}(x, K))^{-\frac{2}{q-1} - \frac{\alpha_+}{2}} \\ (ii) \quad & |\nabla u(x)| \leq c_{76} (d(x))^{\frac{\alpha_+}{2} - 1} (\text{dist}(x, K))^{-\frac{2}{q-1} - \frac{\alpha_+}{2}} \end{aligned} \quad (3.66)$$

for all $x \in \Omega$. As in the proof of Lemma 3.8,

$$\begin{aligned} |u \mathcal{L}_{\kappa} \tilde{\zeta}_n| \leq c_{77} (R[\tilde{\eta}_n])^{2q' - 2} u \left(\phi_{\kappa} R^2[\tilde{\eta}_n] + R[\tilde{\eta}_n] |\nabla \phi_{\kappa} \cdot \nabla R[\tilde{\eta}_n]| \right. \\ \left. + \phi_{\kappa} (R[\tilde{\eta}_n] |\Delta R[\tilde{\eta}_n]| + |\nabla R[\tilde{\eta}_n]|^2) \right). \end{aligned} \quad (3.67)$$

Let \mathcal{O} be a relatively open neighborhood of K such that $\bar{\mathcal{O}} \subset \mathcal{O}_n$. We set $G_{\mathcal{O}, \beta_0} = \{x \in \Omega_{\beta_0} : \sigma(x) \in \mathcal{O}\}$ and $G_{\mathcal{O}^c, \beta_0} = \Omega_{\beta_0} \setminus G_{\mathcal{O}}$. If $x \in G_{\mathcal{O}}$, $\text{dist}(x, K_n^c) \geq \tau > 0$. Then, by (3.66)-(i) and (3.63), $u^q \tilde{\zeta}_n \in L^q(G_{\mathcal{O}})$. Since $u(x) = \circ(W(x))$ in $G_{\mathcal{O}^c}$ it follows that $u^q \tilde{\zeta}_n \in L^1(\Omega_{\beta_0})$ and thus $u^q \tilde{\zeta}_n$ is integrable in Ω . Similarly, using (N22-1)-(i) and (ii), $u \mathcal{L}_{\kappa} \tilde{\zeta}_n \in L^1(\Omega)$. Since $\tilde{\zeta}_n$ does not vanish in a neighborhood of K , we introduce a cut-off function $\theta_{\epsilon} \in C^2(\bar{\Omega})$ for $0 < \epsilon \leq \frac{\beta_0}{2}$, with the following properties,

$$\begin{aligned} 0 \leq \theta_{\epsilon} \leq 1, \quad \theta_{\epsilon}(x) = 0 \quad \forall x \in G_{\mathcal{O}, \epsilon}, \quad \theta_{\epsilon}(x) = 1 \quad \forall x \in \bar{\Omega} \text{ s.t. } \text{dist}(x, G_{\mathcal{O}, \epsilon}) \geq \epsilon \\ |\nabla \theta_{\epsilon}| \leq c_{78} \epsilon^{-1} \chi_{G_{\mathcal{O}, \epsilon, \epsilon} \setminus G_{\mathcal{O}, \epsilon}} \quad \text{and} \quad |D^2 \theta_{\epsilon}| \leq c_{78} \epsilon^{-2} \chi_{G_{\mathcal{O}, \epsilon, \epsilon} \setminus G_{\mathcal{O}, \epsilon}}, \end{aligned}$$

where we have taken ϵ small enough so that

$$G_{\mathcal{O}, \epsilon, \epsilon} := \{x \in \Omega : \text{dist}(x, G_{\mathcal{O}, \epsilon}) \leq \epsilon\} \subset G_{K_n, 2\epsilon} = \{x \in \Omega_{2\epsilon} : \sigma(x) \in K_n\}.$$

Clearly $\theta_{\epsilon} \tilde{\zeta}_n \in \mathbf{X}_{\kappa}^K(\Omega)$, thus

$$\int_{\Omega} \left(u \mathcal{L}_{\kappa} (\theta_{\epsilon} \tilde{\zeta}_n) + u^q \theta_{\epsilon} \tilde{\zeta}_n \right) dx = 0.$$

Next

$$\begin{aligned} \int_{\Omega} \left(u \mathcal{L}_{\kappa} (\theta_{\epsilon} \tilde{\zeta}_n) + u^q \theta_{\epsilon} \tilde{\zeta}_n \right) dx &= \int_{\Omega \setminus G_{\mathcal{O}, \epsilon, \epsilon}} \left(u \mathcal{L}_{\kappa} (\tilde{\zeta}_n) + u^q \tilde{\zeta}_n \right) dx + \int_{G_{\mathcal{O}, \epsilon, \epsilon}} \left(u \mathcal{L}_{\kappa} (\theta_{\epsilon} \tilde{\zeta}_n) + u^q \theta_{\epsilon} \tilde{\zeta}_n \right) dx \\ &= I_{\epsilon} + II_{\epsilon} \end{aligned}$$

Clearly

$$\lim_{\epsilon \rightarrow 0} I_{\epsilon} = \int_{\Omega} \left(u \mathcal{L}_{\kappa} \tilde{\zeta}_n + u^q \tilde{\zeta}_n \right) dx$$

and

$$\lim_{\epsilon \rightarrow 0} \int_{G_{\mathcal{O}, \epsilon, \epsilon}} u^q \theta_{\epsilon} \tilde{\zeta}_n dx = 0.$$

Finally, since $\mathcal{L}_{\kappa}(\theta_{\epsilon} \tilde{\zeta}_n) = \theta_{\epsilon} \mathcal{L}_{\kappa} \tilde{\zeta}_n + \tilde{\zeta}_n \Delta \theta_{\epsilon} + 2 \nabla \theta_{\epsilon} \cdot \nabla \tilde{\zeta}_n$, θ_{ϵ} is constant outside $G_{\mathcal{O}, \epsilon, \epsilon} \setminus G_{\mathcal{O}, \epsilon}$ and $\text{dist}(G_{\mathcal{O}, \epsilon, \epsilon} \setminus G_{\mathcal{O}, \epsilon}, F_n^c) \geq \tau > 0$, independent of ϵ there holds, by (3.63)

$$|\mathcal{L}_{\kappa}(\theta_{\epsilon} \tilde{\zeta}_n)| \leq c_{79} \epsilon^{-N+4} e^{-\frac{\tau}{\epsilon}}.$$

Using by (3.66)-(i) we derive

$$\lim_{\epsilon \rightarrow 0} \int_{G_{\mathcal{O}_\epsilon, \epsilon}} u \mathcal{L}_\kappa(\theta_\epsilon \tilde{\zeta}_n) dx = 0,$$

which yields to (3.65).

Step 2. We claim that

$$\int_{\Omega} u^q \phi_\kappa dx < \infty. \quad (3.68)$$

Using the expression of $\mathcal{L}_\kappa \zeta_n$ in (3.65) where replace η_n by $\tilde{\eta}_n$, we derive

$$\begin{aligned} \int_{\Omega} u^q \tilde{\zeta}_n dx &= \int_{\Omega} \left(-\lambda_\kappa (R[\tilde{\eta}_n])^{2q'} \phi_\kappa + 4q' (R[\tilde{\eta}_n])^{2q'-1} \nabla \phi_\kappa \cdot \nabla R[\tilde{\eta}_n] + \right. \\ &\quad \left. 2q' (R[\tilde{\eta}_n])^{2q'-2} \phi_\kappa (R[\tilde{\eta}_n] \Delta R[\tilde{\eta}_n] + (2q' - 1) |\nabla R[\tilde{\eta}_n]|^2) \right) u dx \\ &\leq c_{79} \left(\int_{\Omega} u^q \tilde{\zeta}_n dx \right)^{\frac{1}{q}} \left(\int_{\Omega} (\tilde{L}[\eta_n])^{q'} dx \right)^{\frac{1}{q'}}, \end{aligned} \quad (3.69)$$

where we have set

$$\tilde{L}[\eta] = (\phi_\kappa)^{-\frac{1}{q}} \nabla \phi_\kappa \cdot \nabla R[\eta_n] + (\phi_\kappa)^{\frac{1}{q'}} |\Delta R[\tilde{\eta}_n]| + (\phi_\kappa)^{\frac{1}{q'}} |\nabla R[\tilde{\eta}_n]|^2 \quad (3.70)$$

By Lemma 3.9 we know that

$$\int_{\Omega} (\phi_\kappa)^{-\frac{q'}{q}} |\nabla \phi_\kappa \cdot \nabla R[\eta_n]|^{q'} + \phi_\kappa |\Delta R[\tilde{\eta}_n]|^{q'} dx \leq (c_{72} + c_{73}) \|\eta_n\|_{B^{2-\frac{2+\alpha_+}{2q'}, 2}(\partial\Omega)}^{q'}. \quad (3.71)$$

The last term is estimated in the following way

$$\begin{aligned} \int_{\Omega} \phi_\kappa |\nabla R[\tilde{\eta}_n]|^{2q'} dx &\leq c_{80} \int_0^{\beta_0^2} \int_{\Sigma} s^{q' + \frac{\alpha_+ + 2}{4}} \left| \frac{\partial H[\eta_n]}{\partial s} \right|^{2q'} dS \frac{ds}{s} \\ &\quad + c_{80} \int_0^{\beta_0^2} \int_{\Sigma} s^{\frac{\alpha_+ + 2}{4}} \left(|\nabla_{\Sigma} H[\eta_n]|^{2q'} + (H[\eta_n])^{2q'} \right) dS \frac{ds}{s}, \end{aligned} \quad (3.72)$$

where ∇_{Σ} denotes the covariant gradient on Σ . Since the following interpolation identity holds

$$\left[W^{2, 2q'}(\Sigma), L^{2q'}(\Sigma) \right]_{1-\frac{\alpha_+ + 2}{8q'}, 2q'} = B^{1-\frac{\alpha_+ + 2}{4q'}, 2q'}(\Sigma)$$

we obtain

$$\int_0^{\beta_0^2} \int_{\Sigma} s^{q' + \frac{\alpha_+ + 2}{4}} \left| \frac{\partial H[\eta_n]}{\partial s} \right|^{2q'} \frac{ds}{s} \leq c_{81} \|\eta_n\|_{B^{1-\frac{\alpha_+ + 2}{4q'}, 2q'}(\Sigma)}^{2q'}$$

By the Gagliardo-Nirenberg inequality

$$\|\eta_n\|_{B^{1-\frac{\alpha_+ + 2}{4q'}, 2q'}(\Sigma)}^{2q'} \leq c_{82} \|\eta_n\|_{B^{2-\frac{\alpha_+ + 2}{2q'}, q'}(\Sigma)}^{q'} \|\eta_n\|_{L^\infty(\Sigma)}^{q'} = c_{82} \|\eta_n\|_{B^{2-\frac{\alpha_+ + 2}{2q'}, q'}(\Sigma)}^{q'}. \quad (3.73)$$

By the same inequality

$$\int_{\Sigma} \left(|\nabla_{\Sigma} H[\eta_n]|^{2q'} + (H[\eta_n])^{2q'} \right) dS \leq c_{82} \|H[\eta_n]\|_{L^\infty(\Sigma)}^{q'} \int_{\Sigma} \left(|\Delta_{\Sigma} H[\eta_n]|^{q'} + (H[\eta_n])^{q'} \right) dS. \quad (3.74)$$

Using the estimates on $L[\eta]$ in Lemma 3.9 and the fact that $0 \leq H[\eta_m] \leq 1$, we conclude that

$$\int_{\Sigma} \left(|\nabla_{\Sigma} H[\eta_m]|^{2q'} + (H[\eta_m])^{2q'} \right) dS \leq c_{83} \|\eta_m\|_{B^{2-\frac{\alpha_++2}{2q'}, q'}(\Sigma)}^{q'}$$

It follows from (3.69)

$$\int_{\Omega_{\frac{\beta_0}{2}}} u^q (R[\tilde{\eta}_m])^{2q'} dx \leq c_{84} \int_{\Omega_{\beta_0}} (\tilde{L}\eta_m)^{q'} dx \leq c_{85} \|\eta_m\|_{B^{2-\frac{\alpha_++2}{2q'}, q'}(\Sigma)}^{q'} \quad (3.75)$$

Letting $n \rightarrow \infty$ and using the fact that $\eta_n \rightarrow 0$, we obtain by Fatou's lemma that

$$\int_{\Omega_{\frac{\beta_0}{2}}} u^q \phi_{\kappa} = 0.$$

Combined with the fact that u is bounded in $\Omega'_{\frac{\beta_0}{2}}$ we obtain (3.68). Notice that $\|u\|_{L^q_{\phi_{\kappa}}(\Omega)}$ is bounded independently of u .

Step 3. End of the proof. Since $u^q \in L^1_{\phi_{\kappa}}(\Omega)$, by Proposition 3.2 there exists a unique weak solution $v \in L^1_{\phi_{\kappa}}(\Omega)$ of

$$\begin{aligned} \mathcal{L}_{\kappa} v &= u^q & \text{in } \Omega \\ v &= 0 & \text{in } \partial\Omega, \end{aligned} \quad (3.76)$$

and $v \geq 0$. Then $w = u + v$ is \mathcal{L}_{κ} -harmonic in Ω , and by Theorem 2.31 there exists a unique positive Radon measure τ on $\partial\Omega$ such that $w = \mathbb{K}_{\mathcal{L}_{\kappa}}[\tau]$. Since v and u vanish respectively on $\partial\Omega$ and $\partial\Omega \setminus K$, it follows from Propositions 2.33 and 2.34 that the support of τ is included in K . By Theorem 3.6, τ vanishes on Borel subsets with zero $C_{2-\frac{2+\alpha_+}{2q'}, q'}^{\mathbb{R}^{N-1}}$ -capacity. Since $C_{2-\frac{2+\alpha_+}{2q'}, q'}^{\mathbb{R}^{N-1}}(K) = 0$, $\tau = 0$. This implies that u is a weak solution of

$$\begin{aligned} \mathcal{L}_{\kappa} u + u^q &= 0 & \text{in } \Omega \\ u &= 0 & \text{in } \partial\Omega, \end{aligned} \quad (3.77)$$

and therefore $u = 0$. \square

Remark. Using the fact that u^+ and u_- are subsolutions of (3.61), it is easy to check that Theorem 3.10 remains valid for any signed solution of (3.61).

Remark. If $1 < q < q_c$ (see (3.37)) it follows from Sobolev imbedding theorem that only the empty set has zero $C_{2-\frac{2+\alpha_+}{2q'}, q'}^{\mathbb{R}^{N-1}}$ -capacity. As a consequence of the previous result, if $q \geq q_c$ any isolated boundary singularity of a solution of (3.61) is removable.

4 Isolated boundary singularities

We denote by $\{e_1, \dots, e_N\}$ the canonical basis in $\mathbb{R}^N = \{x = (x', x_N) \in \mathbb{R}^{N-1} \times \mathbb{R}\}$ and by (r, σ) the spherical coordinates therein. Then $\mathbb{R}_+^N = \{(x', x_N) : x' \in \mathbb{R}^{N-1}, x_N > 0\}$. We although denote by S^{N-1} and S_+^{N-1} the unit sphere and the upper hemisphere of \mathbb{R}_+^N , i.e. $S^{N-1} : \cap \mathbb{R}_+^N$. In this section we study the behavior near 0 of solutions of

$$-\Delta u - \frac{\kappa}{d^2} u + |u|^{q-1} u = 0 \quad (4.1)$$

in a bounded convex domain Ω of \mathbb{R}^N with a smooth boundary containing 0 where d is the distance function to the boundary, κ a constant in $(0, \frac{1}{4}]$ and $q > 1$. Although it is not bounded, the model case is $\Omega = \mathbb{R}_+^N = \{(x', x_N) : x' \in \mathbb{R}^{N-1}, x_N > 0\}$ which is represented by (r, σ) , $r > 0$, $\sigma \in S_+^{N-1}$ in spherical coordinates. Then

$$\mathcal{L}_\kappa u = -u_{rr} - \frac{N-1}{r}u_r - \frac{1}{r^2}\Delta_{S^{N-1}}u - \frac{\kappa}{r^2(\mathbf{e}_N \cdot \sigma)^2}u + |u|^{q-1}u. \quad (4.2)$$

We also denote by ∇' the covariant gradient on S^{N-1} in the metric of S^{N-1} obtained by the imbedding into \mathbb{R}^N .

4.1 The spherical \mathcal{L}_κ -harmonic problem

It is straightforward to check that the Poisson kernel $K_{\mathcal{L}_\kappa}$ of \mathcal{L}_κ in \mathbb{R}_+^N has the following expression

$$K_{\mathcal{L}_\kappa}(x, \xi) = c_{N, \kappa} \frac{x_N^{\frac{\alpha_+}{2}}}{|x - \xi|^{N + \alpha_+ - 2}}. \quad (4.3)$$

In spherical coordinates

$$K_{\mathcal{L}_\kappa}(x, 0) = c_{N, \kappa} r^{2-N-\frac{\alpha_+}{2}} \psi(\sigma) \quad r > 0, \sigma \in S_+^{N-1}$$

where $\psi_\kappa(\sigma) = x_N \lfloor_{S_+^{N-1}} \frac{\alpha_+}{2} = (\mathbf{e}_N \cdot \sigma)^{\frac{\alpha_+}{2}}$ solves

$$\begin{aligned} -\Delta_{S^{N-1}} \psi_\kappa - \mu_\kappa \psi_\kappa - \frac{\kappa}{(\mathbf{e}_N \cdot \sigma)^2} \psi_\kappa &= 0 & \text{in } S_+^{N-1} \\ \psi_\kappa &= 0 & \text{in } \partial S_+^{N-1}, \end{aligned} \quad (4.4)$$

and

$$\mu_\kappa = \frac{\alpha_+}{2}(N + \frac{\alpha_+}{2} - 2) \quad (4.5)$$

Notice that equation (4.4) admits a unique positive solution with supremum 1. We could have defined the first eigenvalue μ_κ of the operator

$$\phi \mapsto \mathcal{L}'_\kappa w := -\Delta_{S^{N-1}} w - \frac{\kappa}{(\mathbf{e}_N \cdot \sigma)^2} w$$

by

$$\mu_\kappa = \inf \left\{ \frac{\int_{S_+^{N-1}} (|\nabla w|^2 - \kappa(\mathbf{e}_N \cdot \sigma)^{-2} w^2) dS}{\int_{S_+^{N-1}} \phi^2 dS} : w \in H_0^1(S_+^{N-1}), w \neq 0 \right\}. \quad (4.6)$$

By [?] the infimum exists since $\rho(\sigma) = x_N \lfloor_{S_+^{N-1}} = \mathbf{e}_N \cdot \sigma$ is the first eigenfunction of $-\Delta_{S^{N-1}}$ in $H_0^1(S_+^{N-1})$. The minimizer ψ_κ belongs to $H_0^1(S_+^{N-1})$ only if $1 < \kappa < \frac{1}{4}$. Furthermore

$$\psi_\kappa \in \mathbf{Y}(S_+^{N-1}) := \{\phi \in H_{loc}^1(S^{N-1}) : \rho^{-\frac{\alpha_+}{2}} \phi \in H^1(S_+^{N-1}, \rho^{\alpha_+})\}. \quad (4.7)$$

We can also define μ_k by

$$\mu_k = \inf \left\{ \frac{\int_{S_+^{N-1}} |\nabla'(\rho^{-\frac{\alpha_+}{2}} \omega)|^2 \rho^{\alpha_+} dS}{\int_{S_+^{N-1}} \omega^2 dS} : \omega \in \mathbf{Y}(S_+^{N-1}) \setminus \{0\} \right\}. \quad (4.8)$$

We can use the symmetry of the operator to obtain the second eigenvalue and eigenfunction of \mathcal{L}'_κ on S_+^{N-1} . We first notice that for $j = 1, \dots, N-1$, the function

$$x \mapsto \frac{x_N^{\frac{\alpha_+}{2}} x_j}{|x|^{N+\alpha_+}} \quad (4.9)$$

is \mathcal{L}_κ -harmonic in \mathbb{R}_+^{N-1} , positive (resp. negative) on $\{x = (x_1, \dots, x_N : x_j > 0, x_N > 0)\}$ (resp. $\{x = (x_1, \dots, x_N : x_j < 0, x_N > 0)\}$) and vanishes on $\{x = (x_1, \dots, x_N : x_j = 0, x_N = 0)\}$.

Proposition 4.1. *For any $j = 1, \dots, N-1$ the function*

$$\sigma \mapsto \psi_{\kappa,j}(\sigma) = (\mathbf{e}_N \cdot \sigma)^{\frac{\alpha_+}{2}} \mathbf{e}_j \cdot \sigma$$

satisfies

$$\mathcal{L}_\kappa \psi_{\kappa,j} = (\mu_\kappa + N - 1 + \alpha_+) \rho_{\kappa,j} \quad (4.10)$$

in S_+^{N-1} . It is positive (resp. negative) on $S_+^{N-1} \cap \{x = (x_1, \dots, x_N) = x_j > 0\}$ (resp. $S_+^{N-1} \cap \{x = (x_1, \dots, x_N) = x_j < 0\}$) and it vanishes on $\partial S_+^{N-1} \cap \{x = (x_1, \dots, x_N) = x_j = 0\}$. The real number

$$\mu_{\kappa,2} = \mu_\kappa + N - 1 + \alpha_+ = \left(\frac{\alpha_+}{2} + 1\right)(N + \frac{\alpha_+}{2} - 1)$$

is the second eigenvalue of \mathcal{L}'_κ in $\mathbf{Y}(S_+^{N-1})$.

Proof. There holds

$$\begin{aligned} \mathcal{L}_\kappa \psi_{\kappa,j} &= \mathbf{e}_j \cdot \sigma \mathcal{L}_\kappa \psi_\kappa + \psi_\kappa \Delta_{S^{N-1}} \mathbf{e}_j \cdot \sigma + 2 \nabla' \psi_\kappa \cdot \nabla' \mathbf{e}_j \cdot \sigma \\ &= (\mu_\kappa + N - 1) \psi_{\kappa,j} - \alpha_+ (\mathbf{e}_N \cdot \sigma)^{\frac{\alpha_+}{2} - 1} \nabla'(\mathbf{e}_j \cdot \sigma) \cdot \nabla'(\mathbf{e}_N \cdot \sigma). \end{aligned}$$

Now

$$\nabla\left(\frac{x_j}{r}\right) = \left(\frac{x_j}{r}\right)_r \frac{x}{r} + \frac{1}{r} \nabla'\left(\frac{x_j}{r}\right) = \frac{1}{r} \nabla'\left(\frac{x_j}{r}\right) = \frac{1}{r} \mathbf{e}_j - \frac{x_j}{r^3} x,$$

thus

$$\nabla\left(\frac{x_j}{r}\right) \cdot \nabla\left(\frac{x_N}{r}\right) = -\frac{x_j x_N}{r^4} = \frac{1}{r^2} \nabla'\left(\frac{x_j}{r}\right) \cdot \nabla'\left(\frac{x_N}{r}\right) = \frac{1}{r^2} \nabla'(\mathbf{e}_j \cdot \sigma) \cdot \nabla'(\mathbf{e}_N \cdot \sigma)$$

which implies

$$\nabla'(\mathbf{e}_j \cdot \sigma) \cdot \nabla'(\mathbf{e}_N \cdot \sigma) = -\frac{x_j x_N}{r^2} = -\frac{(\mathbf{e}_j \cdot \sigma)(\mathbf{e}_N \cdot \sigma)}{r^2}$$

and finally

$$\mathcal{L}_\kappa \psi_{\kappa,j} = (\mu_\kappa + N - 1 + \alpha_+) \psi_{\kappa,j}. \quad (4.11)$$

Since $S_+^{N-1} = \{(\sigma' \sin \theta, \cos \theta) : \sigma' \in S^{N-2}, \theta \in [0, \frac{\pi}{2}]\}$, $\mathbf{e}_N \cdot \sigma = \cos \theta$, $\mathbf{e}_j \cdot \sigma = \mathbf{e}_j \cdot \sigma' \sin \theta$ and $dS = (\sin \theta)^{N-2} dS' d\theta$ where dS and dS' are the volume element of S^{N-1} and S^{N-2} respectively, we derive from the fact that $\sigma' \mapsto \mathbf{e}_j \cdot \sigma'$ is an odd function on S^{N-2} ,

$$\begin{aligned} \int_{S_+^{N-1}} \psi_{\kappa,j} \psi_\kappa dS &= \int_{S_+^{N-1}} (\mathbf{e}_N \cdot \sigma)^{\alpha_+} \mathbf{e}_j \cdot \sigma dS \\ &= \int_0^{\frac{\pi}{2}} \left(\int_{S^{N-2}} \mathbf{e}_j \cdot \sigma' dS' \right) (\cos \theta)^{\alpha_+} (\sin \theta)^{N-1} d\theta \\ &= 0. \end{aligned}$$

Hence $\psi_{\kappa,j}$ is an eigenvalue of \mathcal{L}'_{κ} in $\mathbf{Y}(S_+^{N-1})$ with two nodal domains and the space the $\rho_{\kappa,j}$ span is (N-1)-dimensional and any linear combination of the $\rho_{\kappa,j}$ has exactly two nodal domains since

$$\sum_{j=1}^{N-1} a_j \psi_{\kappa,j} = (\mathbf{e}_N \cdot \sigma)^{\frac{\alpha_+}{2}} \left(\sum_{j=1}^{N-1} a_j \mathbf{e}_j \right) \cdot \sigma.$$

This implies that $\mu_{\kappa,2}$ is the second eigenvalue. \square

4.2 The nonlinear eigenvalue problem

If we look for separable solutions under the form

$$u(x) = u(r, \sigma) = r^{\alpha} \omega(\sigma)$$

then necessarily $\alpha = -\frac{2}{q-1}$ and ω is a solution of

$$\begin{aligned} -\Delta_{S^{N-1}} \omega - \ell_{q,N} \omega - \frac{\kappa}{(\mathbf{e}_N \cdot \sigma)^2} \omega + |\omega|^{q-1} \omega &= 0 & \text{in } S_+^{N-1} \\ \omega &= 0 & \text{in } \partial S_+^{N-1}, \end{aligned} \quad (4.12)$$

$$\ell_{q,N} = \frac{2}{q-1} \left(\frac{2}{q-1} + 2 - N \right) \quad (4.13)$$

and (4.6) is transformed accordingly. We denote by

$$\mathcal{E}_{\kappa} = \{ \omega \in \mathbf{Y}(S_+^{N-1}) \cap L^{q+1}(S_+^{N-1}) \text{ s. t. (4.12) holds} \} \quad (4.14)$$

and by \mathcal{E}_{κ}^+ the set of the nonnegative ones. We also recall that $q_c := \frac{2N + \alpha_+}{2N - 4 + \alpha_+}$ and we define a second critical value $q_e := \frac{2N + 2 + \alpha_+}{2N - 2 + \alpha_+}$.

The following result holds

Theorem 4.2. *Assume $0 < \kappa \leq \frac{1}{4}$ and $q > 1$, then*

(i) *If $q \geq q_c$, $\mathcal{E}_{\kappa} = \{0\}$.*

(ii) *If $1 < q < q_c$, \mathcal{E}_{κ}^+ contains exactly two elements: 0 and ω_{κ} . Furthermore ω_{κ} depends only on the azimuthal angle θ .*

(iii) *If $q_e \leq q < q_c$, \mathcal{E}_{κ} contains three elements: 0, ω_{κ} and $-\omega_{\kappa}$.*

Proof. We recall that $q \geq q_c \iff \ell_{q,N} \leq \mu_{\kappa}$. Then non-existence follows by multiplying by ω and integrating on S_+^{N-1} . For existence, we consider the functional

$$J_{\kappa}(w) = \int_{S_+^{N-1}} \left(|\nabla'(w)|^2 + (\mu_{\kappa} - \ell_{q,N}) w^2 + \frac{2}{q+1} \psi_{\kappa}^{q-1} |w|^{q+1} \right) \psi_{\kappa}^2 dS, \quad (4.15)$$

defined in $H^1(S_+^{N-1}, \psi_{\kappa}^2 dS) \cap L^{q+1}(S_+^{N-1}, \psi_{\kappa}^{q+1} dS)$. Since $\mu_{\kappa} - \ell_{q,N} < 0$, there exists a nontrivial minimum $w_{\kappa} > 0$, which satisfies

$$-\operatorname{div}(\psi_{\kappa}^2 \nabla' w_{\kappa}) + (\mu_{\kappa} - \ell_{q,N}) \psi_{\kappa}^2 w_{\kappa} + \psi_{\kappa}^{q+1} w_{\kappa}^q = 0 \quad (4.16)$$

If we set $\omega_\kappa = \psi_\kappa w_\kappa$, then ω_κ satisfies

$$\mathcal{L}'_\kappa \omega_\kappa - \ell_{q,N} \omega_\kappa + \omega_\kappa^q = 0 \quad \text{in } S_+^{N-1}. \quad (4.17)$$

By monotonicity we derive that $\omega_\kappa \in L^p(S_+^{N-1})$ for any $1 < p < \infty$ and finally, that ω_κ satisfies the regularity estimates of Lemma 2.9 and Lemma 2.10. Moreover $\omega_\kappa > 0$ by the maximum principle.

In the case $q \geq q_c$ or equivalently $\mu_\kappa - \ell_{q,N} \geq 0$, nonexistence of nontrivial solution is clear from (4.16).

Uniqueness. By Proposition 2.8 $\omega_\kappa(x) \leq c_{86}(\rho(x))^{\frac{\alpha_+}{2}}$ and by standard scaling techniques $|\nabla \omega_\kappa(x)| \leq c_{87}(\rho(x))^{\frac{\alpha_+}{2}-1}$. Assume now two different positive solutions of (4.12) ω_κ and ω'_κ exist. Since $\max\{\omega_\kappa, \omega'_\kappa\}$ and $\omega_\kappa + \omega'_\kappa$ are respectively a subsolution and a supersolution and they are ordered, we can assume that $\omega'_\kappa < \omega_\kappa < c\omega'_\kappa$ for some $c > 1$. Let $\epsilon > 0$ and $\epsilon' = c^{-1}\epsilon$, then $\epsilon\omega'_\kappa \geq \epsilon'\omega_\kappa$. Set

$$\vartheta_\epsilon = \frac{((\omega'_\kappa + \epsilon')^2 - (\omega_\kappa + \epsilon)^2)_+}{\omega_\kappa + \epsilon}, \quad \vartheta_{\epsilon'} = \frac{((\omega'_\kappa + \epsilon')^2 - (\omega_\kappa + \epsilon)^2)_+}{\omega'_\kappa + \epsilon'}$$

and $S_{\epsilon, \epsilon'} = \{\sigma \in S_+^{N-1} : \omega'_\kappa + \epsilon' > \omega_\kappa + \epsilon\}$. The assume that $S_{\epsilon, \epsilon'} \neq \emptyset$ for any $\epsilon > 0$. Then

$$\int_{S_{\epsilon, \epsilon'}} \left(\nabla \omega'_\kappa \cdot \nabla \vartheta_{\epsilon'} - \nabla \omega_\kappa \cdot \nabla \vartheta_\epsilon - (\ell_{q,N} + \frac{\kappa}{\rho^2})(\omega'_\kappa \cdot \vartheta_{\epsilon'} - \omega_\kappa \cdot \vartheta_\epsilon) + \omega'^q_\kappa \vartheta_{\epsilon'} - \omega^q_\kappa \vartheta_\epsilon \right) dS = 0$$

The first integrand on the l.h. side is equal to

$$\int_{S_{\epsilon, \epsilon'}} \left(\left| \nabla \omega'_\kappa - \frac{\omega'_\kappa + \epsilon'}{\omega_\kappa + \epsilon} \nabla \omega_\kappa \right|^2 + \left| \nabla \omega_\kappa - \frac{\omega_\kappa + \epsilon}{\omega'_\kappa + \epsilon'} \nabla \omega'_\kappa \right|^2 \right) dS \geq 0$$

Since $\epsilon\omega'_\kappa < \epsilon'\omega_\kappa$ and $(\omega'_\kappa + \epsilon')^2 > (\omega_\kappa + \epsilon)^2$, the second integrand on the l.h. side is equal to

$$-\int_{S_{\epsilon, \epsilon'}} \left(\ell_{q,N} + \frac{\kappa}{\rho^2} \right) \left(\frac{\omega'_\kappa}{\omega'_\kappa + \epsilon'} - \frac{\omega_\kappa}{\omega_\kappa + \epsilon} \right) ((\omega'_\kappa + \epsilon')^2 - (\omega_\kappa + \epsilon)^2) dS \geq 0.$$

At end, the last integrand is

$$\int_{S_{\epsilon, \epsilon'}} \left(\frac{\omega'^q_\kappa}{\omega'_\kappa + \epsilon'} - \frac{\omega^q_\kappa}{\omega_\kappa + \epsilon} \right) ((\omega'_\kappa + \epsilon')^2 - (\omega_\kappa + \epsilon)^2) dS$$

If we let $\epsilon \rightarrow 0$, we derive

$$\int_{S_+^{N-1}} (\omega'^{q-1}_\kappa - \omega^{q-1}_\kappa) (\omega'^2_\kappa - \omega^2_\kappa)_+ dS \leq 0$$

This yields a contradiction. Therefore uniqueness holds.

Case $q_e \leq q < q_c$. Assume ω_κ is a solution. Using the representation of S_+^{N-1} already introduced in the proof of Proposition 4.1, with $\sigma = (\sigma', \theta)$ and

$$\Delta_{S^{N-1}} \omega_\kappa = \frac{1}{(\sin \theta)^{N-2}} \frac{\partial}{\partial \theta} \left((\sin \theta)^{N-2} \frac{\partial \omega_\kappa}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \Delta_{S^{N-2}} \omega_\kappa$$

where $\Delta_{S^{N-2}}$ is the Laplace-Beltrami operator on S^{N-2} , we set

$$\bar{\omega}_\kappa(\theta) = \frac{1}{|S^{N-2}|} \int_{S^{N-2}} \omega_\kappa(\sigma', \theta) dS'(\sigma').$$

Then $\bar{\omega}_\kappa$ is independent of $\sigma' \in S^{N-2}$ and furthermore

$$\int_{S_+^{N-1}} (\omega_\kappa - \bar{\omega}_\kappa) \psi_\kappa dS = \int_0^{\frac{\pi}{2}} \left(\int_{S^{N-2}} (\omega_\kappa - \bar{\omega}_\kappa) dS' \right) (\sin \theta)^{N-2} (\cos \theta)^{\frac{\alpha_\pm}{2}} d\theta = 0,$$

thus $\bar{\omega}_\kappa$ is the projection of ω_κ onto the first eigenspace of \mathcal{L}_κ and

$$\int_{S_+^{N-1}} (\omega_\kappa - \bar{\omega}_\kappa) \mathcal{L}_\kappa (\omega_\kappa - \bar{\omega}_\kappa) dS \geq \mu_{\kappa,2} \int_{S_+^{N-1}} (\omega_\kappa - \bar{\omega}_\kappa)^2 dS.$$

At end, noting that

$$\int_{S_+^{N-2}} (\overline{g_q \circ \omega_\kappa} - g_q \circ \bar{\omega}_\kappa) (\omega_\kappa - \bar{\omega}_\kappa) dS' = 0$$

with $g_q \circ u = |u|^{q-1}u$,

$$\begin{aligned} \int_{S_+^{N-1}} (g_q \circ \omega_\kappa - \overline{g_q \circ \omega_\kappa}) (\omega_\kappa - \bar{\omega}_\kappa) dS &= \int_0^{\frac{\pi}{2}} \int_{S_+^{N-2}} (g_q \circ \omega_\kappa - \overline{g_q \circ \omega_\kappa}) (\omega_\kappa - \bar{\omega}_\kappa) dS' (\sin \theta)^{N-2} d\theta \\ &= \int_0^{\frac{\pi}{2}} \int_{S_+^{N-2}} (g_q \circ \omega_\kappa) - g_q \circ \bar{\omega}_\kappa (\omega_\kappa - \bar{\omega}_\kappa) dS' (\sin \theta)^{N-2} d\theta \\ &\geq 2^{1-q} \int_{S_+^{N-1}} |\omega_\kappa - \bar{\omega}_\kappa|^{q+1} dS, \end{aligned}$$

we derive that $w = \omega_\kappa - \bar{\omega}_\kappa$, satisfies

$$\int_{S_+^{N-1}} ((\mu_{\kappa,2} - \ell_{N,q}) (\omega_\kappa - \bar{\omega}_\kappa)^2 + 2^{1-q} |\omega_\kappa - \bar{\omega}_\kappa|^{q+1}) dS \leq 0$$

which implies $\omega_\kappa = \bar{\omega}_\kappa$ and it satisfies

$$\frac{1}{(\sin \theta)^{N-2}} \frac{d}{d\theta} \left((\sin \theta)^{N-2} \frac{d\omega_\kappa}{d\theta} \right) + \left(\ell_{q,N} + \frac{\kappa}{\cos^2 \theta} \right) \omega_\kappa - g_q \circ \omega_\kappa = 0. \quad (4.18)$$

Since $\mu_{\kappa,1} < \ell_{q,N} \leq \mu_{\kappa,2}$, by [4, Th. 4, Corol. 1] this equation admits three solutions, ω_κ , $-\omega_\kappa$ and 0. \square

Remark. For $\epsilon > 0$ small enough the function $\epsilon \psi_\kappa$ is a subsolution for problem (4.12). This implies

$$\omega_\kappa(\sigma) \geq \epsilon \psi_\kappa(\sigma) \quad \forall \sigma \in S_+^{N-1}. \quad (4.19)$$

4.3 Isolated boundary singularities

Throughout this section we assume that $\Omega \subset \mathbb{R}_+^N$, $0 \in \partial\Omega$ the tangent plane to $\partial\Omega$ at 0 is $\partial\mathbb{R}_+^N$ and that $1 < q < q_c$.

Lemma 4.3. *There holds*

$$\lim_{|x| \rightarrow 0} \frac{\mathbb{G}_{\mathcal{L}_\kappa} [(K_{\mathcal{L}_\kappa}(\cdot, 0))^q](x)}{K_{\mathcal{L}_\kappa}(x, 0)} = 0 \quad (4.20)$$

Proof. We recall the following estimates (??), (2.52)

$$(i) \quad G_{\mathcal{L}_\kappa}(x, y) \leq c_3 \min \left\{ \frac{1}{|x-y|^{N-2}}, \frac{(d(x))^{\frac{\alpha_+}{2}} (d(y))^{\frac{\alpha_+}{2}}}{|x-y|^{N+\alpha_+-2}} \right\}$$

$$(ii) \quad c_3^{-1} \frac{(d(x))^{\frac{\alpha_+}{2}}}{|x|^{N+\alpha_+-2}} \leq K_{\mathcal{L}_\kappa}(x, 0) \leq c_3 \frac{(d(x))^{\frac{\alpha_+}{2}}}{|x|^{N+\alpha_+-2}}.$$

Then

$$\begin{aligned} \frac{\mathbb{G}_{\mathcal{L}_\kappa}[K_{\mathcal{L}_\kappa}(\cdot, 0)](x)}{K_{\mathcal{L}_\kappa}(x, 0)} &\leq c_3^{q+1} |x|^{N+\alpha_+-2} \int_{\Omega} \frac{(d(y))^{\frac{(q+1)\alpha_+}{2}} dy}{|x-y|^{N+\alpha_+-2} |y|^{q(N+\alpha_+-2)}} \\ &\leq c_3^{q+1} |x|^{N+\frac{\alpha_+}{2}-q(N+\frac{\alpha_+}{2}-2)} \int_{\mathbb{R}^N} \frac{d\eta}{|e_x - \eta|^{N+\alpha_+-2} |\eta|^{q(N+\alpha_+-2)}} \end{aligned}$$

where $e_x = |x|^{-1}x$. This last integral is finite and independent of x . Since $q < q_c$, (4.20) follows. \square

Corollary 4.4. *Let $u_{k\delta_0}$ be the unique solution of*

$$\begin{aligned} \mathcal{L}_\kappa u + |u|^{q-1}u &= 0 && \text{in } \Omega \\ u &= k\delta_0 && \text{in } \partial\Omega. \end{aligned} \quad (4.21)$$

Then

$$\lim_{x \rightarrow 0} \frac{u_{k\delta_0}}{K_{\mathcal{L}_\kappa}(x)} = k. \quad (4.22)$$

Proof. This is a consequence of (4.20) and the inequality

$$k\mathbb{K}_{\mathcal{L}_\kappa}[\delta_0](x) - k^q \mathbb{G}[(\mathbb{K}_{\mathcal{L}_\kappa}[\delta_0])^q](x) \leq u_{k\delta_0}(x) \leq k\mathbb{K}_{\mathcal{L}_\kappa}[\delta_0](x).$$

\square

Proposition 4.5. *There exists $u_{\infty,0} = \lim_{k \rightarrow \infty} u_{k\delta_0}$ and there holds*

$$\lim_{\substack{x \rightarrow 0, x \in \Omega \\ |x|^{-1} \rightarrow \sigma}} |x|^{\frac{2}{q-1}} u_{\infty,0}(x) = \omega_\kappa(\sigma), \quad (4.23)$$

uniformly on compact subsets of S_+^{N-1} .

Proof. The correspondence $k \mapsto u_{k\delta_0}$ is increasing and by the Keller-Osserman estimate, it converges, when $k \rightarrow \infty$ to some smooth function $u_{\infty,0}$ defined in Ω where it satisfies (1.1). By Proposition 6.1, for any $0 < R < R_0$, $u_{k\delta_0}$, and therefore $u_{\infty,0}$, vanishes on any compact subset of $\partial\Omega \setminus \{0\}$ and furthermore

$$u_{\infty,0}(x) \leq \begin{cases} c_{K,\gamma,\kappa} (\text{dist}(x, K))^\gamma & \forall \gamma \in (\frac{\alpha_-}{2}, \frac{\alpha_+}{2}) \quad \text{if } 0 < \kappa < \frac{1}{4} \\ c_K \sqrt{\text{dist}(x, K)} \sqrt{\ln \left(\frac{\text{diam}(\Omega)}{\text{dist}(x, K)} \right)} & \text{if } \kappa = \frac{1}{4} \end{cases}$$

for all compact set $K \subset \partial\Omega \setminus \{0\}$. Combining this estimate with Propositions 6.3, 7.5 and 7.6, we obtain

$$c_{91} d(x) |x|^{-\frac{2}{q-1}-1} \leq v_{\infty,0}(x) \leq u_{\infty,0}(x) \leq c_{90} (d(x))^{\frac{\alpha_+}{2}} |x|^{-\frac{2}{q-1}-\frac{\alpha_+}{2}} \quad \forall x \in \Omega, \quad (4.24)$$

and

$$|\nabla u_{\infty,0}(x)| \leq c_{90}(d(x))^{\frac{\alpha_+}{2}-1}|x|^{-\frac{2}{q-1}-\frac{\alpha_+}{2}} \quad \forall x \in \Omega. \quad (4.25)$$

For $\ell > 0$, we put $T_\ell[v](x) = \ell^{\frac{2}{q-1}}v(\ell x)$, $\Omega_\ell = \ell^{-1}\Omega$, $d_\ell(y) = \text{dist}(y, \partial\Omega_\ell)$. If v satisfies (4.1) in Ω and vanishes on $\partial\Omega \setminus \{0\}$, $T_\ell[v]$ vanishes on $\partial\Omega_\ell \setminus \{0\}$ and satisfies

$$-\Delta T_\ell[v] - \frac{\kappa}{d_\ell^2}T_\ell[v] + |T_\ell[v]|^{q-1}T_\ell[v] = 0 \quad \in \Omega_\ell. \quad (4.26)$$

In order to avoid ambiguity, we set $u_{k\delta_0} = u_{k\delta_0}^\Omega$, $v_{k\delta_0} = v_{k\delta_0}^\Omega$, $u_{\infty,0} = u_{\infty,0}^\Omega$ and $v_{\infty,0} = v_{\infty,0}^\Omega$, then $T_\ell[u_{k\delta_0}^\Omega] = u_{\ell^{\frac{2}{q-1}-N+2-\frac{\alpha_+}{2}}k\delta_0}^{\Omega_\ell}$ and $T_\ell[v_{k\delta_0}^\Omega] = v_{\ell^{\frac{2}{q-1}-N+2-\frac{\alpha_+}{2}}k\delta_0}^{\Omega_\ell}$. Since inequalities (4.24) and (4.25) are invariant under the scaling transformation, the standard elliptic equations regularity theory yields the following estimates

$$c_{93}d_\ell(y)|y|^{-\frac{2}{q-1}-1} \leq v_{\infty,0}^{\Omega_\ell}(y) \leq u_{\infty,0}^{\Omega_\ell}(y) \leq c_{92}(d_\ell(y))^{\frac{\alpha_+}{2}}|y|^{-\frac{2}{q-1}-\frac{\alpha_+}{2}} \quad \forall y \in \Omega_\ell, \quad (4.27)$$

and

$$|\nabla u_{\infty,0}^{\Omega_\ell}(y)| \leq c_{92}(d_\ell(y))^{\frac{\alpha_+}{2}-1}|y|^{-\frac{2}{q-1}-\frac{\alpha_+}{2}} \quad \forall y \in \Omega_\ell, \quad (4.28)$$

valid for any $\ell \geq 1$. If we let $k \rightarrow \infty$, we obtain $T_\ell[u_{\infty,0}^\Omega] = u_{\infty,0}^{\Omega_\ell}$ and because of the group property of the transformation T_ℓ , $T_{\ell'}[u_{\infty,0}^{\Omega_\ell}] = u_{\infty,0}^{\Omega_{\ell'\ell}}$ for any $\ell, \ell' > 0$. Estimates (4.27) and (4.28) imply that $\{u_{\infty,0}^{\Omega_\ell}\}$ is relatively compact for the topology of convergence on compact subsets of \mathbb{R}_+^N . Therefore there exists a sequence $\{\ell_n\}$ tending to 0 and a function U such that $\{u_{\infty,0}^{\Omega_{\ell_n}}\}$ converges to U uniformly on any compact subset of \mathbb{R}_+^N . Therefore U is a weak solution of

$$-\Delta U - \frac{\kappa}{y_N^2}U + U^q = 0 \quad \text{in } \mathbb{R}_+^N \quad (4.29)$$

Furthermore

$$c_{93}y_N|y|^{-\frac{2}{q-1}-1} \leq v_{\infty,0}^{\mathbb{R}_+^N}(y) \leq u_{\infty,0}^{\mathbb{R}_+^N}(y) \leq c_{92}y_N^{\frac{\alpha_+}{2}}|y|^{-\frac{2}{q-1}-\frac{\alpha_+}{2}} \quad \forall y \in \mathbb{R}_+^N. \quad (4.30)$$

By Proposition 7.5, $v_{\infty,0}^{\mathbb{R}_+^N}(y) = |y|^{-\frac{2}{q-1}}\omega_\kappa^*(\frac{y}{|y|})$. Since $T_{\ell'}[u_{\infty,0}^{\Omega_{\ell_n}}] = u_{\infty,0}^{\Omega_{\ell'\ell_n}}$ we derive $T_{\ell'}[U] = U$ for any $\ell' > 0$, thus U is self similar. Set $\omega(\frac{y}{|y|}) = U(\frac{y}{|y|})$. If we set $\sigma = \frac{y}{|y|}$ then there holds

$$c_{93}\omega_\kappa^*(\sigma) \leq \omega(\sigma) \leq c_{92}\psi_\kappa(\sigma) \quad \forall \sigma \in S_+^{N-1}. \quad (4.31)$$

Therefore ω satisfies (4.12) and it coincides with the unique positive element ω_κ of \mathcal{E}_κ . Thus $u_{\infty,0}^{\Omega_\ell}$ converges to U on compact subsets of \mathbb{R}_+^N . In particular (4.23) holds on compact subsets of S_+^{N-1} . \square

5 The boundary trace of positive solutions

As before we assume that $0 < \kappa \leq \frac{1}{4}$, $q > 1$ and Ω is a bounded smooth domain, convex if $\kappa = \frac{1}{4}$. Although the construction of the boundary trace can be made in a more general framework, we restrict ourselves to the class $\mathcal{U}_+(\Omega)$ of positive smooth functions u satisfying

$$\mathcal{L}_\kappa u + |u|^{q-1}u = 0 \quad (5.1)$$

in Ω .

Lemma 5.1. *Let $f \in L^1_{\phi_\kappa}(\Omega)$. If u is a nonnegative solution of*

$$\mathcal{L}_\kappa u = f \quad \text{in } \Omega \quad (5.2)$$

there exists $\mu \in \mathfrak{M}_+(\partial\Omega)$ such that u admits μ for boundary trace and

$$u = \mathbb{G}_{\mathcal{L}_\kappa}[f] + \mathbb{K}_{\mathcal{L}_\kappa}[\mu]. \quad (5.3)$$

Proof. Let $v = \mathbb{G}_{\mathcal{L}_\kappa}[f]$, then $u - v$ is \mathcal{L}_κ -harmonic and positive thus the result follows. \square

Definition Let $G \subset \Omega$ be a domain. A function $u \in L^q_{loc}(G)$ is a supersolution (resp. subsolution) of (5.1) if

$$\mathcal{L}_\kappa u + |u|^{q-1}u \geq 0 \quad (\text{resp. } \mathcal{L}_\kappa u + |u|^{q-1}u \leq 0) \quad (5.4)$$

in the sense of distributions in G .

The following comparison principle holds [3, Lemma 3.2]

Proposition 5.2. *Let $G \subset \Omega$ be a smooth domain and \bar{u}, \underline{u} a pair of nonnegative supersolution and subsolution respectively in G .*

(i) *If there holds*

$$\limsup_{\text{dist}(x, \partial G) \rightarrow 0} (\bar{u}(x) - \underline{u}(x)) < 0, \quad (5.5)$$

then $\underline{u} < \bar{u}$ in G .

(ii) *Assume $\bar{G} \subset \Omega$ and \bar{u} and \underline{u} belong to $H^1(G) \cap C(\bar{G})$. If $\underline{u} \leq \bar{u}$ in ∂G , then $\underline{u} \leq \bar{u}$ in G .*

5.1 Construction of the boundary trace

We use the notations of [20]

Proposition 5.3. *Let v be a non-negative function in $C(\Omega)$.*

(i) *If v is a subsolution of (5.1), there exists a minimal solution u_* dominating v , i.e. $v \leq u_* \leq U$ for any solution $U \geq v$.*

(ii) *If v is a supersolution of (5.1), there exists a maximal solution u^* dominated by v , i.e. $U \leq u^* \leq v$ for any solution $U \leq v$.*

Proof. (i) Let $\{\Omega_n\}$ be a smooth exhaustion Ω and for each $n \in \mathbb{N}$, u_n the positive solution of

$$\begin{aligned} \mathcal{L}_\kappa u + |u|^{q-1}u &= 0 && \text{in } \Omega_n \\ u &= v && \text{in } \partial\Omega_n. \end{aligned} \quad (5.6)$$

By the comparison principle $u_n \geq v$, which implies $u_{n+1}(x) \geq u_n(x) \forall x \in \Omega_n$. Since $\{u_n\}$ is uniformly bounded on compact subsets of Ω and thus in C^2 by standard regularity arguments that $u_n \uparrow u_*$ which is a positive solution of (5.1). Furthermore, if U is any solution of (5.1) dominating v , it dominates u_n in Ω_n and thus $u_* \leq U$.

The proof of (ii) is similar: we construct a decreasing sequence $\{u'_n\}$ of nonnegative solutions of (5.1) in Ω_n coinciding with v on $\partial\Omega_n$ and dominated by v . It converges to some u^* which satisfies $U \leq u^* \leq v$ for any solution U dominated by v . \square

Proposition 5.4. Let $0 \leq u, v \in C(\Omega)$.

(i) If u and v are subsolutions (resp. supersolutions) then $\max(u, v)$ is a subsolution (resp. $\min(u, v)$ is a supersolution).

(ii) If u and v are supersolutions then $u + v$ is a supersolution.

(iii) If u is a subsolution and v is a supersolution then $(u - v)_+$ is a subsolution.

Proof. The first two statements follow Kato's inequality. The last statement is verified using that

$$\begin{aligned} -\Delta(u - v)_+ &\leq \text{sign}_+(u - v)(-\Delta(u - v)) \leq -\text{sign}_+(u - v)(u^p - v^p) + \kappa \frac{(u - v)_+}{d^2(x)} \\ &\leq -(u - v)_+^q + \kappa \frac{(u - v)_+}{d^2(x)}. \end{aligned}$$

□

Notation 5.5. Let u, v be nonnegative continuous functions in Ω .

(a) If u is a subsolution, $[u]_{\dagger}$ denotes the smallest solution dominating u .

(b) If u is a supersolution, $[u]^{\dagger}$ denotes the largest solution dominated by u .

(c) If u, v are subsolutions then $u \vee v := [\max(u, v)]_{\dagger}$.

(d) If u, v are supersolutions then $u \wedge v := [\inf(u, v)]^{\dagger}$ and $u \oplus v = [u + v]^{\dagger}$.

(e) If u is a subsolution and v is a supersolution then $u \ominus v := [(u - v)_+]_{\dagger}$.

The next result based upon local uniform estimates is due to Dynkin [10].

Proposition 5.6. (i) Let $\{u_k\} \subset C(\Omega)$ be a sequence of positive subsolutions (resp. supersolutions) of (5.1). Then $U := \sup u_k$ (resp. $U := \inf u_k$) is a subsolution (resp. supersolution).

(ii) Let $\mathcal{T} \subset C(\Omega)$ be a family of positive solutions of (5.1). Suppose that, for every pair $u_1, u_2 \in \mathcal{T}$ there exists $v \in \mathcal{T}$ such that

$$\max(u_1, u_2) \leq v \quad \text{resp.} \quad \min(u_1, u_2) \geq v.$$

Then there exists a monotone sequence $\{u_n\} \subset \mathcal{T}$ such that

$$u_n \uparrow \sup \mathcal{T} \quad \text{resp.} \quad u_n \downarrow \inf \mathcal{T}.$$

Furthermore $\sup \mathcal{T}$ (resp. $\inf \mathcal{T}$) is a solution.

Definition 5.7. Let $F \subset \partial\Omega$ be a closed set. We set

$$U_F := \sup \left\{ u \in \mathcal{U}_+(\Omega) : \lim_{x \rightarrow \xi} \frac{u(x)}{W(x)} = 0, \forall \xi \in \partial\Omega \setminus F \right\}, \quad (5.7)$$

and

$$[u]_F = \sup \left\{ v \in \mathcal{U}_+(\Omega) : v \leq u, \lim_{x \rightarrow \xi} \frac{v(x)}{W(x)} = 0, \forall \xi \in \partial\Omega \setminus F \right\} \quad (5.8)$$

Notice that $F \mapsto U_F$ and $F \mapsto [u]_F$ are increasing with respect to the inclusion order relation in $\partial\Omega$, $[u]_F = u \wedge U_F$. As a consequence of Proposition 6.3, U_F satisfies

$$\lim_{x \rightarrow \xi} \frac{U_F(x)}{W(x)} = 0, \quad \forall \xi \in \partial\Omega \setminus K. \quad (5.9)$$

Proposition 5.8. Let $E, F \subset \partial\Omega$ be closed sets. Then

(i) $U_E \wedge U_F = U_{E \cap F}$.

(ii) If $F_n \subset \partial\Omega$ is a decreasing sequence of closed sets there holds

$$\lim_{n \rightarrow \infty} U_{F_n} = U_F \quad \text{where} \quad F = \bigcap F_n.$$

Proof. (i) $U_E \wedge U_F$ is the largest solution dominated by $\inf(U_E, U_F)$ and therefore, by definition, it is the largest solution which vanishes outside $E \cap F$.

(ii) If $V := \lim U_{F_n}$ then $U_F \leq V$. But $\text{supp}(V) \subset F_n$ for each $n \in \mathbb{N}$ and consequently $V \leq U_F$. \square

For $\beta > 0$, we recall that $\Omega_\beta, \Sigma_\beta$ and the mapping $x \mapsto (d(x), \sigma(x))$ have been defined in the proof of Lemma 3.9. We also set $\Omega'_\beta = \Omega \setminus \overline{\Omega}_\beta$ and, if $Q \subset \partial\Omega$, $\Sigma_\beta(Q) = \{x \in \Omega_\beta : \sigma(x) \in Q\}$.

Proposition 5.9. *Let $u \in \mathcal{U}(\Omega)$.*

(i) *If $A, B \subset \partial\Omega$ are closed sets. Then*

$$[[u]_A]_B = [[u]_B]_A = [u]_{A \cap B}. \quad (5.10)$$

(ii) *If $\{F_n\}$ is a decreasing sequence of closed subsets of $\partial\Omega$ and $F = \cap F_n$, then*

$$[u]_{F_n} \downarrow [u]_F.$$

(iii) *If $A, B \subset \partial\Omega$ be closed sets. Then*

$$[u]_A \leq [u]_{A \cap B} + [u]_{\overline{A \setminus B}}. \quad (5.11)$$

Proof. (i) It follows directly from definition that,

$$[[u]_A]_B \leq \inf(u, U_A, U_B).$$

The largest solution dominated by u and vanishing on $A^c \cup B^c$ is $[u]_{A \cap B}$. Thus

$$[[u]_A]_B \leq [u]_{A \cap B}.$$

On the other hand

$$[u]_{A \cap B} = [[u]_{A \cap B}]_B \leq [[u]_A]_B,$$

this proves (5.10).

(ii) If $F_n \downarrow F$, it follows by Proposition 5.8-(ii) that $U_{F_n} \rightarrow U_F$, thus

$$[u]_F \leq \lim_{n \rightarrow \infty} [u]_{F_n} = \lim_{n \rightarrow \infty} u \wedge U_{F_n} \leq \lim_{n \rightarrow \infty} \inf(u, U_{F_n}) \leq \inf(u, U_F).$$

Since $[u]_F$ is the largest solution dominated by $\inf(u, U_F)$, $[u]_{F_n}$ is the largest solution dominated by $\inf(u, U_{F_n})$ and $U_{F_n} \downarrow U_F$ by Proposition 5.8, the function $v = \lim_{n \rightarrow \infty} [u]_{F_n}$ is a solution of (5.1) dominated by $\inf(u, U_F)$, thus $v \leq [u]_F$ and the proof of (ii) is complete.

(iii) Without loss of generality we assume that $A \cap B \neq \emptyset$. Let $O, O' \subset \partial\Omega$ be a relatively open set such that $A \cap B \subset O$ and $\overline{A \cap B^c} \subset O'$. Set $v = [u]_A$ and let v_β^1 be the solution of

$$\begin{aligned} \mathcal{L}_\kappa w + |w|^{q-1}w &= 0 && \text{in } \Omega'_\beta \\ w &= \chi_{\Sigma_\beta(\overline{O})}v && \text{on } \Sigma_\beta. \end{aligned}$$

Also we denote by v_β^2 and v_β^3 the solutions of the above problem with respective boundary data $\chi_{\Sigma(\overline{O'})}v$ and $\chi_{\Sigma(O^c \cap O'^c)}v$. Then $v_\beta^i \leq v|_{\Omega'_\beta} \leq v_\beta^1 + v_\beta^2 + v_\beta^3$, $i = 1, 2, 3$. Let now $\{\beta_j\}$ be a decreasing sequence converging to 0 and such that

$$v_{\beta_j}^i \rightarrow v^i \leq v \leq v^1 + v^2 + v^3, \quad i = 1, 2, 3 \text{ locally uniformly in } \Omega.$$

By definition of v^i and Proposition 6.1, we have that $v^1 \leq [v]_{\overline{O}}$, $v^2 \leq [v]_{\overline{O'}}$ and $v^3 \leq [v]_{O^c \cap O'^c}$. But by (i) we have

$$[v]_{O^c \cap O'^c} = [[u]_A]_{O^c \cap O'^c} = [u]_{A \cap O^c \cap O'^c} = 0.$$

Thus

$$v \leq [v]_{\overline{O}} + [v]_{\overline{O'}}$$

We can consider decreasing sequences $\{O_n\}$ and $\{O'_n\}$ such that $\cap \overline{O_n} = A \cap B$ and $\cap \overline{O'_n} = \overline{A \cap B^c}$. By (ii) we obtain

$$v \leq [[u]_A]_{A \cap B} + [[u]_A]_{\overline{A \cap B^c}} \leq [u]_{A \cap B} + [u]_{\overline{A \cap B^c}}$$

which is (iii). \square

Remark. Since any $u \in \mathcal{U}_+(\Omega)$ is dominated by $u_{\partial\Omega}$, it follows from (iii) that for any set $A \subset \partial\Omega$, there holds

$$u = [u]_{\partial\Omega} \leq [u]_{\overline{A}} + [u]_{\overline{\partial\Omega \setminus A}} \leq [u]_{\overline{A}} + [u]_{\overline{\partial\Omega \setminus A}}. \quad (5.12)$$

Proposition 5.10. *Let u be a positive solution of (5.1). If $u \in L^q_{\phi_\kappa}(\Omega)$ it possesses a boundary trace $\mu \in \mathfrak{M}(\partial\Omega)$, i.e., u is the solution of the boundary value problem (3.36) with this measure μ .*

Proof. If $v := \mathbb{G}_{\mathcal{L}_\kappa}[u^q]$ then $v \in L^1_{\phi_\kappa}(\Omega)$ and $u + v$ is a positive \mathcal{L}_κ -harmonic function. Hence $u + v \in L^1_{\phi_\kappa}(\Omega)$ and there exists a non-negative measure $\mu \in \mathfrak{M}(\partial\Omega)$ such that $u + v = \mathbb{K}_{\mathcal{L}_\kappa}[\mu]$. By Proposition 3.5 this implies the result. \square

Proposition 5.11. *Let u be a positive solution of (5.1) and $\mu \in \mathfrak{M}(\partial\Omega)$. If for an exhaustion $\{\Omega_n\}$ of Ω , we have*

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} Z(x) u d\omega_{\Omega_n}^{x_0} = \int_{\partial\Omega} Z(x) d\mu, \quad \forall Z \in C(\overline{\Omega}),$$

where $\omega_{\Omega_n}^{x_0}$ is the \mathcal{L}_κ -harmonic measure of Ω_n relative to a point $x_0 \in \Omega_1$, then u and $|u|^p$ belong to $L^1_{\phi_\kappa}(\Omega)$. Furthermore u possesses the boundary trace $\mu \in \mathfrak{M}(\partial\Omega)$, i.e., u is the solution of the boundary value problem (3.36) with this measure μ .

Proof. Let $\mathbb{G}_{\mathcal{L}_\kappa}^n$ be the green function of \mathcal{L}_κ in Ω_n , then

$$\mathbb{G}_{\mathcal{L}_\kappa}^n(x, y) \leq \mathbb{G}_{\mathcal{L}_\kappa}^{n+1}(x, y), \quad \forall x, y \in \Omega_n$$

and

$$\mathbb{G}_{\mathcal{L}_\kappa}^n \uparrow \mathbb{G}_{\mathcal{L}_\kappa}.$$

Since

$$\int_{\partial\Omega_n} u d\omega_{G^{x_0}} = u(x_0) + \int_{\Omega_n} \mathbb{G}_{\mathcal{L}_\kappa}^n(x, x_0) (u(x))^p dx,$$

we derive, as $n \rightarrow \infty$,

$$\mu(\partial\Omega) = u(x_0) + \int_{\Omega} \mathbb{G}_{\mathcal{L}_\kappa}(x, x_0) |u(x)|^p dx.$$

By Proposition 2.1 this implies $|u|^p \in L^1_{\phi_\kappa}(\Omega)$, and the result follows by Proposition 5.10. \square

Proposition 5.12. *If $F \subset \partial\Omega$ is a closed set and u a positive solution of (5.1) with boundary trace $\mu \in \mathfrak{M}(\partial\Omega)$, then $[u]_F$ has boundary trace $\mu \chi_F$.*

Proof. The function $[u]_F$ belongs to $\mathcal{U}_+(\Omega)$ and is dominated by u which satisfies (5.1), thus $[u]_F \in L^q_{\phi_\kappa}(\Omega)$ and $[u]_F$ admits a boundary trace $\mu_F \leq \mu$ by Proposition 5.10. Let v be the solution of (3.36) with boundary data $\mu\chi_F$. Let $O \subset \partial\Omega$ relatively open such that $F \subset O$. By 5.12 we have

$$v \leq [v]_{\overline{O}} + [v]_{\overline{O^c}}.$$

Let A be an open set such that $F \subset A \subset \overline{A} \subset O$, and for exhaustion we take $\Omega_n = \Omega'_n$ which is smooth for n large enough, and $\partial\Omega_n = \Sigma_{\frac{1}{n}}$. Then

$$\int_{\partial\Omega_n} [v]_{\overline{O^c}} d\omega_{\Omega_n}^{x_0} = \int_{\Sigma_{\frac{1}{n}}(A)} [v]_{\overline{O^c}} d\omega_{\Omega_n}^{x_0} + \int_{\partial\Omega_n \setminus \Sigma_{\frac{1}{n}}(A)} [v]_{\overline{O^c}} d\omega_{\Omega_n}^{x_0}$$

But

$$\int_{\Sigma_{\frac{1}{n}}(A)} [v]_{\overline{O^c}} d\omega_{\Omega_n}^{x_0} \leq \int_{\Sigma_{\frac{1}{n}}(A)} v d\omega_{\Omega_n}^{x_0} \rightarrow 0$$

and

$$\int_{\partial\Omega_n \setminus \Sigma_{\frac{1}{n}}(A)} [v]_{\overline{O^c}} d\omega_{\Omega_n}^{x_0} \leq \int_{\partial\Omega_n \setminus \Sigma_{\frac{1}{n}}(A)} U_{\overline{O^c}} d\omega_{\Omega_n}^{x_0} \rightarrow 0,$$

as $n \rightarrow \infty$, thus $[v]_{\overline{O^c}} = 0$ by Proposition 5.11 and therefore $v \leq [v]_{\overline{O}} \leq [u]_{\overline{O}}$. Since O be an arbitrary open set, take a sequence of open set $\{O_n\}$ such that $F \subset O_n \subset \overline{O_n} \subset O_{n-1}$ and $\cap O_n = F$. Using Proposition 5.9 we derive

$$v \leq [u]_F,$$

and thus $\mu\chi_F \leq \mu_F$. Conversely, let $Z \in C(\overline{\Omega})$, $Z \geq 0$,

$$\begin{aligned} \int_{\partial\Omega_n} Z[u]_F d\omega_{\Omega_n}^{x_0} &= \int_{\partial\Omega_n \cap \Sigma_{\frac{1}{n}}(A)} Z[u]_F d\omega_{\Omega_n}^{x_0} + \int_{\partial\Omega_n \setminus \Sigma_{\frac{1}{n}}(A)} Z[u]_F d\omega_{\Omega_n}^{x_0} \\ &\leq \int_{\partial\Omega_n \cap \Sigma_{\frac{1}{n}}(A)} Z u d\omega_{\Omega_n}^{x_0} + \int_{\partial\Omega_n \setminus \Sigma_{\frac{1}{n}}(A)} Z U_F d\omega_{\Omega_n}^{x_0} \\ &\leq I_n + II_n. \end{aligned}$$

Because of (5.9), $II_n \rightarrow 0$ as $n \rightarrow \infty$, thus

$$\int_{\partial\Omega} Z d\mu_F \leq \int_{\partial\Omega} Z \chi_F d\mu \iff \mu_F \leq \mu\chi_O,$$

and the result follow by regularity since O is arbitrary. \square

The next result shows that the boundary trace has a local character.

Proposition 5.13. *Let $u \in \mathcal{U}_+(\Omega)$ and $\xi \in \partial\Omega$. We assume that there exists $\rho > 0$ such that*

$$\int_{B_\rho(\xi) \cap \Omega} (u(x))^q \phi_\kappa(x) dx < \infty.$$

(i) *Then*

$$[u]_F^q \in L^1_{\phi_\kappa}(\Omega) \quad \forall F \subset \partial\Omega \cap B_\rho(\xi), F \text{ closed.}$$

Thus $[u]_F$ possesses a boundary trace $\mu_F \in \mathfrak{M}(\partial\Omega)$, and $\text{supp}(\mu_F) \subset F$.

(ii) There exists a nonnegative Radon measure μ_ρ on $B_\rho(\xi)$ such that for any closed set $F \subset B_\rho(\xi) \cap \partial\Omega$

$$\mu_F = \mu_\rho \chi_F,$$

and for any exhaustion $\{\Omega_n\}$ of Ω and any $Z \in C(\overline{\Omega})$ such that $\text{supp}(Z) \cap \partial\Omega \subset \partial\Omega \cap B_\rho(\xi)$

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} u(x)Z(x)d\omega_{\Omega_n}^{x_0} = \int_{\partial\Omega} u(x)Z(x)d\mu_\rho. \quad (5.13)$$

Proof. (i) Let F be a closed set and $0 < \rho' < \rho$ be such that

$$F \subset \partial\Omega \cap B_{\rho'}(\xi).$$

Since $[u]_F \leq \inf(u, U_F)$ and $U_F \in C(\overline{\Omega} \setminus F)$, we have

$$\int_{\Omega} [u]_F^q \phi_\kappa(x) dx \leq \int_{B_\rho(\xi) \cap \Omega} |u|^p \phi_\kappa(x) dx + \int_{\Omega \setminus B_\rho(\xi)} |U_F|^p \phi_\kappa(x) dx < \infty.$$

(ii) Let $0 < \rho_1 < \rho_2 < \rho$, then

$$[u]_{\overline{B_{\rho_2}(\xi)} \cap \partial\Omega} \leq u \leq [u]_{\overline{B_{\rho_2}(\xi)} \cap \partial\Omega} + U_{\overline{\partial\Omega \setminus \overline{B_{\rho_2}(\xi)}}}.$$

The function $[u]_{\overline{B_{\rho_2}(\xi)} \cap \partial\Omega}$ which belongs $L_{\phi_\kappa}^q(\Omega)$ admits a boundary trace $\nu \in \mathfrak{M}(\partial\Omega)$ and

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} U_{\overline{\partial\Omega \setminus \overline{B_{\rho_2}(\xi)}}} \zeta(x) d\omega_{\Omega_n}^{x_0} = 0,$$

for any $Z \in C(\overline{\Omega})$ such that $\text{supp}(Z) \cap \partial\Omega \subset \partial\Omega \cap B_{\rho_1}(\xi)$. Combined with Proposition 5.12 it follows identity (5.13) and finally statement (ii). \square

Definition 5.14. The set \mathcal{R}_u of boundary points a such that there exists $r > 0$ such that (5.13) holds is relatively open. Using a partition of unity there exists a positive Radon measure μ_u on \mathcal{R}_u such that

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} u(x)Z(x)d\omega_{\Omega_n}^{x_0} = \int_{\partial\Omega} u(x)Zd\mu_u \quad (5.14)$$

for any $Z \in C(\overline{\Omega})$ such that $\text{supp}(Z) \cap \partial\Omega \subset \mathcal{R}_u$. The set $\mathcal{S}_u := \partial\Omega \setminus \mathcal{R}_u$ is closed. The couple (\mathcal{S}_u, μ_u) is the boundary trace of u , denoted by $\text{Tr}_{\partial\Omega}(u)$. The measure μ_u is the regular part of $\text{Tr}_{\partial\Omega}(u)$, the set (\mathcal{S}_u) is its singular part.

Proposition 5.15. Let u be a positive solution in Ω and let $\{\Omega_n\}$ be an exhaustion of Ω . If $y \in \mathcal{S}_u$ then for every nonnegative $Z \in C(\overline{\Omega})$ such that $Z(y) \neq 0$ we have

$$\lim_{n \rightarrow \infty} \int_{\partial\Omega_n} Zud\omega_{\Omega_n}^{x_0} = \infty.$$

Proof. Let $Z \in C(\overline{\Omega})$, $Z \geq 0$, such that $Z(y) \neq 0$ and

$$\liminf_{n \rightarrow \infty} \int_{\partial\Omega_n} Zud\omega_{\Omega_n}^{x_0} < \infty.$$

There exists a subsequence n_j such that

$$\lim_{j \rightarrow \infty} \int_{\partial\Omega_{n_j}} Z u d\omega_{\Omega_n}^{x_0} = M < \infty.$$

Let r be such that $Z(x) > \frac{Z(y)}{2}$, $\forall x \in B_r(y) \cap \overline{\Omega}$, then for any $r' < r$ we have that

$$\limsup_{j \rightarrow \infty} \int_{\partial\Omega_{n_j}} [u]_{\overline{B_{r'}(y) \cap \partial\Omega}} d\omega_{\Omega_n}^{x_0} < \infty.$$

In view of the proposition of 5.11 the last fact implies that $[[u]_{\overline{B_{r'}(y)}}]^p \in L_{\phi_\kappa}(\Omega)$, which implies that

$$|u|^p \in L_{\phi_\kappa}(B_{r''}(y)) \quad \forall r'' < r'.$$

Which is clearly a contradiction, by Proposition 5.13. \square

Proposition 5.16. *Let u be a positive solution of (5.1) in Ω with boundary trace (\mathcal{S}_u, μ_u) . Then*

$$\int_{\Omega} (u \mathcal{L}_\kappa \zeta + u^q \zeta) dx = \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa} [\mu_u \chi_F] \mathcal{L}_\kappa \zeta dx,$$

for any $\zeta \in \mathbf{X}(\Omega)$ such that $\text{supp}(\zeta) \cap \partial\Omega \subset F$.

Proof. Consider the function $\zeta \in \mathbf{X}(\Omega)$ such that $\text{supp}(\zeta) \cap \partial\Omega \subset F$. Set $K = \text{supp}(\zeta)$,

$$O_\varepsilon = \{x \in \mathbb{R}^N : \text{dist}(x, K) < \varepsilon\}$$

and $\varepsilon_0 > 0$ small enough such that

$$\overline{O_\varepsilon} \cap \partial\Omega \subset \mathcal{R}_u, \quad \forall 0 < \varepsilon < \varepsilon_0.$$

Let $\varepsilon < \frac{\varepsilon_0}{4}$ and η be a cut off function such that $\eta \in C_0^\infty(O_\varepsilon)$, $0 \leq \eta \leq 1$ and $\eta \equiv 1$ on $\overline{O_{\frac{\varepsilon}{2}}}$. For $0 < \beta \leq \beta_0$, let v_β be the solution of

$$\begin{aligned} \mathcal{L}_\kappa w + |w|^{q-1} w &= 0 && \text{in } \Omega'_\beta \\ w &= \eta u && \text{on } \Sigma_\beta. \end{aligned}$$

Then there exists a sequence $\{\beta_j\}$ decreasing to 0 such that $v_{\beta_j} \rightarrow v$ locally uniformly, and

$$v \leq [u]_{\partial\Omega \cap \overline{O_\varepsilon}}.$$

Thus v has boundary trace μ_0 such that

$$\mu_0 \leq \mu_u \chi_{\partial\Omega \cap \overline{O_\varepsilon}}.$$

Let v_β^1 and v_β^2 be the solutions of

$$\begin{aligned} \mathcal{L}_\kappa w + |w|^{q-1} w &= 0 && \text{in } \Omega'_\beta \\ w &= \eta [u]_{\partial\Omega \cap \overline{O_{2\varepsilon}}} && \text{on } \Sigma_\beta. \end{aligned}$$

and

$$\begin{aligned} \mathcal{L}_\kappa w + |w|^{q-1} w &= 0 && \text{in } \Omega'_\beta \\ w &= \eta U_{\partial\Omega \setminus O_{2\varepsilon}} && \text{on } \Sigma_\beta, \end{aligned}$$

respectively. Notice that $u \leq [u]_{\partial\Omega \cap \overline{O_{2\varepsilon}}} + U_{\partial\Omega \setminus O_{2\varepsilon}}$ we have that

$$v_\beta \leq v_\beta^1 + v_\beta^2 \leq [u]_{\partial\Omega \cap \overline{O_{2\varepsilon}}} + v_\beta^2.$$

Since $[u]_{\partial\Omega \cap \overline{O_{2\varepsilon}}}^p \in L^1_{\phi_\kappa}(\Omega)$. By (6.20) we have that

$$\eta(x)U_{\partial\Omega \setminus O_{2\varepsilon}}(x) \leq c_{90}d^{\frac{\alpha_+}{2}}(x) \quad \forall x \in \Omega.$$

where $c_{90} > 0$ depends on N, q, κ and $\text{dist}(\text{supp}(\eta), \partial\Omega \setminus O_\varepsilon)$. Thus $v_\beta^2(x) \leq c_{90}d^{\frac{\alpha_+}{2}}(x)$ and

$$v_\beta \leq [u]_{\partial\Omega \cap \overline{O_{2\varepsilon}}} + c_{90}d^{\frac{\alpha_+}{2}}(x), \quad \forall x \in \Omega'_\beta. \quad (5.15)$$

Let w_β be the solution of

$$\begin{aligned} \mathcal{L}_\kappa w + |w|^{q-1}w &= 0 && \text{in } \Omega'_\beta \\ w &= \chi_{\Sigma_\beta(\overline{\partial\Omega \setminus O_{\frac{\varepsilon}{2}}})}[u]_F && \text{on } \Sigma_\beta. \end{aligned}$$

Then

$$[u]_F \leq v_\beta + w_\beta \quad \text{in } \Omega'_\beta.$$

We have that $w_{\beta_j} \rightarrow 0$ locally uniformly in Ω which implies that

$$[u]_F \leq v.$$

Thus we have

$$\mu_u \chi_F \leq \mu_0 \leq \mu_u \chi_{\partial\Omega \cap \overline{O_\varepsilon}}. \quad (5.16)$$

Set $Z = \eta\zeta_\beta$ where ζ_β is the solution of

$$\begin{aligned} \mathcal{L}_\kappa w &= \mathcal{L}_\kappa \zeta && \text{in } \Omega'_\beta \\ w &= 0 && \text{on } \Sigma_\beta. \end{aligned}$$

Since $\zeta \in \mathbf{X}(\Omega)$, there exists a constant c_{91} such that $\zeta_\beta \leq c_{91}\phi_\kappa$ in Ω'_β . Thus there exists a decreasing sequence $\{\beta_j\}$ converging to 0 such that $\zeta_{\beta_j} \rightarrow \zeta$ locally uniformly. Now,

$$\begin{aligned} \int_{\Omega'_\beta} u \mathcal{L}_\kappa Z dx + \int_{\Omega'_\beta} u^q Z dx &= - \int_{\partial\Omega'_\beta} \frac{\partial Z}{\partial \mathbf{n}} u dS \\ &= - \int_{\partial\Omega'_\beta} \frac{\partial \zeta_\beta}{\partial \mathbf{n}} \eta u dS. \\ &= \int_{\Omega'_\beta} v_\beta \mathcal{L}_\kappa \zeta_\beta dx + \int_{\Omega'_\beta} v_\beta^q \zeta_\beta dx \\ &= \int_{\Omega'_\beta} v_\beta \mathcal{L}_\kappa \zeta dx + \int_{\Omega'_\beta} v_\beta^q \zeta dx, \end{aligned} \quad (5.17)$$

We note here that in view of the proof of (6.22), we have

$$|\nabla \zeta_\beta| \leq c_{92}d^{\frac{\alpha_+}{2}}, \quad \forall x \in \Omega'_\beta,$$

where the constant $c_{92} > 0$ does not depend on β . Also by remark ?? and our assumptions we have

$$\int_{\Omega \cap O_{2\varepsilon}} [u]_{\partial\Omega \cap \bar{O}_{3\varepsilon}} d^{\frac{\alpha_+}{2}-1} dx < \infty.$$

By (6.20)

$$\int_{\Omega \cap O_{2\varepsilon}} U_{\partial\Omega \setminus O_{3\varepsilon}} d^{\frac{\alpha_+}{2}-1} dx < \infty.$$

The last two inequalities above implies that

$$\int_{\Omega \cap O_{2\varepsilon}} u d^{\frac{\alpha_+}{2}-1} dx \leq \int_{\Omega \cap O_{2\varepsilon}} U_{\partial\Omega \setminus O_{3\varepsilon}} d^{\frac{\alpha_+}{2}-1} dx + \int_{\Omega \cap O_{2\varepsilon}} [u]_{\partial\Omega \cap \bar{O}_{3\varepsilon}} d^{\frac{\alpha_+}{2}-1} dx < \infty.$$

Combining all above we can choose a decreasing subsequence $\{\beta_j\}$ to the origin such that if we take the limit in (5.17) to obtain

$$\int_{\Omega} u \mathcal{L}_{\kappa} \zeta dx + \int_{\Omega} u^q \zeta dx = \int_{\Omega} v \mathcal{L}_{\kappa} \zeta dx + \int_{\Omega} v^q \zeta dx = \int_{\Omega} \mathbb{K}_{\mathcal{L}_{\kappa}}[\mu] \mathcal{L}_{\kappa} \zeta dx$$

Be (5.16) we have the desired result if we send ε to zero. \square

5.2 Subcritical case

We recall that

$$q_c = \frac{N + \frac{\alpha_+}{2}}{N + \frac{\alpha_+}{2} - 2}$$

is the critical exponent for the equation. If $1 < q < q_c$, we have seen in section 4 that for any $a \in \partial\Omega$ and $k \geq 0$ there exists $u_{k\delta_a}$ and $\lim_{k \rightarrow \infty} u_{k\delta_a} = u_{\infty, a}$. Furthermore, by Proposition 5.15, $Tr_{\partial\Omega}(u_{\infty, a}) = (\{a\}, 0)$.

Theorem 5.17. *Assume $1 < q < q_c$ and $a \in \mathcal{S}_u$. Then*

$$u(x) \geq u_{\infty, a}(x), \quad \forall x \in \Omega. \quad (5.18)$$

For proof of the above uses some ideas of the proof of Theorem 7.1 in [19] and needs several intermediate lemmas.

Lemma 5.18. *Assume $1 < q < q_c$. Let $\{\xi^n\}$ be a sequence of points in Ω converging to $a \in \partial\Omega$ and let $l \in (0, 1)$. we define the sets*

$$\Omega_n := \Omega'_{d(\xi^n)} = \{x \in \Omega : d(x) > d(\xi^n)\} \quad \text{and} \quad \Sigma_n := \partial\Omega_n. \quad (5.19)$$

Let $x_0 \in \Omega'_1$ and denote by $\omega_n := \omega_{\Omega_n}^{x_0}$ the \mathcal{L}_{κ} -harmonic measure in Ω_n relative to x_0 . Put

$$V_n = B_{lr_n}(\xi^n) \cap \partial\Omega_n, \quad r_n = d(\xi^n).$$

Let $h_n \in L^\infty(\Sigma_n)$ $n = 1, 2, \dots$, and suppose that there exist numbers c and k such that

$$\text{supp}(h_n) \subset V_n \quad \text{and} \quad 0 \leq h_n \leq cr_n^{-N - \frac{\alpha_+}{2} + 2} \quad (5.20)$$

and

$$\lim_{n \rightarrow \infty} \int_{\Sigma_n} h_n \phi d\omega_{\Omega_n}^{x_0} = k\phi(a), \quad \forall \phi \in C(\bar{\Omega}).$$

Let w_n be the solution of the problem

$$\begin{aligned} \mathcal{L}_\kappa w_n + |w_n|^{q-1} w_n &= 0 & \text{in } \Omega_n \\ w_n &= h_n & \text{on } \partial\Sigma_n. \end{aligned}$$

Then

$$w_n \rightarrow u_{k,a} \quad \text{locally uniformly in } \Omega.$$

Proof. Let $\eta^n \in \partial\Omega$ be such that $d(\xi^n) = |\xi^n - \eta^n|$. By Corollary 2.29 we have

$$\mathbb{K}_{\mathcal{L}_\kappa}(x, \eta^n) \geq \frac{1}{c_{43}} r_n^{-N - \frac{\alpha_+}{2} + 2} \geq \frac{1}{cc_{43}} h_n(x), \quad \forall x \in \Sigma_n, \quad (5.21)$$

by the maximum principle,

$$\mathbb{K}_{\mathcal{L}_\kappa}(x, \eta^n) \geq \frac{1}{cc_{43}} w_n(x), \quad \forall x \in \Omega_n. \quad (5.22)$$

Moreover

$$\int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}^q(x, y) d^{\frac{\alpha_+}{2}}(x) dx \leq c(q), \quad \forall 1 < q < q_c,$$

where $c(q)$ is a constant independent of y . Since q is subcritical, it follows that the sequences $\{\mathbb{K}_{\mathcal{L}_\kappa}^q(\cdot, \eta^n)\}$ and $\{\mathbb{K}_{\mathcal{L}_\kappa}(\cdot, \eta^n)\}$ are uniformly integrable in $L^1_{\phi_\kappa}(\Omega)$. Let \bar{w}_n denotes the extension of w_n to Ω defined by $\bar{w}_n = 0$ in $\Omega \setminus \Omega_n$. In view of (5.21) we conclude that the sequences $\{\bar{w}_n^q\}$ and $\{\bar{w}_n\}$ are uniformly integrable in $L^1_{\phi_\kappa}(\Omega)$, and locally uniformly bounded in Ω . By regularity results for elliptic equations there exists a subsequence of $\{\bar{w}_n\}$, say again $\{\bar{w}_n\}$ that converges locally uniformly in Ω to a solution w of (5.1). This fact and the uniform integrability mentioned above imply that

$$w_n \rightarrow w \quad \text{in } L^q_{\phi_\kappa}(\Omega) \cap L^1_{\phi_\kappa}(\Omega).$$

Since $w \in L^q_{\phi_\kappa}(\Omega)$ by Proposition 5.10 there exists $\mu \in \mathfrak{M}(\Omega)$ such that

$$\int_{\Omega} w \mathcal{L}_\kappa \eta dx + \int_{\Omega} |w|^{q-1} w \eta dx = - \int_{\Omega} \mathbb{K}_{\mathcal{L}_\kappa}[\mu] \mathcal{L}_\kappa \eta dx \quad \forall \eta \in \mathbf{X}(\Omega).$$

Furthermore, using (5.21) we prove below that measure μ is concentrated at a . Let $\phi_{\kappa,n}$ be the first eigenfunction of \mathcal{L}_κ in Ω_n normalized by $\phi_{\kappa,n}(x_0) = 1$ for some $x_0 \in \Omega_1$. Let $\eta \in \mathbf{X}(\Omega)$ be nonnegative function and let η_n be the solution of the problem

$$\begin{aligned} \mathcal{L}_\kappa \eta_n &= \frac{\phi_{\kappa,n}}{\phi_\kappa} \mathcal{L}_\kappa \eta & \text{in } \Omega_n \\ \eta_n &= 0 & \text{in } \partial\Omega_n. \end{aligned}$$

Then $\eta_n \in C^2(\bar{\Omega}_n)$ and since $\phi_{\kappa,n} \rightarrow \phi_\kappa$,

$$\mathcal{L}_\kappa \eta_n \rightarrow \mathcal{L}_\kappa \eta \quad \text{and } \eta_n \rightarrow \eta \quad \text{as } n \rightarrow \infty.$$

Then we have

$$\int_{\Omega_n} w_n \mathcal{L}_\kappa \eta_n dx + \int_{\Omega} |w_n|^{q-1} w_n \eta dx = - \int_{\Omega} v_n \mathcal{L}_\kappa \eta_n dx, \quad (5.23)$$

where v_n solves

$$\begin{aligned} \mathcal{L}_\kappa v_n &= 0 & \text{in } \Omega_n \\ v_n &= h_n & \text{on } \partial\Sigma_n. \end{aligned}$$

By the same arguments as above there exists a subsequence of $\{v_n \chi_{\Omega_n}\}$, for simplicity $\{v_n \chi_{Gw_n}\}$, converging in $L^1_{\phi_\kappa}(\Omega)$ to a nonnegative \mathcal{L}_κ -harmonic function v . By (5.21) we have

$$cc_{43} K_{\mathcal{K}_\kappa}(x, a) \geq v(x), \quad \forall x \in \Omega. \quad (5.24)$$

Thus there exists a measure $\nu \in \mathfrak{M}(\partial\Omega)$, concentrated at a such that v solves

$$\begin{aligned} \mathcal{L}_\kappa v &= 0 & \text{in } \Omega \\ v &= \nu & \text{on } \partial\Omega. \end{aligned}$$

But

$$k = \lim_{n \rightarrow \infty} \int_{\Sigma_n} h_n d\omega_{\Omega_n}^{x_0} = \lim_{n \rightarrow \infty} v_n(x_0) = v(x_0) = \int_{\partial\Omega} d\nu,$$

the results follows if we sent n to ∞ in (5.23). \square

Lemma 5.19. *For every $l \in (0, 1)$ there exists a constant $c_l = c(N, \kappa, q, l)$ such that, for every positive solution u of (5.1) in Ω and every $x_0 \in \Omega$,*

$$u(x) \leq c_l u(y), \quad \forall x, y \in B_{lr_0}(x_0), \quad r_0 = d(x_0). \quad (5.25)$$

Proof. Put $r_1 = \frac{1+l}{2} r_0$. Then u satisfies

$$\mathcal{L}_\kappa u + u^q = 0, \quad \in B_{r_1}(x_0).$$

Denote by Ω_{r_0} the domain

$$\Omega_{r_0} = \{y \in \mathbb{R}^n : r_0 y \in \Omega\}.$$

Set $v(y) = u(r_0 y)$, and $y_0 = \frac{x_0}{r}$, then $v(y)$ satisfies

$$-\Delta v - \kappa \frac{v}{\text{dist}^2(y, \partial\Omega_{y_0})} + r_0^2 |v|^{q-1} v = 0, \quad \in B_{\frac{1+l}{2}}(y_0).$$

Now note that

$$\frac{1}{\text{dist}^2(y, \partial\Omega_{y_0})} \leq \frac{4}{(1-l)^2}, \quad \forall y \in B_{\frac{1+l}{2}}(y_0)$$

and by Keller Osserman condition

$$r_0^2 |v(y)|^{q-1} = r_0^2 |u(r_0 y)|^{q-1} \leq C(\Omega, \kappa, N) r_0^2 \frac{1}{d^2(r_0 y)} \leq C(\Omega, \kappa, N) B_{\frac{1+l}{2}}(y_0).$$

Thus by Harnack inequality there exists a constant $c_l > 0$ such that

$$v(z) \leq c_l v(y), \quad \forall z, y \in B_l(y_0),$$

and the results follows. \square

For the proof of the next lemma we need some notations. Let $\beta > 0$ and $\xi \in \Sigma_\beta = \partial\Omega'_\beta$. We set $\Delta_r^\beta(\xi) = \Sigma_\beta \cap B_r(\xi)$ and, for $0 < r < \beta < 2r$, $x_r^\beta = x_r^\beta(\xi) \in \bar{\Omega}_\beta$, such that $d(x_r) = |x_r^\beta - \xi| = r$. Also we denote by $\omega_{\Omega'_\beta}^x$ the \mathcal{L}_κ -harmonic measure in $\Omega'_\beta := \Omega \setminus \bar{\Omega}_\beta$ relative to x

Lemma 5.20. *Let $r_0 = r_0(\Omega) > 0$ be small enough and $0 < r \leq \frac{r_0}{4}$. Then there exists a constant c_{95} which depends only on Ω, N such that*

$$\omega_{\Omega'_\beta}^x(\Delta_r(\xi)) > c_{95} \quad \forall x \in \Omega \cap B_{\frac{r}{2}}(\xi). \quad (5.26)$$

Proof. Since $x \mapsto \omega_{\Omega'_\beta}^x$ is a positive \mathcal{L}_κ -harmonic in Ω'_β , it is a positive superharmonic function (relative to the Laplacian) in Ω'_β . Thus

$$\omega_{\Omega'_\beta}^x \geq v_{\Omega'_\beta}^x, \quad \forall x \in \Omega'_\beta,$$

where $v_{\Omega'_\beta}^x$ is the standard harmonic measure in Ω'_β relative to $x \in \Omega'_\beta$. The result follows by Lemma 2.1 in [7]. \square

Lemma 5.21. *Let $\kappa = \frac{1}{4}$, $\varepsilon \in (0, 1)$ and $x_0 \in \Omega_1$. Let $\{\xi^n\}$ be a sequence of points in Ω converging to $a \in \partial\Omega$. Then there exist $n_0 = n_0(\varepsilon, \Omega) \in \mathbb{N}$ and $c_{96} = c_{96}(\Omega, N, \varepsilon)$ such that*

$$\omega_{\Omega_n}^{x_0}(B_{d(\xi^n)}(\xi^n) \cap \partial\Omega_n) \geq c_{96} d(\xi^n)^{N+\frac{1}{2}-2} (-\log d(\xi^n))^{1-\varepsilon} \quad \forall n \geq n_0. \quad (5.27)$$

Proof. We recall that for any $n \in \mathbb{N}$ Ω_n is defined by (5.19), $G_{\mathcal{L}_\frac{1}{4}}^{\Omega_n} \leq G_{\mathcal{L}_\frac{1}{4}}^\Omega := G_{\mathcal{L}_\frac{1}{4}}^\Omega$, and for a fixed point $y_0 \in \Omega_1$

$$G_{\mathcal{L}_\frac{1}{4}}^{\Omega_n} \chi_{\Omega_n}(x) \uparrow G_{\mathcal{L}_\frac{1}{4}}(x, y_0), \quad \text{locally uniformly in } \Omega \setminus y_0. \quad (5.28)$$

Set $x(\xi^n) = x_{r_n}^{2r_n}(\xi^n)$, with $r_n = \frac{d(\xi^n)}{2}$. By (2.9) we have

$$\frac{1}{r_n^{N-2}} G_{\mathcal{L}_\frac{1}{4}}^n(x, x(\xi^n)) < c_{97}, \quad \forall x \in \Omega_n \cap \partial B_{r_n}(\xi^n),$$

and by Lemma 5.20 there exists $r_0 = r_0(\Omega) > 0$ such that for any $r_n \leq \frac{r_0}{4}$

$$\frac{1}{r_n^{N-2}} G_{\mathcal{L}_\frac{1}{4}}^{\Omega_n}(x, x(\xi^n)) \leq c_{98} \omega_{\Omega_n}^x(\partial\Omega_n \cap B_{r_n}(\xi^n)), \quad \forall x \in \Omega_n \cap \partial B_{r_n}(\xi^n).$$

Since if $|x - y| > \varepsilon > 0$ there holds

$$G_{\mathcal{L}_\frac{1}{4}}^{\Omega_n}(x, y) \approx c_{99}(\varepsilon, \Omega_n) \text{dist}(x, \partial\Omega_n) \text{dist}(y, \partial\Omega_n).$$

Thus we have by maximum principle and properties of Green function

$$\frac{1}{r_n^{N-2}} G_{\mathcal{L}_\frac{1}{4}}^{\Omega_n}(x, x(\xi^n)) \leq c_{100} \omega_{\Omega_n}^x(\partial\Omega_n \cap B_{r_n}(\xi^n)), \quad \forall x \in \Omega_n \setminus B_{r_n}(\xi^n). \quad (5.29)$$

By [3, Lemma 2.8] there exists $\beta_0 = \beta_0(\Omega, \varepsilon) > 0$ such that the function

$$h_1(x) = d^{\frac{1}{2}}(x) (-\log d(x)) \left(1 + (-\log d(x))^{-\varepsilon}\right),$$

is a supersolution in Ω_{β_0} and the function

$$h_2(x) = d^{\frac{1}{2}}(x) (-\log d(x)) \left(1 - (-\log d(x))^{-\varepsilon}\right),$$

is a subsolution in Ω_{β_0} . Set

$$c_{101} = \frac{1 - (-\log d(\xi_n))^{-\varepsilon}}{1 + (-\log d(\xi_n))^{-\varepsilon}}$$

and

$$H(x) = h_2(x) - c_{101}h_1(x).$$

Let $n_0 \in \mathbb{N}$ such that $r_n \leq \frac{\beta_0}{4}$, $\forall n \geq n_0$. then the function $H(x)$ is a nonnegative subsolution in $\Omega_n \setminus \Omega'_{\beta_0}$, and $H(x) = 0$, $\forall x \in \partial\Omega_n$. By (5.28) we can choose $n_1 \in \mathbb{N}$ such that

$$G_{\mathcal{L}_{\frac{1}{4}}}(x_0, x) \geq \beta_0^{\frac{1}{2}}, \quad \forall x \in \partial\Omega'_{\beta_0}.$$

Thus we can find a constant $c_{102} = c_{102}(\beta_0) > 0$ such that

$$c_{102}H(x) \leq G_{\mathcal{L}_{\frac{1}{4}}}(x_0, x), \quad \forall x \in \partial\Omega'_{\beta_0}.$$

Since H vanishes on $\partial\Omega_n$ it follows by the by maximum principle that

$$c_{102}H(x) \leq G_{\mathcal{L}_{\frac{1}{4}}}(x_0, x), \quad \forall x \in \overline{\Omega}_n \setminus \Omega'_{\beta_0}. \quad (5.30)$$

But

$$H(x(\xi^n)) \geq c_{103}(\beta_0) \geq c_{104}(\Omega, N)r_n^{N+\frac{1}{2}-2}(-\log r_n)^{1-\varepsilon}$$

and the result follows by the above inequality and inequalities (5.30) and (5.29). \square

Lemma 5.22. *Let $\kappa < \frac{1}{4}$, $\varepsilon \in (0, \sqrt{1-4\kappa})$ and $x_0 \in \Omega_1$. Let $\{\xi^n\}$ be a sequence of points in Ω converging to $a \in \partial\Omega$. Then there exists $n_0 = n_0(\varepsilon, \Omega) \in \mathbb{N}$ such that*

$$\omega_{\Omega_n}^{x_0}(B_{d(\xi^n)}(\xi^n) \cap \partial\Omega'_n) \geq c_{105}(\Omega, N, \kappa, \varepsilon)d(\xi^n)^{N+\frac{\alpha_-}{2}+\varepsilon-2}, \quad \forall n \geq n_0,$$

where Ω_n is defined by (5.19)

Proof. The proof is same as in Lemma 5.21. The only difference is that we use $d^{\alpha_-}(1-d^\varepsilon)$ and the supersolution $d^{\alpha_-}(1+d^\varepsilon)$ as a subsolution. \square

Proof of Theorem 5.17. Step 1: if

$$\limsup_{x \in \Omega, x \rightarrow a} (d(x))^{N+\frac{\alpha_+}{2}-2}u(x) < \infty, \quad (5.31)$$

then $a \in \mathcal{R}_u$. Thus we have to prove that there exists $r_0 > 0$ such that $u \in L^q_{\phi_\kappa}(\Omega \cap B_{r_0}(a))$. By (5.31) there exists $r_1 > 0$ such that

$$\sup_{x \in \Omega \cap B_{r_1}(a)} d^{N+\frac{\alpha_+}{2}-2}(x)u(x) = M < \infty.$$

Let U be a smooth open domain such that

$$\Omega \cap B_{\frac{r_1}{2}}(a) \subset U \subset \Omega \cap B_{r_1}(a),$$

and

$$\overline{U} \cap \partial\Omega \subset \partial\Omega \cap B_{r_1}(a).$$

For $\beta > 0$, set

$$d_U(x) = \text{dist}(x, \partial U) \quad \forall x \in U, \quad U_\beta = \{x \in U : d_U(x) > \beta\}, \quad V_\beta = U \setminus U_\beta.$$

Let $\beta_0 > 0$ be small enough such that $d_U \in C^2(\overline{U}_{\beta_0})$. Let $0 < \beta < \beta_0$ and $\zeta(x) = d_U(x) - \beta$. Then u satisfies

$$\int_{\partial V_\beta} u dS = \int_{V_\beta \setminus V_{\beta_0}} (u \mathcal{L}_\kappa \zeta + u^q \zeta) dx - \int_{\partial V_{\beta_0}} \frac{\partial u}{\partial \mathbf{n}} \zeta dS.$$

Now

$$\left| \int_{\partial V_{\beta_0}} \frac{\partial u}{\partial \mathbf{n}} \zeta dS \right| \leq c_{106}(\beta_0 - \beta),$$

where c_{106} depends on $q, \kappa, \Omega, \beta_0$,

$$\int_{V_\beta \setminus V_{\beta_0}} u \mathcal{L}_\kappa \zeta dx \leq - \int_{V_\beta \setminus V_{\beta_0}} u \Delta \zeta dx \leq c_{107} \int_{V_\beta \setminus V_{\beta_0}} u dx.$$

and by (5.31)

$$u^{q-1}(x) \leq c_{108}(d(x))^{-(q-1)(N+\frac{\alpha_+}{2}-2)} \leq c_{108}(d_U(x))^{-(q-1)(N+\frac{\alpha_+}{2}-2)} \quad \forall x \in U.$$

Combining the above inequalities, we derive

$$\int_{\partial V_\beta} u dS \leq c_{109} \left(\int_\beta^{\beta_0} (\sigma^{1-(q-1)(N+\frac{\alpha_+}{2}-2)} + 1) \int_{\partial V_\sigma} u(x) dS d\sigma + 1 \right).$$

Multiplying the above inequality by $\beta^{\frac{\alpha_+}{2}}$ we get

$$\int_{\partial V_\beta} u d_U^{\frac{\alpha_+}{2}} dS \leq c_{109} \left(\int_\beta^{\beta_0} (\sigma^{1-(q-1)(N+\frac{\alpha_+}{2}-2)} + 1) \int_{\partial V_\sigma} d_U^{\frac{\alpha_+}{2}}(x) u(x) dS d\sigma + 1 \right).$$

Set

$$U(\sigma) = \int_{\partial V_\sigma} d_U^{\frac{\alpha_+}{2}}(x) u(x) dS,$$

Then we have

$$U(\beta) \leq c_{110} \left(\int_\beta^{\beta_0} (\sigma^{1-(q-1)(N+\frac{\alpha_+}{2}-2)} + 1) U(\sigma) d\sigma + 1 \right), \quad (5.32)$$

Set

$$W(\beta) = \int_\beta^{\beta_0} (\sigma^{1-(q-1)(N+\frac{\alpha_+}{2}-2)} + 1) U(\sigma) d\sigma + 1,$$

then

$$W'(\beta) = -(\beta^{1-(p-1)(N+\frac{\alpha_+}{2}-2)} + 1) U(\beta) = -h(\beta) U(\beta).$$

Thus inequality (5.32) becomes

$$-W'(\beta) \leq c_{110} h(\beta) W(\beta) \iff (H(\beta) W(\beta))' \geq 0,$$

where

$$H(\beta) = e^{-c_{110} \int_\beta^{\beta_0} h(s) ds}.$$

Thus we have

$$W(\beta) \leq \frac{1}{H(\beta)} W(\beta_0), \quad \forall 0 < \beta < \beta_0.$$

But

$$\frac{1}{H(\beta)} = e^{c_{110} \int_{\beta}^{\beta_0} h(s) ds} = e^{c_{110} \int_{\beta}^{\beta_0} \sigma^{1-(q-1)(N+\frac{\alpha_+}{2}-2)+1} ds} < \infty$$

if and only if

$$2 - (q-1)(N + \frac{\alpha_+}{2} - 2) > 0 \iff q < q_c.$$

Thus we have proved that

$$\int_U u^q (d_U(x))^{\frac{\alpha_+}{2}} dx < \infty,$$

which implies the existence of a $r_2 > 0$ such that

$$\int_{\Omega \cap B_{r_2}(a)} u^q (d(x))^{\frac{\alpha_+}{2}} dx < \infty,$$

i.e. $a \in \mathcal{R}_u$, which is the claim.

Step 2. Since $a \in \mathcal{S}_u$ the previous statement implies that there exists a sequence $\{\xi^n\} \subset \Omega$ such that

$$\xi^n \rightarrow a \quad \text{and} \quad \limsup_{n \rightarrow \infty} (d(\xi^n))^{N+\frac{\alpha_+}{2}-2} u(\xi^n) = \infty. \quad (5.33)$$

By Lemma 5.19, there exists a constant c_l such that

$$u(x) \leq c_l u(y), \quad \forall x, y \in B_{\frac{r_n}{2}}(\xi^n), \quad r_n = d(\xi^n). \quad (5.34)$$

Put $V_n := B_{\frac{r_n}{2}}(\xi^n) \cap \partial\Omega'_{r_n}$, and, for $k > 0$, $h_{n,k} := \frac{k}{b_n} u \chi_{V_n}$.

Case 1: $\kappa = \frac{1}{4}$. By (5.34) and Lemma 5.21 there exists a constant $c_{111} > 0$ such that

$$b_n := \int_{V_n} u dS \geq c_{111} A_n r_n^{N+\frac{1}{2}-2} (-\log r_n)^{1-\varepsilon}, \quad A_n := \sup_{x \in B_{\frac{r_n}{2}}(\xi^n)} u(x).$$

Then

$$\int_{\partial\Omega'_n} h_{n,k} dS = k, \quad h_{n,k} \leq \frac{k}{c_2} r_n^{2-\frac{\alpha_+}{2}N} \chi_{V_n}, \quad \forall n \geq n_0. \quad (5.35)$$

By (5.33),

$$b_n \rightarrow \infty, \quad r_n \rightarrow 0. \quad (5.36)$$

Hence, for every $k > 0$ there exists n_k such that

$$u \geq h_{n,k} \quad \text{on} \quad \partial\Omega'_n \quad \forall n \geq n_k. \quad (5.37)$$

Let $w_{n,k}$ be defined as in Lemma 5.18 with h_n replaced by $h_{n,k}$. By (5.35) and (5.36), the sequence $\{h_{n,k}\}_{n=1}^{\infty}$ satisfies (5.20) for every fixed $k > 0$. Therefore by Lemma 5.18

$$\lim_{n \rightarrow \infty} w_{n,k} = u_{k\delta_a} \quad \text{locally uniformly in } \Omega.$$

By (5.37), $u \geq w_{n,k}$ in $x \in \Omega : d(x) > r_n$. Hence $u \geq u_{k\delta_a}$ for every $k > 0$. The proof in the case $0 < \kappa < \frac{1}{4}$ is similar but for the definition of b_n . \square

As a consequence we provide a full classification of positive solution of (4.1) with a boundary isolated singularity.

Theorem 5.23. Assume $1 < q < q_c$ and $u \in C(\overline{\Omega} \setminus \{0\})$ is a positive solution of (4.1) in Ω which vanishes on $\partial\Omega \setminus \{0\}$. Then the following alternative holds

(i) Either there exists $k \geq 0$ such that

$$\lim_{\substack{x \rightarrow 0, x \in \Omega \\ |x|^{-1} \rightarrow \sigma}} |x|^{N + \frac{\alpha_+}{2} - 2} u(x) = k\psi_1(\sigma) \quad (5.38)$$

and u solves

$$\begin{aligned} -\Delta u - \frac{\kappa}{d^2} u + u^q &= 0 && \text{in } \Omega \\ u &= k\delta_0 && \text{in } \partial\Omega. \end{aligned} \quad (5.39)$$

(ii) Or

$$\lim_{\substack{x \rightarrow 0, x \in \Omega \\ |x|^{-1} \rightarrow \sigma}} |x|^{\frac{2}{q-1}} u(x) = \omega_\kappa(\sigma) \quad (5.40)$$

locally uniformly on S_+^{N-1} .

The result is a consequence of the following result

Lemma 5.24. Assume $1 < q < q_c$, $a \in \partial\Omega$ and $F_\epsilon(a) = \partial\Omega \cap \overline{B_\epsilon(a)}$. Then

$$\lim_{\epsilon \rightarrow 0} U_{F_\epsilon(a)} = u_{\infty, a}. \quad (5.41)$$

Proof. Without loss of generality, we can assume $a = 0$. Clearly, $U_{\{0\}} := \lim_{\epsilon \rightarrow 0} U_{F_\epsilon(0)}$ is a solution of (5.1) which satisfies

$$\lim_{x \rightarrow \xi} \frac{U_{\{0\}}}{W(x)} = 0 \quad \forall \xi \in \partial\Omega \setminus \{0\}$$

locally uniformly on $\partial\Omega \setminus \{0\}$. By (6.20) it verifies

$$U_{\{0\}}(x) \leq c|x|^{-\frac{2}{q-1}} \left(\frac{d(x)}{|x|} \right)^{\frac{\alpha_+}{2}}. \quad (5.42)$$

□

Using (4.24) we see that there exists $c = c_{112}(N, \kappa, q) > 1$ such that

$$U_{\{0\}}(x) \leq cu_{\infty, 0}(x) \quad \forall x \in \Omega. \quad (5.43)$$

Assume $U_{\{0\}} \neq u_{\infty, 0}$, thus $U_{\{0\}}(x) > u_{\infty, 0}(x)$ for all $x \in \Omega$ and put $\tilde{u} = u_{\infty, 0} - \frac{1}{2c}(U_{\{0\}} - u_{\infty, 0})$. By convexity \tilde{u} is a supersolution of (5.1) which is smaller than $u_{\infty, 0}$. Now $\frac{c+1}{2c}u_{\infty, 0}$ is a subsolution, thus there exists a solution \underline{u} of (5.1) in Ω which satisfies

$$\frac{c+1}{2c}u_{\infty, 0}(x) \leq \underline{u}(x) \leq \tilde{u}(x) < u_{\infty, 0}(x) \quad \forall x \in \Omega. \quad (5.44)$$

This implies that $Tr_{\partial\Omega}(\underline{u}) = (\{0\}, 0)$, and by Theorem 5.17, $\underline{u} \geq u_{\infty, 0}$, which is a contradiction.

Proof of Theorem 5.23 Assume $a = 0$ without loss of generality. If $a \in \mathcal{S}_u$, then for any $\epsilon > 0$, $u \leq U_{F_\epsilon(0)}$ which is a maximal solution which vanishes on $\partial\Omega \setminus F_\epsilon(0)$. Thus, using (5.41)

$$u \leq \lim_{\epsilon \rightarrow 0} U_{F_\epsilon(0)} = U_{\{0\}} = u_{\infty, 0}.$$

If $0 \in \mathcal{R}_u$, this implies that $Tr_{\partial\Omega}(u) = (\emptyset, k\delta_0)$ for some $k \geq 0$ and we conclude with Corollary 4.4. □

The next result can be proven by using the same approximation methods as in [19, Th 9.6].

Theorem 5.25. . Assume $S \subset \partial\Omega$ is closed and ν is a positive Radon measure on $\mathcal{R} = \partial\Omega \setminus S$. Then there exists a positive solution of (4.1) in Ω with boundary trace (S, μ) .

6 Appendix I: barriers and a priori estimates

6.1 Barriers

Following a localization principle introduced in [19] we the following lemma is at the core of the a priori estimates construction

Proposition 6.1. Let $\Omega \subset \mathbb{R}^N$ be a C^2 domain $0 < \kappa \leq \frac{1}{4}$ and $p > 1$. Then there exists $R_0 > 0$ such that for any $z \in \partial\Omega$ and $0 < R \leq R_0$, there exists a super solution $f := f_{R,z}$ of (4.1) in $\Omega \cap B_R(z)$ such that $f \in C(\overline{\Omega} \cap B_R(z))$, $f(x) \rightarrow \infty$ when $\text{dist}(x, K) \rightarrow 0$, for any compact subset $K \subset \Omega \cap \partial B_R(z)$ and which vanishes on $\partial\Omega \cap B_R(z)$, and more precisely

$$f(x) = \begin{cases} c_{\beta,\gamma,\kappa,q}(R^2 - |x - z|^2)^{-\beta} d^\gamma(x) & \forall \gamma \in (\frac{\alpha_-}{2}, \frac{\alpha_+}{2}) \quad \text{if } 0 < \kappa < \frac{1}{4} \\ c_{\beta,\gamma,q}(R^2 - |x - z|^2)^{-\beta} \sqrt{d(x)} \sqrt{\ln\left(\frac{\text{diam}(\Omega)}{d(x)}\right)} & \text{if } \kappa = \frac{1}{4} \end{cases} \quad (6.1)$$

for $\beta \geq \max\{\frac{2}{q-1} + \gamma, \frac{N-2}{2}, 1\}$.

Proof. We assume $z = 0$

Step 1: $\kappa < \frac{1}{4}$. Set $f(x) = \Lambda(R^2 - |x|^2)^{-\beta} (d(x))^\gamma$ where $\beta, \gamma > 0$ to be chosen later on. Then, with $r = |x|$,

$$\begin{aligned} & \Lambda^{-1} \mathcal{L}_\kappa f \\ &= -(R^2 - r^2)^{-\beta} (\Delta d^\gamma + \kappa d^{\gamma-2}) - d^\gamma \Delta(R^2 - r^2)^{-\beta} - 2\nabla(R^2 - r^2)^{-\beta} \nabla d^\gamma \end{aligned}$$

Since $\Delta d(x) = (N-1)H_d$ where H_d is the mean curvature of the foliated set $\Sigma_d := \{x \in \Omega : d(x) = d\}$ and $|\nabla d|^2 = 1$,

$$\begin{aligned} \Delta d^\gamma &= (N-1)\gamma H_d d^{\gamma-1} + \gamma(\gamma-1)d^{\gamma-2} \\ \Delta d^\gamma + \kappa d^{\gamma-2} &= (N-1)\gamma H_d d^{\gamma-1} + (\gamma(\gamma-1) + \kappa) d^{\gamma-2} \\ \nabla d^\gamma &= \gamma \phi_1^{\gamma-1} \nabla d, \end{aligned}$$

$$\nabla(R^2 - r^2)^{-\beta} = 2\beta(R^2 - r^2)^{-\beta-1} x,$$

thus

$$\begin{aligned} \nabla(R^2 - r^2)^{-\beta} \nabla d^\gamma &= 2\beta\gamma d^{\gamma-1} (R^2 - r^2)^{-\beta-1} x \nabla d \\ \Delta(R^2 - r^2)^{-\beta} &= 2N\beta(R^2 - r^2)^{-\beta-1} + 4\beta(\beta+1)(R^2 - r^2)^{-\beta-2} r^2 \\ &= 2\beta(R^2 - r^2)^{-\beta-2} (NR^2 + (2\beta+2-N)r^2) \end{aligned}$$

Then

$$\begin{aligned} \Lambda^{-1} \mathcal{L}_\kappa f &= -(R^2 - r^2)^{-\beta-2} d^{\gamma-2} \left[(R^2 - r^2)^2 ((N-1)\gamma H_d d + \gamma(\gamma-1) + \kappa) \right. \\ &\quad \left. + 2\beta d^2 (NR^2 + (2\beta+2-N)r^2) + 4\beta\gamma d(R^2 - r^2)x\nabla d \right] \end{aligned}$$

Therefore

$$\begin{aligned} \mathcal{L}_\kappa f + f^q &= \Lambda(R^2 - r^2)^{-\beta-2} d^{\gamma-2} \left[\Lambda^{q-1} (R^2 - r^2)^{-(q-1)\beta+2} d^{(q-1)\gamma+2} \right. \\ &\quad \left. -(R^2 - r^2)^2 ((N-1)\gamma H_d d + \gamma(\gamma-1) + \kappa) \right. \\ &\quad \left. - 2\beta d^2 (NR^2 + (2\beta+2-N)r^2) + 4\beta\gamma d(R^2 - r^2)x\nabla d \right] \end{aligned} \quad (6.2)$$

If we fix $\beta \geq \max\{\frac{2}{q-1} + \gamma, \frac{N-2}{2}, 1\}$, there holds

$$2\beta d^2 (NR^2 + (2\beta+2-N)r^2) + 4\beta\gamma d(R^2 - r^2)x\nabla d \leq 4d^2\beta(\beta+1)NR^2 + 4\beta dR(R^2 - r^2)$$

We choose $\frac{\alpha_-}{2} < \gamma < \frac{\alpha_+}{2}$ so that $\gamma(\gamma-1) + \kappa < 0$. There exist $\delta_0, \epsilon_0 > 0$ such that

$$(N-1)\gamma H_d d + \gamma(\gamma-1) + \kappa < -\epsilon_0 < -1$$

provided $d(x) \leq \delta_0$. We set

$$A = \left\{ x \in \Omega \cap B_R : d(x) \leq \frac{\epsilon_0(R^2 - r^2)}{16\beta R} \right\} \quad \text{and} \quad B := A \cap \left\{ x \in \Omega \cap B_R : d(x) \leq \delta_0 \right\}$$

Then, if $x \in B$, there holds

$$\begin{aligned} -(R^2 - r^2)^2 ((N-1)\gamma H_d d + \gamma(\gamma-1) + \kappa) - 2\beta d^2 (NR^2 + (2\beta+2-N)r^2) \\ + 4\beta\gamma d(R^2 - r^2)x\nabla d \geq \frac{(R^2 - r^2)^2 \epsilon_0}{2} \end{aligned}$$

Finally, assume $x \in A^c \cap \left\{ x \in \Omega \cap B_R : d(x) \leq \delta_0 \right\}$ and thus

$$d \geq c_1 \frac{R^2 - r^2}{R}$$

In order to have

$$\begin{aligned} (i) \quad & \Lambda^{q-1} (R^2 - r^2)^{2-(q-1)\beta} d^{(q-1)\gamma+2} \geq d^2 R^2 \\ (ii) \quad & \Lambda^{q-1} (R^2 - r^2)^{2-(q-1)\beta} d^{(q-1)\gamma+2} \geq dR(R^2 - r^2) \end{aligned} \quad (6.3)$$

or equivalently

$$\begin{aligned} (i) \quad & \iff \Lambda^{\frac{1}{\gamma}} d \geq (R^2 - r^2)^{\frac{\beta}{\gamma}} \\ (ii) \quad & \iff \Lambda^{\frac{q-1}{(q-1)\gamma+1}} d \geq R^{\frac{1}{(q-1)\gamma+1}} (R^2 - r^2)^{\frac{(q-1)\beta-1}{(q-1)\gamma+1}} \end{aligned} \quad (6.4)$$

it is sufficient to have, for (i)

$$c_1 \Lambda^{\frac{1}{\gamma}} \frac{R^2 - r^2}{R} \geq (R^2 - r^2)^{\frac{\beta}{\gamma}} \quad \forall r \in (0, R) \iff \Lambda \geq c_2 R^{2\beta-\gamma} \quad (6.5)$$

and for (ii)

$$\begin{aligned} c_1 \Lambda \frac{R^{q-1} R^2 - r^2}{R} &\geq R^{\frac{1}{(q-1)\gamma+1}} (R^2 - r^2)^{\frac{(q-1)\beta-1}{(q-1)\gamma+1}} \quad \forall r \in (0, R) \\ \iff \Lambda &\geq c_2 R^{2\beta-\gamma-\frac{2}{q-1}} \end{aligned} \quad (6.6)$$

where $c_2 = c_2(N, \gamma, \beta) > 0$ since $\beta > \gamma + \frac{2}{q-1}$.

At end, in the set $C := \{x \in \Omega : d(x) \geq \delta_0\}$, it suffices that

$$\Lambda \geq c_3 \max \left\{ R^{2\beta}, R^{2\beta-\frac{1}{q-1}} \right\} \quad (6.7)$$

for some $c_3 = c_3(N, \gamma, \beta, \max |H_d|, \delta_0) > 0$ in order to insure

$$\begin{aligned} (i) \quad &\Lambda^{q-1} (R^2 - r^2)^{-(q-1)\beta+2} d^{(q-1)\gamma+2} \geq (R^2 - r^2)^2 (N-1)\gamma |H_d| d \\ (ii) \quad &\Lambda^{q-1} (R^2 - r^2)^{-(q-1)\beta+2} d^{(q-1)\gamma+2} \geq 4d^2 \beta(\beta+1) N R^2 \\ (iii) \quad &\Lambda^{q-1} (R^2 - r^2)^{-(q-1)\beta+2} d^{(q-1)\gamma+2} \geq 4\beta d R (R^2 - r^2). \end{aligned} \quad (6.8)$$

Noticing that $2\beta > 2\beta - \frac{1}{q-1}$, $2\beta - \gamma > 2\beta - \gamma - \frac{1}{q-1}$, we conclude that there exists a constant $c_4 = c_4(N, \gamma, \beta, \max |H_d|, \delta_0) > 0$ such that if

$$\Lambda \geq c_4 \max \left\{ R^{2\beta}, R^{2\beta-\gamma-\frac{1}{q-1}} \right\} \quad (6.9)$$

there holds

$$\mathcal{L}_\kappa(f) + f^q \geq 0 \quad \text{in } \Omega. \quad (6.10)$$

Step 2: $\kappa = \frac{1}{4}$. Set $f(x) = \Lambda (R^2 - r^2)^{-\beta} \sqrt{d} (\ln \frac{eR}{d})^{\frac{1}{2}}$ for some Λ, β to be fixed. Then

$$\begin{aligned} \Delta \sqrt{d} (\ln \frac{eR}{d})^{\frac{1}{2}} &= \frac{1}{\sqrt{d}} \left(\frac{1}{2} (\ln \frac{eR}{d})^{\frac{1}{2}} - \frac{1}{2} (\ln \frac{eR}{d})^{-\frac{1}{2}} \right) \Delta d \\ &\quad + \frac{1}{d^{\frac{3}{2}}} \left(-\frac{1}{4} (\ln \frac{eR}{d})^{\frac{1}{2}} - \frac{1}{4} (\ln \frac{eR}{d})^{-\frac{3}{2}} \right) \\ &= \frac{N-1}{\sqrt{d}} \left(\frac{1}{2} (\ln \frac{eR}{d})^{\frac{1}{2}} - \frac{1}{2} (\ln \frac{eR}{d})^{-\frac{1}{2}} \right) H_d \\ &\quad + \frac{1}{d^{\frac{3}{2}}} \left(-\frac{1}{4} (\ln \frac{eR}{d})^{\frac{1}{2}} - \frac{1}{4} (\ln \frac{eR}{d})^{-\frac{3}{2}} \right) \end{aligned}$$

Thus

$$\begin{aligned} \Delta \sqrt{d} (\ln \frac{eR}{d})^{\frac{1}{2}} + \frac{\kappa}{d^2} \sqrt{d} (\ln \frac{eR}{d})^{\frac{1}{2}} &= \frac{N-1}{\sqrt{d}} \left(\frac{1}{2} (\ln \frac{eR}{d})^{\frac{1}{2}} - \frac{1}{2} (\ln \frac{eR}{d})^{-\frac{1}{2}} \right) H_d - \frac{1}{4d^{\frac{3}{2}}} (\ln \frac{eR}{d})^{-\frac{3}{2}} \\ &= \frac{1}{d^{\frac{3}{2}}} (\ln \frac{eR}{d})^{-\frac{3}{2}} \left[(N-1) d H_d \left(\frac{1}{2} (\ln \frac{eR}{d})^2 - \frac{1}{2} (\ln \frac{eR}{d}) \right) - \frac{1}{4} \right] \end{aligned}$$

Further

$$\nabla (R^2 - r^2)^{-\beta} \nabla \sqrt{d} (\ln \frac{eR}{d})^{\frac{1}{2}} = \frac{\beta (R^2 - r^2)^{-\beta-1} (\ln \frac{eR}{d})^{-\frac{1}{2}}}{\sqrt{d}} \left((\ln \frac{eR}{d}) - 1 \right) x \nabla d.$$

Therefore

$$\begin{aligned} \Lambda^{-1} \mathcal{L}_\kappa f &= -(R^2 - r^2)^{-\beta-2} d^{-\frac{3}{2}} (\ln \frac{eR}{d})^{-\frac{3}{2}} \\ &\quad \left[(R^2 - r^2)^2 \left[(N-1) d H_d \left(\frac{1}{2} (\ln \frac{eR}{d})^2 - \frac{1}{2} (\ln \frac{eR}{d}) \right) - \frac{1}{4} \right] \right. \\ &\quad \left. + 2\beta (R^2 - r^2) d \left[(\ln \frac{eR}{d})^2 - (\ln \frac{eR}{d}) \right] x \nabla d + 2\beta d^2 (\ln \frac{eR}{d})^2 \left[N R^2 + (2\beta + 2 - N) r^2 \right] \right] \end{aligned}$$

Finally

$$\begin{aligned} \mathcal{L}_\kappa f + f^q &= \Lambda(R^2 - r^2)^{-\beta-2} d^{-\frac{3}{2}} (\ln \frac{eR}{d})^{-\frac{3}{2}} \left[\Lambda^{q-1} (R^2 - r^2)^{(1-q)\beta+2} d^{\frac{q+3}{2}} (\ln \frac{eR}{d})^{\frac{1}{2}(q-1)+2} \right. \\ &\quad \left. - (R^2 - r^2)^2 \left[(N-1)dH_d \left(\frac{1}{2} (\ln \frac{eR}{d})^2 - \frac{1}{2} (\ln \frac{eR}{d}) \right) - \frac{1}{4} \right] \right. \\ &\quad \left. - 2\beta(R^2 - r^2)d \left[(\ln \frac{eR}{d})^2 - (\ln \frac{eR}{d}) \right] x \nabla d - 2\beta d^2 (\ln \frac{eR}{d})^2 [NR^2 + (2\beta + 2 - N)r^2] \right]. \end{aligned} \quad (6.11)$$

Notice that $\frac{eR}{d} \geq e$ thus $-\frac{1}{2} \leq (\ln \frac{eR}{d})^2 - (\ln \frac{eR}{d}) \leq (\ln \frac{eR}{d})^2$. If β is large enough, as in Step 1, there holds

$$\begin{aligned} &|2\beta(R^2 - r^2)d \left[(\ln \frac{eR}{d})^2 - (\ln \frac{eR}{d}) \right] x \cdot \nabla d + 2\beta d^2 (\ln \frac{eR}{d})^2 [NR^2 + (2\beta + 2 - N)r^2]| \\ &\leq 4N\beta(\beta + 1) (\ln \frac{R}{d})^2 ((R^2 - r^2)dR + d^2 R^2). \end{aligned}$$

There exists $\delta_0 > 0$ such that

$$(N-1)dH_d \left(\frac{1}{2} (\ln \frac{eR}{d})^2 - \frac{1}{2} (\ln \frac{eR}{d}) \right) - \frac{1}{4} \leq -\frac{1}{8} < -1$$

if $d(x) \leq \delta_0$. If we define A, B by

$$A = \left\{ x \in \Omega \cap B_R : d(x) \leq \frac{\epsilon_0(R^2 - r^2)}{16\beta R (\ln \frac{eR}{d})^2} \right\} \quad \text{and} \quad B := A \cap \left\{ x \in \Omega \cap B_R : d(x) \leq \delta_0 \right\}$$

there holds if $x \in B$

$$\begin{aligned} &-2\beta(R^2 - r^2)d \left[(\ln \frac{eR}{d})^2 - (\ln \frac{eR}{d}) \right] x \cdot \nabla d - 2\beta d^2 (\ln \frac{eR}{d})^2 [NR^2 + (2\beta + 2 - N)r^2] \\ &\quad - (R^2 - r^2)^2 \left[(N-1)dH_d \left(\frac{1}{2} (\ln \frac{eR}{d})^2 - \frac{1}{2} (\ln \frac{eR}{d}) \right) - \frac{1}{4} \right] \geq \frac{(R^2 - r^2)^2}{16}. \end{aligned}$$

If $x \in A^c \cap \{x \in \Omega \cap \Omega : d(x) \leq \delta_0\}$, then

$$d(x) \geq c_1 \frac{R^2 - r^2}{R (\ln \frac{eR}{d})^2}. \quad (6.12)$$

In order to have

$$\begin{aligned} (i) \quad &\Lambda^{q-1} (R^2 - r^2)^{(1-q)\beta+2} d^{\frac{q+3}{2}} (\ln \frac{eR}{d})^{\frac{q+3}{2}} \geq (\ln \frac{eR}{d})^2 (R^2 - r^2) dR \\ (ii) \quad &\Lambda^{q-1} (R^2 - r^2)^{(1-q)\beta+2} d^{\frac{q+3}{2}} (\ln \frac{eR}{d})^{\frac{q+3}{2}} \geq (\ln \frac{eR}{d})^2 d^2 R^2 \end{aligned} \quad (6.13)$$

or equivalently

$$\begin{aligned} (i) \quad &\Lambda^{\frac{2q-2}{q+1}} d (\ln \frac{eR}{d})^{\frac{q-1}{q+1}} \geq (R^2 - r^2)^{\frac{2(q-1)\beta-2}{q+1}} R^{\frac{2}{q+1}} \\ (ii) \quad &\Lambda^2 d \ln \frac{eR}{d} \geq R^{\frac{4}{q-1}} (R^2 - r^2)^{2\beta - \frac{4}{q-1}} \end{aligned} \quad (6.14)$$

Up to taking c_1 small enough, (6.12) is fulfilled if

$$\frac{eR}{d} \leq \frac{R^2}{R^2 - r^2} \left(\ln \left(\frac{R^2}{R^2 - r^2} \right) \right)^2 \iff d \geq \frac{e(R^2 - r^2)}{R} \left(\ln \left(\frac{R^2}{R^2 - r^2} \right) \right)^{-2}. \quad (6.15)$$

Inequality (6.13)-(i) will be insured if

$$\Lambda^{\frac{2q-2}{q+1}} \geq \frac{1}{\epsilon} (R^2 - r^2)^{2\frac{(q-1)\beta-1}{q+1}-1} R^{\frac{2}{q+1}+1} (\ln(\frac{R^2}{R^2-r^2}))^{\frac{2}{q+1}}$$

which holds if, for any $\epsilon > 0$, we have for any $r \in (0, R)$

$$\Lambda^{\frac{2q-2}{q+1}} \geq C_\epsilon (R^2 - r^2)^{2\frac{(q-1)\beta-1}{q+1}-1} R^{\frac{2}{q+1}+1} \left(\frac{R^2}{R^2 - r^2} \right)^\epsilon.$$

A sufficient condition for such a task is, with the help of (6.15),

$$\Lambda \geq c_3 R^{3\beta - \frac{2}{q-1}}. \quad (6.16)$$

As for (6.13)-(ii), it will be insured if

$$\Lambda \geq c_4 R^{2\beta - \frac{2}{q-1} - \frac{1}{2}} \quad (6.17)$$

Thus, if

$$\Lambda \geq c_5 \max\{R^{2\beta - \frac{2}{q-1} - \frac{1}{2}}, R^{3\beta - \frac{2}{q-1}}\} \quad (6.18)$$

for some $c_5 > 0 = c_5(N, \gamma, \beta, \delta_0, |H_d|)$, the function f satisfies (6.10). \square

6.2 A priori estimates

By the Keller-Osserman estimate, it is clear that any solution u of 4.1 in Ω satisfies

$$u(x) \leq C(q, \Omega, N) d^{-\frac{2}{q-1}}(x), \quad \forall x \in \Omega. \quad (6.19)$$

This estimate is also a consequence of the following result [3, Prop 3.4]

Proposition 6.2. *Let ϕ_* be the first positive eigenfunction of $-\Delta$ in $H_0^1(\Omega)$. For $q > 1$, there exists $\gamma > 0$ and $\epsilon_0 > 0$ such that for any $0 \leq \epsilon \leq \epsilon_0$ the function $h_{+\epsilon} = \gamma(\phi_* - \epsilon)^{-\frac{2}{q-1}}$ is a supersolution of 4.1 in $\Omega_{\epsilon, \phi_*} := \{x \in \Omega : \phi_*(x) > \epsilon\}$.*

We recall here that

$$W(x) = \begin{cases} d^{\frac{\alpha_-}{2}}(x) & \text{if } \kappa < \frac{1}{4} \\ d^{\frac{1}{2}}(x) |\log d(x)| & \text{if } \kappa = \frac{1}{4} \end{cases}$$

Proposition 6.3. *Let Ω be a bounded open domain uniformly of class C^2 and let F be a compact subset of the boundary. Let u be a nonnegative solution of 5.1 in Ω such that*

$$\lim_{x \in \Omega, x \rightarrow \xi} \frac{u(x)}{W(x)} = 0, \quad \forall \xi \in \partial\Omega \setminus F,$$

locally uniformly in $\partial\Omega \setminus F$. Then there exists a constant C depending only on q, κ and Ω such that,

$$|u(x)| \leq C d^{\frac{\alpha_+}{2}}(x) (\text{dist}(x, F))^{-\frac{2}{q-1} - \frac{\alpha_+}{2}}, \quad \forall x \in \Omega, \quad (6.20)$$

$$\left| \frac{u(x)}{d^{\frac{\alpha_+}{2}}(x)} - \frac{u(y)}{d^{\frac{\alpha_+}{2}}(y)} \right| \leq C |x - y|^\beta (\text{dist}(x, F))^{-\frac{2}{q-1} - \beta - \frac{\alpha_+}{2}} \quad \forall (x, y) \in \Omega \times \Omega \quad (6.21)$$

such that $\text{dist}(x, F) \leq \text{dist}(y, F)$,

$$|\nabla u(x)| \leq C d^{\frac{\alpha_+}{2}-1}(x) (\text{dist}(x, F))^{-\frac{2}{q-1} - \frac{\alpha_+}{2}}, \quad \forall x \in \Omega. \quad (6.22)$$

Proof. The proof is based on the proof of Proposition 3.4.3 in [22]. Let $\xi \in \partial\Omega \setminus F$ and put $d_F(\xi) = \frac{1}{2}\text{dist}(\xi, F)$. Denote by Ω^ξ the domain

$$\Omega^\xi = \{y \in \mathbb{R}^n : d_F(\xi)y \in \Omega\}.$$

If u is a positive solution of (5.1) in Ω , denote by u^ξ the function

$$u^\xi(y) = |d_F(\xi)|^{\frac{2}{q-1}} u(d_F(\xi)y), \quad \forall y \in \Omega^\xi.$$

Then,

$$-\Delta u^\xi - \kappa \frac{u}{|\text{dist}(y, \partial\Omega^\xi)|^2} + |u^\xi|^q = 0 \quad \text{in } \Omega^\xi.$$

Let R_0 be the constant in Proposition 6.1. First, we assume that

$$\text{dist}(\xi, F) \leq \frac{1}{1 + R_0}.$$

Set $r_0 = \frac{3R_0}{4}$, then the solution $W_{r_0, \xi}$ mentioned in Proposition 6.1 satisfies

$$u^\xi(y) \leq W_{r_0, \xi}(y), \quad \forall y \in B_{\frac{3R_0}{4}}(\xi) \cap \Omega^\xi.$$

Thus u^ξ is bounded in $B_{\frac{3R_0}{5}}(\xi) \cap \Omega^\xi$ by a constant $C > 0$ depending only on n, q, κ and the C^2 characteristic of Ω^ξ . As $d_F(\xi) \leq 1$ a C^2 characteristic of Ω is also a C^2 characteristic of Ω^ξ therefore the constant C can be taken to be independent of ξ . We note here that the constant $0 < R_0 < 1$ depends on C^2 characteristic of Ω .

Now we note that

$$\lim_{y \in \Omega^\xi, y \rightarrow P} \frac{u^\xi(y)}{W(x)} = 0, \quad \forall P \in \partial\Omega^\xi \cap B_{\frac{3R_0}{5}}(\xi).$$

Thus in view of the proof of Lemmas 2.11 and 2.12, by the above inequality and in view of the proof of Theorem 2.12 in [12], we have that there exists $C > 0$ depending only on n, p, κ such that

$$u^\xi(y) \leq |\text{dist}(y, \partial\Omega^\xi)|^{\frac{\alpha_+}{2}}, \quad \forall y \in B_{\frac{R_0}{2}}(\xi) \cap \Omega^\xi. \quad (6.23)$$

Hence

$$u(x) \leq d^{\frac{\alpha_+}{2}}(x) d_F(\xi)^{-\frac{2}{q-1} - \frac{\alpha_+}{2}}, \quad \forall x \in B_{d_F(\xi)\frac{R_0}{2}}(\xi) \cap \Omega.$$

Let $x \in \Omega_{\frac{R_0}{2}}$ and assume that

$$d(x) \leq \frac{R_0}{2} d_F(x).$$

Let ξ be the unique point in $\partial\Omega \setminus F$ such that $|x - \xi| = d(x)$. Then we have

$$d_F(\xi) \leq d(x) + d_F(x) \leq (1 + R_0)d_F(x) < 1$$

and

$$|u(x)| \leq C d^{\frac{\alpha_+}{2}}(x) ((1 + R_0)\text{dist}(x, F))^{-\frac{2}{q-1} - \frac{\alpha_+}{2}}.$$

If $d(x) > \frac{R_0}{4} d_F(x)$, then by (6.19) we have that

$$|u(x)| \leq C d^{-\frac{2}{q-1}}(x) \leq C d^{\frac{\alpha_+}{2}}(x) \left(\frac{R_0}{2} \text{dist}(x, F) \right)^{-\frac{2}{q-1} - \frac{\alpha_+}{2}}.$$

Thus (6.20) holds for every $x \in \Omega_{\frac{R_0}{2}}$ such that $\text{dist}(x, F) < \frac{1}{1+R_0}$.

Now we assume that $x \in \Omega_{\frac{R_0}{2}}$ and

$$\text{dist}(x, F) \geq \frac{1}{1+R_0}.$$

Let ξ be the unique point in $\partial\Omega \setminus F$ such that $|x - \xi| = d(x)$. Similarly with the proof of 6.23 we can prove that

$$u(x) \leq C d^{\frac{\alpha_+}{2}}(x) \leq d^{\frac{\alpha_+}{2}}(x) C ((1+R_0)\text{dist}(x, F))^{-\frac{2}{q-1}-\frac{\alpha_+}{2}}, \quad \forall x \in B_{\frac{R_0}{2}}(\xi) \cap \Omega.$$

Now if $x \in \Omega \setminus \Omega_{\frac{R_0}{2}}$, the proof of (6.20) follows by (6.19).

(ii) Let $x_0 \in \Omega$. Set

$$\Omega^{x_0} = \{y \in \mathbb{R}^n : d(x_0)y \in \Omega\},$$

and $d_{x_0}(y) = \text{dist}(y, \partial\Omega^{x_0})$. If $x \in B_{\frac{d(x_0)}{2}}(x_0)$ then $y = \frac{x}{d(x_0)}$ belongs to $B_{\frac{1}{2}}(y_0)$, where $y_0 = \frac{x_0}{d(x_0)}$. Also we have that $\frac{1}{2} \leq d_{x_0}(y) \leq \frac{3}{2}$ for each $y \in B_{\frac{1}{2}}(y_0)$. Set now $v(y) = u(d(x_0)y)$, $\forall y \in B_{\frac{1}{2}}(y_0)$. Then v satisfies

$$-\Delta v - \kappa \frac{u}{|d_{x_0}(y)|^2} + d^2(x_0)|v|^q = 0 \quad \text{in } B_{\frac{1}{2}}(y_0).$$

By standard elliptic estimate we have

$$\sup_{y \in B_{\frac{1}{4}}(y_0)} |\nabla v| \leq C \left(\sup_{y \in B_{\frac{1}{3}}(y_0)} |v| + \sup_{y \in B_{\frac{1}{3}}(y_0)} d^2(x_0)|v|^q \right),$$

Now since $\nabla v(y) = d(x_0)\nabla u(d(x_0)y)$, by above inequality and (6.20) we have that

$$|\nabla u(x_0)| \leq C \left(d^{\frac{\alpha_+}{2}-1}(x_0) (\text{dist}(x_0, F))^{-\frac{2}{q-1}-\frac{\alpha_+}{2}} + d^{\frac{q\alpha_+}{2}+1}(x_0) (\text{dist}(x_0, F))^{-q(\frac{2}{q-1}-\frac{\alpha_+}{2})} \right).$$

Using $\frac{2q}{q-1} = \frac{2}{q-1} + 2$ and the fact that x_0 is arbitrary the result follows. \square

Proposition 6.4. *Let $O \subset \partial\Omega$ be a relatively open subset and $F = \bar{O}$. Let U_F be defined by (5.7) be the maximal solution of (5.1) which vanishes on $\partial\Omega \setminus F$. Then for any compact set $K \subset O$, there holds*

$$\lim_{\xi \rightarrow x} (d(\xi))^{\frac{2}{q-1}} U_F(\xi) = \ell_\kappa = \left(\frac{2(q+1)}{(q-1)^2} + \kappa \right)^{\frac{1}{q-1}} \quad \text{uniformly with respect to } x \in K. \quad (6.24)$$

Proof. Step 1. We claim that for any $\epsilon > 0$ there exists $C_\epsilon, \tau_\epsilon > 0$ such that for any $z \in O$ such that $\bar{B}_{2\tau_\epsilon}(z) \subset O$, there holds

$$u(x) \leq (\epsilon + \ell_\kappa^{q-1})^{\frac{1}{q-1}} \tau^{-\frac{2}{q-1}} + C_\epsilon \quad \forall \tau \in (0, \tau_\epsilon], \forall x \in \Sigma_\tau(\bar{B}_{\tau_\epsilon}(z)). \quad (6.25)$$

We recall that $\Sigma_\tau(\bar{B}_{\tau_\epsilon}(z)) = \{x \in \Omega, x \approx (d(x), \sigma(x)), d(x) = \tau, \sigma(x) \in \bar{B}_{\tau_\epsilon}(z)\}$. Set $g(x) = \ell d^{-\frac{2}{q-1}}(x)$, then

$$\mathcal{L}_\kappa g + g^q = \frac{2(N-1)}{q-1} H_d d^{-\frac{q+1}{q-1}} + (\ell^{q-1} - \ell_\kappa^{q-1}) d^{-\frac{2q}{q-1}}, \quad (6.26)$$

where H_d is the mean curvature of Σ_d . If Ω is convex we take $\ell = \ell_\kappa$ and g is a supersolution for $d(x) \leq R_0$ for some R_0 . In the general case, we take $\ell = \ell(\epsilon) = (\epsilon + \ell_\kappa^{q-1})^{\frac{1}{q-1}}$, and $g = g_\epsilon = \ell(\epsilon)d^{-\frac{2}{q-1}}$ is a supersolution in the set Ω_{τ_ϵ} where

$$\tau_\epsilon = \max \left\{ \tau : 0 < \tau \leq \frac{R_0}{2}, \frac{2(N-1)}{q-1} \|H_\tau\|_{L^\infty(\Sigma_\tau)} + \epsilon > 0 \right\}.$$

Then $f_{2\tau_\epsilon, z} + g_\epsilon$ is a supersolution of (5.1) in $B_{2\tau_\epsilon}(z) \cap \Omega$ which tends to infinity on $\partial(B_{2\tau_\epsilon}(z) \cap \Omega) = \partial\Omega \cap B_{2\tau_\epsilon}(z) \cup \Omega \cap \partial B_{2\tau_\epsilon}(z)$. Since we can replace $g_\epsilon(x)$ by $g_{\epsilon, \tau}(x) = \ell(d(x) - \tau)^{-\frac{2}{q-1}}$ for $\tau \in (0, \rho_\epsilon)$, any positive solution u of (5.1) in Ω is bounded from above by $f_{2\tau_\epsilon, z} + g_{\epsilon, \tau}$ and therefore by $f_{2\tau_\epsilon, z} + g_\epsilon$. This implies (6.25) with $C_\epsilon = \max\{f_{2\tau_\epsilon, z}(y) : |y - z| \leq \tau_\epsilon\}$, and it can be made explicit thanks to (6.1).

Step 2. With the same constants as in step 1, we claim that

$$U_F(x) \geq (\ell_\kappa^{q-1} - \epsilon)^{\frac{1}{q-1}} \tau^{-\frac{2}{q-1}} - C_\epsilon \quad \forall \tau \in (0, \tau_\epsilon], \forall x \in \Sigma_\tau(\overline{B_{\tau_\epsilon}(z)}). \quad (6.27)$$

If in the definition of the function g , we take $\ell = \ell(\epsilon) = (\ell_\kappa^{q-1} - \epsilon)^{\frac{1}{q-1}}$, then g is a subsolution in the same set Ω_{τ_ϵ} . Since $U_F + f_{2\tau_\epsilon, z}$ is a supersolution of (5.1) in $B_{2\tau_\epsilon}(z) \cap \Omega$ which tends to infinity on the boundary, it dominates the subsolution $g_{\epsilon, -\tau} = \ell(d(\cdot) + \tau)^{-\frac{2}{q-1}}$ for $\tau \in (0, \rho_\epsilon)$ and thus, as $\tau \rightarrow 0$, $g_\epsilon(x) \leq U_F(x) + f_{2\tau_\epsilon, z}(x)$. This implies (6.27) with the same constant C_ϵ .

Step 3. End of the proof. Since $K \subset O$ is precompact, for any $\epsilon > 0$, there exists a finite number of points $z_j, j = 1, \dots, k$ such that $K \subset \cup_{j=1}^k \overline{B_{\tau_\epsilon}(z_j)}$ with $\overline{B_{2\tau_\epsilon}(z_j)} \subset O$. Therefore

$$(\ell_\kappa^{q-1} - \epsilon)^{\frac{1}{q-1}} \tau^{-\frac{2}{q-1}} - C_\epsilon \leq U_F(x) \leq (\epsilon + \ell_\kappa^{q-1})^{\frac{1}{q-1}} \tau^{-\frac{2}{q-1}} + C_\epsilon \quad \forall \tau \in (0, \tau_\epsilon], \forall x \in \Sigma_\tau(K). \quad (6.28)$$

Since ϵ is arbitrary, it yields to

$$\lim_{\tau \rightarrow 0} \|\tau^{\frac{2}{q-1}} U_F - \ell_\kappa\|_{L^\infty(\Sigma_\tau(K))} = 0 \quad (6.29)$$

which is (6.24). \square

Corollary 6.5. *Let $U_{\partial\Omega}$ be the maximal solution of (5.1) in Ω , then*

$$\lim_{d(x) \rightarrow 0} (d(x))^{\frac{2}{q-1}} U_{\partial\Omega}(x) = \ell_\kappa. \quad (6.30)$$

7 Appendix II: The associated semilinear equation with singular potential

In the section we consider the following operator

$$v \mapsto \mathcal{M}_\tau v := -\Delta v - \frac{\tau}{|x|^2} v \quad (7.1)$$

defined in a bounded C^2 domain such that $0 \in \partial\Omega$, and the associated semilinear equation

$$\mathcal{M}_\tau w + |w|^{q-1} w = 0 \quad (7.2)$$

in Ω . The study of this operator is straightforward and we refer to [14] and [?] for verifications of the next statements. The critical value of κ is $\frac{N^2}{4}$ and since $\frac{N^2}{4} > \frac{1}{4}$ we will always impose $\kappa < \frac{N^2}{4}$.

Proposition 7.1. Assume $\kappa < \frac{N^2}{4}$, then there exist a positive solution $v_s \in C^2(\bar{\Omega} \setminus \{0\})$ of the following problem

$$\begin{aligned} \mathcal{M}_\kappa v_s &= 0 & \text{in } \Omega \\ v_s &= \delta_0 & \text{in } \partial\Omega \end{aligned} \quad (7.3)$$

satisfying

$$c_{a1}^{-1} \frac{d(x)}{|x|^{\frac{N+\sqrt{N^2-4\kappa}}{2}}} \leq v_s(x) \leq c_{a1} \frac{d(x)}{|x|^{\frac{N+\sqrt{N^2-4\kappa}}{2}}}. \quad (7.4)$$

We notice that $v_s(x) = o(K_{\mathcal{L}_\kappa}(x, 0))$ as $x \rightarrow 0$. The associated semilinear problem with a singularity at 0

$$\begin{aligned} \mathcal{M}_\tau w + |w|^{q-1}w &= 0 & \text{in } \Omega \\ w &= 0 & \text{in } \partial\Omega \end{aligned} \quad (7.5)$$

admits a critical value $q_{c^*} = \frac{N+2+\sqrt{N^2-4\kappa}}{N-2+\sqrt{N^2-4\kappa}}$. If we assume that the tangent plane to $\partial\Omega$ at 0 is $\partial\mathbb{R}_+^N := \{x = (x', x_N) : x_N = 0\}$ and the normal unit outward normal vector at 0 is $-\mathbf{e}_N$, then we can look for solutions of (7.2) in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ under the form $v(r, \sigma) = r^{-\frac{2}{q-1}}\omega(\sigma)$. Then ω satisfies

$$\begin{aligned} -\Delta_{S^{N-1}}\omega - (\ell_{q,N} + \kappa)\omega + |\omega|^{q-1}\omega &= 0 & \text{in } S_+^{N-1} \\ \omega &= 0 & \text{in } \partial S_+^{N-1}, \end{aligned} \quad (7.6)$$

where $\ell_{q,N}$ is given by (4.13). The following result is straightforward.

Proposition 7.2. If $1 < q < q_{c^*}$ there exists a unique positive function $\omega \in C^2(\bar{S}_+^{N-1})$ solution of (7.6). We denote it by ω_κ^* .

The set \mathcal{F}_κ of the solutions of (7.6) has the following structure

Proposition 7.3. Assume $\kappa < \frac{N^2}{4}$ and $q > 1$, then

I- If $q \geq q_{c^*}$, $\mathcal{F}_\kappa = \{0\}$.

II- If $1 < q < q_{c^*}$, then $\mathcal{F}_\kappa^+ = \{0, \omega_\kappa^*\}$.

III- If $q_{e^*} \leq q < q_{c^*}$, $\mathcal{F}_\kappa = \{0, \pm\omega_\kappa^*\}$, where $q_{e^*} = \frac{N+2+\sqrt{(N+2)^2-4\kappa}}{N-2+\sqrt{(N+2)^2-4\kappa}}$.

The next two results characterize the solutions of

Proposition 7.4. Assume $\kappa < \frac{N^2}{4}$ and $q \geq q_{c^*}$, then any function $v \in C^2(\bar{\Omega} \setminus \{0\})$ which satisfies (7.5) is identically zero.

Proposition 7.5. Assume $\kappa < \frac{N^2}{4}$ and $1 < q < q_{c^*}$. If $v \in C^2(\bar{\Omega} \setminus \{0\})$ is a positive solution of problem (7.5), then the following alternative holds.

(i) Either

$$\lim_{x \rightarrow 0} |x|^{\frac{2}{q-1}} v(x) = \omega_\kappa^*\left(\frac{x}{|x|}\right) \quad (7.7)$$

(ii) Or there exist $\kappa \geq 0$ such that

$$\lim_{x \rightarrow 0} |x|^{\frac{N-2+\sqrt{N^2-4\kappa}}{2}} v(x) = k \frac{x_N}{|x|} \quad (7.8)$$

and $v = v_{k\delta_0}$ is the unique solution of

$$\begin{aligned} \mathcal{M}_\tau v + |v|^{q-1}v &= 0 & \text{in } \Omega \\ v &= c_{N,\kappa} k \delta_0 & \text{in } \partial\Omega. \end{aligned} \quad (7.9)$$

Remark. Notice that in the above statement, the limit when $k \rightarrow \infty$ of a solution $v_{k\delta_0}$ is the unique positive solution $v_{\infty,0}$ satisfying (7.7). Actually $v_{\infty,0}$ is the unique positive solution of (7.5) such that $\limsup_{x \rightarrow 0} |x|^{\frac{N-2+\sqrt{N^2-4\kappa}}{2}} v(x) = \infty$.

Any positive solution of (7.5) is a subsolution for (3.61). Therefore the next comparison result which follows from the maximum principle, (7.4) and (2.52) provides a useful estimate from below of positive solutions of (3.61) with a boundary isolated singularities.

Proposition 7.6. Assume $\kappa < \frac{1}{4}$ and $1 < q < q_e$. Then there exists $c = c(a, 2) > 0$ depending on Ω , q and κ such that $v_{k\delta_0} \leq u_{ck\delta_0}$ for any $k > 0$, and $v_{\infty,0} \leq u_{\infty,0}$.

References

- [1] D.R. Adams & L.I. Hedberg, *Function Spaces and Potential Theory*, Grundlehren Math. Wiss., Vol. 314, Springer (1996).
- [2] A. Ancona, *Negatively curved manifolds, elliptic operators and the Martin boundary*. Annals Math. 2nd Series 125 (1987), 495-536.
- [3] C. Bandle, V. Moroz & W. Reichel, *Boundary blow up type sub-solutions to semilinear elliptic equations with Hardy potential*, J. London Math. Soc. 77 (2008), 503-523.
- [4] H. Berestycki, *Le nombre de solutions de certains problèmes semi-linéaires elliptiques*, J. Funct. Anal. 40 (1981), 1-29.
- [5] H. Brezis & M. Marcus, *Hardy's inequalities revisited*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 25 (1997), 217-237.
- [6] G. Barbatis, S. Filippas & A. Tertikas, *A unified approach to improved L^p Hardy inequalities with best constants*, Trans. Amer. Math. Soc. 356 (2003), 2169-2196.
- [7] L. Caffarelli, E. Fabes, S. Mortola & S. Salsa, *Boundary behavior of nonnegative solutions of elliptic operators in divergence form*, Indiana Univ. Math. J 30, 621-640.
- [8] G. Dal Maso, *On the integral representation of certain local functionals*, Ricerche Mat. 32 (1983), 85-113.
- [9] J. Davila & L. Dupaigne, *Hardy-type inequalities*, J. Eur. Math. Soc. (JEMS) 6 (2004), 335-365.
- [10] E.B. Dynkin, *Superdiffusions and partial differential equations*, American Mathematical Society Colloquium Publications 50. American Mathematical Society, Providence, RI, 2002.
- [11] D. Feyel & A. de la Pradelle, *Topologies fines et compactifications associées à certains espaces de Dirichlet*, Ann. Inst. Fourier (Grenoble) 27 (1977), 121-146.
- [12] S. Filippas, L. Moschini & A. Tertikas, *Sharp two-sided heat kernel estimates for critical Schrödinger operators on bounded domains*. Comm. Math. Phys. 273 (2007), 237-281.
- [13] A. Gmira & L. Véron, *Boundary singularities of solutions of some nonlinear elliptic equations*, Duke Math. J. 64 (1991), 271-324.
- [14] B. Guerch & L. Véron, *Local properties of stationary solutions of singular Schrödinger equations*, Revista Mat. Iberoamericana 7 (1991), 65-114.
- [15] R.A. Hunt & R. L. Wheeden, *Positive harmonic functions on Lipschitz domains*. Transactions of the American Mathematical Society, 147 (1970), 507-527.
- [16] M. Marcus, V. J. Mizel & Y. Pinchover, *On the best constant for Hardy's inequality in \mathbb{R}^n* , Trans. Amer. Math. Soc. 350 (1998), 3237-3255.

- [17] M. Marcus, *Complete classification of the positive solutions of $-\Delta u + u^q = 0$* . J. Anal. Math. 117, 187-220 (2012).
- [18] M. Marcus & L. Véron, *Removable singularities and boundary trace*. J. Math. Pures Appl. 80 (2001), 879-900.
- [19] M. Marcus & L. Véron, *The boundary trace and generalized boundary value problem for semilinear elliptic equations with coercive absorption*. Comm. Pure Appl. Math. 56 (2003), 689-731.
- [20] M. Marcus & L. Véron, *The precise boundary trace of the positive solutions of the equations $\Delta u = u^q$ in the supercritical case*. Cont. Math. 446 (2007), 345-383.
- [21] M. Marcus & L. Véron, *Boundary trace of positive solutions of semilinear elliptic equations in Lipschitz domains: the subcritical case*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 10 (2011), 913-984.
- [22] M. Marcus & L. Véron, *Nonlinear second order elliptic equations involving measures*. De Gruyter Series in Nonlinear Analysis and Applications, 21. De Gruyter, Berlin, 2014. xiv+248 pp. ISBN: 978-3-11-030515-9; 978-3-11-030531-9 35-0
- [23] M. Marcus & L. Véron, *Boundary trace of positive solutions of supercritical semilinear elliptic equations in dihedral domains*. Ann. Sc. Norm. Super. Pisa Cl. Sci., to appear.
- [24] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press, 1970.
- [25] L. Véron, *Singularities of Solutions of Second Order Quasilinear Equations*, Pitman Research Notes in Math. 353, Addison-Wesley-Longman, 1996.
- [26] L. Véron, *Elliptic Equations Involving Measures*, Handbook of Differential Equations, M. Chipot, P. Quittner, eds. Elsevier: Stationary Partial Differential Equations volume 1, 593-712, 2004.
- [27] L. Véron & C. Yarur, *Boundary value problems with measures for elliptic equations with singular potentials*. J. Funct. Anal. 262 (2012) 733-772.