



HAL
open science

A Fitting Theorem for Simple Theories

Daniel Palacin, Frank Olaf Wagner

► **To cite this version:**

Daniel Palacin, Frank Olaf Wagner. A Fitting Theorem for Simple Theories. Bulletin of the London Mathematical Society, 2016. hal-01073018v3

HAL Id: hal-01073018

<https://hal.science/hal-01073018v3>

Submitted on 28 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

A FITTING THEOREM FOR SIMPLE THEORIES

DANIEL PALACÍN AND FRANK O. WAGNER

ABSTRACT. The Fitting subgroup of a type-definable group in a simple theory is relatively definable and nilpotent. Moreover, the Fitting subgroup of a supersimple hyperdefinable group has a normal hyperdefinable nilpotent subgroup of bounded index, and is itself of bounded index in a hyperdefinable subgroup.

1. INTRODUCTION

The *Fitting subgroup* $F(G)$ of a group G is the group generated by all normal nilpotent subgroups. Since the product of two normal nilpotent subgroups of class c and c' respectively is again a normal nilpotent subgroup of class $c + c'$, it is clear that the Fitting subgroup of a finite group is nilpotent. In general, this need not be the case, and some additional finiteness conditions are needed. For groups with the chain condition on centralisers (\mathfrak{M}_c), nilpotency of the Fitting subgroup was shown by Bryant [3] for periodic groups, by Poizat and Wagner [9, 12] in the stable case (stability being a model-theoretic condition implying, in particular, that definable groups are \mathfrak{M}_c), and generally by Derakhshan and Wagner [4].

In this paper, we shall prove nilpotency of the Fitting subgroup in the model-theoretic context of groups type-definable in simple structures (Theorem 3.9), and virtual nilpotency of the Fitting subgroup for groups hyperdefinable in supersimple structures (Theorem 4.7). Simplicity is a fundamental broadening of stability mentioned above; algebraic examples of simple structures include ω -free bounded pseudo-algebraically closed fields, generic (differential) difference fields, or vector spaces over a finite field with a non-degenerate bilinear form.

In this context, a weaker chain condition on centralisers one might call (uniform) $\widetilde{\mathfrak{M}}_c$ arises naturally [13, Theorem 4.2.12]: The (uniform) chain condition on centralisers up to finite index. More precisely, we shall assume that there are natural numbers $n, d < \omega$ such that any chain $(C_G(a_j : j < i) : i < k)$ of centralizers, each of index at least d in its predecessor, has length $k < n$.

Similarly to the approach in [9, 12] we shall also need this chain condition on quotients by relatively definable subgroups, which again follows from simplicity, as the quotients are again type-definable. In this context, in contrast to \mathfrak{M}_c -groups, the centralizer of an infinite set need not be definable, and the centralizer of a relatively definable set (or even subgroup) need not even be type-definable. Nevertheless, the general theory of simplicity (Fact 2.6) yields the (relative) definability

2010 *Mathematics Subject Classification.* 03C45, 20F18 (primary), 20F24, 20F12, 20F19 (secondary).

Key words and phrases. simple theory; group; nilpotent; Fitting subgroup.

The first author was supported by the project SFB 878 and the project MTM 2011-26840. The second author was partially supported by ValCoMo (ANR-13-BS01-0006).

of a weaker, still purely group theoretic notion introduced by Haimo [5] under the name of FC-centralizer: For $H \leq G$ put

$$\tilde{C}_G(H) = \{g \in G : |H : C_H(g)| < \infty\}.$$

However, whereas for subgroups $H, K \leq G$ trivially $H \leq C_G(K)$ if and only if $K \leq C_G(H)$, no such symmetry has to hold for the FC-centralisers, even if one only asks for inclusion up to finite index. Nevertheless, for type-definable groups in a simple theory, symmetry does hold (Proposition 2.7). Note that Fact 2.6 (and more generally Fact 3.1) and symmetry are the only consequences of simplicity of the ambient theory we will use until we study hyperdefinable groups in section four. In fact, Hempel [6] has recently shown that symmetry of the FC-centraliser holds in general for type-definable groups; adapting our methods she deduces nilpotency of the Fitting subgroup of any group with the uniform $\widetilde{\mathfrak{M}}_c$ -condition in definable sections.

In the course of the paper, we shall also need that soluble and nilpotent groups are contained (up to finite index) in relatively definable soluble and nilpotent supergroups. In the definable simple case this has been shown by Milliet [7]. We follow his proofs (which adapt ideas from the hyperdefinable case [13]) in the type-definable context, adding some details about the existence of a suitable normal relatively definable abelian/central series.

Some words about definability: A subset of a structure is *definable* if it is the set of realizations of some first-order formula, which may contain parameters, such as

$$\begin{aligned} Z(G) &= \{x : G(x) \wedge \forall y [G(y) \rightarrow xy = yx]\}, \quad \text{or} \\ C_G(a) &= \{x : G(x) \wedge xa = ax\}, \end{aligned}$$

where the group G is given by a predicate $G(x)$, or by a formula abbreviated as $G(x)$. A set is definable *relative* to a superset X if it is the intersection with X of a definable set; it is *type-definable* if it is given as the intersection of definable sets. Finally, X is *hyperdefinable* if it is the quotient of a type-definable set in countably many variables by a type-definable equivalence relation. In particular, a type-definable group (G, \cdot) is a type-definable set G together with a definable subset of $G \times G \times G$ which is the graph of the group operation. A subgroup H of a group G is relatively definable if the underlying set H is definable relative to G . Note that the quotient of a type-definable group by a relatively definable normal subgroup is still type-definable¹; however the quotient of a type-definable group by a type-definable normal subgroup is only hyperdefinable, which in a way makes hyperdefinable groups a more natural category to work with.

Recall that if κ is an infinite cardinal at least the size of the language, we call a structure κ -saturated if any intersection of less than κ definable sets is non-empty as soon as all finite subintersections are. In the presence of type-definable sets, we shall always assume that the ambient structure is more highly saturated than the size of the intersections used in the type-definitions.

Although we make no assumption on the language used, we can in fact assume that it is reduced to the pure group language $\{\cdot, {}^{-1}, =\}$ (in addition to the predicate(s) (type-)defining the ambient group). All our definable sets have an easy

¹or *type-interpretable*, if we do not consider classes modulo \emptyset -definable equivalence relations as elements

group-theoretic interpretation (except for Section four where the model theory becomes more involved).

2. ALMOST JUST DEFINITIONS

Definition 2.1. A subgroup H of G is *almost contained* in another subgroup K , denoted by $H \lesssim K$, if $H \cap K$ has finite index in H . Then H and K are *commensurable*, denoted by $H \sim K$, if $H \lesssim K$ and $K \lesssim H$.

Observe that \lesssim is a transitive relation among subgroups of G , and that \sim is an equivalence relation.

Definition 2.2. Let H be a subgroup of a group G . The *almost normalizer* of H is defined as

$$\tilde{N}_G(H) = \{g \in G : H \sim H^g\}.$$

Note that if H and K are commensurable, then $\tilde{N}_G(H) = \tilde{N}_G(K)$. By compactness, if G is type-definable and H relatively definable, then $\tilde{N}_G(H)$ is relatively definable if and only if all $\tilde{N}_G(H)$ -conjugates of H are uniformly commensurable with H ; otherwise $\tilde{N}_G(H)$ is not even type-definable.

We shall usually work in a context where commensurability is uniform. Then a theorem by Schlichting [11], generalized by Bergmann and Lenstra [2] (see also [13, Theorem 4.2.4] for definability questions), provides an invariant object:

Fact 2.3. Let \mathcal{H} be a family of uniformly commensurable subgroups of a group G , i.e. the index $|H : H \cap H^*|$ is finite and bounded independently of $H, H^* \in \mathcal{H}$. Then there is a subgroup N commensurable with any $H \in \mathcal{H}$, which is invariant under any automorphism of G stabilizing \mathcal{H} setwise. Moreover, N is a finite extension of a finite intersection of groups in \mathcal{H} ; if the latter are relatively definable, so is N .

It follows that if H is uniformly commensurable with all its $\tilde{N}_G(H)$ -conjugates, then there is a normal subgroup $\tilde{H} \trianglelefteq \tilde{N}_G(H)$ commensurable with H . Clearly, any two choices for \tilde{H} will be commensurable, and $N_G(\tilde{H}) = \tilde{N}_G(\tilde{H}) = \tilde{N}_G(H)$. We call \tilde{H} a *conjugacy-connected component* of H . Clearly, any two conjugacy-connected components of H are commensurable, and have the same normalizer, which is equal to their almost normalizer.

Definition 2.4. Let K and H be subgroups of a group G with $H \lesssim \tilde{N}_G(K)$, and suppose K has a conjugacy-connected component \tilde{K} . The *almost centralizer of H modulo K* is given by

$$\tilde{C}_G(H/K) = \{g \in \tilde{N}_G(K) : |H : C_H(g/\tilde{K})| \text{ is finite}\}.$$

For $n < \omega$ the n -th iterated almost centralizer of H modulo K is defined inductively by $\tilde{C}_G^0(H/K) = K$, and if $\tilde{C}_G^n(H/K)$ has a conjugacy-connected component, then

$$\tilde{C}_G^{n+1}(H/K) = \bigcap_{i \leq n} \tilde{N}_G(\tilde{C}_G^i(H/K)) \cap \tilde{C}_G(H/\tilde{C}_G^n(H/K)).$$

If $K = \{1\}$ it is omitted. Moreover, we define the *almost centre* of G as $\tilde{Z}(G) = \tilde{C}_G(G)$ and for $n < \omega$ the n -th iterated almost centre of G as $\tilde{Z}_n(G) = \tilde{C}_G^n(G)$.

Thus $\tilde{C}_G(H/K) = \tilde{C}_G^1(H/K)$ and $\tilde{Z}(G) = \tilde{C}_G^1(G)$. For any subgroup L we put $\tilde{C}_L^n(H/K) = \tilde{C}_G^n(H/K) \cap L$ and $\tilde{N}_L(K) = \tilde{N}_G(K) \cap L$.

Remark 2.5. One easily sees that $\tilde{C}_G(H/K)$ is an H -invariant subgroup of $\tilde{N}_G(K)$, as are all $\tilde{C}_G^n(H/K)$ for $n > 0$. Moreover, if $H^* \sim H$ and $K^* \sim K$, then $\tilde{C}_G^n(H/K) = \tilde{C}_G^n(H^*/K^*)$ for all $n > 0$.

Fact 2.6. *If H is type-definable and K a relatively definable subgroup of H in a simple theory, then commensurability among conjugates is uniform. In particular, K has a conjugacy-connected component, and both $\tilde{N}_G(K)$ and $\tilde{C}_G^n(H/K)$ are relatively definable subgroups of G .*

Proof: This follows immediately from [13, Lemma 4.2.6]. \square

In a simple theory, the existence of generic elements yields the following symmetry property, which plays an essential role throughout the paper.

Proposition 2.7. *Let G be type-definable in a simple theory, and H and K be type-definable subgroups of G . The following are equivalent:*

- (1) $H \lesssim \tilde{C}_G(K)$.
- (2) *There are independent generic elements $h \in H$ and $k \in K$ with $[h, k] = 1$.*

In particular, $H \lesssim \tilde{C}_G(K)$ if and only if $K \lesssim \tilde{C}_G(H)$.

Proof: Suppose $H \lesssim \tilde{C}_G(K)$. So there is a generic $h \in H$ with $h \in \tilde{C}_G(K)$. Thus $C_K(h)$ has finite index in K , and there is generic $k \in K$ over h with $k \in C_G(h)$. Then h and k are independent, and $[h, k] = 1$.

Conversely, suppose $h \in H$ and $k \in K$ are independent generics with $[h, k] = 1$. As $k \in C_K(h)$ and k is generic over h , the index of $C_K(h)$ in K is finite. Thus $h \in \tilde{C}_G(K)$; as h is generic, we get $H \lesssim \tilde{C}_G(K)$. \square

Of course, if $H, K \leq N_G(N)$ we also have $H \lesssim \tilde{C}_G(K/N)$ if and only if $K \lesssim \tilde{C}_G(H/N)$, by working in the group $N_G(N)/N$. Thus symmetry also holds for relative almost centralizers.

We shall finish this section by recalling two group-theoretic facts.

Fact 2.8 ([8, Theorem 3.1]). *There is a finite bound of the size of conjugacy classes in a group G if and only if the derived subgroup G' is finite.*

Fact 2.9 ([1, 10]). *Let H and N be subgroups of G with N normalized by H . If the set of commutators*

$$\{[h, n] : h \in H, n \in N\}$$

is finite, then the group $[H, N]$ is finite.

3. NILPOTENCY IN TYPE-DEFINABLE GROUPS IN A SIMPLE THEORY

In this section, we shall generalize the results of Milliet [7] to the relatively definable context. For this we need the following result.

Fact 3.1 ([13, Theorem 4.2.12]). *Let G be type-definable in a simple theory, and \mathcal{H} a family of uniformly relatively definable subgroups. Then there are $n, d < \omega$ such that any chain $(\bigcap_{j < i} H_j : i < k)$ of intersections of groups $H_i \in \mathcal{H}$, each of index at least d in its predecessor, has length $k < n$.*

Lemma 3.2. *Let G be a type-definable group in a simple theory, and H a soluble subgroup of G . Then there is a relatively definable soluble subgroup S containing H , and a series of relatively definable S -invariant subgroups*

$$\{1\} = S_0 < S_1 < \cdots < S_n = S,$$

all normalized by $N_G(H)$, such that S_i/S_{i+1} is abelian for all $i < n$. The derived length n of S is at most three times the derived length of H . Moreover, S_1 and S_n/S_{n-1} are finite.

Proof: Suppose first that H is abelian. By Fact 3.1 there is a finite tuple $\bar{h} \in H$ and $d < \omega$ such that for any $h \in H$

$$|C_G(\bar{h}) : C_G(\bar{h}, h)| < d.$$

Hence the $N_G(H)$ -conjugates of $C_G(\bar{h})$ are all commensurable, and $N_G(H) \leq \tilde{N}_G(C_G(\bar{h})) =: N$. Now Fact 2.6 yields that N is relatively definable, and $C_G(\bar{h})$ has a relatively definable conjugacy-connected component C normalized by N . Then

$$H = C_H(\bar{h}) \leq C_G(\bar{h}) \lesssim C.$$

Now, as

$$C \sim C_G(\bar{h}) \sim C_G(\bar{h}, h) = C_{C_G(\bar{h})}(h) \sim C_C(h)$$

for any $h \in H$, the relatively definable subgroup $\tilde{Z}(C)$ of G contains $H \cap C$, whence $H \lesssim \tilde{Z}(C)$. By compactness there is a finite bound on the size of conjugacy classes in $\tilde{Z}(C)$, so $\tilde{Z}(C)'$ is finite by Fact 2.8 and hence definable. Put $S_2 = C_{\tilde{Z}(C)}(\tilde{Z}(C)')$, a relatively definable characteristic subgroup of C , which must be normalized by N . Since $\tilde{Z}(C)'$ is finite, S_2 has finite index in $\tilde{Z}(C)$, so $H \lesssim S_2$. If $S_1 = S_2 \cap \tilde{Z}(C)'$, then S_1 is finite, abelian (even central in S_2) and normalized by N , and S_2/S_1 is abelian. Put $S_3 = HS_2$, a finite extension of S_2 and thus relatively definable. Then S_3/S_2 is abelian as well; moreover, if \bar{h}' is a system of representatives of S_3/S_2 , then

$$N_G(H) \leq N_1 = \{g \in N : h^g \in S_3 \text{ for all } h \in \bar{h}'\} \leq N_G(S_3).$$

So N_1 is a relatively definable subgroup of G normalizing S_3 and containing $N_G(H)$. Replace G by N_1/S_3 , we finish by induction. \square

Note that the above proof merely uses the $\tilde{\mathfrak{M}}_c$ -condition for G and for certain relatively definable sections of G , but not symmetry of the almost centraliser. This is different for nilpotency, where the following lemma is used.

Lemma 3.3. *Let G be a type-definable group in a simple theory. Then there is a characteristic relatively definable subgroup G_0 of finite index and a finite characteristic subgroup $N \leq Z(G_0)$ such that $\tilde{Z}(G) \leq C_G(G_0/N)$.*

Proof: As trivially $\tilde{C}_G(G) \lesssim \tilde{C}_G(G)$, Proposition 2.7 yields

$$G \lesssim \tilde{C}_G(\tilde{C}_G(G)) = \tilde{C}_G(\tilde{Z}(G)),$$

and so $\tilde{C}_G(\tilde{Z}(G))$ is a characteristic relatively definable subgroup of finite index in G . For independent $g \in \tilde{Z}(G)$ and $h \in \tilde{C}_G(\tilde{Z}(G))$ we have

$$[g, h] \in \text{acl}(g) \cap \text{acl}(h) = \text{acl}(\emptyset).$$

As every element in $\tilde{Z}(G)$ is the product of two generic elements g, g' each of which can be chosen independently of $h \in \tilde{C}_G(\tilde{Z}(G))$, and

$$[gg', h] = [g, h][[g, h], g'][g', h] \in \text{acl}(\emptyset),$$

the set of commutators

$$\{[g, h] : g \in \tilde{Z}(G), h \in \tilde{C}_G(\tilde{Z}(G))\}$$

is bounded, whence finite by compactness. By Fact 2.9 the characteristic group $Z = [\tilde{Z}(G), \tilde{C}_G(\tilde{Z}(G))]$ is finite. We put $G_0 = \tilde{C}_G(\tilde{Z}(G)) \cap C_G(Z)$ and $N = Z \cap G_0$. \square

Lemma 3.4. *Let G be a type-definable group in a simple theory, and H a nilpotent subgroup of G . Then there is a relatively definable nilpotent subgroup N with $H \lesssim N$, and a series of relatively definable $N_G(H)$ -invariant subgroups*

$$\{1\} = N_0 < N_1 < \cdots < N_n = N$$

such that $N_{i+1} \leq C_G(N/N_i)$ for all $i < n$. The nilpotency class n of N is at most two times the nilpotency class of H .

The same conclusion holds if H is merely FC-nilpotent, i.e. $H \lesssim \tilde{Z}_k(H)$ for some k .

Proof: We use induction on the (FC-)nilpotency class of H . Consider the family $\mathcal{H} = \{C_G(g) : g \in \tilde{C}_G(H)\}$. By Fact 3.1 there is a finite intersection C of groups in \mathcal{H} such that any further intersection has boundedly finite index. As \mathcal{H} is clearly $N_G(H)$ -invariant, we obtain $\tilde{N}_G(C) \geq N_G(H)$. Fact 2.6 yields that $\tilde{N}_G(C)$ is relatively definable, and C has a relatively definable conjugacy-connected component \tilde{C} . So $N_G(\tilde{C}) = \tilde{N}_G(C) \geq N_G(H)$. As $H \lesssim C_G(g)$ for any $C_G(g) \in \mathcal{H}$, we get

$$H \lesssim C \sim \tilde{C}.$$

By construction

$$\tilde{C}_G(H \cap \tilde{C}) = \tilde{C}_G(H) \leq \tilde{C}_G(C) = \tilde{C}_G(\tilde{C}),$$

whence $\tilde{Z}(H \cap \tilde{C}) \leq \tilde{Z}(\tilde{C})$. Now consider the characteristic relatively definable groups G_0 and N given by Lemma 3.3 applied to \tilde{C} . Then $H \lesssim G_0$. We put $N_1 = N$ and $N_2 = \tilde{Z}(\tilde{C}) \cap G_0$. Note that N_1, N_2 and G_0 are all relatively definable and normalized by $N_G(\tilde{C}) \geq N_G(H)$, and $G_0 \cap H$ is normalized by $N_G(H)$; moreover $N_1 \leq C_G(G_0 \cap H)$ and $N_2 \leq C_G(G_0 \cap H/N_1)$. Now $(H \cap G_0)/N_2$ is a nilpotent subgroup of $N_G(\tilde{C})/N_2$ of smaller (FC-)nilpotency class, and we finish by induction. \square

Remark 3.5. If in addition H is normalized by N , then NH is nilpotent of class at most three times the class of H ; if c is the nilpotency class of H and $\bar{h} \in H$ is a system of representatives of NH/N , then

$$\begin{aligned} \{1\} &\leq C_{N_1}(\bar{h}) \leq C_{N_1}^2(\bar{h}) \leq \cdots \leq C_{N_1}^{c+1}(\bar{h}) = N_1 \\ &\leq C_{N_2}(\bar{h}/N_1) \leq C_{N_2}^2(\bar{h}/N_1) \leq \cdots \leq C_{N_2}^{c+1}(\bar{h}) = N_2 \\ &\leq \cdots \leq C_{N_n}^{c+1}(\bar{h}/N_{n-1}) = N_n = N \\ &\leq NZ(H) \leq NZ_2(H) \leq \cdots \leq NZ_c(H) = NH \end{aligned}$$

is a relatively definable central series for NH normalised by $N_G(H)$.

Lemma 3.6. *Let G be a type-definable group in a simple theory and \mathcal{H} a directed system of nilpotent subgroups. Then $H = \bigcup \mathcal{H}$ is soluble.*

Proof: Let (n, d) be the bounds given by Fact 3.1 for chains of centralizers, and (n', d') the bounds for chains of centralizers modulo $\tilde{Z}(G)$. The proof is by induction on n .

Consider G_0 and N as given by Lemma 3.3. As HG_0/G_0 is finite and nilpotent, we may assume that $H \leq G_0$. If $H \leq \tilde{Z}(G)$ we are done, as $(\tilde{Z}(G) \cap G_0)/N$ and N are abelian.

If $H \not\leq \tilde{Z}(G)$, then consider some $H_0 \in \mathcal{H}$ with $H_0 \not\leq \tilde{Z}(G)$ and we take $h_0 \in C_{H_0}(H_0/\tilde{Z}(G)) \setminus \tilde{Z}(G)$; note that such an element exists as $H_0/\tilde{Z}(G)$ is nilpotent.

If $C_H(h_0/\tilde{Z}(G))$ has index greater than d' in H , then there is some $H_1 > H_0$ in \mathcal{H} such that $C_{H_1}(h_0/\tilde{Z}(G))$ has index greater than d' in H_1 . Choose $h_1 \in C_{H_1}(H_1/\tilde{Z}(G)) \setminus \tilde{Z}(G)$, and note that $C_G(h_1, h_0/\tilde{Z}(G))$ has index greater than d' in $C_G(h_1/\tilde{Z}(G))$. If $C_H(h_1/\tilde{Z}(G))$ has index more than d' in H , then we can iterate this process, which must stabilize after at most n' steps. It follows that there is some $h \in H \setminus \tilde{Z}(G)$ such that $C_H(h/\tilde{Z}(G))$ has index at most d' in H .

Since $C_G(h)$ has infinite index in G , the induction hypothesis for $n - 1$ yields that $C_H(h)$ is soluble. Moreover, as N is central in G_0 the map from $C_H(h/N)$ to N given by $x \mapsto [h, x]$ is a homomorphism with abelian image and kernel $C_H(h)$. Thus $C_H(h/N)/C_H(h)$ is abelian. Similarly, as $\tilde{Z}(G)$ is centralised by H modulo N , the map $x \mapsto [h, x]N$ is a homomorphism from $C_H(h/\tilde{Z}(G))$ to $\tilde{Z}(G)/N$ with abelian image and kernel $C_H(h/N)$. Therefore, $C_H(h/\tilde{Z}(G))$ is soluble. Finally, as $C_H(h/\tilde{Z}(G))$ contains a normal subgroup K of H with H/K finite, whence nilpotent, we see that H must be soluble. \square

Corollary 3.7. *A locally nilpotent subgroup H of a type-definable group in a simple theory is soluble.*

Proof: The collection of finitely generated subgroups of H satisfies the hypotheses of Lemma 3.6. \square

Lemma 3.8. *Let G be a type-definable group acting definably on a type-definable abelian group A , in a simple theory. Suppose that $H \leq G$ is abelian, and that there are $\bar{g} = (g_i : i < k)$ in H and $m_i < \omega$ for $i < k$ such that $(g_i - 1)^{m_i}A$ is finite for all $i < k$, and for any $g \in H$ the index of $C_A(\bar{g}, g)$ in $C_A(\bar{g})$ is finite. Put $m = 1 + \sum_{i < k} (m_i - 1)$. Then there is a relatively definable supergroup \bar{H} of H such that $\bar{C}_A^m(\bar{H})$ has finite index in A .*

Proof: By [13, Lemma 4.2.6] the group

$$\bar{H} = \{g \in C_G(\bar{g}) : |C_A(\bar{g}) : C_A(\bar{g}, g)| \text{ finite}\}$$

is relatively definable, and it clearly contains H . By the pigeonhole principle, for any m indices $(i_j : j < m) \in k^m$, there must be at least one $i < k$ such that m_i of the indices are equal to i . As the group ring $\mathbb{Z}(H)$ is commutative, this implies that

$$(g_{i_0} - 1)(g_{i_1} - 1) \cdots (g_{i_{m-1}} - 1)A$$

is finite. Hence there is a subgroup A_0 of finite index in A such that

$$(g_{i_0} - 1)(g_{i_1} - 1) \cdots (g_{i_{m-1}} - 1)A_0 = 0$$

for all of the finitely many choices $(i_j : j < m) \in k^m$. It follows that for all choices of $(i_j : 1 \leq j < m) \in k^{m-1}$ we have

$$(g_{i_1} - 1) \cdots (g_{i_{m-1}} - 1)A_0 \leq C_A(g_i : i < k).$$

As $C_A(h_0, g_i : i < k)$ has finite index in $C_A(g_i : i < k)$ for all $h_0 \in \bar{H}$, the group

$$(h_0 - 1)(g_{i_1} - 1) \cdots (g_{i_{m-1}} - 1)A_0$$

is finite for all choices of $(i_j : 1 \leq j < m) \in k^{m-1}$, as is

$$(h_0 - 1)(g_{i_1} - 1) \cdots (g_{i_{m-1}} - 1)A.$$

By the same argument (keeping h_0 fixed) and the fact that $\bar{H} \leq C_G(\bar{g})$ we see that for any h_1 in G and all choices of $(i_j : 2 \leq j < m) \in k^{m-2}$ the group

$$(h_1 - 1)(h_0 - 1)(g_{i_2} - 1) \cdots (g_{i_{m-1}} - 1)A$$

is finite, and inductively that

$$(h_{m-1} - 1) \cdots (h_1 - 1)(h_0 - 1)A$$

is finite for any $(h_j : j < m)$ in \bar{H} . It follows that

$$\bar{H} \leq \tilde{C}_{\bar{H}}((h_{m-2} - 1) \cdots (h_0 - 1)A)$$

for all $(h_j : j < m - 1)$ in G , whence by symmetry

$$(h_{m-2} - 1) \cdots (h_0 - 1)A \lesssim \tilde{C}_A(\bar{H}).$$

But $\tilde{C}_A(\bar{H})$ is relatively definable; we may divide out and note that

$$(h_{m-2} - 1) \cdots (h_0 - 1)A / \tilde{C}_A(\bar{H})$$

is finite for all choices of $(h_j : j < m - 1)$ in \bar{H} . Hence

$$\bar{H} \leq \tilde{C}_{\bar{H}}((h_{m-3} - 1) \cdots (h_0 - 1)A / \tilde{C}_A(\bar{H}))$$

and by symmetry

$$(h_{m-3} - 1) \cdots (h_0 - 1)A \lesssim \tilde{C}_A(\bar{H} / \tilde{C}_A(\bar{H})) = \tilde{C}_A^2(\bar{H}).$$

Inductively, we see that $A \lesssim \tilde{C}_A^m(\bar{H})$. \square

Theorem 3.9. *Let G be a type-definable group in a simple theory. Then the Fitting subgroup $F(G)$ is nilpotent.*

Proof: As $F(G)$ is soluble by Lemma 3.6, by Lemma 3.2 there is a chain

$$\{1\} = S_0 < S_1 < \cdots < S_d = S$$

of relatively definable normal subgroups of G such that S contains $F(G)$ and all quotients S_{i+1}/S_i for $i < d$ are abelian. Since $F(S_i) = F(G) \cap S_i$, we may assume by induction on d that $F(G)' \leq F(S_{d-1})$ is nilpotent. By Lemma 3.4 and Remark 3.5 there is a relatively definable normal nilpotent group N containing $F(G)'$, and a relatively definable series

$$\{1\} = N_0 < N_1 < \cdots < N_k = N$$

of normal subgroups of G with $[N, N_{i+1}] \leq N_i$ for all $i < k$.

Fix $i > 0$. Any $g \in F(G)$ is contained in a normal nilpotent subgroup H_g . Since $N_i H_g$ is again nilpotent, there is $m_g < \omega$ such that

$$(g - 1)^{m_g} N_i \leq N_{i-1}.$$

By Fact 3.1 there is a finite tuple $\bar{g} \in F(G)$ such that for any $g \in F(G)$ the index $|C_{N_i}(\bar{g}/N_{i-1}) : C_{N_i}(\bar{g}, g/N_{i-1})|$ is finite. Furthermore, by Lemma 3.8 (applied to G/N with its abelian subgroup $F(G)/N$ acting on N_i/N_{i-1} by conjugation) there is $m_i < \omega$ and a relatively definable group $H_i \geq F(G)$ such that $N_i \lesssim \tilde{C}_G^{m_i}(H_i/N_{i-1})$. Then, the finite intersection $\bigcap_i H_i$ is a relatively definable supergroup of $F(G)$. By Facts 3.1 and 2.3 there is a relatively definable normal subgroup H which is a finite extension of a finite intersection of G -conjugates of $\bigcap_i H_i$. Thus $H \geq F(G)$, and $H \lesssim H_i$ for all i . By Lemma 3.2 applied to the abelian normal subgroup $F(G)/N$

of G/N , we may restrict H and assume that there are relatively definable normal subgroups

$$N \leq Z \leq A \leq H$$

of G with Z/N and H/A finite and A/Z abelian. Then

$$N_i \lesssim \tilde{C}_G^{m_i}(H_i/N_{i-1}) \leq \tilde{C}_G^{m_i}(H_i \cap H/N_{i-1}) = \tilde{C}_G^{m_i}(H/N_{i-1}),$$

and inductively

$$\begin{aligned} N_i &\lesssim \tilde{C}_G^{m_i}(H/N_{i-1}) \\ &\leq \tilde{C}_G^{m_i}(H/\tilde{C}_G^{m_{i-1}}(H/N_{i-2})) = \tilde{C}_G^{m_i+m_{i-1}}(H/N_{i-2}) \\ &\vdots \\ &\leq \tilde{C}_G^{m_i+m_{i-1}+\dots+m_1}(H). \end{aligned}$$

Thus $N \lesssim \tilde{C}_N^m(H)$ for $m = m_1 + m_2 + \dots + m_k$. Since $Z \lesssim N$ and $N \leq A \leq H$ we obtain $Z \lesssim \tilde{C}_A^m(A)$, whence $A = \tilde{Z}^{m+1}(A)$. So A is nilpotent-by-finite by Lemma 3.4, as is $F(G)$, since $F(G) \lesssim A$. If F is a normal nilpotent subgroup of finite index in $F(G)$ and K a normal nilpotent group containing representatives for $F(G)/F$, then FK is a nilpotent, as is $F(G)$. \square

4. HYPERDEFINABLE GROUPS

In the previous section we have systematically used the fact that type-definability is preserved under quotients whenever we divide out by a relatively definable subgroup. However, type-definability is not preserved when quotienting by a type-definable subgroup, and in fact such quotients (and even the slightly more general ones defined below) arise naturally from model-theoretic considerations in simplicity theory. We are thus led to the following definition.

Definition 4.1. A *hyperdefinable* group is a group whose domain is given by a partial type π modulo a type-definable equivalence relation E , and whose group law is induced by an E -invariant type-definable relation on π^3 .

Note that a quotient of a hyperdefinable group by a hyperdefinable group is again hyperdefinable. In this context, we have to replace *finite index* by *bounded index*, i.e. the index remains bounded even in a very saturated elementary extension. With this replacement in the definitions of Section 2, almost containment is transitive, commensurability is an equivalence relation, and we still have good definability properties in a simple theory.

Fact 4.2 ([13, Proposition 4.4.10 and Corollary 4.5.16]). *Let K and H be hyperdefinable subgroups of a hyperdefinable group G in a simple theory, with $H \lesssim \tilde{N}_G(K)$. Then:*

- (1) $\tilde{N}_G(K)$ is hyperdefinable, and a conjugacy-connected component \tilde{K} exists.
- (2) $\tilde{C}_G^n(H/K)$ is hyperdefinable for all $n < \omega$.

Moreover, the proof of Proposition 2.7 remains valid in the hyperdefinable context.

In contrast to the type-definable case, simplicity does not necessarily yield a finite chain condition on centralizers (even though there is an ordinal α such that any descending chain of hyperdefinable subgroups having unbounded index in its

predecessor stabilizes, up to bounded index, after α many steps). In order to adapt the arguments from the previous section we shall make a stronger assumption, supersimplicity. More precisely, we shall assume the following consequence of supersimplicity: There is no infinite descending chain of hyperdefinable subgroups, each of unbounded index in its predecessor. In particular, we obtain a minimal condition on centralizers, up to bounded index.

As a consequence, all proofs of the previous section adapt to this wider context and therefore we obtain the same result, up to bounded index. Note that Remark 3.5 need no longer hold, as a system of representatives for a subgroup of bounded index can now be infinite.

Alternatively, we offer a distinct proof of virtual nilpotency of the Fitting subgroup of a hyperdefinable group of ordinal SU-rank in a simple theory, which in addition provides a bound on the nilpotency class. For the rest of the section, the ambient theory will be simple. We first recall some facts starting with the Lascar inequalities for SU-rank.

Fact 4.3 ([13, Theorem 5.1.6 (1)]). *If H and K are hyperdefinable subgroups of a common hyperdefinable group, then*

$$\mathrm{SU}(K) + \mathrm{SU}(HK/K) \leq \mathrm{SU}(HK) \leq \mathrm{SU}(K) \oplus \mathrm{SU}(HK/K),$$

where \oplus is the least symmetric increasing function on ordinals satisfying $f(\alpha, \beta + 1) = f(\alpha, \beta) + 1$.

Fact 4.4 ([13, Proposition 5.4.3]). *Let G be an \emptyset -hyperdefinable group of rank $\mathrm{SU}(G) = \omega^\alpha \cdot n + \gamma$ with $\gamma < \omega^\alpha$. Then G has an \emptyset -hyperdefinable normal subgroup H of SU-rank $\omega^\alpha \cdot n$.*

Corollary 4.5. *Let G be an \emptyset -hyperdefinable group of rank $\mathrm{SU}(G) = \omega^{\alpha_1} \cdot n_1 + \dots + \omega^{\alpha_k} \cdot n_k$ with $\alpha_i > \alpha_{i+1}$ for $i < k$ and $n_i > 0$ for $i \leq k$. Then there exists a series of \emptyset -hyperdefinable G -invariant subgroups*

$$\{1\} = G_0 \trianglelefteq G_1 \trianglelefteq \dots \trianglelefteq G_\ell = G$$

with $\ell \leq n_1 + \dots + n_k$ such that each quotient G_{i+1}/G_i is unbounded of monomial SU-rank $\omega^{\beta_i} \cdot m_i$ and its \emptyset -hyperdefinable G -invariant subgroups of unbounded index have SU-rank strictly smaller than ω^{β_i} .

Proof: By Fact 4.4 there is an \emptyset -hyperdefinable normal subgroup G_1 of G of minimal monomial Lascar rank of the form $\mathrm{SU}(G_1) = \omega^{\alpha_1} \cdot m$ with positive $m \leq n_1$. By minimality, G_1 is as required. If $\mathrm{SU}(G_1) = \mathrm{SU}(G)$ we are done. Otherwise, $\mathrm{SU}(G/G_1) < \mathrm{SU}(G)$ by the Lascar inequalities, so we finish by induction on $\mathrm{SU}(G)$. \square

Next, we recall the supersimple version of Zilber's Indecomposability Theorem.

Fact 4.6 ([13, Theorem 5.4.5 and Remark 5.4.7]). *Let G be an \emptyset -hyperdefinable group of rank $\mathrm{SU}(G) < \omega^{\alpha+1}$. If \mathfrak{X} is a family of hyperdefinable subsets of G , then there exists a hyperdefinable subgroup $K \leq X_1^{\pm 1} \dots X_m^{\pm 1}$ for some $X_1, \dots, X_m \in \mathfrak{X}$, such that $\mathrm{SU}(XK) < \mathrm{SU}(K) + \omega^\alpha$ for all $X \in \mathfrak{X}$. Moreover, $\mathrm{SU}(K) = \omega^\alpha \cdot n$, and K is unique up to commensurability. In particular, if \mathfrak{X} is invariant under all automorphisms we can choose K hyperdefinable over \emptyset , and if \mathfrak{X} is G -invariant, we can take K to be normal in G .*

Finally, we state the hyperdefinable version of our main result in the supersimple case. Recall that the \emptyset -connected component N_\emptyset^0 of a hyperdefinable group N is the intersection of all \emptyset -hyperdefinable subgroups of N of bounded index in N .

Theorem 4.7. *Let G be an \emptyset -hyperdefinable group of rank $\text{SU}(G) = \omega^{\alpha_1} \cdot n_1 + \dots + \omega^{\alpha_k} \cdot n_k$. Then $F(G)$ has bounded index in an \emptyset -hyperdefinable FC-nilpotent normal subgroup N of class $\ell \leq n_1 + \dots + n_k$. In particular, the hyperdefinable normal group N_\emptyset^0 is nilpotent of class 2ℓ and has bounded index in $F(G)$.*

Proof: By Lemma 4.5 there is a finite series of \emptyset -hyperdefinable G -invariant subgroups

$$\{1\} = G_0 \trianglelefteq G_1 \trianglelefteq \dots \trianglelefteq G_\ell = G$$

with $\ell \leq n_1 + \dots + n_k$, such that each quotient G_{i+1}/G_i is unbounded of monomial Lascar rank $\omega^{\beta_i} \cdot m_i$, and its \emptyset -hyperdefinable G -invariant subgroups of unbounded index have SU-rank strictly smaller than ω^{β_i} . Clearly, we may assume that all G_i are \emptyset -connected, i.e. have no \emptyset -hyperdefinable subgroup of bounded index.

Let N be the intersection $\bigcap_{i < \ell} \tilde{C}_G(G_{i+1}/G_i)$, an \emptyset -hyperdefinable normal subgroup of G . Note that $N \leq \tilde{C}_G((G_{i+1} \cap N)/(G_i \cap N))$. Hence by symmetry we get

$$G_{i+1} \cap N \lesssim \tilde{C}_G(N/(G_i \cap N))$$

for all $i < \ell$. Inductively,

$$\begin{aligned} N &= G_\ell \cap N \lesssim \tilde{C}_G(N/(G_{\ell-1} \cap N)) \\ &\leq \tilde{C}_G(N/(\tilde{C}_G(N/(G_{\ell-2} \cap N)))) = \tilde{C}_G^2(N/(G_{\ell-2} \cap N)) \\ &\leq \tilde{C}_G^2(N/\tilde{C}_G(N/(G_{\ell-3} \cap N))) = \tilde{C}_G^3(N/(G_{\ell-3} \cap N)) \\ &\leq \dots \leq \tilde{C}_G^\ell(N/(G_0 \cap N)) = \tilde{C}_G^\ell(N). \end{aligned}$$

It follows that $N = \tilde{Z}_\ell(N)$ is FC-nilpotent of class at most ℓ . Then N_\emptyset^0 is nilpotent of class 2ℓ by [13, Proposition 4.4.10 (3)].

In order to finish, we shall show $F(G) \leq N$. Fix $i \leq \ell$, and consider the \emptyset -invariant family \mathfrak{X} formed by the hyperdefinable sets $X_a = [a, G_{i+1}]/G_i$ for $a \in F(G)$. Suppose, towards a contradiction, that the SU-rank of some of these sets is greater than ω^{β_i} . By Fact 4.6 applied to G_{i+1}/G_i we obtain an \emptyset -hyperdefinable G -invariant subgroup $H \leq G_{i+1}/G_i$ of monomial SU-rank which is contained in a finite product of sets $X_{a_0}^{\pm 1}, \dots, X_{a_m}^{\pm 1}$ from \mathfrak{X} ; moreover, $\text{SU}(H) \geq \omega^{\beta_i}$. Thus H has bounded index in G_{i+1}/G_i and must be equal by \emptyset -connectivity. But every $a_j G_i$ is contained in a normal nilpotent subgroup of G/G_i which must also contain X_{a_j} , so $K = \langle a_j G_i, X_{a_j} : j \leq m \rangle$ is a nilpotent subgroup of G/G_i . However,

$$(\dagger) \quad H \leq X_{a_0}^{\pm 1} \dots X_{a_m}^{\pm 1} = [a_0, H]^{\pm 1} \dots [a_m, H]^{\pm 1} \subseteq H$$

and we must have equality, contradicting nilpotency of K : If H is in the k -th element $\gamma_k(K)$ of the lower central series of K , then equation (\dagger) implies that $H \leq \gamma_{k+1}(K)$. Thus $H \leq \bigcap_{k < \omega} \gamma_k(K) = \{1\}$, a contradiction.

It follows that $\text{SU}(X_a) < \omega^{\beta_i}$ for all $X_a \in \mathfrak{X}$. As X_a is in bijection with $G_{i+1}/C_{G_{i+1}}(a/G_i)$, the Lascar inequalities imply that X_a is bounded and so $a \in \tilde{C}_G(G_{i+1}/G_i)$. Thus $F(G) \leq N$, as required. \square

REFERENCES

- [1] Reinhold Baer. Endlichkeitskriterien für Kommutatorgruppen. *Math. Ann.* **124** (1952).
- [2] George M. Bergman and Hendrik W. Lenstra, Jr. Subgroups close to normal subgroups. *J. Alg.* **127** (1989) 80–97.
- [3] Roger M. Bryant. Groups with the minimal condition on centralizers. *J. Alg.* **60** (1979), 371–383.
- [4] Jamshid Derakhshan and Frank O. Wagner. Nilpotency in groups with chain condition on centralizers. *Quart. J. Math. Oxford (2)* **48** (1997), 453–468.
- [5] Franklin Haimo. The *FC*-chain of a group. *Can. J. Math.* **5** (1953), 498–511.
- [6] Nadja Hempel. Almost group theory. Preprint.
- [7] Cédric Milliet. Definable envelopes in groups with simple theory. Preprint. HAL:00657716
- [8] Bernhard Hermann Neumann. Groups covered with permutable subsets. *J. London Math. Soc.* **29** (1954), 236–248.
- [9] Bruno Poizat and Frank O. Wagner. Sous-groupes périodiques d’un groupe stable. *J. Symb. Logic* **58** (1993), 385–400.
- [10] Maxwell Rosenlicht. On a result of Baer. *Proc. Amer. Math. Soc.* **13** (1962), 99–101.
- [11] G. Schlichting. Operationen mit periodischen Stabilisatoren. *Arch. Math. (Basel)* **34** (1980) 97–99.
- [12] Frank O. Wagner. The Fitting subgroup of a stable group. *J. Alg.* **174** (1995), 599–609.
- [13] Frank O. Wagner. *Simple Theories*. Mathematics and Its Applications 503. Kluwer Academic Publishers, Dordrecht, 2000.

UNIVERSITÄT MÜNSTER; INSTITUT FÜR MATHEMATISCHE LOGIK UND GRUNDLAGENFORSCHUNG,
EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY
E-mail address: `daniel.palacin@uni-muenster.de`

UNIVERSITÉ DE LYON; CNRS; UNIVERSITÉ LYON 1; INSTITUT CAMILLE JORDAN UMR5208, 43
BD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE
E-mail address: `wagner@math.univ-lyon1.fr`