

Automatic detection of microemboli by means of a synchronous linear prediction technique

Sébastien Ménigot, Latifa Dreibine, Nawal Meziati, Jean-Marc Girault

▶ To cite this version:

Sébastien Ménigot, Latifa Dreibine, Nawal Meziati, Jean-Marc Girault. Automatic detection of microemboli by means of a synchronous linear prediction technique. IEEE. IEEE International Ultrasonics Symposium (IUS), Sep 2009, Rome, Italy. IEEE, pp.2371 - 2374, 2009, 10.1109/ULT-SYM.2009.5441701. hal-01075514

HAL Id: hal-01075514 https://hal.science/hal-01075514

Submitted on 17 Oct 2014 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



1. Introduction

the most commonly used systems cannot detect all of microemboli events whose energy is lower than the systolic energy, new techniques are proposed.

could detect very small microemboli. If we periodically take and compare the values of the energy of the prediction error (or autoregressive parameters) at different time points in the cardiac cycle, we can therefore detect the presence of non-periodic events such as microemboli.

2. Materials

TansCranial Doppler system (TCD)

- Velocity of blood flow through the brain's arteries
- Pulsed Doppler probe
- Examination in the temporal region



Doppler signals

- The Doppler signals are cyclo-stationary
- Synthetized Embolus Doppler signal proposed by Wedling
- severity

E-Mail: sebastien.menigot@etu.univ-tours.fr jean-marc.girault@univ-tours.fr

Automatic Detection of microemboli by means of a synchronous linear prediction technique Sébastien Ménigot, Latifa Dreibine, Nawal Meziati and Jean-Marc Girault

Úniversité François Rabelais de Tours, Tours, France INSÉRM, U930, Tours, France CNRS, ERL 3106, Tours, France



Discussion

- inaudible up to now
- error does not vary with the cardiac cycle

5. Conclusion

- Large microemboli are all detected
- incorporate "on line" technique



old	standard	SLP	
andard		<i>a</i> ₁	error
)0 %	100 %	100 %	100 %
%	32.37 %	7.49 %	5.80 %

	Gold	standard	SLP
	standard		
ן	100 %	67 %	100 %
ion	0 %	0 %	0 %

Best detection with SLP : embolus can be detected even if it was

Synchronous error autocorrelation does not improve the detection: the

Small microemboli are only detected with our new technique