



HAL
open science

Error-Correction Capability of Reed-Muller codes

Stéphanie Dib, François Rodier

► **To cite this version:**

Stéphanie Dib, François Rodier. Error-Correction Capability of Reed-Muller codes. 2014. hal-01082431v1

HAL Id: hal-01082431

<https://hal.science/hal-01082431v1>

Preprint submitted on 13 Nov 2014 (v1), last revised 2 Feb 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Error-Correction Capability of Reed-Muller codes

Stéphanie Dib, François Rodier *

Abstract

We present an asymptotic limit between correctable and uncorrectable errors on the Reed-Muller codes of any order. This limit is theoretical and does not depend of any decoding algorithm.

1 Introduction

Let \mathbb{F}_2 be the field with 2 elements, and let $RM = RM(n, r)$ be the Reed-Muller code of length 2^n and of order r that is the set of Boolean function with n variables of algebraic degree not more than r .

Decoding an error correcting code beyond half of the minimum distance has been a challenge for the one who study error correcting codes. Here we propose a theoretical bound for decoding almost all errors of Reed-Muller code on any order by a maximum likelihood decoding.

Indeed, the decoder will often be able to recover the correct codeword using an algorithm that generates for each received word the closest codeword even if the received word is more distant than half of the minimum distance. On the contrary, when the number of errors exceeds a certain value, the received vector will be rarely closer to the correct codeword than to any other one. Here we give a proof for that.

It is interesting to compare that fact with the phenomenon of concentration of the nonlinearity of Boolean functions which have been studied by several authors ([3, 4, 9, 11, 12, 13]. The r -nonlinearity of a Boolean function is its Hamming distance to the set of Boolean functions with n variables of algebraic degree not more than r .

*Aix Marseille Université, CNRS, Centrale Marseille, Institut de Mathématiques de Marseille, UMR 7373, 13288 Marseille, France, stephania.dib@gmail.com, francois.rodier@univ-amu.fr

Claude Carlet [2], proved that the density of the set of Boolean functions satisfying

$$NL_r(f) > 2^{n-1} - c\sqrt{2^{n-1} \binom{n}{r} \log 2}$$

tends to 1 when n tends to infinity, if $c > 1$. The authors of the present paper proved that this yield a concentration of the nonlinearities of almost all Boolean functions around

$$2^{n-1} - \sqrt{2^{n-1} \binom{n}{r} \log 2} \tag{1}$$

when $r \leq 2$ but missed the greater values by lack of knowledge of weight distributions ([3, 4, 11, 12]. Kai-Uwe Schmidt generalized this result for all r thanks to a result of Kaufman, Lovett, and Porat [8] helping him to find a better bound for the weights of a RM code [13].

On the other hand, in Helleseth et al. [7] the error-correction capability of Reed-Muller codes is shown for order 1 or 2 Reed-Muller codes giving the same bound as (1). Stéphanie Dib [4, Chapter 3] proved by the same method as for the concentration of the nonlinearities of almost all Boolean functions that the relation (1) was the bound for correcting most of the values of codewords for 1-order RM codes.

We show here that the value given in relation (1) is also the bound for correcting most of the values of codewords for RM codes for any order.

For RM codes, the present work improves the paper by Helleseth et al. *Error correction capability of binary linear codes* [7] where they just prove the fact that the codes $RM(n, r)$ are asymptotically optimal for $r = 1$ (cf. note after inequality (54) of [7]) or $r = 2$ (example 7 of [7]).

2 Presentation

Let $d(e, f)$ be the Hamming distance between the elements e and f in $\mathbb{F}_2^{2^n}$. We denote by $wt(e)$ the weight of an element e in $\mathbb{F}_2^{2^n}$. Let C be a linear code of length m , of dimension k . The Reed-Muller code of length 2^n and of order r has dimension $\sum_0^r \binom{n}{r}$ and minimum distance 2^{n-r} .

2.1 Correctable and uncorrectable errors

Let $\mathbb{F}_2^{2^n}$ be the set of all binary vectors of length 2^n . For any vector $f \in \mathbb{F}_2^{2^n}$, the set

$$f + C = \{f + g \mid g \in C\}$$

is called a coset of C and contains 2^k vectors. One can easily check that two cosets are either disjoint or coincide. This means

$$f \in h + C \implies f + C = h + C.$$

Therefore, the set $\mathbb{F}_2^{2^n}$ can be partitioned into 2^{2^n-k} cosets of C :

$$\mathbb{F}_2^{2^n} = \bigcup_{i=0}^{2^{2^n-k}-1} (f_i + C), \quad f_i \in \mathbb{F}_2^{2^n}$$

where $(f_i + C) \cap (f_j + C) = \emptyset$ for $i \neq j$.

If you send a word g and the decoder receive the word h , we will call $e = g - h$ the error. Thus, the possible error vectors are the vectors in the coset containing h . In maximum-likelihood decoding, the decoder's strategy is, given h , to choose a minimum weight vector f in $h + C$, and to decode h as $h - f$.

The minimum weight vector in a coset is called the coset leader, and when there is more than one vector of minimum weight in a coset, any one of them can be selected as the coset leader.

We denote the set of all coset leaders by $E_0(C)$ (note that $\#E_0(C) = 2^{2^n-k}$). The elements of $E_0(C)$ are called correctable errors, and the elements of $E_1(C) = \mathbb{F}_2^{2^n} - E_0(C)$ are called uncorrectable errors. Only coset leaders are correctable errors, which means that 2^{2^n-k} errors can be corrected with this decoding.

A codeword is an unambiguous correctable error if it is a coset leader, and it is the only vector of minimum weight in this coset.

Proposition 1 .

The following statements are equivalent.

- 1- *A codeword e is an unambiguous correctable error if and only if any of the following statements are true;*
- 2- $\forall e' \in e + C$ if $e \neq e'$ then $wt(e) < wt(e')$;
- 3- $\forall g \in C - \{0\}$, $wt(e) < wt(g + e)$;
- 4- $\forall g \in C - \{0\}$, $d(e, 0) < d(g, e)$.

Proof

The first assertion implies the second because if $e' \in e + C$ and $e \neq e'$ then e' is not the coset leader, so $wt(e) < wt(e')$.

The second assertion implies the first because if $e' \in e + C$ and $e \neq e'$ then $wt(e) < wt(e')$ so e' is not the coset leader and e is the only vector of minimum weight in this coset.

The other statement are clear.

2.2 The probability

We take $\mathbb{F}_2^{2^n}$ as the probability space.

We endow it with the uniform probability P .

3 The results

Let $\lambda_n = c \times 2^{n/2} \sqrt{2 \binom{n}{r} \log 2}$ and $\delta = 2^{n-1} - \lambda_n/2$ where c is a positive real.

We will show that if $c > 1$ then almost all error of weight smaller than δ are correctable, when n tends to infinity. And that if $c < 1$ then almost all error of weight higher than δ are uncorrectable, when n tends to infinity. More precisely we will show the following two theorems.

Theorem 1 . *Let $c > 1$. Then*

$$P_{wt(e) \leq \delta} \left(d(e, 0) < d(e, g) \text{ for all } g \in RM(r) - 0 \right) \rightarrow 1 \text{ when } n \rightarrow \infty.$$

and

Theorem 2 . *Let $c < 1$. Then*

$$P_{wt(e) \geq \delta} \left(\text{there exists } g \in RM(r) - 0 \text{ such that } d(e, 0) \geq d(e, g) \right) \rightarrow 1 \text{ when } n \rightarrow \infty.$$

4 Proof of the Theorem 1. Decoding a large number of errors

We intend to prove that almost all error of weight smaller than δ for $c > 1$ are correctable, when n tends to infinity. It is enough to prove

$$P_{wt(e) \leq \delta} \left(d(e, 0) < d(e, g) \text{ for all } g \in RM(r) - 0 \right) \rightarrow 1 \text{ when } n \rightarrow \infty.$$

We have just to show

$$P_{wt(e) \leq \delta} \left(\delta < d(e, g) \text{ for all } g \in RM(r) - 0 \right) \rightarrow 1 \text{ when } n \rightarrow \infty$$

or

$$P_{wt(e) \leq \delta} \left(\exists g \in RM(r) - 0, \delta \geq d(e, g) \right) \rightarrow 0 \text{ when } n \rightarrow \infty$$

that is

$$P_{wt(e) \leq \delta} \left(\bigcup_{g \in RM(r) - 0} \left(\delta \geq d(e, g) \right) \right) \rightarrow 0 \text{ when } n \rightarrow \infty.$$

It is enough to prove that

$$\sum_{g \in RM(r)-0} P_{wt(e) \leq \delta} (\delta \geq d(e, g)) \rightarrow 0 \text{ when } n \rightarrow \infty.$$

By expressing the conditional probabilities we have to show that

$$\sum_{g \in RM(r)-0} \frac{P\left((wt(e) \leq \delta) \cap (\delta \geq d(e, g))\right)}{P(wt(e) \leq \delta)} \rightarrow 0 \text{ when } n \rightarrow \infty.$$

Let $B_\delta(g)$ be the ball of center g and of radius δ that is the set of e such that $d(e, g) \leq \delta$. The event $B_\delta(g)$ is the set of words f in $\mathbb{F}_2^{2^n}$ such $f \in B_\delta(g)$, that is $d(f, g) \leq \delta$.

Hence Theorem 1 is a consequence of the following proposition.

Proposition 2 . *If $c > 1$ then*

$$\sum_{g \in RM(r)-0} \frac{P(B_\delta(0) \cap B_\delta(g))}{P(B_\delta(0))} \rightarrow 0 \text{ when } n \rightarrow \infty$$

Before the proof of this Proposition we have to evaluate the terms in the sum.

Lemma 1 . *For every real s , one has*

$$P(B_\delta(0) \cap B_\delta(g)) \leq \exp\left(2s^2(2^n - wt(g)) - 2s\lambda\right)$$

Proof.

Replace δ by its value.

$$\begin{aligned} P(B_\delta(0) \cap B_\delta(g)) &= P\left((wt(f) \leq \delta) \cap (wt(f+g) \leq \delta)\right) \\ &= P\left((2^{n-1} - wt(f) \geq \lambda/2) \cap (2^{n-1} - wt(f+g) \geq \lambda/2)\right) \end{aligned}$$

One knows that

$$2^n - 2wt(f) = \sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)}, \quad 2^n - 2wt(f+g) = \sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)+g(x)}.$$

Hence this gives using Markov's inequality.

$$\begin{aligned}
P(B_\delta(0) \cap B_\delta(g)) &= P\left(\left(\sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)} \geq \lambda\right) \cap \left(\sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)+g(x)} \geq \lambda\right)\right) \\
&= P\left(\left(\exp\left(s \sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)}\right) \geq \exp(s\lambda)\right) \cap \right. \\
&\quad \left.\left(\exp\left(s \sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)+g(x)}\right) \geq \exp(s\lambda)\right)\right) \\
&\leq E\left(\exp\left(s \sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)}\right) \exp\left(s \sum_{x \in \mathbb{F}_2^n} (-1)^{f(x)+g(x)}\right)\right) / \exp(s\lambda)^2
\end{aligned}$$

Since the random values $f(x)$ are independant

$$\begin{aligned}
P(B_\delta(0) \cap B_\delta(g)) &\leq E\left(\exp\left(\sum_{x \in \mathbb{F}_2^n} s(-1)^{f(x)}(1 + (-1)^{g(x)})\right)\right) / \exp(s\lambda)^2 \\
&\leq \prod_{x \in \mathbb{F}_2^n} E\left(\exp\left(s(-1)^{f(x)}(1 + (-1)^{g(x)})\right)\right) / \exp(s\lambda)^2
\end{aligned}$$

Because the random values $f(x)$ takes the values ± 1 with probability $1/2$, the calculation of the expectation gives

$$P(B_\delta(0) \cap B_\delta(g)) \leq \prod_{x \in \mathbb{F}_2^n} \cosh\left(s(1 + (-1)^{g(x)})\right) / \exp(s\lambda)^2$$

As $\cosh(t) \leq \exp(t^2/2)$

$$\begin{aligned}
P(B_\delta(0) \cap B_\delta(g)) &\leq \prod_{x \in \mathbb{F}_2^n} \exp\left(s^2(1 + (-1)^{g(x)})^2/2\right) / \exp(s\lambda)^2 \\
&\leq \exp\left(s^2\left(2^n + \sum_{\mathbb{F}_2^n} (-1)^{g(x)}\right)\right) / \exp(s\lambda)^2 \\
&\leq \exp\left(2s^2(2^n - wt(g)) - 2s\lambda\right).
\end{aligned}$$

4.1 Case where the distances are close to 2^{n-1} .

We give a bound for $P(B_\delta(0) \cap B_\delta(g))$ when the distance to 0 of the center g is rather close to 2^{n-1} .

Lemma 2 . *If*

$$|2^{n-1} - d(g, 0)| \leq 2^{n-1} / \binom{n}{r}$$

then:

$$P(B_\delta(0) \cap B_\delta(g)) \leq \frac{1}{2^{c^2 2^{\binom{n}{r}-1}}}.$$

Proof.

From lemma 1 we have

$$\begin{aligned} P(B_\delta(0) \cap B_\delta(g)) &\leq \exp\left(s^2(2^n + 2^n - 2wt(g))\right) / \exp(s\lambda)^2 \\ &\leq \exp\left(s^2 2^n \left(1 + 1/\binom{n}{r}\right)\right) / \exp(s\lambda)^2 \end{aligned}$$

We take $s = \lambda/2^n$.

$$\begin{aligned} P(B_\delta(0) \cap B_\delta(g)) &\leq \exp\left(\lambda^2 2^{-n} \left(1 + 1/\binom{n}{r}\right)\right) / \exp(\lambda^2/2^{n-1}) \\ &\leq \exp\left(2c^2 \binom{n}{r} \log 2 \left(\left(1 + 1/\binom{n}{r}\right)\right)\right) / \exp(4c^2 \binom{n}{r} \log 2) \end{aligned}$$

Simplifying the two members of this fraction by $\exp\left(2c^2 \binom{n}{r} \log 2\right)$ you get

$$P(B_\delta(0) \cap B_\delta(g)) \leq \frac{\exp(2c^2 \log 2)}{\exp(2c^2 \binom{n}{r} \log 2)} \leq \frac{2^{2c^2}}{2^{c^2 \times 2 \binom{n}{r}}}$$

4.2 Case where the distances are away from 2^{n-1} .

We use the following lemma, which is an application of a result by Kaufman, Lovett, and Porat [8].

Lemma 3 . *Let α be a strictly positive real number. The number $B_{r,n}$ of functions g in $RM(r, n)$ satisfying*

$$|wt(g) - 2^{n-1}| \geq 2^{n-1} / \binom{n}{r}$$

fulfills

$$B_{r,n} \leq 2^{\alpha \binom{n}{r}}$$

if n is large enough.

Proof

We use the relation (6) in the proof of Lemma 3 in K.-U. Schmidt's article [13].

We use this lemma to evaluate $\Pi = \sum P(B_\delta(0) \cap B_\delta(g))$ where the sum is on the nonzero g in $RM(n, r)$ fulfilling

$$|wt(g) - 2^{n-1}| \geq 2^{n-1} / \binom{n}{r}.$$

Lemma 4 . *Let α be a strictly positive real number. Then*

$$\Pi < 2^\alpha \binom{n}{r} 2^{-\frac{c^2}{1-2^{-r}} \binom{n}{r}}$$

Proof.

From lemma 1, for all s , we have

$$P(B_\delta(0) \cap B_\delta(g)) \leq \exp(2s^2(2^n - wt(g)) - 2s\lambda)$$

Let us take $s = \frac{\lambda}{2^{n+1} - 2wt(g)}$. We have, expressing the value of λ and noting that $wt(g)$ is not less than the minimum distance 2^{n-r} of $RM(n, r)$:

$$\begin{aligned} P(B_\delta(0) \cap B_\delta(g)) &\leq \exp\left(-\frac{\lambda^2}{2^{n+1} - 2wt(g)}\right) \\ &\leq \exp\left(-\frac{c^2 \times 2^{n+1} \binom{n}{r} \log 2}{2^{n+1} - 2^{n-r+1}}\right) \\ &\leq 2^{-\frac{c^2 \times \binom{n}{r}}{1-2^{-r}}}. \end{aligned}$$

Therefore

$$\Pi \leq B_{r,n} 2^{-\frac{c^2}{1-2^{-r}} \binom{n}{r}} \leq 2^\alpha \binom{n}{r} 2^{-\frac{c^2}{1-2^{-r}} \binom{n}{r}}.$$

4.3 Evaluation of $P(B_\delta(0))$

Proposition 3 . *Let r be a fixed integer, $\delta = 2^{n-1} - c \sqrt{2^{n-1} \binom{n}{r} \log 2}$ where c is a positive constant. We have*

$$P(B_\delta(0)) = \frac{1}{2\pi} \frac{2^{-c^2 \binom{n}{r}}}{2c \sqrt{\binom{n}{r} \log 2}} (1 + o(1)) \quad (2)$$

when n tends to infinity.

This is proved in Stéphanie Dib's thesis [4]. We recall briefly the proof for completeness.

The following lemma (see [2, lemma 1]) gives well-known asymptotic estimate of the sum of binomial coefficients.

Lemma 5 . Let n be a positive integer and $k \leq n$. Then

$$\sum_{i=0}^k \binom{2n}{i} < 2^{2n} \cdot \exp\left(-\frac{(n-k)^2}{n}\right).$$

When k is sufficiently close to n , the following lemma (see [6, chapter IX, (9.98)], [5, chapter VII]) gives an asymptotic estimation for $\binom{2n}{k}$:

Lemma 6 . Let n be a positive integer and $|n - k| \leq n^{\frac{5}{8}}$. Then

$$\binom{2n}{k} = \frac{2^{2n}}{\sqrt{\pi \cdot n}} \cdot \exp\left(-\frac{(n-k)^2}{n}\right) \cdot (1 + o(1)), \quad (3)$$

where the term $o(1)$ is independent of the choice of k .

Proof of the Proposition.

The number of Boolean functions whose Hamming distance to 0 is bounded from above by some number δ equals $\sum_{0 \leq i \leq \delta} \binom{2^n}{i}$. Thus we have

$$\sum_{0 \leq i \leq \delta} \binom{2^n}{i} = \sum_{0 \leq i < 2^{n-1} - 2^{(n-1)\frac{5}{8}}} \binom{2^n}{i} + \sum_{2^{n-1} - 2^{(n-1)\frac{5}{8}} \leq i \leq \delta} \binom{2^n}{i}.$$

The first sum on the right hand side is taken care of by lemma 5 which will show that it is negligible with respect of the second sum.

To estimate a lower bound of the second sum (which we denote S), we use (3)

$$S = \frac{2^{2^n}}{\sqrt{\pi} 2^{n-1}} \cdot (1 + o(1)) \cdot \sum_{2^{n-1} - 2^{(n-1)\frac{5}{8}} \leq i \leq \delta} \exp\left(-\frac{(2^{n-1} - i)^2}{2^{n-1}}\right).$$

We use that the function in the sum is monotonous to replace the sum by an integral.

$$\begin{aligned} S &= \frac{2^{2^n}}{\sqrt{\pi} 2^{n-1}} \cdot (1 + o(1)) \cdot \int_{2^{n-1} - 2^{(n-1)\frac{5}{8}} + 1 \leq i \leq \delta} \exp\left(-\frac{(2^{n-1} - i)^2}{2^{n-1}}\right) di. \\ &= \frac{2^{2^n}}{\sqrt{\pi}} \cdot (1 + o(1)) \cdot \int_c \sqrt{\binom{n}{r} \log 2} \leq v \leq 2^{\frac{n-1}{8}} - 2^{\frac{1-n}{2}} \exp(-v^2) dv. \end{aligned}$$

By [5, chapter VII, Lemma 2] and the fact that

$$c^2 \binom{n}{r} \log 2 - \left(2^{\frac{n-1}{8}} - 2^{\frac{1-n}{2}}\right)^2 \rightarrow -\infty$$

which implies that

$$\int_{2^{\frac{n-1}{8}} - 2^{\frac{1-n}{2}}}^{\infty} \exp(-v^2) dv = o\left(\int_{c\sqrt{\binom{n}{r}\log 2}}^{\infty} \exp(-v^2) dv\right)$$

the last integral is equivalent to

$$\frac{\exp\left(-c^2 \binom{n}{r} \log 2\right)}{2c\sqrt{\binom{n}{r}\log 2}} = \frac{2^{-c^2 \binom{n}{r}}}{2c\sqrt{\binom{n}{r}\log 2}}.$$

Thus

$$\sum_{0 \leq i \leq \delta} \binom{2^n}{i} = \frac{2^{2^n}}{\sqrt{\pi}} \frac{2^{-c^2 \binom{n}{r}}}{2c\sqrt{\binom{n}{r}\log 2}} (1 + o(1)).$$

4.4 Proof of Theorem 1

Therefore

$$\begin{aligned} \sum_{g \in RM(r)-0} \frac{P(B_\delta(0) \cap B_\delta(g))}{P(B_\delta(0))} \\ \leq O(n^{r/2}) \left(2^{\binom{n}{r}} 2^{-c^2 2 \left(\binom{n}{r} - 1\right)} 2^{c^2 \binom{n}{r}} + 2^\alpha \binom{n}{r} 2^{-\frac{c^2}{1-2^{-r}} \binom{n}{r}} 2^{c^2 \binom{n}{r}} \right). \end{aligned}$$

This tends to 0 because the exponent of 2 is, for the left term

$$\binom{n}{r} - c^2 2 \left(\binom{n}{r} - 1\right) + c^2 \binom{n}{r} = -\binom{n}{r} c^2 + 2c^2 \rightarrow -\infty$$

and for the right term

$$\alpha \binom{n}{r} - \frac{c^2}{1-2^{-r}} \binom{n}{r} + c^2 \binom{n}{r} = \binom{n}{r} \left(\alpha - \frac{2^{-r} c^2}{1-2^{-r}} \right).$$

So just take

$$\alpha < \frac{2^{-r} c^2}{1-2^{-r}}$$

so that this term tends to $-\infty$.

5 The error correction capability function

Let $\epsilon_C(t)$ the ratio of the number of errors correctables of weight t to the number of possible errors of weight t . An important property of this ratio is that for any t in the range from half the minimum distance to the covering radius, $\epsilon_C(t)$ decreases with the growing t as the next lemma says.

Lemma 7 .

For any $[n, k]$ code C and any $t = 0, 1, \dots, n - 1$

$$\epsilon_C(t + 1) \leq \epsilon_C(t)$$

with strict inequality for $t_C \leq t \leq r_C$ where we set $t_C = \lfloor (d_C - 1)/2 \rfloor$ and denote the covering radius of C by r_C .

Proof

See Helleseth et al. [7, Lemma 2]. This property is due to the fact that the sets of correctable and uncorrectable errors form a monotone structure, (see, for example, [10, p. 58, Theorem 3.11]) and a result of Bollobas about shadows [1, Theorem 3]

5.1 A corollary of Theorem 1

Corollary 1 . If $c > 1$, then $\epsilon_C(t) \rightarrow 1$ when $n \rightarrow \infty$.

Proof.

Suppose that $\epsilon_C(t_c) \not\rightarrow 1$. Then there exists $\eta < 1$ such that $\epsilon_C(t_c) < \eta$ for an infinity of n .

For an $RM(n, r)$ code, let

$$\frac{2^{2^n}}{\sqrt{\pi}} \frac{2^{-c^2} \binom{n}{r}}{2c \sqrt{\binom{n}{r} \log 2}} = A(c)$$

For $c_1 > c_2 > 1$ fixed, the area of weights between c_1 and c_2 contains only a proportion η of correctable words as the function ϵ_C decreases. From Proposition 3 there are about

$$\sum_{0 \leq i \leq \delta_1} \binom{2^n}{i} = A(c_1)(1 + o(1)).$$

words of weight in $[0, c_1]$ and in the area of weight between c_1 et c_2

$$\sum_{\delta_1 \leq i \leq \delta_2} \binom{2^n}{i} = A(c_2)(1 + o(1)).$$

As $A(c_1) = o(A(c_2))$ there are at most

$$A(c_1)(1 + o(1)) + \eta A(c_2)(1 + o(1)) = \eta A(c_2)(1 + o(1))$$

correctable words of weights $[0, c_2]$, which shows that it is impossible that almost all words are correctable as says Theorem 1.

5.2 Proof of the Theorem 2. An asymptotic decoding upper bound

In the case of RM codes we have a simplification of the proof of the Theorem 3 b in [7].

For Reed-Muller codes of order r , that is to say $RM(n, r)$ we take

$$t = 2^{n-1} - c\sqrt{2^{n-1} \binom{n}{r} \ln 2}.$$

Proposition 4 . *If $c < 1$, then $\epsilon_C(t) \rightarrow 0$ when $n \rightarrow \infty$.*

For every t , one has (cf. Lemma 3 of [7])

$$\begin{aligned} \epsilon_C(t) \times \sum_{i=0}^t \binom{2^n}{i} &\leq \sum_{i=0}^t \epsilon_C(i) \binom{2^n}{i} \\ &= \text{number of correctable errors of weight smaller than } t \\ &\leq \text{total number of correctable errors} \\ &\leq 2^{2^n - k}. \end{aligned}$$

We have, from Proposition 3

$$\sum_{i=0}^t \binom{2^n}{i} = \#B_t = \frac{2^{2^n - c^2 \binom{n}{r}}}{2c\sqrt{\pi \binom{n}{r} \ln 2}}(1 + o(1)).$$

Whence

$$\begin{aligned} \epsilon_C(t) &\geq \frac{2^{2^n - k}}{\sum_{i=0}^t \binom{2^n}{i}} \\ &= \frac{2c\sqrt{\pi \binom{n}{r} \ln 2}}{2^{2^n - c^2 \binom{n}{r}} \times 2^{\sum_{i=0}^r \binom{n}{i} - 2^n}}(1 + o(1)) \\ &= \frac{2c\sqrt{\pi \binom{n}{r} \ln 2}}{2^{\sum_{i=0}^r \binom{n}{i} - c^2 \binom{n}{r}}}(1 + o(1)). \end{aligned}$$

If $c < 1$, when $n \rightarrow \infty$ then the denominator tends toward infinity, so $\epsilon_C(t) \rightarrow 0$.

Remark.

The theorem is still true if we take $t = 2^{n-1} - c\sqrt{2^{n-1}k \ln 2}$ with $c < 1$ if $k \leq 2^{(n-1)/4}$ (to be able to use Proposition 3).

5.3 End of the proof of Theorem 2.

If $wt(t) > \delta$, then $\epsilon(t) \rightarrow 0$. The proportion of correctable errors of weight t tends to 0, whence so is the proportion of correctable errors of weight greater than t . Therefore Theorem 2 is true.

5.4 Asymptotically optimality of RM codes

A sequence $(C_m)_m$ of $[m, k]$ codes where $k = o(m)$ as $m \rightarrow \infty$ is called asymptotically optimal if for any fixed ϵ , $0 < \epsilon < 1$

$$m - 2t_{C_m}(\epsilon) \sim \sqrt{mk \ln 4}$$

where the error correction capability function $t_C(\epsilon)$ is the maximum t such that $\epsilon_C(t) \geq \epsilon$.

Theorem 3 . *The sequence $RM(n, r)_n$ is asymptotically optimal.*

Proof.

Let us take ϵ and try to find $t_C(\epsilon)$. Let $c < 1$.

If $2t = 2^n - c\sqrt{2^n k \log 4}$ then the words of weight t are almost all uncorrectable, therefore $\epsilon_{C_{2^n}}(t) \simeq 0$. And we have $\epsilon_{C_{2^n}}(t) < \epsilon$ for n big enough (and consequently $t \geq t_{C_{2^n}}(\epsilon)$).

As a result $t_{C_{2^n}}(\epsilon) \leq 2^{n-1} - c\sqrt{2^{n-1}k \log 4}$.

Let now $c > 1$.

If $2t = 2^n - c\sqrt{2^n k \log 4}$ then the words of weight t are almost all correctable, therefore $\epsilon_{C_{2^n}}(t) \simeq 1$. And we have $\epsilon_{C_{2^n}}(t) > \epsilon$ for n big enough (and consequently $t < t_{C_{2^n}}(\epsilon)$).

As a result, if $c_1 < 1 < c_2$, one has

$$2^n - c_2\sqrt{2^n k \log 4} < 2t_{C_{2^n}}(\epsilon) < 2^n - c_1\sqrt{2^n k \log 4}$$

or

$$c_1\sqrt{2^n k \log 4} < 2^n - 2t_{C_{2^n}}(\epsilon) < c_2\sqrt{2^n k \log 4}$$

or

$$c_1 < \frac{2^n - 2t_{C_{2^n}}(\epsilon)}{\sqrt{2^n k \log 4}} < c_2.$$

As c_1 and c_2 may be as close to 1 as we wish, we have

$$\frac{2^n - 2t_{C_{2^n}}(\epsilon)}{\sqrt{2^n k \log 4}} \rightarrow 1 \quad \text{quand } n \rightarrow \infty.$$

References

- [1] B. Bollobas: *Combinatorics: Set Systems, Hypergraphs, Families of Vectors, and Combinatorial Probability*. Cambridge, U.K.: Cambridge Univ. Press, 1986.
- [2] C. Carlet: The complexity of boolean functions from cryptographic viewpoint. In Matthias Krause, Pavel Pudlk, Rüdiger Reischuk, and Dieter van Melkebeek, editors, *Complexity of Boolean Functions*, volume 06111 of Dagstuhl Seminar Proceedings. Internationales Begegnungs- und Forschungszentrum fuer Informatik (IBFI), Schloss Dagstuhl, Germany, 2006.
- [3] S. Dib: Distribution of boolean functions according to the second-order nonlinearity. In M. Anwar Hasan and Tor Hellesteth, editors, *WAIFI*, volume 6087 of *Lecture Notes in Computer Science*, pages 8696. Springer, 2010.
- [4] S. Dib: Thèse. *Distribution de la non-linéarité des fonctions Booléennes*. Université d'Aix-Marseille, 2013.
- [5] W. Feller: *An introduction to probability theory and its applications*. Vol. I. Third edition. John Wiley & Sons Inc., New York, 1968.
- [6] R. L. Graham, D. E. Knuth, and O. Patashnik: *Concrete mathematics*. Addison-Wesley Publishing Company Advanced Book Program, Reading, MA, 1989. A foundation for computer science.
- [7] T. Hellesteth, T. Kløve, and Vladimir I. Levenshtein: Error correction capability of binary linear codes. *IEEE Trans. Inform. Theory*, 51(4):14081423, 2005.
- [8] T. Kaufman, S. Lovett, E. Porat: Weight distribution and list-decoding size of Reed-Muller codes. *IEEE Trans. Inform. Theory* 58 (2012), no. 5, 26892696.
- [9] S. Litsyn and A. Shpunt.: On the distribution of Boolean function nonlinearity. *SIAM J. Discrete Math.*, 23(1):79 95, 2008/09.

- [10] W. W. Peterson and E. J. Weldon Jr. Error-Correcting Codes. Cambridge, MA: MIT Press, 1972.
- [11] F. Rodier: Sur la non-linéarité des fonctions booléennes, Acta Arithmetica, vol 115, (2004), 1-22, ArXiv : math.NT/0306395.
- [12] F. Rodier: Asymptotic nonlinearity of Boolean functions, Designs, Codes and Cryptography, 40 :1 2006.
- [13] K. U. Schmidt: Nonlinearity measures of random Boolean functions, preprint, 2013, arXiv:1308.3112.