



HAL
open science

Asymptotic equivalence of discretely observed diffusion processes and their Euler scheme: small variance case

Ester Mariucci

► **To cite this version:**

Ester Mariucci. Asymptotic equivalence of discretely observed diffusion processes and their Euler scheme: small variance case. 2014. hal-01082603v1

HAL Id: hal-01082603

<https://hal.science/hal-01082603v1>

Preprint submitted on 13 Nov 2014 (v1), last revised 3 Mar 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymptotic equivalence of discretely observed diffusion processes and their Euler scheme: small variance case

Ester Mariucci*

*Laboratoire LJK, Université Joseph Fourier UMR 5224
51, Rue des Mathématiques, Campus de Saint Martin d'Hères
BP 53 38041 Grenoble Cedex 09*

Abstract

This paper establishes the global asymptotic equivalence, in the sense of the Le Cam Δ -distance, between scalar diffusion models with unknown drift function and small variance on the one side, and nonparametric autoregressive models on the other side. The time horizon T is kept fixed and both the cases of discrete and continuous observation of the path are treated. We allow non constant diffusion coefficient, bounded but possibly tending to zero. The asymptotic equivalences are established by constructing explicit equivalence mappings.

KEYWORDS. Nonparametric experiments, deficiency distance, asymptotic equivalence, diffusion processes, autoregression.

AMS 2010 subject classification: Primary 62B15; Secondary 62G20, 60G51.

1 Introduction

Diffusion processes obtained as small random perturbations of deterministic dynamical systems have been widely studied and have proved fruitful in applied problems (see e.g. [15]). Among other subjects, they have been applied to contingent claim pricing, see [44] and the references therein, to filtering problems, see e.g. [39, 40] more recently to epidemic data [23]. From a statistical point of view, these models have first been considered by Kutoyants [29] in the framework of continuous observation on a fixed time interval $[0, T]$. However, statistical inference for discretely observed diffusion processes has first been treated several years after, see [16]. In a nonparametric framework we may quote [28], among many others.

In this paper we consider the problem of estimating the drift function f associated with a scalar diffusion process (y_t) continuously or discretely observed on a time interval $[0, T]$, with $T < \infty$ kept fixed. More precisely, we consider the one-dimensional diffusion process (y_t) given by

$$dy_t = f(y_t)dt + \varepsilon\sigma(y_t)dW_t, \quad t \in [0, T], \quad y_0 = w \in \mathbb{R}, \quad (1)$$

*Ester.Mariucci@imag.fr

where $(W_t)_{t \geq 0}$ is a standard $(\mathcal{A}_t)_{t \geq 0}$ -brownian motion defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$. The diffusion coefficient $\varepsilon\sigma(\cdot)$, with $0 < \varepsilon < 1$, is supposed to be known and satisfying the following conditions:

(H1) $\sigma(\cdot)$ is a K -Lipschitz function on \mathbb{R} bounded away from infinity and zero, i.e. there exist strictly positive constants σ_0, σ_1, K with

$$\sigma_0^2 \leq \sigma^2(y) \leq \sigma_1^2 \quad \text{and} \quad |\sigma(z) - \sigma(y)| \leq K|z - y|, \quad \forall z, y \in \mathbb{R}. \quad (2)$$

When (y_t) is discretely observed we will also require the following assumption:

(H2) $\sigma(\cdot)$ is a differentiable function on \mathbb{R} with K -Lipschitz derivative, i.e.

$$|\sigma'(z) - \sigma'(y)| \leq K|z - y| \quad \forall z, y \in \mathbb{R}.$$

More in details, we consider two experiments, the continuous one associated with (y_t) and the discrete one given by the observations $(y_{t_1}, \dots, y_{t_n})$, where $t_i = \frac{i}{n}T$. Our aim is to prove that these nonparametric experiments are both equivalent to an autoregressive model given by Euler type discretizations of y with sampling interval $1/n$, $n \in \mathbb{N}^*$:

$$Z_0 = w, \quad Z_i = Z_{i-1} + \frac{f(Z_{i-1})}{n} + \varepsilon\sigma(Z_{i-1})\xi_i, \quad i = 1, \dots, n, \quad (3)$$

with independent standard normal variables ξ_i .

The concept of asymptotic equivalence that we shall adopt is based on the Le Cam Δ -distance between statistical experiments. Roughly speaking, saying that two statistical models, or experiments, are equivalent in the Le Cam's sense means that any statistical inference procedure can be transferred from one model to the other in such a way that the asymptotic risk remains the same, at least for bounded loss functions. One can use this property in order to obtain asymptotic results working in a simpler but equivalent setting. For the basic concepts and a detailed description of the notion of asymptotic equivalence, we refer to [31, 32]. A short review of this topic will be given in Appendix 4.

In recent years, the Le Cam's theory on the asymptotic equivalence between statistical models has aroused great interest and a large number of works has been published on this subject. In parametric statistics, Le Cam's theory has successfully been applied to a huge variety of experiments. Proving an asymptotic equivalence for nonparametric experiments is more demanding but, nowadays, several works in this subject have appeared. The first results of global asymptotic equivalence for nonparametric experiments date from 1996 and are due to Brown and Low [3] and Nussbaum [37]. A non-exhausting list of subsequent works in this domain includes [1, 21, 43, 8, 7, 41, 9, 35] for nonparametric regression, [6, 26, 2] for nonparametric density estimation models, [20] for generalized linear models, [22] for time series, [5] for GARCH model, [34] for functional linear regression, [19] for spectral density estimation and [33] for inhomogeneous jumps diffusion models. Negative results are somewhat harder to come by; the most notable ones among them are [14, 4, 46].

Asymptotic equivalence results have also been obtained for diffusion models. References concern nonparametric drift estimation with known diffusion coefficient. Among these one can quote [13, 18, 11, 12, 42]. However, the most relevant results to our purposes are due to Milstein and Nussbaum [36] and to Genon-Catalot and Larédo [17]. The former authors have shown the asymptotic equivalence of a diffusion process continuously observed until time $T = 1$ having unknown drift function and constant small known diffusion coefficient, with the corresponding Euler scheme. They also proved the asymptotic sufficiency of the discretized observation of the diffusion with small sampling interval. Hence, our work is a generalization of [36]. It can also be seen as a complement to [17], the difference being that in our case the time horizon is kept fixed and the

diffusion coefficient goes to zero. This setting allows for weaker hypotheses than those assumed by Genon-Catalot and Larédo (for example, we do not need the drift function f to be uniformly bounded).

The interest in proving the asymptotic equivalence between the statistical model associated with the discretization of (1) and (3) lies in the difficulty of making inferences in the discretely observed diffusion model. On the other hand, inference for model (3) is well understood and in practice one often reduces to working with the latter (see e.g. [16, 30, 24, 10]). The result in the present paper can thus be seen as a theoretical justification for such a practice.

The scheme of the proof is to prove both an asymptotic equivalence between the continuous and the discrete observation of (1) and one between the continuous model (1) and the Euler scheme (3). By the triangular inequality, the result will follow. The main difficulty lies in the model (3) being equivalent to a diffusion process with a diffusion coefficient $\bar{\sigma}$ different from σ . In particular, this means that the total variation distance between (3) and (1) is always 1. Thus, to prove the equivalence between these models it is necessary to construct an appropriate randomization. This is made possible by using random time changed experiments. Indeed, one can use random time changes in order to reduce to new diffusion models with diffusion coefficient equal to ε . However, these randomizations do not allow to apply the result of Milstein and Nussbaum directly since the changes of clock oblige to observe the new diffusion processes until different random times. Some care is then needed to overcome this technical obstacle (see Lemma 3.10).

The paper is organized as follows: Section 2 contains the notations and the statement of the main results while Section 3 is devoted to the proofs. Appendix 4 is devoted to background material.

2 Assumptions and notations

To formulate our results we need to assume the standard conditions for existence and uniqueness of a strong solution y for the SDE (1) ([38], Theorem 5.5, page 45). We shall thus work with parameter spaces included in \mathcal{F}_M , the set of all functions f defined on \mathbb{R} and satisfying

$$|f(0)| \leq M \text{ and } |f(z) - f(y)| \leq M|z - y|, \quad \forall z, y \in \mathbb{R}. \quad (4)$$

In particular, observe that every element of \mathcal{F}_M satisfies a condition of linear growth: $|f(z)| \leq M(1 + |z|)$, $\forall z \in \mathbb{R}$. Let $C = C(\mathbb{R}^+, \mathbb{R})$ be the space of continuous mappings ω from \mathbb{R}^+ into \mathbb{R} . Define the *canonical process* $x : C \rightarrow C$ by

$$\forall \omega \in C, \quad x_t(\omega) = \omega_t, \quad \forall t \geq 0.$$

Let \mathcal{C}^0 be the smallest σ -algebra of parts of C that makes x_s , $s \geq 0$, measurable. Further, for any $t \geq 0$, let \mathcal{C}_t^0 be the smallest σ -algebra that makes x_s , s in $[0, t]$, measurable. Finally, set $\mathcal{C}_t := \bigcap_{s>t} \mathcal{C}_s^0$ and $\mathcal{C} := \sigma(\mathcal{C}_t; t \geq 0)$. Let us denote by $P_f^{n,y}$ the distribution induced on (C, \mathcal{C}_T) by the law of y , solution to (1) and by $Q_f^{n,y}$ the distribution defined on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$ by the law of $(y_{t_1}, \dots, y_{t_n})$, $t_i = T \frac{i}{n}$. We call \mathcal{P}_y^T the experiment associated with the continuous observation of y until the time T and \mathcal{Q}_y^n the discrete one, based on the grid values of y :

$$\mathcal{P}_y^T = (C, \mathcal{C}_T, \{P_f^y, f \in \mathcal{F}\}), \quad (5)$$

$$\mathcal{Q}_y^n = (\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n), \{Q_f^{n,y}, f \in \mathcal{F}\}). \quad (6)$$

Finally, let us consider the experiment associated with the Euler scheme corresponding to (1). We denote by $Q_f^{n,Z}$ the distribution of $(Z_i, i = 1, \dots, n)$ defined by (3). Then:

$$\mathcal{Q}_Z^n = (\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n), \{Q_f^{n,Z}, f \in \mathcal{F}\}). \quad (7)$$

Let us now state our main results.

Theorem 2.1. *Suppose that for some $M > 0$ the parameter space \mathcal{F} fulfills $\mathcal{F} \subset \mathcal{F}_M$ and that $\sigma(\cdot)$ satisfies Assumption (H1) with $K = M$. Then, if $\varepsilon n \rightarrow \infty$ as $n \rightarrow \infty$ and $\varepsilon \rightarrow 0$, the experiments \mathcal{P}_y^T and \mathcal{Q}_Z^n are asymptotically equivalent. More precisely we have*

$$\Delta(\mathcal{P}_y^T, \mathcal{Q}_Z^n) = O\left(\frac{1}{\varepsilon n} + (n^{-1} + \varepsilon)^{1/4}\right).$$

Theorem 2.2. *Suppose that for some $M > 0$ the parameter space \mathcal{F} fulfills $\mathcal{F} \subset \mathcal{F}_M$ and that $\sigma(\cdot)$ satisfies assumptions (H1) and (H2), with $K = M$. Then, the sampled values y_{t_1}, \dots, y_{t_n} are an asymptotically sufficient statistic for the experiment \mathcal{P}_y^T .*

Corollary 2.3. *Under the same hypotheses as in Theorem 2.2, the statistical model associated with the sampled values y_{t_1}, \dots, y_{t_n} is asymptotically equivalent to \mathcal{Q}_Z^n , as n goes to infinity.*

Remark 2.4. The conditions on ε and n of Theorem 2.1 are the same as in [36].

3 Proofs

3.1 Random time substitutions for Markov processes

A key tool in establishing the asymptotic equivalence between the diffusion model continuously observed and its Euler scheme is given by random time changes for Markov processes. More in details we will need the following results.

Theorem 3.1. *(see [45]) Let (Y, \mathbb{P}_y) be a (càdlàg) strong (\mathcal{A}_t) -Markov process on $(\Omega, \mathcal{A}, \mathbb{P})$ with state space $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ and let $v : \mathbb{R}^d \rightarrow (0, \infty)$ be a positive continuous function. Define the additive functional*

$$F_t = \int_0^t \frac{ds}{v(Y_s)}, \quad t \geq 0$$

and assume that

$$\int_0^\infty \frac{ds}{v(Y_s)} = \infty, \quad \mathbb{P}_y - \text{a.s.}, \quad \forall y \in \mathbb{R}^d,$$

so that the right continuous inverse

$$T_t = \inf\{s \geq 0 : F_s > t\}, \quad t \geq 0$$

of the functional F is well defined on $[0, \infty)$. Then the process

$$J_t = Y_{T_t}, \quad t \geq 0,$$

is a càdlàg strong (\mathcal{A}_{T_t}) -Markov process on the probability space $(\Omega, \mathcal{A}, \mathbb{P})$.

Assume moreover that (Y, \mathbb{P}_y) is a Feller process with infinitesimal generator \mathcal{L}^Y and domain \mathcal{D} . Then J is also a Feller process whose infinitesimal generator, with domain \mathcal{D} , is given by

$$\mathcal{L}^J h(z) = v(z) \mathcal{L}^Y h(z), \quad h \in \mathcal{D}, \quad z \in \mathbb{R}^d.$$

Property 3.2. For all $\omega \in C$, $s, t > 0$ define:

$$\begin{aligned}\rho_s(\omega) &= \int_0^s \sigma^2(\omega_r) dr; & \eta_t(\omega) &= \inf \{s \geq 0, \rho_s(\omega) \geq t\}, \\ \theta_s(\omega) &= \int_0^s \frac{1}{\sigma^2(\omega_r)} dr; & A_t(\omega) &= \inf \{s \geq 0, \theta_s(\omega) \geq t\}.\end{aligned}$$

Then, the following hold:

1. $\rho_T(x) = A_T(x_{\eta_T(x)})$,
2. $A_t(x) = \int_0^t \sigma^2(x_{A_s(x)}) ds, \quad \forall t \in [0, T]$.

Proof. 1. It is enough to show that $\theta_T(x_{\eta_T(x)}) = \eta_T^n(x)$ since $t \mapsto A_t(x)$ and $t \mapsto \rho_t(x)$ are, respectively, the inverses of the applications $t \mapsto \theta_t(x)$ and $t \mapsto \eta_t(x)$. To prove the last assertion compute:

$$\theta_T(x_{\eta_T(x)}) = \int_0^T \frac{1}{\sigma^2(x_{\eta_r(x)})} dr = \int_0^{\eta_T(x)} \frac{\sigma^2(x_s)}{\sigma^2(x_s)} ds = \eta_T(x);$$

where in the second equality we have performed the change of variable $s = \eta_r(x) \Leftrightarrow r = \rho_s(x)$ that yields $dr = \sigma^2(x_s) ds$.

2. Again, we use that $t \mapsto \theta_t(x)$ is the inverse of the function $t \mapsto A_t(x)$ combined with the following elementary fact:

Let h and g be two differentiable functions on \mathbb{R} such that $h(0) = 0 = g(0)$ and their derivatives never vanish. Then, $h'(z) = \frac{1}{g'(h(z))}$ for all z in \mathbb{R} if and only if h is the inverse of g .

To show the assertion in 2. it is enough to apply this fact to $h(t) = A_t(x)$ and $g(t) = \theta_t(x)$. \square

3.2 Proof of Theorem 2.1

We will proceed in four steps. More precisely, in Step 1 we consider a random time change on the diffusion (1) in order to obtain an experiment equivalent to \mathcal{P}_y^T but associated with a diffusion having diffusion coefficient equal to ε . In Step 2 we build a continuous time discretization of the process (y_t) and, applying a second random time change, we prove an equivalence result between a second experiment associated again with a diffusion having diffusion coefficient equal to ε . In Step 3 we compare, in term of the Le Cam Δ -distance, the two experiments with the diffusion coefficient equal to ε constructed in Steps 1-2. Finally, in Step 4, we prove the equivalence between the experiment associated with the continuous time discretization of (y_t) and the one with the Euler scheme. By means of the triangular inequality we are able to bound the Le Cam Δ -distance between \mathcal{P}_y^T and \mathcal{Q}_Z^n .

Step 1. We start by proving the Le Cam equivalence between \mathcal{P}_y^T and a corresponding diffusion model with coefficient diffusion equal to ε . Recall that P_f^y is the law on (C, \mathcal{C}_T) of a diffusion process with infinitesimal generator \mathcal{L}_1 given by

$$\mathcal{L}_1 = f \nabla + \varepsilon^2 \frac{\sigma^2}{2} \Delta, \tag{8}$$

Define P_f^ε as the law on (C, \mathcal{C}) of a diffusion process with infinitesimal generator \mathcal{L}_2 given by

$$\mathcal{L}_2 = \frac{f}{\sigma^2} \nabla + \varepsilon^2 \frac{1}{2} \Delta \tag{9}$$

and initial condition $\xi_0 = \omega$. Moreover, for all $A > 0$ a \mathcal{C}_t -stopping time, define the experiment

$$\mathcal{P}_\xi^A = (C, \mathcal{C}_A, (P_f^\xi|_{\mathcal{C}_A}, f \in \mathcal{F})).$$

Proposition 3.3. *Suppose that for some $M > 0$ the parameter space \mathcal{F} fulfills $\mathcal{F} \subset \mathcal{F}_M$ and that $\sigma(\cdot)$ satisfies assumption (H1), with $K = M$. Then, the statistical models \mathcal{P}_y^T and $\mathcal{P}_\xi^{A_T(x)}$ are equivalent.*

Proof. Let us prove that $\delta(\mathcal{P}_y^T, \mathcal{P}_\xi^{A_T(x)}) = 0$. Note that (x_t) under P_f^y is a (\mathcal{C}_t) -Markov process with infinitesimal generator as in (8). Define a new process ξ as a change of time of (x_t) with stochastic clock $(\eta_t(x))_t$: $\xi_0 = \omega$, $\xi_t := x_{\eta_t(x)}$, $\forall t > 0$. Theorem 3.1 ensures that the process $(\xi_t)_{t \geq 0}$ is a diffusion process with infinitesimal generator given by (9). Also, remark that, as (x_t) is defined on $[0, T]$, then the trajectories of (ξ_t) are defined until the time $\rho_T(x)$. In order to produce a randomization transforming the family of measures $\{P_f^y, f \in \mathcal{F}\}$ in $\{P_f^\xi|_{\mathcal{C}_{A_T(x)}}, f \in \mathcal{F}\}$, let us consider the following application:

$$\Phi : \{\omega_t : t \in [0, T]\} \rightarrow \{\omega_{\eta_t(\omega)} : t \in [0, \rho_T(\omega)]\}.$$

Observe that the process $\Phi(x)$ is defined until the time $\rho_T(x)$ that is equal to $A_T(\Phi(x))$ (see Property 3.2), so that any set of paths of $\Phi(x)$ belongs to $\mathcal{C}_{A_T(x)}$. Introduce the Markov kernel K defined by $K(\omega, \Gamma) = \mathbb{I}_\Gamma(\Phi(\omega))$, $\forall \omega \in C, \forall \Gamma \in \mathcal{C}_{A_T(x)}$, then:

$$K P_f^y(\Gamma) = \int \mathbb{I}_\Gamma(\Phi(\omega)) P_f^y(d\omega) = P_f^y(\Phi(x) \in \Gamma) = P_f^\xi|_{\mathcal{C}_{A_T(x)}}(\Gamma).$$

Therefore $\delta(\mathcal{P}_y^T, \mathcal{P}_\xi^{A_T(x)}) = 0$.

The same type of computations imply that $\delta(\mathcal{P}_\xi^{A_T(x)}, \mathcal{P}_y^T) = 0$ through use of the application $\Psi : (\omega_t : t \in [0, A_T(\omega)]) \rightarrow (\omega_{A_t(\omega)} : t \in [0, T])$. \square

Step 2. We now introduce a statistical model that approximates the model \mathcal{P}_y^T . Given a path ω in C and a time grid $t_i = T \frac{i}{n}$, we define

$$\bar{f}_n(t, \omega) = \sum_{i=1}^{n-1} f(\omega(t_i)) \mathbb{I}_{[t_i, t_{i+1})}(t), \quad \bar{\sigma}_n(t, \omega) = \sum_{i=1}^{n-1} \sigma(\omega(t_i)) \mathbb{I}_{[t_i, t_{i+1})}(t), \quad \forall t \in [0, T].$$

Then, we denote by $P_f^{n, \bar{y}}$ the law on (C, \mathcal{C}_T) of a diffusion process with infinitesimal generator $\bar{\mathcal{L}}^n$ given by

$$\bar{\mathcal{L}}_t^n(\omega) h(z) = \bar{f}_n(t, \omega) \nabla h(z) + \varepsilon^2 \frac{\bar{\sigma}_n^2(t, \omega)}{2} \Delta h(z), \quad \forall \omega \in C, h \in C^2(\mathbb{R}), z \in \mathbb{R} \quad (10)$$

and initial condition $\bar{y}_0 = \omega$. Consider the experiment

$$\mathcal{P}_{\bar{y}}^{n, T} = (C, \mathcal{C}_T, (P_f^{n, \bar{y}}|_{\mathcal{C}_T}, f \in \mathcal{F}))$$

Again, we want to introduce the diffusion model with diffusion coefficient equal to ε associated to $\mathcal{P}_{\bar{y}}^{n, T}$. To that aim, for all $\omega \in C$, define

$$\bar{A}_0^n(\omega) = 0, \quad \bar{A}_t^n(\omega) = \bar{A}_{t_{i-1}}(\omega) + \sigma^2(\omega_{\bar{A}_{t_{i-1}}(\omega)})(t - t_{i-1}), \quad t \in (t_{i-1}, t_i]; \quad (11)$$

$$\bar{g}_n(t, \omega) = \sum_{i=1}^n \frac{f}{\sigma^2}(\omega_{\bar{A}_{t_i}^n(\omega)}) \mathbb{I}_{(\bar{A}_{t_i}^n(\omega), \bar{A}_{t_{i+1}}^n(\omega)]}(t), \quad t \geq 0, \quad i = 0, \dots, n-1. \quad (12)$$

Lemma 3.4. $\bar{A}_{t_i}^n(x)$ is a \mathcal{C}_t -stopping time for all $i = 1, \dots, n$.

Proof. By (11), $\bar{A}_{t_1}^n(x) = \sigma^2(x(0))t_1$, so the set $\{\bar{A}_{t_1}^n(x) \leq t\} = \begin{cases} \emptyset & \text{if } \sigma^2(x(0))t_1 > t \\ C & \text{otherwise.} \end{cases}$

belongs to \mathcal{C}_t , for all t . By induction, assume that $\bar{A}_{t_{i-1}}^n(x)$ is a (\mathcal{C}_t) -stopping time and remark that (11) implies $\{\bar{A}_{t_i}^n(x) \leq t\} = \{\bar{A}_{t_i}^n(x) \leq t\} \cap \{\bar{A}_{t_{i-1}}^n(x) \leq t\}$. Since (x_t) is (\mathcal{C}_t) -adapted and continuous, in particular it is progressively measurable with respect to (\mathcal{C}_t) . By the induction hypothesis it follows that $x_{\bar{A}_{t_{i-1}}^n(x)}$ is $\mathcal{C}_{\bar{A}_{t_{i-1}}^n(x)}$ -measurable, hence, using (11), $\{\bar{A}_{t_i}^n(x) \leq t\} \in \mathcal{C}_{\bar{A}_{t_{i-1}}^n(x)}$, as $\bar{A}_{t_{i-1}}^n(x)$ is already $\mathcal{C}_{\bar{A}_{t_{i-1}}^n(x)}$ -measurable, again by the induction hypothesis. By the definition of the σ -algebra $\mathcal{C}_{\bar{A}_{t_{i-1}}^n(x)}$ and the induction hypothesis, we then conclude that $\{\bar{A}_{t_i}^n(x) \leq t\} \in \mathcal{C}_t$. Therefore the result. \square

Denote by $P_f^{n,\bar{\xi}}$ the law on (C, \mathcal{C}) of a diffusion process with infinitesimal generator $\tilde{\mathcal{L}}^n$ given by

$$\tilde{\mathcal{L}}_t^n(\omega)h(z) = \bar{g}_n(t, \omega)\nabla h(z) + \frac{\varepsilon^2}{2}\Delta h(z), \quad \forall \omega \in C, h \in C^2(\mathbb{R}), z \in \mathbb{R} \quad (13)$$

and initial condition $\bar{\xi}_0 = \omega$. Thanks to Lemma 3.4 we can define the statistical model associated with the observation of $\bar{\xi}$ until the stopping time $\bar{A}_T^n(x)$:

$$\mathcal{P}_{\bar{\xi}}^{n, \bar{A}_T^n(x)} = (C, \mathcal{C}_{\bar{A}_T^n(x)}, \{P_f^{n, \bar{\xi}}|_{\mathcal{C}_{\bar{A}_T^n(x)}}, f \in \mathcal{F}\}).$$

As in Step 1, one can prove the following proposition. There are, however, some technical points that need to be taken care of; for more details, we refer to [17], Proposition 5.4.

Proposition 3.5. *Under the same hypotheses as in Proposition 3.3, the statistical models $\mathcal{P}_{\bar{y}}^{n, T}$ and $\mathcal{P}_{\bar{\xi}}^{n, \bar{A}_T^n(x)}$ are equivalent.*

Step 3. We shall prove that $\Delta(\mathcal{P}_{\xi}^{A_T(x)}, \mathcal{P}_{\bar{\xi}}^{n, \bar{A}_T^n(x)}) \rightarrow 0$ as $n \rightarrow \infty$. To that aim we will prove that, setting $S_T^n(x) := A_T(x) \wedge \bar{A}_T^n(x)$, $\Delta(\mathcal{P}_{\xi}^{A_T(x)}, \mathcal{P}_{\xi}^{S_T^n(x)}) \rightarrow 0$, $\Delta(\mathcal{P}_{\xi}^{S_T^n(x)}, \mathcal{P}_{\xi}^{\bar{A}_T^n(x)}) \rightarrow 0$ and $\Delta(\mathcal{P}_{\xi}^{\bar{A}_T^n(x)}, \mathcal{P}_{\bar{\xi}}^{n, \bar{A}_T^n(x)}) \rightarrow 0$ as $n \rightarrow \infty$. We shall start by showing that $\Delta(\mathcal{P}_{\xi}^{\bar{A}_T^n(x)}, \mathcal{P}_{\bar{\xi}}^{n, \bar{A}_T^n(x)}) \rightarrow 0$; we need the following lemmas:

Lemma 3.6. *According to the notation above, the law of $(x_{A_t(x)})$ under P_f^{ξ} is the same as the law of (x_t) under $P_f^{\bar{y}}$. Moreover, let $P_f^{\bar{\xi}}$ be the distribution induced on (C, \mathcal{C}) by the law of a diffusion process $(\bar{\zeta}_t)$ satisfying*

$$d\bar{\zeta}_t = \frac{f(\bar{\zeta}_t)}{\sigma^2(\bar{\zeta}_t)}\bar{\sigma}_n^2(t, \bar{\zeta})dt + \varepsilon dW_t, \quad \bar{\zeta}_0 = w.$$

Then, the law of $(x_{\bar{A}_t^n(x)})$ under P_f^{ξ} is the same as the law of (x_t) under $P_f^{\bar{\xi}}$.

Proof. We shall only prove the first assertion, the proof of the second one being very similar. We have $\xi_0 = w = y_0$ and, for all $t > 0$:

$$x_{A_t(x)} = w + \int_0^{A_t(x)} \frac{f(x_s)}{\sigma^2(x_s)} ds + \varepsilon \tilde{W}_{A_t^n(x)},$$

where the process (\tilde{W}_t) is a standard brownian motion under P_f^{ξ} . The change of variable $s = A_u(x)$ implies that $ds = \sigma^2(x_{A_u(x)})du$, hence one can write

$$x_{A_t(x)} = w + \int_0^t \frac{f(x_{A_u(x)})}{\sigma^2(x_{A_u(x)})} \sigma^2(x_{A_u(x)}) du + \varepsilon \int_0^t \sigma(x_{A_u(x)}) B_u,$$

where the process (B_t) is defined by

$$B_t = \int_0^t \frac{d\tilde{W}_{A_u(x)}}{\sigma(x_{A_u(x)})}.$$

Classical results (see e.g. [27], 5.5) ensure that (B_t) is a $\mathcal{C}_{A_t(x)}$ standard brownian motion under P_f^y . It follows that the law of $x_{A_t(x)}$ under P_f^ξ is the same as the law of x under P_f^y . \square

Lemma 3.7. *Let p be an even positive integer and (t_n) a sequence of times bounded by CT for some constant C independent of f ; then $\mathbb{E}_{P_f^\xi} |x_{t_n}|^p = O(1)$, uniformly on \mathcal{F} .*

Proof. In order to bound $\mathbb{E}_{P_f^\xi} |x_{t_n}|^p$ we will use the following facts:

- $(z_1 + \dots + z_m)^p \leq m^{p-1}(z_1^p + \dots + z_m^p)$, $\forall z_1, \dots, z_m \in \mathbb{R}$;
- $\mathbb{E}_{P_f^\xi} \left| \int_0^v h(x_r) dr \right|^p \leq v^{p-1} \int_0^v \mathbb{E}_{P_f^\xi} h^p(x_r) dr$, for any integrable function h ;
- If X is a centered Gaussian random variable with variance σ^2 then $\mathbb{E}[X^p] = \sigma^p(p-1)!!$;
- (Gronwall lemma) Let $I = [0, a]$ be an interval of the real line, α a constant and let β and u continuous real valued functions defined on I . If β is non-negative and if u satisfies the integral inequality:

$$u(t) \leq \alpha + \int_0^t \beta(s)u(s) ds, \quad \forall t \in I,$$

then

$$u(t) \leq \alpha \exp\left(\int_0^t \beta(s) ds\right), \quad t \in I.$$

As one can always construct a brownian motion (\bar{B}_t) under P_f^ξ such that $dx_t = \frac{f(x_t)}{\sigma^2(x_t)} \bar{\sigma}_n^2(t, x) dt + \varepsilon d\bar{B}_t$, applying the first three facts combined with the linear growth of f one can write:

$$\begin{aligned} \mathbb{E}_{P_f^\xi} |x_{t_n}|^p &\leq 3^{p-1} w^p + 3^{p-1} \mathbb{E}_{P_f^\xi} \left(\int_0^{t_n} \frac{f(x_s)}{\sigma^2(x_s)} \bar{\sigma}_n^2(s, x) ds \right)^p + 3^{p-1} \varepsilon^p \mathbb{E}_{P_f^\xi} \bar{B}_{t_n}^p \\ &\leq 3^{p-1} w^p + 3^{p-1} \frac{\sigma_1^{2p}}{\sigma_0^{2p}} (CT)^{p-1} \int_0^{t_n} \mathbb{E}_{P_f^\xi} [f^p(x_s)] ds + 3^{p-1} \varepsilon^p (CT)^{\frac{p}{2}} (p-1)!! \\ &\leq C' \left(1 + \int_0^{t_n} \mathbb{E}_{P_f^\xi} |x_s|^p ds \right), \end{aligned}$$

for some constant C' independent of f . Applying the Gronwall lemma, we obtain

$$\mathbb{E}_{P_f^\xi} |x_{t_n}|^p \leq C' e^{C't_n} \leq C' e^{C'CT} = O(1).$$

\square

Lemma 3.8. *Under the same hypotheses as in Proposition 3.3 and with the same notation as in Steps 1 and 2, we have*

$$\mathbb{E}_{P_f^\xi} \int_0^{A_T^n(x)} \left(\frac{f(x_s)}{\sigma^2(x_s)} - \bar{g}_n(s, x) \right)^2 ds = O(n^{-2} + \varepsilon n^{-1}),$$

uniformly on \mathcal{F} .

Proof. For the sake of brevity, in this proof we will omit the superscript n in each occurrence of \bar{A}_t^n . We start by observing that, for all $y, z \in \mathbb{R}$

$$\begin{aligned} \left| \frac{f(z)}{\sigma^2(z)} - \frac{f(y)}{\sigma^2(y)} \right| &\leq |f(z)| \left| \frac{1}{\sigma^2(z)} - \frac{1}{\sigma^2(y)} \right| + \frac{|f(z) - f(y)|}{\sigma_0^2} \\ &\leq \frac{2M^2\sigma_1}{\sigma_0^4} (1 + |z|) |z - y| + \frac{M}{\sigma_0^2} |z - y|, \end{aligned}$$

hence there exists some constant C such that $\left(\frac{f(z)}{\sigma^2(z)} - \frac{f(y)}{\sigma^2(y)} \right)^2 \leq C(z - y)^2(1 + z^2)$.

Applying this inequality we can write:

$$\begin{aligned} \int_0^{\bar{A}_T(x)} \left(\frac{f(x_s)}{\sigma^2(x_s)} - \bar{g}_n(s, x) \right)^2 ds &\leq \sum_{i=0}^{n-1} \int_{\bar{A}_{t_i}(x)}^{\bar{A}_{t_{i+1}}(x)} C(x_s - x_{\bar{A}_{t_i}(x)})^2 (1 + x_{\bar{A}_{t_i}(x)}^2) ds \\ &= C \sum_{i=0}^{n-1} (1 + x_{\bar{A}_{t_i}(x)}^2) \int_{\bar{A}_{t_i}(x)}^{\bar{A}_{t_{i+1}}(x)} (x_r - x_{\bar{A}_{t_i}(x)})^2 dr \\ &\leq C\sigma_1^2 \sum_{i=0}^{n-1} (1 + x_{\bar{A}_{t_i}(x)}^2) \int_0^{t_{i+1} - t_i} (x_{\bar{A}_{t_i+s}(x)} - x_{\bar{A}_{t_i}(x)})^2 ds, \end{aligned}$$

where in the last step we have performed the change of variables $r = \bar{A}_{t_i+s}(x)$. Thanks to the Cauchy-Schwarz inequality and Lemma 3.7 we obtain

$$\begin{aligned} \mathbb{E}_{P_f^\xi} \int_0^{\bar{A}_T(x)} \left(\frac{f(x_s)}{\sigma^2(x_s)} - \bar{g}_n(s, x) \right)^2 ds &\leq \\ &\leq C\sigma_1^2 \sum_{i=0}^{n-1} \sqrt{\mathbb{E}_{P_f^\xi} (1 + x_{\bar{A}_{t_i}(x)}^2)^2} \sqrt{\mathbb{E}_{P_f^\xi} \left(\int_0^{t_{i+1} - t_i} (x_{\bar{A}_{t_i+r}(x)} - x_{\bar{A}_{t_i}(x)})^2 dr \right)^2} \\ &= C\sigma_1^2 \sum_{i=0}^{n-1} \sqrt{\mathbb{E}_{P_f^\xi} (1 + x_{t_i}^2)^2} \sqrt{\mathbb{E}_{P_f^\xi} \left(\int_0^{\frac{T}{n}} (x_{r+t_i} - x_{t_i})^2 dr \right)^2} \\ &\leq C\sigma_1^2 \sum_{i=0}^{n-1} \sqrt{\mathbb{E}_{P_f^\xi} (2 + 2x_{t_i}^4)} \sqrt{\mathbb{E}_{P_f^\xi} \left(\frac{T}{n} \int_0^{\frac{T}{n}} (x_{r+t_i} - x_{t_i})^4 dr \right)} \\ &= O \left(\sum_{i=0}^{n-1} \sqrt{\mathbb{E}_{P_f^\xi} \left(\frac{T}{n} \int_0^{\frac{T}{n}} (x_{r+t_i} - x_{t_i})^4 dr \right)} \right). \end{aligned}$$

Using the same arguments as in the proof of Lemma 3.7, we can write

$$\begin{aligned} \mathbb{E}_{P_f^\xi} (x_{r+t_i} - x_{t_i})^4 &= \mathbb{E}_{P_f^\xi} \left| \int_{t_i}^{t_i+r} \frac{f(x_s)\sigma^2(x_{t_i})}{\sigma^2(x_s)} ds + \varepsilon\sigma(x_{t_i})(\bar{B}_{t_i+r} - \bar{B}_{t_i}) \right|^4 \\ &\leq 8\mathbb{E}_{P_f^\xi} \left| \int_{t_i}^{t_i+r} \frac{f(x_s)\sigma^2(x_{t_i})}{\sigma^2(x_s)} ds \right|^4 + 8\varepsilon^4\sigma_1^4\mathbb{E}_{P_f^\xi} |\bar{B}_{t_i+r} - \bar{B}_{t_i}|^4 \\ &\leq 8\frac{\sigma_1^8}{\sigma_0^8} r^3 \int_0^r \mathbb{E}_{P_f^\xi} [f^4(x_{s+t_i})] ds + 8\varepsilon^4\sigma_1^4 r^2 6! \\ &= O \left(r^4 + r^3 \int_0^r \mathbb{E}_{P_f^\xi} [x_{s+t_i}^4] ds + \varepsilon^4 r^2 \right) \\ &= O(r^4 + \varepsilon^4 r^2) = O \left(\frac{1}{n^4} + \frac{\varepsilon^4}{n^2} \right). \end{aligned}$$

Putting all the pieces together we get:

$$\begin{aligned} \int_0^{\bar{A}_T(x)} \left(\frac{f(x_s)}{\sigma^2(x_s)} - \bar{g}_n(s, x) \right)^2 ds &= O\left(\sum_{i=0}^{n-1} \sqrt{\left(\frac{T}{n} \int_0^{\frac{T}{n}} O\left(\frac{1}{n^4} + \frac{\varepsilon^4}{n^2} \right) dr \right)} \right) \\ &= O\left(\frac{1}{n^2} + \frac{\varepsilon^2}{n} \right). \end{aligned}$$

□

Proposition 3.9. *Under the same hypotheses as in Proposition 3.3, we have*

$$\Delta(\mathcal{P}_\xi^{\bar{A}_T^n(x)}, \mathcal{P}_\xi^{n, \bar{A}_T^n(x)}) = O(\sqrt{\varepsilon^{-2}n^{-2} + n^{-1}}).$$

Proof. We use an inequality involving the Hellinger process in order to bound $\left\| P_f^\xi | \mathcal{C}_{\bar{A}_T^n(x)} - P_f^{n, \bar{\xi}} | \mathcal{C}_{\bar{A}_T^n(x)} \right\|_{\text{TV}}$ and hence $\Delta(\mathcal{P}_\xi^{\bar{A}_T^n(x)}, \mathcal{P}_\xi^{n, \bar{A}_T^n(x)})$. More precisely, let h_f be the Hellinger process of order 1/2 between the measures $P_f^\xi | \mathcal{C}_{\bar{A}_T^n(x)}$ and $P_f^{n, \bar{\xi}} | \mathcal{C}_{\bar{A}_T^n(x)}$, that is, (see Jacod and Shiryaev, [25], page 239)

$$h_f(t)(x) = \frac{1}{8\varepsilon^2} \int_0^t \left(\frac{f(x_s)}{\sigma^2(x_s)} - \bar{g}_n(s, x) \right)^2 ds.$$

Then:

$$\left\| P_f^\xi | \mathcal{C}_{\bar{A}_T^n(x)} - P_f^{n, \bar{\xi}} | \mathcal{C}_{\bar{A}_T^n(x)} \right\|_{\text{TV}} \leq 4 \sqrt{\mathbb{E}_{P_f^\xi} h_f(\bar{A}_T^n(x))(x)},$$

as in [25], 4b, Theorem 4.21, page 279. Hence we conclude thanks to Lemma 3.8. □

We now prove that $\Delta(\mathcal{P}_\xi^{A_T^n(x)}, \mathcal{P}_\xi^{S_T^n(x)}) \rightarrow 0$ and $\Delta(\mathcal{P}_\xi^{S_T^n(x)}, \mathcal{P}_\xi^{\bar{A}_T^n(x)}) \rightarrow 0$ as $n \rightarrow \infty$. Again, we start with a lemma:

Lemma 3.10. *Under the hypotheses of Proposition 3.3 and with the same notation as in Steps 1 and 2, we have*

$$\mathbb{E}_{P_f^\xi} |A_T(x) - \bar{A}_T^n(x)| = O\left(\frac{1}{n} + \varepsilon \right), \quad (14)$$

uniformly over \mathcal{F} .

Proof. The crucial point in proving (14) is to use the convergence of diffusion processes with small variance to some deterministic solution. To that aim, let us introduce the following ODEs:

$$\frac{dz_t}{dt} = f(z_t), \quad \frac{d\bar{z}_t}{dt} = \frac{f(\bar{z}_t)}{\sigma^2(\bar{z}_t)} \bar{\sigma}_n^2(t, \bar{z}), \quad z_0 = w = \bar{z}_0.$$

By means of Property 3.2, the Lipschitz character of $\sigma^2(\cdot)$ and the linear growth of f , we get,

$$\begin{aligned} \mathbb{E}_{P_f^\xi} |A_T(x) - \bar{A}_T^n(x)| &= \mathbb{E}_{P_f^\xi} \left| \int_0^T (\sigma^2(x_{A_t(x)}) - \bar{\sigma}_n^2(t, x_{\bar{A}_t^n(x)})) dt \right| \\ &\leq 2\sigma_1 M \mathbb{E}_{P_f^\xi} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} |x_{A_t(x)} - x_{\bar{A}_t^n(x)}| dt \\ &\leq 2\sigma_1 M \mathbb{E}_{P_f^\xi} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} (|x_{A_t(x)} - z_t| + |z_t - \bar{z}_{t_i}| + |\bar{z}_{t_i} - x_{\bar{A}_t^n(x)}|) dt. \end{aligned}$$

For all $t \in [t_i, t_{i+1}]$, we shall analyze the terms $I = \mathbb{E}_{P_f^\xi} |x_{A_t(x)} - z_t|$, $II = |z_t - \bar{z}_{t_i}|$ and $III = \mathbb{E}_{P_f^\xi} |\bar{z}_{t_i} - x_{\bar{A}_{t_i}^n(x)}|$, separately.

- Term I: By means of Lemma 3.6 and some standard calculations one can write

$$\begin{aligned} \mathbb{E}_{P_f^\xi} |x_{A_t(x)} - z_t| &= \mathbb{E}_{P_f^y} |x_t - z_t| = \mathbb{E}_{P_f^y} \left| \int_0^t (f(x_s) - f(z_s)) ds + \varepsilon \int_0^t \sigma(x_s) dW_s \right| \\ &\leq M \mathbb{E}_{P_f^y} \int_0^t |x_s - z_s| ds + \varepsilon \sigma_1 \sqrt{t}; \end{aligned}$$

hence, an application of the Gronwall lemma yields $\mathbb{E}_{P_f^\xi} |x_{A_t(x)} - z_t| \leq \varepsilon \sqrt{t} \exp(MT)$.

- Term II: By the triangular inequality it is enough to bound $|z_t - z_{t_j}|$ and $|z_{t_i} - \bar{z}_{t_j}|$, separately. It is easy to see that $|z_s - z_{t_j}|$ is a $O(n^{-1})$ as well as $|\bar{z}_s - \bar{z}_{t_j}|$ for all $s \in [t_j, t_{j+1}]$, $j = 0, \dots, n-1$. Moreover, observe that there exists a constant C , independent of f , such that $\left| f(x) - \frac{f(y)}{\sigma^2(y)} \sigma^2(z) \right| \leq C(|x - y| + (1 + |y|)|y - z|)$. We get:

$$\begin{aligned} |z_{t_i} - \bar{z}_{t_i}| &= \left| \int_0^{t_i} \left(f(z_s) - \frac{f(\bar{z}_s)}{\sigma^2(\bar{z}_s)} \sigma^2(s, \bar{z}) \right) ds \right| \\ &= \left| \sum_{j=0}^{i-1} \int_{t_j}^{t_{j+1}} \left(f(z_s) - \frac{f(\bar{z}_s)}{\sigma^2(\bar{z}_s)} \sigma^2(\bar{z}_{t_j}) \right) ds \right| \\ &\leq C \sum_{j=0}^{i-1} \int_{t_j}^{t_{j+1}} (|z_s - \bar{z}_s| + (1 + |\bar{z}_s|)|\bar{z}_s - \bar{z}_{t_j}|) ds \\ &\leq C \int_0^{t_i} |z_s - \bar{z}_s| ds + \frac{C' t_i}{n}, \end{aligned}$$

for some constant C' , independent of f . Therefore, applying the Gronwall lemma one obtains

$$|z_{t_i} - \bar{z}_{t_i}| \leq \frac{C' t_i}{n} e^{C t_i}$$

that allows us to conclude $|z_t - \bar{z}_{t_i}| = O(n^{-1})$.

- Term III: By means of Lemma 3.6 we know that $\mathbb{E}_{P_f^\xi} |\bar{z}_{t_i} - x_{\bar{A}_{t_i}^n(x)}| = \mathbb{E}_{P_f^\xi} |\bar{z}_{t_i} - x_{t_i}|$.

$$\begin{aligned} \mathbb{E}_{P_f^\xi} |\bar{z}_{t_i} - x_{t_i}| &= \mathbb{E}_{P_f^\xi} \left| \sum_{j=0}^{i-1} \int_{t_j}^{t_{j+1}} \left[\left(\frac{f(\bar{z}_s)}{\sigma^2(\bar{z}_s)} \sigma^2(\bar{z}_{t_j}) - \frac{f(x_s)}{\sigma^2(x_s)} \sigma^2(x_{t_j}) \right) ds + \varepsilon \sigma(x_{t_j}) dW_s \right] \right| \\ &\leq \mathbb{E}_{P_f^\xi} \left| \sum_{j=0}^{i-1} \int_{t_j}^{t_{j+1}} \left(\frac{f(\bar{z}_s)}{\sigma^2(\bar{z}_s)} \sigma^2(\bar{z}_{t_j}) - \frac{f(x_s)}{\sigma^2(x_s)} \sigma^2(x_{t_j}) \right) ds \right| + \varepsilon \sigma_1 \sqrt{t_i} \\ &\leq \mathbb{E}_{P_f^\xi} \sum_{j=0}^{i-1} \int_{t_j}^{t_{j+1}} \left(\frac{M \sigma_1^2}{\sigma_0^2} |\bar{z}_s - x_s| + \frac{2 \sigma_1 M}{\sigma_0^4} (1 + |\bar{z}_s|)(|\bar{z}_{t_j} - \bar{z}_s| + |x_s - x_{t_j}|) \right) ds + \varepsilon \sigma_1 \sqrt{t_i} \\ &\leq \mathbb{E}_{P_f^\xi} \int_0^{t_i} \frac{M \sigma_1^2}{\sigma_0^2} |\bar{z}_s - x_s| ds + C n^{-1} t_i + \varepsilon \sigma_1 \sqrt{t_i}, \end{aligned}$$

for some constant C independent of f . An application of the Gronwall lemma gives

$$\mathbb{E}_{P_f^\xi} |\bar{z}_{t_i} - x_{t_i}| \leq (C n^{-1} t_i + \varepsilon \sigma_1 \sqrt{t_i}) \exp\left(\frac{M \sigma_1^2}{\sigma_0^2} t_i\right).$$

Putting all the pieces together we obtain $\mathbb{E}_{P_f^\xi} |A_T(x) - \bar{A}_T^n(x)| = O\left(\frac{1}{n} + \varepsilon\right)$. \square

Proposition 3.11. *Under the same hypotheses of Proposition 3.3, $\Delta(\mathcal{P}_\xi^{A_T(x)}, \mathcal{P}_\xi^{S_T^n(x)}) \rightarrow 0$ and $\Delta(\mathcal{P}_\xi^{S_T^n(x)}, \mathcal{P}_\xi^{\bar{A}_T^n(x)}) \rightarrow 0$ as $n \rightarrow \infty$.*

Proof. We shall prove only the first statement, the proof of the second one being identical. Since $\mathcal{C}_{S_T^n(x)} \subset \mathcal{C}_{A_T(x)}$, it is clear that $\delta(\mathcal{P}_\xi^{A_T(x)}, \mathcal{P}_\xi^{S_T^n(x)}) = 0$. To control $\delta(\mathcal{P}_\xi^{S_T^n(x)}, \mathcal{P}_\xi^{A_T(x)})$ we will introduce the following Markov kernel K^n :

$$K^n(\omega, A) := \mathbb{E}_{P_0^\xi}(\mathbb{1}_A | \mathcal{C}_{S_T^n(x)})(\omega), \quad \forall A \in \mathcal{C}_{A_T(x)}, \omega \in C,$$

where P_0^ξ is defined as P_f^ξ with $f \equiv 0$. Remark that the Markov kernel K^n thus constructed coincides with the Markov kernel N defined in [17], Proposition 6.2, when $\varepsilon \equiv 1$. Making the same computations as in the cited proposition, we obtain that

$$\begin{aligned} \|K^n P_f^\xi | \mathcal{C}_{S_T^n(x)} - P_f^\xi | \mathcal{C}_{A_T(x)}\|_{TV} &\leq \frac{1}{2} \sqrt{\mathbb{E}_{P_f^\xi | \mathcal{C}_{A_T(x)}} \int_{S_T^n(x)}^{A_T(x)} \frac{f^2(x_r)}{\sigma^4(x_r)} dr} \\ &\leq \frac{M}{\sqrt{2}\sigma_0^2} \sqrt{\mathbb{E}_{P_f^\xi | \mathcal{C}_{A_T(x)}} \left(|A_T(x) - \bar{A}_T^n(x)| + \int_0^{|A_T(x) - \bar{A}_T^n(x)|} x_r^2 dr \right)} \\ &= O\left(\mathbb{E}_{P_f^\xi | \mathcal{C}_{A_T(x)}} |A_T(x) - \bar{A}_T^n(x)|^{1/4}\right). \end{aligned}$$

We then conclude that $\Delta(\mathcal{P}_\xi^{A_T(x)}, \mathcal{P}_\xi^{S_T^n(x)}) \rightarrow 0$ by means of Lemma 3.10. \square

Step 4. Using Steps 1–3 and the triangular inequality, one can find that $\Delta(\mathcal{P}_y^T, \mathcal{P}_y^{n,T}) = O\left(\frac{1}{\varepsilon n} + (n^{-1} + \varepsilon)^{1/4}\right)$. Hence, to conclude the proof of Theorem 2.1, we only need to show the following proposition:

Proposition 3.12. *Under the same hypotheses of Proposition 3.3, $\Delta(\mathcal{P}_y^{n,T}, \mathcal{Q}_Z^n) = 0$, for all n .*

Proof. Note that, by using the Girsanov theorem, we can show that the measure $P_f^{n,\bar{y}} | \mathcal{C}_T$ is absolutely continuous with respect to $P_0^{n,\bar{y}}$ and the density is given by

$$\frac{dP_f^{n,\bar{y}}}{dP_0^{n,\bar{y}}} | \mathcal{C}_T(\omega) = \exp\left(\sum_{i=0}^{n-1} \left(\frac{f(\omega_{t_i})}{\varepsilon^2 \sigma^2(\omega_{t_i})} (\omega_{t_{i+1}} - \omega_{t_i}) - \frac{f^2(\omega_{t_i})}{2n\varepsilon^2 \sigma^2(\omega_{t_i})}\right)\right).$$

Hence, by means of the Fisher's factorization theorem, we can deduce that the application $S : \omega \rightarrow (\omega_{t_1}, \dots, \omega_{t_n})$ is a sufficient statistic for the family of probability measures $\{P_f^{n,\bar{y}} | \mathcal{C}_T; f \in \mathcal{F}\}$. We complete the proof remarking that the distribution of $(x_{t_1}, \dots, x_{t_n})$ under $P_f^{n,\bar{y}}$ is the same as the one of (Z_1, \dots, Z_n) under \mathbb{P} and finally invoking the following property of the Le Cam distance (see Le Cam [31]):

Let $\mathcal{P}_i = (\mathcal{X}_i, \mathcal{A}_i, \{P_{i,\theta}, \theta \in \Theta\})$, $i = 1, 2$, be two statistical models and let $(\mathcal{X}_1, \mathcal{A}_1)$ be a Polish space. Let $S : \mathcal{X}_1 \rightarrow \mathcal{X}_2$ be a sufficient statistics such that the distribution of S under $P_{1,\theta}$ is equal to $P_{2,\theta}$. Then $\Delta(\mathcal{P}_1, \mathcal{P}_2) = 0$. \square

3.3 Proof of Theorem 2.2

We will proceed in three Steps.

Step 1. Let us consider the application $F : \mathbb{R} \rightarrow \mathbb{R}$ defined as $F(x) = \int_0^x \frac{1}{\varepsilon\sigma(u)} du$. Remark that F is well defined and one to one. Using the Itô formula, we have that

$$F(y_t) = F(w) + \int_0^t \left(\frac{f(y_s)}{\varepsilon\sigma(y_s)} - \frac{\sigma'(y_s)}{2\varepsilon} \right) ds + W_t.$$

Thus, if we set $\mu_t := F(y_t)$, the new process (μ_t) satisfies the following SDE:

$$\mu_0 = F(w); \quad d\mu_t = \left(\frac{f(F^{-1}(\mu_t))}{\varepsilon\sigma(F^{-1}(\mu_t))} - \frac{\sigma'(F^{-1}(\mu_t))}{2\varepsilon} \right) dt + dW_t, \quad t \in [0, T]. \quad (15)$$

Observe that, under hypotheses (H1) and (H2), the drift function $b(x) := \frac{f(F^{-1}(x))}{\varepsilon\sigma(F^{-1}(x))} - \frac{\sigma'(F^{-1}(x))}{2\varepsilon}$ is clearly bounded and also Lipschitz:

$$\begin{aligned} \varepsilon|b(x) - b(y)| &= \left| \left(\frac{f(F^{-1}(x))}{\sigma(F^{-1}(x))} - \frac{\sigma'(F^{-1}(x))}{2} \right) - \left(\frac{f(F^{-1}(y))}{\sigma(F^{-1}(y))} - \frac{\sigma'(F^{-1}(y))}{2} \right) \right| \\ &\leq \left| \frac{f(F^{-1}(x))}{\sigma(F^{-1}(x))} - \frac{f(F^{-1}(y))}{\sigma(F^{-1}(y))} \right| + \left| \frac{\sigma'(F^{-1}(x))}{2} - \frac{\sigma'(F^{-1}(y))}{2} \right| \\ &\leq \frac{\sigma_1 M + M^2}{\sigma_0^2} |F^{-1}(x) - F^{-1}(y)| + \frac{M}{2} |F^{-1}(x) - F^{-1}(y)| \\ &= \left| \int_x^y \varepsilon\sigma(F(u)) du \right| \left(\frac{M}{2} + \frac{M(\sigma_1 + M)}{\sigma_0^2} \right) \leq \left(\frac{M}{2} + \frac{M(\sigma_1 + M)}{\sigma_0^2} \right) \sigma_1 \varepsilon |x - y|; \end{aligned}$$

In particular the existence and the uniqueness of a strong solution μ for the SDE (15) are guaranteed. Let us denote by P_f^μ (resp. $Q_f^{n,\mu}$) the law of μ (resp. $(\mu_{t_1}, \dots, \mu_{t_n})$) and introduce the statistical models

$$\mathcal{P}_\mu^T = (C, \mathcal{C}_T, (P_f^\mu, f \in \mathcal{F})), \quad \mathcal{Q}_\mu^n = (\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n), (Q_f^{n,\mu}, f \in \mathcal{F})).$$

By construction, \mathcal{P}_μ^T (resp. \mathcal{Q}_μ^n) is the image experiment of \mathcal{P}_y^T (resp. \mathcal{Q}_y^n) by F . Thus we have:

Proposition 3.13. *Under the hypotheses of Theorem 2.2, the statistical models \mathcal{P}_y^T (resp. \mathcal{Q}_y^n) and \mathcal{P}_μ^T (resp. \mathcal{Q}_μ^n) are equivalent.*

Step 2. Using the same notations as above, define a new drift function \bar{b}_n :

$$\bar{b}_n(t, \omega) = \sum_{i=0}^{n-1} b(\omega_{t_i}) \mathbb{1}_{(t_i, t_{i+1}]}(t), \quad \forall \omega \in C, \quad t \in [0, T]$$

and consider the diffusion process $(\bar{\mu}_t)$ on (C, \mathcal{C}_T) having drift function given by \bar{b}_n and diffusion coefficient equal to 1, i.e.

$$\bar{\mu}_0 = F(w); \quad d\bar{\mu}_t = \bar{b}_n(t, \bar{\mu}) dt + dW_t, \quad t \in [0, T]. \quad (16)$$

Denote by $P_f^{n, \bar{\mu}}$ the law of the solution of (16) and introduce the corresponding statistical model:

$$\mathcal{P}_\mu^{n, T} = (C, \mathcal{C}_T, (P_f^{n, \bar{\mu}}, f \in \mathcal{F})).$$

Proposition 3.14. *Under the hypotheses of Theorem 2.2, the statistical models $\mathcal{P}_\mu^{n, T}$ and \mathcal{P}_μ^T are asymptotically equivalent as n goes to infinity.*

Proof. One can use the same arguments as in the proof of Proposition 3.9 obtaining a first bound given by

$$\Delta(\mathcal{P}_\mu^T, \mathcal{P}_{\bar{\mu}}^{n,T}) \leq \sup_{f \in \mathcal{F}} \|P_f^\mu - P_f^{n, \bar{\mu}}\|_{TV} \leq 4 \sup_{f \in \mathcal{F}} \sqrt{\mathbb{E}_{P_f^\mu} \frac{1}{8} \int_0^T (b(x_s) - \bar{b}_n(s, x))^2 ds}.$$

Now, thanks to the L -Lipschitz character of b , one can write $\int_0^T (b(x_s) - \bar{b}_n(s, x))^2 ds \leq L \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} (x_s - x_{t_i})^2 ds$ so that the usual computations yield $\Delta(\mathcal{P}_\mu^T, \mathcal{P}_{\bar{\mu}}^{n,T}) = O(n^{-1})$. \square

Step 3. Consider now the statistical model associated with the discrete observations $(\bar{\mu}_{t_1}, \dots, \bar{\mu}_{t_n})$:

$$\mathcal{Q}_{\bar{\mu}}^n = (\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n), (Q_f^{n, \bar{\mu}}, f \in \mathcal{F})),$$

where $Q_f^{n, \bar{\mu}}$ denotes the law of the vector $(\bar{\mu}_{t_1}, \dots, \bar{\mu}_{t_n})$.

Proposition 3.15. *Under the hypotheses of Theorem 2.2, we have*

$$\Delta(\mathcal{P}_{\bar{\mu}}^{n,T}, \mathcal{Q}_{\bar{\mu}}^n) = 0, \quad \Delta(\mathcal{Q}_{\bar{\mu}}^n, \mathcal{P}_{\bar{\mu}}^n) = 0, \quad \forall n.$$

Proof. The first equivalence can be proved by means of a sufficient statistic as in the proof of Proposition 3.12; the second one follows directly from Step 2 since $\|Q_f^{n, \bar{\mu}} - Q_f^{n, \bar{\mu}}\|_{TV} \leq \|P_f^\mu - P_f^{n, \bar{\mu}}\|_{TV}$ as we are only restricting to a smaller σ -algebra. \square

4 Appendix

4.1 Asymptotic equivalence in the sense of Le Cam

A *statistical model* is a triplet $\mathcal{P}_j = (\mathcal{X}_j, \mathcal{A}_j, \{P_{j, \theta}; \theta \in \Theta\})$ where $\{P_{j, \theta}; \theta \in \Theta\}$ is a family of probability distributions all defined on the same σ -field \mathcal{A}_j over the *sample space* \mathcal{X}_j and Θ is the *parameter space*. The *deficiency* $\delta(\mathcal{P}_1, \mathcal{P}_2)$ of \mathcal{P}_1 with respect to \mathcal{P}_2 quantifies “how much information we lose” by using \mathcal{P}_1 instead of \mathcal{P}_2 and is defined as $\delta(\mathcal{P}_1, \mathcal{P}_2) = \inf_K \sup_{\theta \in \Theta} \|KP_{1, \theta} - P_{2, \theta}\|_{TV}$, where TV stands for “total variation” and the infimum is taken over all “transitions” K (see [31], page 18). In our setting, however, the general notion of “transitions” can be replaced with the notion of Markov kernels. Indeed, when the model \mathcal{P}_1 is dominated and the sample space $(\mathcal{X}_2, \mathcal{A}_2)$ of the experiment \mathcal{P}_2 is a Polish space, the infimum appearing on the definition of the deficiency δ can be taken over all Markov kernels K on $\mathcal{X}_1 \times \mathcal{A}_2$ (see [37], Proposition 10.2), i.e.

$$\delta(\mathcal{P}_1, \mathcal{P}_2) = \inf_K \sup_{\theta \in \Theta} \sup_{A \in \mathcal{A}_2} \left| \int_{\mathcal{X}_1} K(x, A) P_{1, \theta}(dx) - P_{2, \theta}(A) \right|. \quad (17)$$

The experiment $KP_{1, \theta} = (\mathcal{X}_1, \mathcal{A}_1, \{KP_{1, \theta}\}_{\theta \in \Theta})$ is called a *randomization* of \mathcal{P}_1 by the kernel K . If the kernel is deterministic, i.e. for $T : (\mathcal{X}_1, \mathcal{A}_1) \rightarrow (\mathcal{X}_2, \mathcal{A}_2)$ a random variable, $T(x, A) := \mathbb{1}_A(T(x))$, the experiment $T\mathcal{P}_1$ is called the *image experiment by the random variable* T . Closely associated with the notion of deficiency is the so called Δ -distance, i.e. the pseudo metric defined by:

$$\Delta(\mathcal{P}_1, \mathcal{P}_2) := \max(\delta(\mathcal{P}_1, \mathcal{P}_2), \delta(\mathcal{P}_2, \mathcal{P}_1)).$$

The sufficiency of a statistic can be expressed in terms of the Δ -distance. More precisely, the following holds (see [17], Proposition 8.1, page 23). *Let $T : (\mathcal{X}_1, \mathcal{A}_1) \rightarrow (\mathcal{X}_2, \mathcal{A}_2)$ be a random variable. The statistic T is sufficient for \mathcal{P}_1 if and only if $\Delta(\mathcal{P}_1, T\mathcal{P}_1) = 0$.*

Also, remark that thanks to (17), if $\mathcal{P}_1 = (\mathcal{X}, \mathcal{A}_1, \{P_\theta; \theta \in \Theta\})$ and $\mathcal{P}_2 = (\mathcal{X}, \mathcal{A}_2, \{P_\theta; \theta \in \Theta\})$ with $\mathcal{A}_2 \subset \mathcal{A}_1$, then $\delta(\mathcal{P}_1, \mathcal{P}_2) = 0$.

Two sequences of statistical models $(\mathcal{P}_1^n)_{n \in \mathbb{N}}$ and $(\mathcal{P}_2^n)_{n \in \mathbb{N}}$ are called *asymptotically equivalent* if $\Delta(\mathcal{P}_1^n, \mathcal{P}_2^n)$ tends to zero as n goes to infinity. Similarly, the statistic T^n is *asymptotically sufficient* for \mathcal{P}_1^n if $\Delta(\mathcal{P}_1^n, T^n \mathcal{P}_1^n)$ tends to zero as n goes to infinity.

Acknowledgements

I would like to thank Valentine Genon-Catalot for several interesting discussions, especially in suggesting to taking into account the relation between diffusion processes with small variance and deterministic limits. Also, I would like to give a special thank to Pierre Étoré, with whom a lot of hours were spent discussing different approaches to the proof of Lemma 3.10. More generally, I am very grateful for all the time he has invested in supervising this project.

References

- [1] Lawrence D. Brown, T. Tony Cai, Mark G. Low, and Cun-Hui Zhang. Asymptotic equivalence theory for nonparametric regression with random design. *Ann. Statist.*, 30(3):688–707, 2002. Dedicated to the memory of Lucien Le Cam.
- [2] Lawrence D. Brown, Andrew V. Carter, Mark G. Low, and Cun-Hui Zhang. Equivalence theory for density estimation, Poisson processes and Gaussian white noise with drift. *Ann. Statist.*, 32(5):2074–2097, 2004.
- [3] Lawrence D. Brown and Mark G. Low. Asymptotic equivalence of nonparametric regression and white noise. *Ann. Statist.*, 24(6):2384–2398, 1996.
- [4] Lawrence D. Brown and Cun-Hui Zhang. Asymptotic nonequivalence of nonparametric experiments when the smoothness index is $1/2$. *Ann. Statist.*, 26(1):279–287, 1998.
- [5] Boris Buchmann and Gernot Müller. Limit experiments of GARCH. *Bernoulli*, 18(1):64–99, 2012.
- [6] Andrew V. Carter. Deficiency distance between multinomial and multivariate normal experiments. *Ann. Statist.*, 30(3):708–730, 2002. Dedicated to the memory of Lucien Le Cam.
- [7] Andrew V. Carter. A continuous Gaussian approximation to a nonparametric regression in two dimensions. *Bernoulli*, 12(1):143–156, 2006.
- [8] Andrew V. Carter. Asymptotic approximation of nonparametric regression experiments with unknown variances. *Ann. Statist.*, 35(4):1644–1673, 2007.
- [9] Andrew V. Carter. Asymptotically sufficient statistics in nonparametric regression experiments with correlated noise. *J. Probab. Stat.*, pages Art. ID 275308, 19, 2009.
- [10] Fabienne Comte, Valentine Genon-Catalot, and Yves Rozenholc. Penalized nonparametric mean square estimation of the coefficients of diffusion processes. *Bernoulli*, pages 514–543, 2007.
- [11] Arnak Dalalyan and Markus Reiß. Asymptotic statistical equivalence for scalar ergodic diffusions. *Probab. Theory Related Fields*, 134(2):248–282, 2006.

- [12] Arnak Dalalyan and Markus Reiß. Asymptotic statistical equivalence for ergodic diffusions: the multidimensional case. *Probab. Theory Related Fields*, 137(1-2):25–47, 2007.
- [13] Sylvain Delattre and Marc Hoffmann. Asymptotic equivalence for a null recurrent diffusion. *Bernoulli*, 8(2):139–174, 2002.
- [14] Sam Efromovich and Alex Samarov. Asymptotic equivalence of nonparametric regression and white noise model has its limits. *Statist. Probab. Lett.*, 28(2):143–145, 1996.
- [15] Mark I. Freidlin and Alexander D. Wentzell. *Random perturbations of dynamical systems*, volume 260 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, third edition, 2012. Translated from the 1979 Russian original by Joseph Szücs.
- [16] V Genon-Catalot. Maximum contrast estimation for diffusion processes from discrete observations. *Statistics*, 21(1):99–116, 1990.
- [17] Valentine Genon-Catalot and Catherine Laredo. Asymptotic equivalence of nonparametric diffusion and Euler scheme experiments. *The Annals of Statistics*, 42(3):1145–1165, 2014.
- [18] Valentine Genon-Catalot, Catherine Laredo, and Michael Nussbaum. Asymptotic equivalence of estimating a Poisson intensity and a positive diffusion drift. *Ann. Statist.*, 30(3):731–753, 2002. Dedicated to the memory of Lucien Le Cam.
- [19] Georgi K. Golubev, Michael Nussbaum, and Harrison H. Zhou. Asymptotic equivalence of spectral density estimation and Gaussian white noise. *Ann. Statist.*, 38(1):181–214, 2010.
- [20] Ion Grama and Michael Nussbaum. Asymptotic equivalence for nonparametric generalized linear models. *Probab. Theory Related Fields*, 111(2):167–214, 1998.
- [21] Ion Grama and Michael Nussbaum. Asymptotic equivalence for nonparametric regression. *Math. Methods Statist.*, 11(1):1–36, 2002.
- [22] Ion G. Grama and Michael H. Neumann. Asymptotic equivalence of nonparametric autoregression and nonparametric regression. *Ann. Statist.*, 34(4):1701–1732, 2006.
- [23] Romain Guy. Inférence dans le cadre des maladies transmissibles par des processus de diffusion. *Ph.D thesis*, 2013.
- [24] Marc Hoffmann. Adaptive estimation in diffusion processes. *Stochastic processes and their Applications*, 79(1):135–163, 1999.
- [25] Jean Jacod and Albert N Shiryaev. *Limit theorems for stochastic processes*, volume 288. Springer-Verlag Berlin, 1987.
- [26] Michael Jähnisch and Michael Nussbaum. Asymptotic equivalence for a model of independent non identically distributed observations. *Statist. Decisions*, 21(3):197–218, 2003.
- [27] Ioannis Karatzas and Steven E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2000.

- [28] Yu A Kutoyants. On nonparametric estimation of trend coefficients in a diffusion process. *Statistics and Control of Stochastic Processes*, 230:250, 1984.
- [29] Yu A Kutoyants. *Parameter estimation for stochastic processes*, volume 6. Heldermann, 1984.
- [30] Catherine F Laredo. A sufficient condition for asymptotic sufficiency of incomplete observations of a diffusion process. *The Annals of Statistics*, pages 1158–1171, 1990.
- [31] Lucien Le Cam. *Asymptotic methods in statistical decision theory*. Springer Series in Statistics. Springer-Verlag, New York, 1986.
- [32] Lucien Le Cam and Grace Lo Yang. *Asymptotics in statistics*. Springer Series in Statistics. Springer-Verlag, New York, second edition, 2000. Some basic concepts.
- [33] Ester Mariucci. Asymptotic equivalence for inhomogeneous jump diffusion processes, 2014.
- [34] Alexander Meister. Asymptotic equivalence of functional linear regression and a white noise inverse problem. *Ann. Statist.*, 39(3):1471–1495, 2011.
- [35] Alexander Meister and Markus Reiß. Asymptotic equivalence for nonparametric regression with non-regular errors. *Probab. Theory Related Fields*, 155(1-2):201–229, 2013.
- [36] Grigori Milstein and Michael Nussbaum. Diffusion approximation for nonparametric autoregression. *Probab. Theory Related Fields*, 112(4):535–543, 1998.
- [37] Michael Nussbaum. Asymptotic equivalence of density estimation and Gaussian white noise. *Ann. Statist.*, 24(6):2399–2430, 1996.
- [38] Bernt Øksendal. *Stochastic differential equations*. Universitext. Springer-Verlag, Berlin, 1985. An introduction with applications.
- [39] Jean Picard. Nonlinear filtering of one-dimensional diffusions in the case of a high signal-to-noise ratio. *SIAM Journal on Applied Mathematics*, 46(6):1098–1125, 1986.
- [40] Jean Picard. Efficiency of the extended Kalman filter for nonlinear systems with small noise. *SIAM Journal on Applied Mathematics*, 51(3):843–885, 1991.
- [41] Markus Reiß. Asymptotic equivalence for nonparametric regression with multivariate and random design. *Ann. Statist.*, 36(4):1957–1982, 2008.
- [42] Markus Reiß. Asymptotic equivalence for inference on the volatility from noisy observations. *Ann. Statist.*, 39(2):772–802, 2011.
- [43] Angelika Rohde. On the asymptotic equivalence and rate of convergence of nonparametric regression and Gaussian white noise. *Statist. Decisions*, 22(3):235–243, 2004.
- [44] Masayuki Uchida and Nakahiro Yoshida. Asymptotic expansion for small diffusions applied to option pricing. *Stat. Inference Stoch. Process.*, 7(3):189–223, 2004.
- [45] VA Volkonskii. Random substitution of time in strong Markov processes. *Theory of Probability & Its Applications*, 3(3):310–326, 1958.
- [46] Yazhen Wang. Asymptotic nonequivalence of Garch models and diffusions. *Ann. Statist.*, 30(3):754–783, 2002. Dedicated to the memory of Lucien Le Cam.