



HAL
open science

A collision process responsible for widespread formation of H anions

Zoltán Juhász, Béla Sulik, Sándor T S Kovács, Elie Lattouf, Jimmy Rangama, Erika Bene, Bernd Huber, François Frémont, Alain Méry, Jean-Christophe Pouilly, et al.

► To cite this version:

Zoltán Juhász, Béla Sulik, Sándor T S Kovács, Elie Lattouf, Jimmy Rangama, et al.. A collision process responsible for widespread formation of H anions. *Journal of Physics: Conference Series*, 2014, XXVIII International Conference on Photonic, Electronic and Atomic Collisions (ICPEAC 2013) 24–30 July 2013, Lanzhou, China, 488 (10), pp.102024. 10.1088/1742-6596/488/10/102024 . hal-01084494

HAL Id: hal-01084494

<https://hal.science/hal-01084494v1>

Submitted on 3 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

OPEN ACCESS

A collision process responsible for widespread formation of H anions

To cite this article: Zoltán Juhász *et al* 2014 *J. Phys.: Conf. Ser.* **488** 102024

View the [article online](#) for updates and enhancements.

Related content

- [Energy distribution of slow fragments. OH⁺ and O⁺, in water dissociation by electron impact](#)
Natalia Ferreira, L Sigaud and E C Montenegro
- [Anion and cation emission via binary-encounters in OH⁺ + Ar collisions: The role of dissociative excitation and statistical aspects](#)
Z Juhász, E Lattouf, J-Y Chesnel *et al.*
- [Electron capture and single ionization in H⁺ + Ar collisions: classical calculations](#)
F Frémont

Recent citations

- [Two- and many-body effects in cation emission from H₂O molecules by O⁺ impact at keV energies: Similarities between ionization of atoms and proton emission from molecules](#)
B Sulik *et al*
- [Anion emission from H₂O molecules colliding with positive O⁺ ions at keV energies: the role of dissociative excitation](#)
J-Y Chesnel *et al*
- [Anion and cation emission via binary-encounters in OH⁺ + Ar collisions: The role of dissociative excitation and statistical aspects](#)
Z Juhász *et al*



The banner features a decorative top border with a repeating pattern of red, white, and blue diagonal stripes. On the left, the ECS logo is displayed in green and blue, followed by the text 'The Electrochemical Society' and 'Advancing solid state & electrochemical science & technology'. In the center, the text '239th ECS Meeting with IMCS18' is written in a large, bold, dark blue font. Below this, 'DIGITAL MEETING • May 30-June 3, 2021' and 'Live events daily • Free to register' are written in a smaller, dark blue font. On the right side, there is a stylized graphic of a person's head with glowing blue lines representing neural connections or data flow. A red button with white text 'Register now!' is positioned at the bottom right. A small logo with the number '18th' is also visible in the upper right area of the banner.

A collision process responsible for widespread formation of H^- anions

Zoltán Juhász^{*1}, Béla Sulik^{*}, Sándor T. S. Kovács^{*}, Elie Lattouf[†], Jimmy Rangama[†], Erika Bene^{*},
Bernd Huber[†], François Frémont[†], Alain Méry[†], Jean-Christophe Pouilly[†], Patrick Rousseau[†],
Péter Herczku^{*}, and Jean-Yves Chesnel[†]

^{*}MTA Institute for Nuclear Research (Atomki), Bem tér 18/c, H-4026 Debrecen, Hungary

[†]Centre de Recherche sur les Ions, les Matériaux et la Photonique (CIMAP), Unité Mixte CEA-CNRS-EnsiCaen-
Université de Caen Basse-Normandie, UMR 6252, Bd Maréchal Juin, F-14000 Caen, France

Synopsis The hydride ion H^- is widely understood to play a major role in many processes important for physics, cosmology, astrophysics and chemistry. Here we report that H^- ions can be formed by collision-induced fragmentation of molecules containing hydrogen and heavier nuclei. We observed H^- emission from $\text{OH}^+ + \text{Ar}$ and $\text{OH}^+ + \text{acetone}$ collisions at the energy of 7 keV. Our data reveal a process of electron-grabbing by fast hydrogen fragments emitted in hard collisions with large momentum transfer between atomic cores.

The formation of negative ions in various environments has been a subject of great interest over the past decades. Even a small concentration of anions has an appreciable effect on the physical and chemical properties of their environment.

In this work we observed H^- formation in collisions of OH^+ projectiles with Ar atoms and acetone molecules ($\text{CH}_3\text{-CO-CH}_3$). The experiments were conducted at the 14.5 GHz ECR ion source of the ARIBE facility, at GANIL in Caen, France. The molecular OH^+ beam with incident energy of 7 keV crossed an effusive gas jet of either argon atoms or acetone molecules. Electrons and negative ions produced in the collision were measured by an electrostatic spectrometer. The spectra were taken from 2° to 150° with respect to the beam direction.

The most remarkable feature of the spectra was the appearance of well-defined peaks [1]. These structures could not be attributed to electron emission. A kinematic analysis has shown that one of the peaks is due to the emission of H^- ions moving with nearly the velocity of the OH^+ projectile ion. The other peak corresponds to H^- emission from the acetone target in direct two-body collisions with the oxygen center of the projectile. Moreover, we compared the measured cross sections with a classical two-body potential-scattering calculation.

Figure 1 shows the measured cross sections for H^- emission in $\text{OH}^+ + \text{Ar}$ collisions. Our new data recorded at small observation angles are indicated by full symbols, while our earlier data at larger angles are shown by open circle symbols. It is a notable feature that the H^- production cross section is proportional to the calculated $\text{H} + \text{Ar}$ elastic scattering cross section. This fact can be explained by a two step process: (i) A large momentum transfer collision liberates a proton. (ii) The receding proton

grabs two electrons in a double capture process with a practically observation-angle-independent probability. The agreement of the theoretical curve with earlier experimental data for $\text{H}^{2+} + \text{Ar}$ collisions at the same velocity supports this picture and its generalization [2]. These earlier data are shown in Fig. 1 by open square symbols. Since the process is likely in any collisions of H-containing molecules at few-keV energies, it is important for both its generality and its fundamental understanding.

This work was supported by the Transnational Access Program ITS-LEIF, granted by the European Project HPRI-CT-2005-026015, the Hungarian OTKA Foundation (Grant N° K73703) and the French-Hungarian Cooperation Program PHC Balaton (N° 27860ZL / TÉT_11-2-2012-028)




Figure 1. Measured cross sections for H^- ejection from the OH^+ projectile as a function of the observation angle for an Ar target. The curves represent 2-body elastic scattering cross sections multiplied by a factor of 0.012. Squares: data of Ref. [2] for H^{2+} impact on Ar.

References

- [1] Z Juhász *et al* 2012 *J Phys: Conf Ser* **388** 102051, and *in press* in *Phys. Rev. A* (2013).
- [2] F B Alarcón *et al* 2006 *Int. J. Mass Spectr.* **248** 21

¹E-mail: zjuhasz@atomki.hu