



HAL
open science

On p-almost direct products and residual properties of pure braid groups of nonorientable surfaces

Paolo Bellingeri, Sylvain Gervais

► **To cite this version:**

Paolo Bellingeri, Sylvain Gervais. On p-almost direct products and residual properties of pure braid groups of nonorientable surfaces. *Algebraic & Geometric Topology*, 2016, 16 (1), pp.547-568. hal-01088951

HAL Id: hal-01088951

<https://hal.science/hal-01088951>

Submitted on 29 Nov 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On p -almost direct products and residual properties of pure braid groups of nonorientable surfaces

Paolo Bellingeri and Sylvain Gervais

Abstract

We prove that the n^{th} pure braid group of a nonorientable surface (closed or with boundary, but different from \mathbb{RP}^2) is residually 2-finite. Consequently, this group is residually nilpotent. The key ingredient in the closed case is the notion of p -almost direct product, which is a generalization of the notion of almost direct product. We prove therefore also some results on lower central series and augmentation ideals of p -almost direct products.

1 Introduction

Let M be a surface (orientable or not, possibly with boundary) and $F_n(M) = \{(x_1, \dots, x_n) \in M^n / x_i \neq x_j \text{ for } i \neq j\}$ its n^{th} configuration space. The fundamental group $\pi_1(F_n(M))$ is called the n^{th} *pure braid group* of M and shall be denoted by $P_n(M)$.

The mapping class group $\Gamma(M)$ of M is the group of isotopy classes of homeomorphisms $h : M \rightarrow M$ which are identity on the boundary. Let $\mathcal{X}_n = \{z_1, \dots, z_n\}$ a set of n distinguished points in the interior of M ; the pure mapping class group $\text{P}\Gamma(M, \mathcal{X}_n)$ relatively to \mathcal{X}_n is the group of the isotopy classes of homeomorphisms $h : M \rightarrow M$ satisfying $h(z_i) = z_i$ for all i : since this group does not depend on the choice of the set \mathcal{X}_n but only on its cardinality we can write $\text{P}_n\Gamma(M)$ instead of $\text{P}\Gamma(M, \mathcal{X}_n)$. Forgetting the marked points, we get a morphism $\text{P}_n\Gamma(M) \rightarrow \Gamma(M)$ whose kernel is known to be isomorphic to $P_n(M)$ when M is not a sphere, a torus, a projective plane or a Klein bottle (see [Sc, GJ]).

Now, recall that if \mathcal{P} is a group-theoretic property, then a group G is said to be *residually* \mathcal{P} if, for all $g \in G$, $g \neq 1$, there exists a group homomorphism $\varphi : G \rightarrow H$ such that H is \mathcal{P} and $\varphi(g) \neq 1$. We are interested in this paper to the following properties: nilpotence, being free and being a finite p -group for a prime number p (mostly $p = 2$). Recall that, if for subgroups H and K of G , $[H, K]$ is the subgroup generated by $\{[h, k] / (h, k) \in H \times K\}$ where $[h, k] = h^{-1}k^{-1}hk$, the lower central series of G , $(\Gamma_k G)_{k \geq 1}$, is defined inductively by $\Gamma_1 G = G$ and $\Gamma_{k+1} G = [G, \Gamma_k G]$. It is well known that G is residually nilpotent if, and only if, $\bigcap_{k=1}^{+\infty} \Gamma_k G = \{1\}$. From the lower central series of G one can define another filtration $D_1(G) \supseteq D_2(G) \supseteq \dots$ setting $D_1(G) = G$, and for $i \geq 2$, defining $D_i(G) = \{x \in G \mid \exists n \in \mathbb{N}^*, x^n \in \Gamma_i(G)\}$. After Garoufalidis and Levine [GLe], this filtration is called *rational lower central series* of G and a group G is residually torsion-free nilpotent if, and only if, $\bigcap_{i=1}^{\infty} D_i(G) = \{1\}$.

When M is an orientable surface of positive genus (possibly with boundary) or a disc with holes, it is proved in [BGG] and [BB] that $P_n(M)$ is residually torsion-free nilpotent for all $n \geq 1$. The fact that a

Date: 17th November 2014

2010 AMS Mathematics Subject Classification: 20F36, 20F14, 20D15, 57M05

Key words: braid groups, pure braid groups, nonorientable surfaces, mod p Torelli group, p -almost direct product, lower central series, residually nilpotent, residually 2-finite

group is residually torsion-free nilpotent has several important consequences, notably that the group is bi-orderable [MR] and residually p -finite [Gr]. The goal of this article is to study the nonorientable case and, more precisely, to prove the following:

THEOREM 1 *The n^{th} pure braid group of a nonorientable surface different from \mathbb{RP}^2 is residually 2-finite.*

In the case of $P_n(\mathbb{RP}^2)$ we give some partial results at the end of Section 4. Since a finite 2-group is nilpotent, a residually 2-finite group is residually nilpotent. Thus, we have

COROLLARY 1 *The n^{th} pure braid group of a nonorientable surface different from \mathbb{RP}^2 is residually nilpotent.*

Remark that in [Go] it was shown that the n^{th} pure braid group of a nonorientable surface is not bi-orderable and therefore it is not residually torsion-free nilpotent. Let us notice also that if pure braid groups of nonorientable surfaces with boundary are residually p for a prime $p \neq 2$ therefore pure braid groups of nonorientable closed surfaces are also residually p (Remark 3); however since the technique proposed in the nonorientable case applies only for $p = 2$, the question if pure braid groups of nonorientable surfaces are residually p for $p \neq 2$ is still open (recall that there are groups residually p for infinitely many primes p which are not residually torsion-free nilpotent, see [H]).

Remark that one can prove that finite type invariants separate classical braids using the fact that the pure braid group P_n is residually nilpotent without torsion (see [Pa]). Moreover using above residual properties it is possible to construct algebraically a universal finite type invariant over \mathbb{Z} on the classical braid group B_n (see [Pa]). Similar constructions were afterwards proposed for braids on orientable surfaces (see [BF, GP]): in a further paper we will explore the relevance of Theorem 1 in the realm of finite type invariants over $\mathbb{Z}/2\mathbb{Z}$ for braids on non orientable surfaces.

From now on, $M = N_{g,b}$ is a nonorientable surface of genus g with b boundary components, simply denoted by N_g when $b = 0$. We will see N_g as a sphere S^2 with g open discs removed and g Möbius strips glued on each circle (see figure 2 where each crossed disc represents a Möbius strip). The surface $N_{g,b}$ is obtained from N_g by removing b open discs. The mapping class groups $\Gamma(N_{g,b})$ and pure mapping class group $P_n\Gamma(N_{g,b})$ will be denoted respectively $\Gamma_{g,b}$ and $\Gamma_{g,b}^n$.

The paper is organized as follows. In Section 2, we prove Theorem 1 for surfaces with boundary: following what the authors did in the orientable case (see [BGG]), we embed $P_n(N_{g,b})$ in a Torelli group. The difference here is that we must consider mod 2 Torelli groups. In Section 3 we introduce the notion of p -almost direct product, which generalizes the notion of almost direct product (see Definition 1) and we prove some results on lower central series and augmentations ideals of p -almost direct products (Theorems 4 and 5) that can be compared with similar results on almost direct products (Theorem 3.1 in [FR] and Theorem 3.1 in [Pa]).

In Section 4, the existence of a split exact sequence

$$1 \longrightarrow P_{n-1}(N_{g,1}) \longrightarrow P_n(N_g) \longrightarrow \pi_1(N_g) \longrightarrow 1$$

and results from Section 2 and 3 are used to prove Theorem 1 in the closed case (Theorem 7). The method is similar to the one developed for orientable surface in [BB]: the difference will be that the semi-direct product $P_{n-1}(N_{g,1}) \rtimes \pi_1(N_g)$ is a 2-almost-direct product (and not an almost-direct product as in the case of closed oriented surfaces).

For proving the main result of the paper, we will also need a group presentation for $P_n(N_{g,b})$ when $b \geq 1$. Although generators of this group seem to be known, we could not find a group presentation in the literature. Thus, we give one in the Appendix (Theorem A).

Acknowledgments. The research of the first author was partially supported by French grant ANR-11-JS01-002-01. The authors are grateful to Carolina de Miranda e Pereiro and John Guaschi for useful discussions and comments.

2 The case of non-empty boundary

In this section, $N = N_{g,b}$ is a nonorientable surface of genus $g \geq 1$ with boundary (ie $b \geq 1$). In this case, one has $P_n(N) = \text{Ker}(\Gamma_{g,b}^n \rightarrow \Gamma_{g,b})$ for all $n \geq 1$.

2.1 Notations

We will follow notations from [PS]. A simple closed curve in N is an embedding $\alpha : S^1 \rightarrow N \setminus \partial N$. Such a curve is said two-sided (resp. one-sided) if it admits a regular neighborhood homeomorphic to an annulus (resp. a Möbius strip). We shall consider the following elements in $\Gamma_{g,b}$.

- If α is a two-sided simple closed curve in N , τ_α is a Dehn twist along α .
- Let μ and α be two simple closed curves such that μ is one-sided, α is two-sided and $|\alpha \cap \mu| = 1$. A regular neighborhood K (resp. M) of $\alpha \cup \mu$ (resp. μ) is diffeomorphic to a Klein bottle with one hole (resp. a Möbius strip). Pushing M once along α , we get a diffeomorphism of K fixing the boundary: it can be extended via the identity to N . Such a diffeomorphism is called a crosscap slide, and denoted by $Y_{\mu,\alpha}$.
- Consider a one-sided simple closed curve μ containing exactly one marked point z_i . Sliding z_i once along μ , we get a diffeomorphism S_μ of N which is identity outside a regular neighborhood of μ . Such a diffeomorphism will be called puncture slide along μ .

2.2 Blowup homomorphism

In this section, we recall the construction of the Blowup homomorphism $\eta_{g,b}^n : \Gamma_{g,b}^n \rightarrow \Gamma_{g+n,b}$ given in [Sz1], [Sz2] and [PS].

Let $U = \{z \in \mathbb{C} / |z| \leq 1\}$ and, for $i = 1, \dots, n$, fix an embedding $e_i : U \rightarrow N$ such that $e_i(0) = z_i$, $e_i(U) \cap e_j(U) = \emptyset$ if $i \neq j$ and $e_i(U) \cap \partial N = \emptyset$ for all i . If we remove the interior of each $e_i(U)$ (thus getting the surface $N_{g,b+n}$) and identify, for each $z \in \partial U$, $e_i(z)$ with $e_i(-z)$, we get a nonorientable surface of genus $g + n$ with b boundary components, that is to say a surface homeomorphic to $N_{g+n,b}$. Let us denote by $\gamma_i = e_i(S^1)$ the boundary of $e_i(U)$, and by μ_i its image in $N_{g+n,b}$: it is a one-sided simple closed curve.

Now, let h be an element of $\Gamma_{g,b}^n$. It can be represented by a homeomorphism $N_{g,b} \rightarrow N_{g,b}$, still denoted h , such that

- (1) $h(e_i(z)) = e_i(z)$ if h preserves local orientation at z_i ;
- (2) $h(e_i(z)) = e_i(\bar{z})$ if h reverses local orientation at z_i .

Such a homeomorphism h commutes with the identification leading to $N_{g+n,b}$ and thus induces an element $\eta(h) \in \Gamma_{g+n,b}$. It is proved in [Sz2] that the map $\eta_{g,b}^n = \eta : \Gamma_{g,b}^n \rightarrow \Gamma_{g+n,b}$ which sends h to $\eta(h)$ is well defined for $n = 1$, but the proof also works for $n > 1$. This homomorphism is called *blowup* homomorphism.

PROPOSITION 1 *The blowup homomorphism $\eta_{g,b}^n : \Gamma_{g,b}^n \rightarrow \Gamma_{g+n,b}$ is injective if $(g + n, b) \neq (2, 0)$.*

REMARK 1 This result is proved in [Sz1] for $(g, b) = (0, 1)$, but the proof can be adapted in our case as follows.

Proof. Suppose that $h : N_{g,b} \rightarrow N_{g,b}$ is a homeomorphism satisfying $h(z_i) = z_i$ for all i and $\eta(h) : N_{g+n,b} \rightarrow N_{g+n,b}$ is isotopic to identity. Then h is isotopic to a map equal to identity on $e_i(U)$ for all i (otherwise, μ_i is isotopic to μ_i^{-1} since $\eta(h)(\mu_i)$ is isotopic to μ_i) and its restriction to $N_{g,b+n}$ is an element of the kernel of the natural map $\Gamma_{g,b+n} \rightarrow \Gamma_{g+n,b}$ induced by glueing a Möbius strip on n boundary components. However, this kernel is generated by the Dehn twists along the curves γ_i (see [St, theorem 3.6]). Now, any γ_i bounds a disc with one marked point in $N_{g,b}$: the corresponding Dehn twist is trivial in $\Gamma_{g,b}$ and therefore h is isotopic to identity. \square

2.3 Embedding $P_n(N_{g,b})$ in $\Gamma_{g+n+2(b-1),1}$

Gluing a one-holed torus onto $b-1$ boundary components of $N_{g,b}$, we get $N_{g,b}$ as a subsurface of $N_{g+2(b-1),1}$. This inclusion induces a homomorphism $\chi_{g,b} : \Gamma_{g,b} \rightarrow \Gamma_{g+2(b-1),1}$ which is injective (see [St]). Thus, the composed map $\lambda_{g,b}^n = \chi_{g+n,b} \circ \eta_{g,b}^n : \Gamma_{g,b}^n \rightarrow \Gamma_{g+n+2(b-1),1}$ is also injective.

Recall that the mod p Torelli group $I_p(N_{g,1})$ is the subgroup of $\Gamma_{g,1}$ defined as the kernel of the action of $\Gamma_{g,1}$ on $H_1(N_{g,1}; \mathbb{Z}/p\mathbb{Z})$. In the following we will consider in particular the case of the mod 2 Torelli group $I_2(N_{g,1})$.

PROPOSITION 2 *If $b \geq 1$, $\lambda_{g,b}^n(P_n(N_{g,b}))$ is a subgroup of the Torelli subgroup $I_2(N_{g+n+2(b-1),1})$.*




Figure 1: Image of the generators of $P_n(N_{g,b})$ in $\Gamma_{g+n+2(b-1),1}$

Proof. The image of the generators (see figures 2, 6 and theorem A) $(B_{i,j})_{1 \leq i < j \leq n}$, $(\rho_{k,l})_{\substack{1 \leq k \leq n \\ 1 \leq l \leq g}}$ and $(x_{u,t})_{\substack{1 \leq u \leq n \\ 1 \leq t \leq b-1}}$ of $P_n(N_{g,b})$ under $\lambda_{g,b}^n$ are respectively (see figure 1):

- * Dehn twist along curves $\beta_{i,j}$, which bound a subsurface homeomorphic to $N_{2,1}$;
- * crosscap slides $Y_{\mu_k, \alpha_{k,l}}$;
- * the product $\tau_{\xi_{u,t}} \tau_{\delta_t}^{-1}$ of Dehn twists along the bounding curves $\xi_{u,t}$ and δ_t .

According to [Sz2], all these elements are in the mod 2 Torelli subgroup $I_2(N_{g+n+2(b-1),1})$. \square

REMARK 2 The embedding provided in Proposition 2 does not hold for $I_p(N_{g+n+2(b-1),1})$, when $p \neq 2$: for example, the cross slide $Y_{\mu_k, \alpha_{k,t}}$ is not in the mod p Torelli subgroup since it sends μ_k to μ_k^{-1} .

2.4 Conclusion of the proof

We shall use the following result, which is a straightforward consequence of a similar result for mod p Torelli groups of orientable surfaces due to L. Paris [P]:

THEOREM 2 *Let $g \geq 1$. The mod p Torelli group $I_p(N_{g,1})$ is residually p -finite.*

Proof. We use Dehn-Nielsen-Baer Theorem (see for instance Theorem 5.15.3 of [CVZ]) which states that $\Gamma_{g,1}$ embeds in $\text{Aut}(\pi_1(N_{g,1}))$. Since $\pi_1(N_{g,1})$ is free we can apply Theorem 1.4 in [P] which claims that if G is a free group, its mod p Torelli group (i.e. the kernel of the canonical map from $\text{Aut}(G)$ to $\text{GL}(H_1(G, \mathbb{F}_p))$) is residually p -finite. Therefore $I_p(N_{g,1})$ is residually p -finite. \square

THEOREM 3 *Let $g \geq 1$, $b > 0$, $n \geq 1$. $P_n(N_{g,b})$ is residually 2-finite.*

Proof. The group $P_n(N_{g,b})$ is a subgroup of $I_2(N_{g+n+2(b-1),1})$ by Proposition 2 and by injectivity of the map $\lambda_{g,b}^n$. Then by Theorem 2 it follows that $P_n(N_{g,b})$ is residually 2-finite. \square

3 p -almost direct products

3.1 On residually p -finite groups

Let p be a prime number and G a group. If H is a subgroup of G , we denote by H^p the subgroup generated by $\{h^p / h \in H\}$. Following [P], we define the lower \mathbb{F}_p -linear central filtration $(\gamma_n^p G)_{n \in \mathbb{N}^*}$ of G by $\gamma_1^p G = G$ and, for $n \geq 1$, $\gamma_{n+1}^p G$ is the subgroup of G generated by $[G, \gamma_n^p G] \cup (\gamma_n^p G)^p$. Note that the subgroups $\gamma_n^p G$ are characteristic in G and that the quotient group $G/\gamma_2^p G$ is nothing but the first homology group $H_1(G; \mathbb{F}_p)$. The followings are proved in [P]:

- for $m, n \geq 1$, $[\gamma_m^p G, \gamma_n^p G] \subset \gamma_{m+n}^p G$;
- a finitely generated group G is a finite p -group if, and only if, there exists some $N \geq 1$ such that $\gamma_N^p G = \{1\}$;
- a finitely generated group G is residually p -finite if, and only if, $\bigcap_{n=1}^{+\infty} \gamma_n^p G = \{1\}$;

and clearly, if $f : G \rightarrow G'$ is a group homomorphism, then $f(\gamma_n^p G) \subset \gamma_n^p G'$ for all $n \geq 1$.

DEFINITION 1 *Let $1 \longrightarrow A \xrightarrow{\quad} B \xrightarrow{\quad \lambda \quad} C \longrightarrow 1$ be a split exact sequence.*

- If the action of C induced on $H_1(A; \mathbb{Z})$ is trivial (i.e. the action is trivial on $A^{\text{Ab}} = A/[A, A]$), we say that B is an almost direct product of A and C .
- If the action of C induced on $H_1(A; \mathbb{F}_p)$ is trivial (i.e. the action is trivial on $A/\gamma_2^p A$), we say that B is a p -almost direct product of A and C .

Let us remark that, as in the case of almost direct products (Proposition 6.3 of [BGoGu]), the fact to be a p -almost direct product does not depend on the choice of the section.

PROPOSITION 3 *Let $1 \longrightarrow A \longrightarrow B \xrightarrow{\lambda} C \longrightarrow 1$ be a split exact sequence of groups. Let σ, σ' be sections for λ , and suppose that the induced action of C on A via σ on $H_1(A; \mathbb{F}_p)$ is trivial. Then the same is true for the section σ' .*

Proof. Let $a \in A$ and $c \in C$. By hypothesis, $\sigma(c) a (\sigma(c))^{-1} \equiv a \pmod{\gamma_2^p A}$. Let σ' be another section for λ . Then $\lambda \circ \sigma'(c) = \lambda \circ \sigma(c)$, and so $\sigma'(c) (\sigma(c))^{-1} \in \text{Ker}(\lambda)$. Thus there exists $a' \in A$ such that $\sigma'(c) = a' \sigma(c)$, and hence

$$\sigma'(c) a (\sigma'(c))^{-1} \equiv a' \sigma(c) a (\sigma(c))^{-1} a'^{-1} \equiv a' a a'^{-1} \equiv a \pmod{\gamma_2^p A}.$$

Thus the induced action of C on $H_1(A; \mathbb{F}_p)$ via σ' is also trivial. \square

The first goal of this section is to prove the following Theorem (see Theorem 3.1 in [FR] for an analogous result for almost direct products).

THEOREM 4 *Let $1 \longrightarrow A \longrightarrow B \xrightarrow{\lambda} C \longrightarrow 1$ be a split exact sequence where B is a p -almost direct product of A and C . Then, for all $n \geq 1$, one has a split exact sequence*

$$1 \longrightarrow \gamma_n^p A \longrightarrow \gamma_n^p B \xrightarrow{\lambda_n} \gamma_n^p C \longrightarrow 1$$

$\swarrow \sigma_n$

where λ_n and σ_n are restrictions of λ and σ .

We shall need the following preliminary result.

LEMMA 1 *Under the hypotheses of Theorem 4, one has, for all $m, n \geq 1$*

$$[\gamma_m^p C', \gamma_n^p A] \subset \gamma_{m+n}^p A$$

where C' denotes $\sigma(C)$.

Proof. First, we prove by induction on n that $[C', \gamma_n^p A] \subset \gamma_{n+1}^p A$ for all $n \geq 1$. The cas $n = 1$ corresponds to the hypotheses: the action of C on $H_1(A; \mathbb{F}_p) = A/\gamma_2^p A$ is trivial if, and only if, $[C', A] \subset \gamma_2^p A$. Thus, suppose that $[C', \gamma_n^p A] \subset \gamma_{n+1}^p A$ for some $n \geq 1$ and let us prove that $[C', \gamma_{n+1}^p A] \subset \gamma_{n+2}^p A$. In view of the definition of $\gamma_{n+1}^p A$, we have to prove that $[C', [A, \gamma_n^p A]] \subset \gamma_{n+2}^p A$ and $[C', (\gamma_n^p A)^p] \subset \gamma_{n+2}^p A$. For the first case, we use a classical result (see [MKS], theorem 5.2) which says

$$[C', [A, \gamma_n^p A]] = [\gamma_n^p A, [C', A]] [A, [\gamma_n^p A, C']].$$

We have just seen that $[C', A] \subset \gamma_2^p A$ thus $[\gamma_n^p A, [C', A]] \subset [\gamma_n^p A, \gamma_2^p A] \subset \gamma_{n+2}^p A$. Then, the induction hypotheses says that $[\gamma_n^p A, C'] \subset \gamma_{n+1}^p A$ thus $[A, [\gamma_n^p A, C']] \subset [A, \gamma_{n+1}^p A] \subset \gamma_{n+2}^p A$. The second case works as follows: for $c \in C'$ and $x \in \gamma_n^p A$, one has, using the fact that $[u, vw] = [u, w][u, v][[u, v], w]$ (see [MKS])

$$[c, x^p] = [c, x][c, x^{p-1}][[c, x^{p-1}], x] = \dots = [c, x]^p [[c, x], x] [[c, x^2], x] \dots [[c, x^{p-1}], x].$$

Since $c \in C'$ and $x \in \gamma_n^p A$, one has $[c, x^i] \in [C', \gamma_n^p A] \subset \gamma_{n+1}^p A$ for all i , $1 \leq i \leq p-1$, which leads to $[c, x]^p \in (\gamma_{n+1}^p A)^p \subset \gamma_{n+2}^p A$ and $[[c, x^i], x] \in [\gamma_{n+1}^p A, A] \subset \gamma_{n+2}^p A$.

Now, we suppose that $[\gamma_m^p C', \gamma_n^p A] \subset \gamma_{m+n}^p A$ for some $m \geq 1$ and all $n \geq 1$ and prove that $[\gamma_{m+1}^p C', \gamma_n^p A] \subset \gamma_{m+n+1}^p A$. As above, there are two cases which work on the same way:

$$(i) \quad \begin{aligned} [[C', \gamma_m^p C'], \gamma_n^p A] &= [[\gamma_n^p A, C'], \gamma_m^p C'] [[\gamma_m^p C', \gamma_n^p A], C'] \\ &\subset [\gamma_{n+1}^p A, \gamma_m^p C'] [\gamma_{m+n}^p A, C'] \\ &\subset \gamma_{m+n+1}^p A. \end{aligned}$$

(ii) For $c \in \gamma_m^p C'$ and $x \in \gamma_n^p A$, one has

$$[c^p, x] = [c, [x, c^{p-1}]] [c^{p-1}, x] [c, x] = \cdots = [c, [x, c^{p-1}]] \cdots [c, [x, c]] [c, x]^p$$

which is an element of $\gamma_{m+n+1}^p A$ by induction hypotheses. \square

Proof of Theorem 4. Restrictions of λ and σ give rise to morphisms $\lambda_n : \gamma_n^p B \rightarrow \gamma_n^p C$ and $\sigma_n : \gamma_n^p C \rightarrow \gamma_n^p B$ satisfying $\lambda_n \circ \sigma_n = \text{Id}_{\gamma_n^p C}$, σ_n is onto and $\text{Ker}(\lambda_n) = A \cap \gamma_n^p B$. Thus, we need to prove that $A \cap \gamma_n^p B = \gamma_n^p A$ for all $n \geq 1$. Clearly one has $\gamma_n^p A \subset A \cap \gamma_n^p B$. In order to prove the converse inclusion, we follow the method developed in [FR] for almost semi-direct product and define $\tau : B \rightarrow B$ by $\tau(b) = (\sigma\lambda(b))^{-1}b$. This map has the following properties:

(i) since $\lambda\sigma = \text{Id}_C$, $\tau(B) \subset A$;

(ii) for $x \in B$, $\tau(x) = x$ if, and only if, $x \in A$;

(iii) for $(b_1, b_2) \in B^2$, $\tau(b_1 b_2) = [\sigma\lambda(b_2), \tau(b_1)^{-1}] \tau(b_1) \tau(b_2)$;

(iv) for $b \in B$, setting $a = \tau(b)$ and $c = \sigma\lambda(b)$, we get $b = ca$ with $c \in C' = \sigma(C)$ and $a \in A$, this decomposition being unique.

We claim that $\tau(\gamma_n^p B) \subset \gamma_n^p A$ for all $n \geq 1$. From this, we conclude easily the proof: if $x \in A \cap \gamma_n^p B$, then $x = \tau(x) \in \gamma_n^p A$.

One has $\tau(\gamma_1^p B) \subset \gamma_1^p A$. Suppose inductively that $\tau(\gamma_n^p B) \subset \gamma_n^p A$ for some $n \geq 1$ and let us prove that $\tau(\gamma_{n+1}^p B) \subset \gamma_{n+1}^p A$. Suppose first that x is an element of $\gamma_n^p B$. Then using (iii) we get:

$$\begin{aligned} \tau(x^p) &= [\sigma\lambda(x), \tau(x^{p-1})^{-1}] \tau(x^{p-1}) \tau(x) \\ &\vdots \\ &= [\sigma\lambda(x), \tau(x^{p-1})^{-1}] [\sigma\lambda(x), \tau(x^{p-2})^{-1}] \cdots [\sigma\lambda(x), \tau(x)^{-1}] \tau(x)^p. \end{aligned}$$

Since $\sigma\lambda(x) \in \gamma_n^p C'$ and, by induction hypotheses, $\tau(x^i) \in \gamma_n^p A$ for $1 \leq i \leq p-1$, we get $\tau(x^p) \in [\gamma_n^p C', \gamma_n^p A] \cdot (\gamma_n^p A)^p \subset \gamma_{n+1}^p A$ by lemma 1 : this prove that $\tau((\gamma_n^p B)^p) \subset \gamma_{n+1}^p A$. Next, let $b \in B$ and $x \in \gamma_n^p B$. Setting $a = \tau(b) \in A$, $y = \tau(x) \in \gamma_n^p A$ by induction hypotheses, $c = \sigma\lambda(b) \in C'$ and $z = \sigma\lambda(x) \in \gamma_n^p C'$, we get

$$\begin{aligned} \tau([b, x]) &= \left(\sigma\lambda([b, x]) \right)^{-1} [b, x] \\ &= [\sigma\lambda(b), \sigma\lambda(x)]^{-1} [b, x] \\ &= [c, z]^{-1} [ca, zy] = [z, c] a^{-1} c^{-1} y^{-1} z^{-1} cazy \\ &= [z, c] (a^{-1} c^{-1} y^{-1} cya) (a^{-1} y^{-1} c^{-1} z^{-1} cza) (a^{-1} y^{-1} z^{-1} azy) \\ &= [z, c] (a^{-1} [c, y] a) (a^{-1} y^{-1} [c, z] ya) (a^{-1} y^{-1} ay) (y^{-1} a^{-1} z^{-1} azy) \\ &= [z, c] (a^{-1} [c, y] a) (a^{-1} y^{-1} [c, z] ya) [a, y] (y^{-1} [a, z] y) \\ &= \left[[c, z], (a^{-1} [y, c] a) \right] (a^{-1} [c, y] a) [z, c] (a^{-1} y^{-1} [c, z] ya) [a, y] (y^{-1} [a, z] y) \end{aligned}$$

$$= \left[[c, z], (a^{-1}[y, c]a) \right] (a^{-1}[c, y]a) \left[[c, z], ya \right] [a, y] (y^{-1}[a, z]y).$$

Now, $[c, z] \in [C', \gamma_n^p C'] \subset \gamma_{n+1}^p C'$, $[y, c] \in [\gamma_n^p A, C'] \subset \gamma_{n+1}^p A$ (lemma 1) thus $[c, z], (a^{-1}[y, c]a) \in \gamma_{n+1}^p A$. Then, $[c, z], ya \in [\gamma_{n+1}^p C', A] \subset \gamma_{n+1}^p A$, $[a, y] \in [A, \gamma_n^p A] \subset \gamma_{n+1}^p A$ and $[a, z] \in [A, \gamma_n^p C'] \subset \gamma_{n+1}^p A$. Thus, $\tau([b, x]) \in \gamma_{n+1}^p A$ and $\tau([B, \gamma_n^p B]) \subset \gamma_{n+1}^p A$. \square

COROLLARY 2 *Let $1 \longrightarrow A \xrightarrow{\quad} B \xrightarrow{\quad \lambda \quad} C \longrightarrow 1$ be a split exact sequence such that B is a p -almost direct product of A and C . If A and C are residually p -finite, then B is residually p -finite.*

3.2 Augmentation ideals

Given a group G and $\mathbb{K} = \mathbb{Z}$ or \mathbb{F}_2 we will denote by $\mathbb{K}[G]$ the group ring of G over \mathbb{K} and by $\overline{\mathbb{K}[G]}$ the augmentation ideal of G . The group ring $\mathbb{K}[G]$ is filtered by the powers $\overline{\mathbb{K}[G]}^j$ of $\overline{\mathbb{K}[G]}$ and we can define the associated graded algebra $gr(\mathbb{K}[G]) = \bigoplus \overline{\mathbb{K}[G]}^j / \overline{\mathbb{K}[G]}^{j+1}$.

The following theorem provides a decomposition formula for the augmentation ideal of a 2-almost direct product (see Theorem 3.1 in [Pa] for an analogous in the case of almost direct products).

Let $A \rtimes C$ be a semi-direct product between two groups A and C . It is a classical result that the map $a \otimes c \mapsto ac$ induces a \mathbb{K} -isomorphism from $\mathbb{K}[A] \otimes \mathbb{K}[C]$ to $\mathbb{K}[A \rtimes C]$. Identifying these two \mathbb{K} -modules, we have the following:

THEOREM 5 *If $A \rtimes C$ is a 2-almost direct product then :*

$$\overline{\mathbb{F}_2[A \rtimes C]}^k = \sum_{i+h=k} \overline{\mathbb{F}_2[A]}^i \otimes \overline{\mathbb{F}_2[C]}^h \quad \text{for all } k.$$

Proof. We sketch the proof which is almost verbatim the same as the proof of Theorem 3.1 in [Pa]. Let $R_k = \sum_{i+h=k} \overline{\mathbb{F}_2[A]}^i \otimes \overline{\mathbb{F}_2[C]}^h$; R_k is a descending filtration on $\mathbb{F}_2[A] \otimes \mathbb{F}_2[C]$, and with the above identification,

we get that $R_k \subset \overline{\mathbb{F}_2[A \rtimes C]}^k$. To verify the other inclusion we have to check that $\prod_{j=1}^k (a_j c_j - 1) \in R_k$

for every a_1, \dots, a_k in A and c_1, \dots, c_k in C . Actually it is enough to verify that $e = \prod_{j=1}^k (e_j - 1) \in R_k$

either $e_j \in A$ or $e_j \in C$ (see Theorem 3.1 in [Pa] for a proof of this fact): we call e a *special* element. We associate to a special element e an element in $\{0, 1\}^k$: let $type(e) = (\delta(e_1), \dots, \delta(e_k))$ where $\delta(e_j) = 0$ if $e_j \in A$ and $\delta(e_j) = 1$ if $e_j \in C$. We will say that the special element e is *standard* if

$$type(e) = (\overbrace{0, \dots, 0}^i, \overbrace{1, \dots, 1}^h)$$

In this case $e \in \overline{\mathbb{F}_2[A]}^i \otimes \overline{\mathbb{F}_2[C]}^h \subset R_k$ and we are done. We claim that we can reduce all special elements to linear combinations of standard elements. If e is not standard, then it must be of the form

$$e = \prod_{i=1}^r (a_i - 1) \prod_{j=1}^s (c_j - 1)(c - 1)(a - 1) \prod_{l=1}^t (e_l - 1)$$

where $a_1, \dots, a_r, a \in A$, $c_1, \dots, c_s, c \in A$, $\tilde{e} = \prod_{l=1}^t (e_l - 1)$ is special and $r + s + t + 2 = k$. Therefore

$$\text{type}(e) = (\overbrace{0, \dots, 0}^r, \overbrace{1, \dots, 1}^s, 1, 0, \delta(e_1), \dots, \delta(e_t)).$$

Now we can use the assumption that $A \rtimes C$ is a 2-almost direct product to claim that one has commutation relations in $\mathbb{Z}[A \rtimes C]$ expressing the difference $(c - 1)(a - 1) - (a - 1)(c - 1)$ as a linear combination of terms of the form

$$(a' - 1)(a'' - 1)c \quad \text{with } a', a'' \in A$$

for any $a \in A$ and $c \in C$. In fact,

$$(c - 1)(a - 1) - (a - 1)(c - 1) = ca - ac = (cac^{-1}a^{-1} - 1)ac = (f - 1)ac$$

where $f = [c^{-1}, a^{-1}] \in [C, A] \subset \gamma_2^2(A)$ by lemma 1. We can decompose f as $f = h_1 k_1 \cdots h_m k_m$ where, for $j = 1, \dots, m$, h_j belongs to $[A, A]$ and $k_j = (k'_j)^2$ for some $k'_j \in A$. One knows (see for instance [Ch] p. 194) that for $j = 1, \dots, m$ $(h_j - 1)$ is a linear combination of terms of the form

$$(h'_j - 1)(h''_j - 1)\alpha_j \quad \text{with } h'_j, h''_j, \alpha_j \in A.$$

On the other hand for $j = 1, \dots, m$ we have also that

$$(k_j - 1) = (k'_j - 1)(k'_j - 1) \quad \text{with } k'_j \in A \quad \text{since the coefficients are } \mathbb{F}_2.$$

Then, recalling that $(hk - 1) = (h - 1)k + (k - 1)$ for any $h, k \in A$, we can conclude that $f - 1$ can be rewritten as a linear combination of terms of the form

$$(f' - 1)(f'' - 1)\alpha \quad \text{with } f', f'', \alpha \in A$$

and that $(c - 1)(a - 1) - (a - 1)(c - 1)$ is a linear combination of terms of the form

$$(f' - 1)(f'' - 1)\alpha c \quad \text{with } f', f'', \alpha \in A.$$

Rewriting $(f'' - 1)\alpha$ as $(f''\alpha - 1) - (\alpha - 1)$ we obtain that the difference $(c - 1)(a - 1) - (a - 1)(c - 1)$ can be seen as a linear combination of terms of the form

$$(a' - 1)(a'' - 1)c \quad \text{with } a', a'' \in A.$$

Therefore e can be rewritten as a sum whose first term is the special element

$$e' = \prod_{i=1}^r (a_i - 1) \prod_{j=1}^s (c_j - 1) (a - 1)(c - 1) \prod_{l=1}^t (e_l - 1)$$

and whose second term is a linear combination of elements of the form $e''c$ where

$$e'' = \prod_{i=1}^r (a_i - 1) \prod_{j=1}^s (c_j - 1) (a' - 1)(a'' - 1) \prod_{l=1}^t (ce_l c^{-1} - 1)c.$$

Using the lexicographic order from the left, one has $\text{type}(e) > \text{type}(e')$ and $\text{type}(e) > \text{type}(e'')$.

By induction on the lexicographic order we infer that e' and e'' belong to R_k : since $R_k \cdot c \subset R_k$ for any $c \in C$ it follows that e belongs to R_k and we are done. \square

4 The closed case

4.1 A presentation of $P_n(N_g)$ and induced identities

We recall a group presentation of $P_n(N_g)$ given in [GG3]: the geometric interpretation of generators is provided in Figure 2.

THEOREM 6 ([GG3]) *For $g \geq 2$ and $n \geq 1$, $P_n(N_g)$ has the following presentation:*

generators: $(B_{i,j})_{1 \leq i < j \leq n}$ and $(\rho_{k,l})_{\substack{1 \leq k \leq n \\ 1 \leq l \leq g}}$.

relations: (a) for all $1 \leq i < j \leq n$ and $1 \leq r < s \leq n$,

$$B_{r,s} B_{i,j} B_{r,s}^{-1} = \begin{cases} B_{i,j} & \text{if } i < r < s < j \text{ or } r < s < i < j & \text{(a}_1\text{)} \\ B_{i,j}^{-1} B_{r,j}^{-1} B_{i,j} B_{r,j} B_{i,j} & \text{if } r < i = s < j & \text{(a}_2\text{)} \\ B_{s,j}^{-1} B_{i,j} B_{s,j} & \text{if } i = r < s < j & \text{(a}_3\text{)} \\ B_{s,j}^{-1} B_{r,j}^{-1} B_{s,j} B_{r,j} B_{i,j} B_{r,j}^{-1} B_{s,j}^{-1} B_{r,j} B_{s,j} & \text{if } r < i < s < j & \text{(a}_4\text{)} \end{cases}$$

(b) for all $1 \leq i < j \leq n$ and $1 \leq k, l \leq g$,

$$\rho_{i,k} \rho_{j,l} \rho_{i,k}^{-1} = \begin{cases} \rho_{j,l} & \text{if } k < l & \text{(b}_1\text{)} \\ \rho_{j,k}^{-1} B_{i,j}^{-1} \rho_{j,k}^2 & \text{if } k = l & \text{(b}_2\text{)} \\ \rho_{j,k}^{-1} B_{i,j}^{-1} \rho_{j,k} B_{i,j}^{-1} \rho_{j,l} B_{i,j} \rho_{j,k}^{-1} B_{i,j} \rho_{j,k} & \text{if } k > l & \text{(b}_3\text{)} \end{cases}$$

(c) for all $1 \leq i \leq n$, $\rho_{i,1}^2 \cdots \rho_{i,g}^2 = T_i$ where $T_i = B_{1,i} \cdots B_{i-1,i} B_{i,i+1} \cdots B_{i,n}$ (c)

(d) for all $1 \leq i < j \leq n$, $1 \leq k \leq n$, $k \neq j$ and $1 \leq l \leq g$,

$$\rho_{k,l} B_{i,j} \rho_{k,l}^{-1} = \begin{cases} B_{i,j} & \text{if } k < i \text{ or } j < k & \text{(d}_1\text{)} \\ \rho_{j,l}^{-1} B_{i,j}^{-1} \rho_{j,l} & \text{if } k = i & \text{(d}_2\text{)} \\ \rho_{j,l}^{-1} B_{k,j}^{-1} \rho_{j,l} B_{k,j}^{-1} B_{i,j} B_{k,j} \rho_{j,l}^{-1} B_{k,j} \rho_{j,l} & \text{if } i < k < j & \text{(d}_3\text{)} \end{cases}$$




Figure 2: Generators of $P_n(N_g)$

For $1 \leq k \leq g$, let us consider the element a_k in $P_n(N_g)$ given by $a_k = \rho_{k,g-1} \rho_{k,g}$ and set $U = a_n \cdots a_2$.

LEMMA 2 *The following relations holds in $P_n(N_g)$:*

- (1) $[\rho_{i,k}, \rho_{j,k}^{-1}] = B_{i,j}^{-1}$ for $1 \leq i < j \leq n$ and $1 \leq k \leq g$; (e)
- (2) U commutes with $\rho_{1,l}$ for $1 \leq l \leq g - 2$; (f₁)
- (3) $[\rho_{1,g-1}, U^{-1}] = T_1^{-1}$; (f₂)
- (4) $a_k a_j a_k^{-1}$ commutes with $B_{i,k}$ for $1 \leq i < j < k \leq n$; (g)
- (5) $a_n a_{n-1} \cdots a_1$ commutes with $B_{j,k}$ for $1 \leq j < k \leq n$; (h)
- (6) U commutes with $B_{i,j}$ for $2 \leq i < j \leq n$; (i)
- (7) $a_n a_{n-1} \cdots a_1$ commutes with T_1 ; (j)
- (8) T_1 commutes with $B_{j,k}$ for $2 \leq j < k \leq n$; (k)

Proof. Some of these identities can easily be verified drawing corresponding braids. This is the case for example for the first, the fourth and the eighth ones (see figure 3, 4 and 5). Let us give an algebraic proof for the others.




Figure 3: identity (e)




Figure 4: identity (g)

- (2) This is a direct consequence of the definitions of the a_i 's and U , and relations (b₁).




Figure 5: identity (k)

(3) By relation (b₁), $\rho_{1,g-1}$ commutes with $\rho_{j,g}$ for $2 \leq j \leq n$. Thus, using relation (e), we get:

$$\begin{aligned}
\rho_{1,g-1}^{-1} U \rho_{1,g-1} &= \rho_{1,g-1}^{-1} a_n \cdots a_2 \rho_{1,g-1} \\
&= \rho_{1,g-1}^{-1} (\rho_{n,g-1} \rho_{n,g}) \cdots (\rho_{2,g-1} \rho_{2,g}) \rho_{1,g-1} \\
&= (B_{1,n}^{-1} \rho_{n,g-1} \rho_{n,g}) \cdots (B_{1,2}^{-1} \rho_{2,g-1} \rho_{2,g}) \\
&= B_{1,n}^{-1} B_{1,n-1}^{-1} \cdots B_{1,2}^{-1} (\rho_{n,g-1} \rho_{n,g}) \cdots (\rho_{2,g-1} \rho_{2,g}) \quad \text{by (d}_1\text{)} \\
&= T_1^{-1} U.
\end{aligned}$$

(5) Let j and k be integers such that $1 \leq j < k \leq n$. By (d₁), a_1, \dots, a_{j-1} commute with $B_{j,k}$. Then, one has

$$\begin{aligned}
a_j B_{j,k} a_j^{-1} &= \rho_{j,g-1} \rho_{j,g} B_{j,k} \rho_{j,g}^{-1} \rho_{j,g-1}^{-1} \\
&= \rho_{j,g-1} \rho_{k,g}^{-1} B_{j,k}^{-1} \rho_{k,g} \rho_{j,g-1}^{-1} \quad \text{by (d}_2\text{)} \\
&= \rho_{k,g}^{-1} \rho_{j,g-1} B_{j,k}^{-1} \rho_{j,g-1}^{-1} \rho_{k,g} \quad \text{by (b}_1\text{)} \\
&= \rho_{k,g}^{-1} \rho_{k,g-1}^{-1} B_{j,k} \rho_{k,g-1} \rho_{k,g} \quad \text{by (d}_2\text{)} \\
&= a_k^{-1} B_{j,k} a_k
\end{aligned}$$

and we get

$$\begin{aligned}
a_n \cdots a_1 B_{j,k} a_1^{-1} \cdots a_n^{-1} &= a_n \cdots a_{k+1} a_k a_{k-1} \cdots a_{j+1} a_k^{-1} B_{j,k} a_k a_{j+1}^{-1} \cdots a_{k-1}^{-1} a_k^{-1} a_{k+1}^{-1} \cdots a_n^{-1} \\
&= a_n \cdots a_{k+1} B_{j,k} a_{k+1}^{-1} \cdots a_n^{-1} \quad \text{by (g)} \\
&= B_{j,k} \quad \text{by (d}_1\text{)}.
\end{aligned}$$

(6) By (d₁), $a_1 = \rho_{1,g-1} \rho_{1,g}$ commutes with $B_{i,j}$ for $2 \leq i < j \leq n$. Thus, relation (i) is a direct consequence of (h).

(7) A direct consequence of (h) since $T_1 = B_{1,2} \cdots B_{1,n}$. □

4.2 The pure braid group $P_n(N_g)$ is residually 2-finite

Following [GG1], one has, for $g \geq 2$, a split exact sequence

$$1 \longrightarrow P_{n-1}(N_{g,1}) \xrightarrow{\mu} P_n(N_g) \xrightarrow{\lambda} P_1(N_g) = \pi_1(N_g) \longrightarrow 1 \quad (1)$$

where λ is induced by the map which forgets all strands except the first one, and μ is defined by capping the boundary component by a disc with one marked point (the first strand in $P_n(N_g)$). According to the definition of μ and to Theorem A, $\text{Im}(\mu)$ is generated by $\{\rho_{i,k}, 2 \leq i \leq n, 1 \leq k \leq g\} \cup \{B_{i,j}, 2 \leq i < j \leq n\}$.

The section given in [GG1] is geometric, *i.e.* it is induced by a crossed section at the level of fibrations. In order to study the action of $\pi_1(N_g)$ on $P_{n-1}(N_{g,1})$, we need an algebraic one. Recall that $\pi_1(N_g)$ has a group presentation with generators p_1, \dots, p_g and the single relation $p_1^2 \cdots p_g^2 = 1$. We define the set map $\sigma : \pi_1(N_g) \rightarrow P_n(N_g)$ by setting

$$\sigma(p_i) = \begin{cases} \rho_{1,i} & \text{for } 1 \leq i \leq g-3, \\ \rho_{1,g-2}U^{-1} & \text{for } i = g-2, \\ U\rho_{1,g-1} & \text{for } i = g-1, \\ \rho_{1,g}T_1^{-1} & \text{for } i = g. \end{cases}$$

PROPOSITION 4 *The map σ is a well defined homomorphism satisfying $\lambda \circ \sigma = \text{Id}_{\pi_1(N_g)}$.*

Proof. Since $\lambda(\rho_{1,i}) = p_i$ for all $1 \leq i \leq g$ and $\lambda(U) = \lambda(T_1) = 1$, one has clearly $\lambda\sigma = \text{Id}_{\pi_1(N_g)}$ if σ is a group homomorphism. Thus, we have just to prove that $\sigma(p_1)^2 \cdots \sigma(p_g)^2 = 1$:

$$\begin{aligned} \sigma(p_1)^2 \cdots \sigma(p_g)^2 &= (\rho_{1,1}^2 \cdots \rho_{1,g-3}^2)(\rho_{1,g-2}U^{-1})^2(U\rho_{1,g-1})^2(\rho_{1,g}T_1^{-1})^2 \\ &= \rho_{1,1}^2 \cdots \rho_{1,g-3}^2 \rho_{1,g-2}^2 \underbrace{U^{-1}\rho_{1,g-2}\rho_{1,g-1}Ua_1T_1^{-1}}_{\rho_{1,g}T_1^{-1}} \rho_{1,g}T_1^{-1} \\ &= \rho_{1,1}^2 \cdots \rho_{1,g-3}^2 \rho_{1,g-2}^2 \underbrace{U^{-1}\rho_{1,g-1}Ua_1T_1^{-1}}_{\rho_{1,g}T_1^{-1}} \rho_{1,g}T_1^{-1} \quad \text{by (f}_1\text{)} \\ &= \rho_{1,1}^2 \cdots \rho_{1,g-3}^2 \rho_{1,g-2}^2 \rho_{1,g-1}^2 U^{-1}T_1 \underbrace{Ua_1T_1^{-1}}_{\rho_{1,g}T_1^{-1}} \rho_{1,g}T_1^{-1} \quad \text{by (f}_2\text{)} \\ &= \rho_{1,1}^2 \cdots \rho_{1,g-3}^2 \rho_{1,g-2}^2 \rho_{1,g-1}^2 U^{-1}T_1T_1^{-1}Ua_1\rho_{1,g}T_1^{-1} \quad \text{by (j)} \\ &= \rho_{1,1}^2 \cdots \rho_{1,g-3}^2 \rho_{1,g-2}^2 \rho_{1,g-1}^2 a_1\rho_{1,g}T_1^{-1} \\ &= \rho_{1,1}^2 \cdots \rho_{1,g-3}^2 \rho_{1,g-2}^2 \rho_{1,g-1}^2 \rho_{1,g}^2 T_1^{-1} \\ &= 1 \quad \text{by (c)}. \end{aligned}$$

□

So, the exact sequence (1) splits. In order to apply Theorem 4, we have to prove that the action of $\pi_1(N_g)$ on $P_{n-1}(N_{g,1})$ is trivial on $H_1(P_{n-1}(N_{g,1}); \mathbb{F}_2)$. This is the claim of the following proposition.

PROPOSITION 5 *For all $x \in \text{Im}(\sigma)$ and $a \in \text{Im}(\mu)$, one has $[x^{-1}, a^{-1}] = xax^{-1}a^{-1} \in \gamma_2^2(\text{Im}(\mu))$.*

Proof. It is enough to prove the result for $a \in \{B_{j,k}, 2 \leq j < k \leq n\} \cup \{\rho_{j,l}, 2 \leq j \leq n \text{ and } 1 \leq l \leq g\}$ and $x \in \{\sigma(p_1), \dots, \sigma(p_g)\}$, respectively sets of generators of $\text{Im}(\mu)$ and $\text{Im}(\sigma)$. Suppose first that $2 \leq j < k \leq n$. One has:

- $[\sigma(p_i)^{-1}, B_{j,k}^{-1}] = [\rho_{1,i}^{-1}, B_{j,k}^{-1}] = 1$ for $1 \leq i \leq g-3$ by (d₁);
- $[\sigma(p_{g-2})^{-1}, B_{j,k}^{-1}] = [U\rho_{1,g-2}^{-1}, B_{j,k}^{-1}] = 1$ by (d₁) and (i);
- $[\sigma(p_{g-1})^{-1}, B_{j,k}^{-1}] = [\rho_{1,g-1}^{-1}U^{-1}, B_{j,k}^{-1}] = 1$ by (d₁) and (i);
- $[\sigma(p_g)^{-1}, B_{j,k}^{-1}] = [T_1\rho_{1,g}^{-1}, B_{j,k}^{-1}] = 1$ by (d₁) and (k).

Now, let j and l be integers such that $2 \leq j \leq n$ and $1 \leq l \leq g$ and let us first prove that $[\rho_{1,i}^{-1}, \rho_{j,l}^{-1}] \in \gamma_2^2(\text{Im}(\mu))$ for all $i, 1 \leq i \leq n$:

- this is clear for $i < l$ by (b₁);
- for $i = l$, the relation (b₂) gives $[\rho_{1,l}^{-1}, \rho_{j,l}^{-1}] = \rho_{j,l}^{-1} B_{1,j}^{-1} \rho_{j,l}$. But

$$B_{1,j}^{-1} = B_{2,j} \cdots B_{j-1,j} B_{j,j+1} \cdots B_{j,n} \rho_{j,g}^{-2} \cdots \rho_{j,1}^{-2} \quad (\text{relation (c)})$$

is an element of $\gamma_2^2(\text{Im}(\mu))$ by (e), thus we get $[\rho_{1,l}^{-1}, \rho_{j,l}^{-1}] \in \gamma_2^2(\text{Im}(\mu))$.

- If $l < i$ then $[\rho_{1,i}^{-1}, \rho_{j,i}^{-1}] = [B_{1,j} \rho_{j,i}^{-1} B_{1,j} \rho_{j,i}, \rho_{j,i}^{-1}]$ by (b₃) so $[\rho_{1,i}^{-1}, \rho_{j,i}^{-1}] \in \gamma_2(\text{Im}(\mu))$ since $\rho_{j,i}$, $\rho_{j,i}^{-1}$ and $B_{1,j}$ are elements of $\text{Im}(\mu)$.

From this, we deduce the following facts.

$$(1) \quad [\sigma(p_i)^{-1}, \rho_{j,i}^{-1}] \in \gamma_2^2(\text{Im}(\mu)) \text{ for } i \leq g-3 \text{ since } \sigma(p_i) = \rho_{1,i}.$$

(2) $[\sigma(p_{g-2})^{-1}, \rho_{j,l}^{-1}] = [U \rho_{1,g-2}^{-1}, \rho_{j,l}^{-1}] = \rho_{1,g-2} [U, \rho_{j,k}^{-1}] \rho_{1,g-2}^{-1} [\rho_{1,g-2}^{-1}, \rho_{j,l}^{-1}]$. But U and $\rho_{j,l}^{-1}$ are elements of $\text{Im}(\mu)$ thus $[U, \rho_{j,k}^{-1}] \in \Gamma_2(\text{Im}(\mu)) \subset \gamma_2^2(\text{Im}(\mu))$. Consequently, $\rho_{1,g-2} [U, \rho_{j,k}^{-1}] \rho_{1,g-2}^{-1} \in \gamma_2^2(\text{Im}(\mu))$ since $\gamma_2^2(\text{Im}(\mu))$ is a characteristic subgroup of $\text{Im}(\mu)$ and $\text{Im}(\mu)$ is normal in $P_n(N_g)$. Thus, we get $[\sigma(p_{g-2})^{-1}, \rho_{j,l}^{-1}] \in \gamma_2^2(\text{Im}(\mu))$. In the same way, one has

$$[\sigma(p_{g-1})^{-1}, \rho_{j,l}^{-1}] = [\rho_{1,g-1}^{-1} U^{-1}, \rho_{j,l}^{-1}] = U [\rho_{1,g-1}^{-1}, \rho_{j,l}^{-1}] U^{-1} [U, \rho_{j,l}^{-1}] \in \gamma_2^2(\text{Im}(\mu)).$$

(4) At last,

$$[\sigma(p_g)^{-1}, \rho_{j,l}^{-1}] = [T_1 \rho_{1,g}^{-1}, \rho_{j,l}^{-1}] = \rho_{1,g} [T_1, \rho_{j,l}^{-1}] \rho_{1,g}^{-1} [\rho_{1,g}, \rho_{j,l}^{-1}] \in \gamma_2^2(\text{Im}(\mu))$$

since $T_1, \rho_{j,l} \in \text{Im}(\mu)$ and $[\rho_{1,g}, \rho_{j,l}^{-1}] \in \gamma_2^2(\text{Im}(\mu))$. □

We are now ready to prove the main result of this section

THEOREM 7 *For all $g \geq 2$ and $n \geq 1$, the pure braid group $P_n(N_g)$ is residually 2-finite.*

Proof. Proposition 4 says that the sequence

$$1 \longrightarrow P_{n-1}(N_{g,1}) \longrightarrow P_n(N_g) \longrightarrow \pi_1(N_g) \longrightarrow 1$$

splits. Now $P_{n-1}(N_{g,1})$ is residually 2-finite (Theorem 3). It is proved in [B1] and [B2] that $\pi_1(N_g)$ is residually free for $g \geq 4$, so it is residually 2-finite. This result is proved in [LM] (lemma 8.9) for $g = 3$. When $g = 2$, $\pi_1(N_2)$ has presentation $\langle a, b \mid aba^{-1} = b^{-1} \rangle$ so is a 2-almost direct product of \mathbb{Z} by \mathbb{Z} . Since \mathbb{Z} is residually 2-finite, $\pi_1(N_2)$ is residually 2-finite by corollary 2. So, using Proposition 5 and Corollary 2, we can conclude that $P_n(N_g)$ is residually 2-finite. □

REMARK 3 It follows from the proof of Theorem 7 that, when $g > 2$, if $P_n(N_{g,1})$ is residually p -finite for some $p \neq 2$ then the pure braid group $P_n(N_g)$ is also residually p -finite.

4.3 The case $P_n(\mathbb{RP}^2)$

The main reason to exclude $N_1 = \mathbb{RP}^2$ in Theorem 7 is that the exact sequence (1) doesn't exist in this case, but forgetting at most $n - 2$ strands we get the following exact sequence ($1 \leq m \leq n - 2$; see [VB])

$$1 \longrightarrow P_m(N_{1,n-m}) \longrightarrow P_n(\mathbb{RP}^2) \longrightarrow P_{n-m}(N_g) \longrightarrow 1 .$$

This sequence splits if, and only if $n = 3$ and $m = 1$ (see [GG2]). Thus, what we know is the following:

- $P_1(\mathbb{RP}^2) = \pi_1(\mathbb{RP}^2) = \mathbb{Z}/2\mathbb{Z}$: $P_1(\mathbb{RP}^2)$ is a 2-group.
- $P_2(\mathbb{RP}^2) = Q_8$, the quaternion group (see [VB]): $P_2(\mathbb{RP}^2)$ is a 2-group.
- One has the exact sequence

$$1 \longrightarrow P_1(N_{1,2}) \longrightarrow P_3(\mathbb{RP}^2) \longrightarrow P_2(\mathbb{RP}^2) \longrightarrow 1$$

where $P_1(N_{1,2}) = \pi_1(N_{1,2})$ is a free group of rank 2, thus is residually 2-finite. Since $P_2(\mathbb{RP}^2)$ is 2-finite, we can conclude that $P_3(\mathbb{RP}^2)$ is residually 2-finite using lemma 1.5 of [Gr].

Appendix: a group presentation for $P_n(N_{g,b})$

Here we apply classical method (see [B, GG3]) to give a presentation of the n^{th} pure braid group of a nonorientable surface with boundary. Since $b \geq 1$, we'll see $N_{g,b}$ as a disc D^2 with $g + b - 1$ open discs removed and g Möbius strips glued on g boundary components so obtained (see figure 6).




Figure 6: Generators $x_{k,t}$ for $P_n(N_{g,b})$, $b \geq 1$

THEOREM A For $g \geq 1$, $b \geq 1$ and $n \geq 1$, $P_n(N_{g,b})$ has the following presentation:

generators: $(B_{i,j})_{1 \leq i < j \leq n}$, $(\rho_{k,t})_{\substack{1 \leq k \leq n \\ 1 \leq t \leq g}}$ and $(x_{u,t})_{\substack{1 \leq u \leq n \\ 1 \leq t \leq b-1}}$.

relations : (a) for all $1 \leq i < j \leq n$ and $1 \leq r < s \leq n$,

$$B_{r,s}B_{i,j}B_{r,s}^{-1} = \begin{cases} B_{i,j} & \text{if } i < r < s < j \text{ or } r < s < i < j & \text{(a}_1\text{)} \\ B_{i,j}^{-1}B_{r,j}^{-1}B_{i,j}B_{r,j}B_{i,j} & \text{if } r < i = s < j & \text{(a}_2\text{)} \\ B_{s,j}^{-1}B_{i,j}B_{s,j} & \text{if } i = r < s < j & \text{(a}_3\text{)} \\ B_{s,j}^{-1}B_{r,j}^{-1}B_{s,j}B_{r,j}B_{i,j}B_{r,j}^{-1}B_{s,j}^{-1}B_{r,j}B_{s,j} & \text{if } r < i < s < j & \text{(a}_4\text{)} \end{cases}$$

(b) for all $1 \leq i < j \leq n$ and $1 \leq k, l \leq g$,

$$\rho_{i,k}\rho_{j,l}\rho_{i,k}^{-1} = \begin{cases} \rho_{j,l} & \text{if } k < l & \text{(b}_1\text{)} \\ \rho_{j,k}^{-1}B_{i,j}^{-1}\rho_{j,k}^2 & \text{if } k = l & \text{(b}_2\text{)} \\ \rho_{j,k}^{-1}B_{i,j}^{-1}\rho_{j,k}B_{i,j}^{-1}\rho_{j,l}B_{i,j}\rho_{j,k}^{-1}B_{i,j}\rho_{j,k} & \text{if } k > l & \text{(b}_3\text{)} \end{cases}$$

(d) for all $1 \leq i < j \leq n$, $1 \leq k \leq n$, $k \neq j$ and $1 \leq l \leq g$,

$$\rho_{k,l}B_{i,j}\rho_{k,l}^{-1} = \begin{cases} B_{i,j} & \text{if } k < i \text{ or } j < k & \text{(d}_1\text{)} \\ \rho_{j,l}^{-1}B_{i,j}^{-1}\rho_{j,l} & \text{if } k = i & \text{(d}_2\text{)} \\ \rho_{j,l}^{-1}B_{k,j}^{-1}\rho_{j,l}B_{k,j}^{-1}B_{i,j}B_{k,j}\rho_{j,l}^{-1}B_{k,j}^{-1}\rho_{j,l} & \text{if } i < k < j & \text{(d}_3\text{)} \end{cases}$$

(l) for all $1 \leq i < j \leq n$, $1 \leq u \leq n$, $u \neq j$ and $1 \leq t \leq b-1$,

$$x_{u,t}B_{i,j}x_{u,t}^{-1} = \begin{cases} B_{i,j} & \text{if } u < i \text{ or } j < u & \text{(l}_1\text{)} \\ x_{j,t}^{-1}B_{i,j}x_{j,t} & \text{if } u = i & \text{(l}_2\text{)} \\ x_{j,t}^{-1}B_{u,j}x_{j,t}B_{u,j}^{-1}B_{i,j}B_{u,j}x_{j,t}^{-1}B_{u,j}^{-1}x_{j,t} & \text{if } i < u < j & \text{(l}_3\text{)} \end{cases}$$

(m) for all $1 \leq k, u \leq n$, $k \neq u$, $1 \leq l \leq g$ and $1 \leq t \leq b-1$,

$$x_{u,t}\rho_{k,l}x_{u,t}^{-1} = \begin{cases} \rho_{k,l} & \text{if } k < u & \text{(m}_1\text{)} \\ x_{k,t}^{-1}B_{u,k}x_{k,t}B_{u,j}^{-1}\rho_{k,l}B_{u,k}x_{k,t}^{-1}B_{u,k}^{-1}x_{k,t} & \text{if } u < k & \text{(m}_2\text{)} \end{cases}$$

(n) for all $1 \leq i < j \leq n$ and $1 \leq s, t \leq b-1$,

$$x_{i,t}x_{j,s}x_{i,t}^{-1} = \begin{cases} x_{j,s} & \text{if } t < s & \text{(n}_1\text{)} \\ x_{j,t}^{-1}B_{i,j}x_{j,t}B_{i,j}^{-1}x_{j,t} & \text{if } t = s & \text{(n}_2\text{)} \\ x_{j,t}^{-1}B_{i,j}x_{j,t}B_{i,j}^{-1}x_{j,s}B_{i,j}x_{j,t}^{-1}B_{i,j}^{-1}x_{j,t} & \text{if } s < t & \text{(n}_3\text{)} \end{cases}$$

Proof. The proof works by induction and generalizes those of [GG3] (closed non orientable case) and [B] (orientable case, possibly with boundary components). It uses the following short exact sequence obtained by forgetting the last strand (see [FN]):

$$1 \longrightarrow \pi_1(N_{g,b} \setminus \{z_1, \dots, z_n\}, z_{n+1}) \xrightarrow{\alpha} P_{n+1}(N_{g,b}) \xrightarrow{\beta} P_n(N_{g,b}) \longrightarrow 1.$$

The presentation is correct for $n = 1$: $P_1(N_{g,b}) = \pi_1(N_{g,b})$ is free on the $\rho_{1,l}$'s and $x_{1,t}$'s for $1 \leq l \leq g$ and $1 \leq t \leq b-1$. Suppose inductively that $P_n(N_{g,b})$ has the given presentation. Then, observe that $\{B_{i,n+1} / 1 \leq i \leq n\} \cup \{\rho_{n+1,l} / 1 \leq l \leq g\} \cup \{x_{n+1,t} / 1 \leq t \leq b-1\}$ is a free generators set of $\text{Im}(\alpha)$ and $(B_{i,j})_{1 \leq i < j \leq n}$, $(\rho_{k,l})_{1 \leq k \leq n}$ and $(x_{u,t})_{1 \leq u \leq n}$ are coset representative for the considered generators of $P_n(N_{g,b})$.

There are three types of relations for $P_{n+1}(N_{g,b})$. The first one comes from the relations in $\text{Im}(\alpha)$: there are none here, since this group is free. The second type comes from the relations in $P_n(N_{g,b})$: they lift to the same relations in $P_{n+1}(N_{g,b})$. Finally, the third type arrives by studying the action of $P_n(N_{g,b})$ on $\text{Im}(\alpha)$ by conjugation. We leave to the reader to verify that this action corresponds to the given relations. \square

REMARK 4 What precedes proves that $P_{n+1}(N_{g,b})$ is a semidirect product of the free group

$\pi_1(N_{g,b} \setminus \{z_1, \dots, z_n\}, z_{n+1})$ by $P_n(N_{g,b})$. Therefore, by recurrence, we get that $P_{n+1}(N_{g,b})$ is an iterated semidirect product of (finitely generated) free groups.

References

- [B1] B. Baumslag, Residual free groups, *Proc. London Math. Soc. (3)* **17** (1967), 402-418.
- [B2] G. Baumslag, On generalised free products. *Math. Z.* **78** (1962), 423-438.
- [BB] V.G. Bardakov and P. Bellingeri, On residual properties of pure braid groups of closed surfaces, *Communications in Algebra* **37** (2009), no 5, 1481-1490.
- [B] P. Bellingeri, On presentations of surface braid groups, *J. Algebra* **274** (2004), no 2, 543-563.
- [BF] P. Bellingeri and L. Funar, Braids on surfaces and finite type invariants, *C. R. Math. Acad. Sci. Paris* **338** (2004), 157-162.
- [BGG] P. Bellingeri, S. Gervais and J. Guaschi, Lower central series of Artin Tits and surface braid groups, *J. Algebra* **319** (2008), 1409-1427.
- [BGoGu] P. Bellingeri, E. Godelle and J. Guaschi, Exact sequences, lower central series and representations of surface braid groups, arxiv:1106.4982.
- [Ch] K.-T. Chen, Extensions of C^∞ function algebra by integrals and Malcev completion of π_1 , *Adv. Math.* **23** (1977), 181-210.
- [CVZ] H.-D. Coldewey, E. Vogt, H. Zieschang . Surfaces and planar discontinuous groups, Lecture Notes in Mathematics 835 (Springer, Berlin, 1980).
- [FN] E. Fadell and L. Neuwirth, Configuration spaces, *Math. Scandinavica* **10** (1962), 111-118.
- [FR] M. Falk and R. Randell, The lower central series of fiber-type arrangement, *Inv. Math.* **82** (1985), 77-88.
- [GLe] S. Garoufalidis and J. Levine, Finite type 3-manifold invariants and the structure of the Torelli group I, *Invent. Math.* **131** (1998), 541-594.
- [GG1] D. Gonçalves and J. Guaschi, On the structure of surface pure braid groups, *J. Pure Appl. Algebra* **186** (2004), 187-218.
- [GG2] D. Gonçalves and J. Guaschi, The braid groups of the projective plane and the Fadell-Neuwirth short exact sequence, *Geom. Dedicata* **130** (2007), 93-107.
- [GG3] D. Gonçalves and J. Guaschi, Braid groups of non-orientable surfaces and the Fadell-Neuwirth short exact sequence, *J. Pure Appl. Algebra* **214** (2010), 667-677.
- [Go] J. González-Meneses, Ordering pure braid groups on closed surfaces, *Pacific J. Math.* **203** (2002), nº 2, 369-378.
- [GP] J. González-Meneses and L. Paris, Vassiliev invariants for braids on surfaces, *Trans. Amer. Math. Soc.* **356** (2004), 219-243.
- [GJ] J. Guaschi and D. Juan-Pineda, A survey of surface braid groups and the lower algebraic K-theory of their group rings, to appear in "Handbook of Group Actions" *Kunming Conference Proceedings*

- [Gr] K.W. Gruenberg, Residual properties of infinite soluble groups. *Proc. London Math. Soc.* **7** (1957), 29-62.
- [H] B. Hartley, On residually finite p -groups. Symposia Mathematica, Vol. XVII (Convegno sui Gruppi Infiniti, INDAM, Roma, 1973), 225–234. Academic Press, London, 1976.
- [LM] G. Levitt and A. Minasyan, Residual properties of automorphism groups of (relatively) hyperbolic groups, *arXiv:1305.5403v2* (2013).
- [MKS] W. Magnus, A. Karrass and D. Solitar, Combinatorial group theory, Second revised edition, *Dover Publications Inc., New York*, 1976.
- [MR] R. Botto Mura and A. Rhemtulla, Orderable groups, Lecture Notes in Pure and Applied Mathematics **27**, Marcel Dekker, New York, 1977.
- [Pa] S. Papadima, The universal finite-type invariant for braids, with integer coefficients, *Topology Appl.* **118** (2002), 169-185.
- [P] L. Paris, Residual p properties of mapping class groups and surface groups, *Trans. Amer. Math. Soc.* **361** (2009), no. 5, 2487-2507.
- [PS] L. Paris, B. Szepietowski, A presentation for the mapping class groups and surface group of a nonorientable surface, *arXiv:1308.5856v1* (2013).
- [Sc] G. P. Scott, Braid groups and the group of homeomorphisms of a surface, *Proc. Cambridge Philos. Soc.* **68** (1970), 605-617.
- [St] M. Stukow, Commensurability of geometric subgroups of mapping class groups, *Geom. Dedicata* **143** (2009), 117-142.
- [Sz1] B. Szepietowski, Embedding the braid group in mapping class group, *Publ. Mat.* **54** (2010), 359-368.
- [Sz2] B. Szepietowski, Crosscap slides and the level 2 mapping class group of a nonorientable surface, *Geom. Dedicata* **160** (2012), 169-183.
- [VB] J. Van Buskirk, Braid groups of compact 2-manifolds with elements of finite order, *Trans. Amer. Math. Soc.* **122** (1966), 81-97.

PAOLO BELLINGERI, Université de Caen, CNRS UMR 6139, LMNO, Caen, 14000 (France). Email: paolo.bellingeri@math.unicaen.fr

SYLVAIN GERVAIS, Université de Nantes, CNRS-UMR 6629, Laboratoire Jean Leray, 2, rue de la Houssinière, F-44322 NANTES cedex 03 (France). Email: sylvain.gervais@univ-nantes.fr