



HAL
open science

The Li-Yau inequality and applications under a curvature-dimension condition

Dominique Bakry, François Bolley, Ivan Gentil

► **To cite this version:**

Dominique Bakry, François Bolley, Ivan Gentil. The Li-Yau inequality and applications under a curvature-dimension condition. 2014. hal-01094046v1

HAL Id: hal-01094046

<https://hal.science/hal-01094046v1>

Preprint submitted on 11 Dec 2014 (v1), last revised 21 Jul 2016 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Li-Yau inequality and applications under a curvature-dimension condition

Dominique Bakry*, François Bolley† and Ivan Gentil‡

December 12, 2014

Abstract

We prove a global Li-Yau inequality for a general Markov semigroup under a curvature-dimension condition. This inequality implies all classical Li-Yau type inequalities known to us. Moreover, on a Riemannian manifold, it proves to be equivalent to a new parabolic Harnack inequality, both in negative and positive curvature, and giving new subsequent bounds on the heat kernel of the semigroup.

Key words: Li-Yau inequality, Harnack inequality, heat kernel bounds, curvature-dimension condition, logarithmic Sobolev inequality.

1 Introduction

In their seminal paper [LY86], P. Li and S.-T. Yau proved that on a Riemannian manifold M with dimension n and non-negative Ricci curvature, for any positive function f and any $t > 0$,

$$-\Delta(\log P_t f) \leq \frac{n}{2t}, \quad (1)$$

where Δ is the Laplace-Beltrami operator on M and $(t, x) \mapsto P_t f(x)$ is the solution to the heat equation $\partial_t u = \Delta u$ with initial condition f . Equivalently, the Li-Yau inequality can be written :

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} \leq \frac{\Delta P_t f}{P_t f} + \frac{n}{2t},$$

where $|\nabla P_t f|$ stands for the length of $\nabla P_t f$. This gradient estimate is a crucial step towards parabolic Harnack inequalities and various subsequent on and off-diagonal bounds on heat kernels. It is optimal since the equality is achieved for the heat kernel on the Euclidean space, that is, when f converges to a Dirac mass.

*Institut de mathématiques de Toulouse, Umr Cnrs 5219, Université Toulouse 3. bakry@math.univ-toulouse.fr

†Laboratoire de probabilités et modèles aléatoires, Umr Cnrs 7599, Université Paris 6. francois.bolley@upmc.fr

‡Institut Camille Jordan, Umr Cnrs 5208, Université Claude Bernard Lyon 1. gentil@math.univ-lyon1.fr

This inequality has been generalised to Riemannian manifolds with a Ricci curvature bounded from below by a real constant ρ . Still in [LY86], P. Li and S.-T. Yau proved that if the Ricci curvature is bounded from below by $\rho = -K < 0$ then for any $\alpha > 1$

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} \leq \alpha \frac{\Delta P_t f}{P_t f} + \frac{n\alpha^2 K}{2(\alpha - 1)} + \frac{n\alpha^2}{2t}.$$

Improving this inequality into a form which would be optimal for small, large as well as intermediate times has been the purpose of many subsequent works. In this direction and still in the negative curvature case, let us mention the diverse bounds

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} \leq \alpha \frac{\Delta P_t f}{P_t f} + \frac{n\alpha^2 K}{4(\alpha - 1)} + \frac{n\alpha^2}{2t} \quad (2)$$

for any $\alpha > 1$, derived by B. Davies in [Dav89, Chapter 5.3], and

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} - \frac{\Delta P_t f}{P_t} \leq \sqrt{2nK} \sqrt{\frac{|\nabla P_t f|^2}{P_t f} + \frac{n}{2t} + 2nK} + \frac{n}{2t}$$

derived by S.-T. Yau [Yau95], itself improved by the first author and Z. Qian [BQ99], as

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} - \frac{\Delta P_t f}{P_t} \leq \sqrt{nK} \sqrt{\frac{|\nabla P_t f|^2}{P_t f} + \frac{n}{2t} + \frac{nK}{4}} + \frac{n}{2t}. \quad (3)$$

In the meanwhile, R. Hamilton [Ham93] had proved the inequality

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} \leq e^{2Kt} \frac{\Delta P_t f}{P_t} + \frac{n}{2t} e^{4Kt} \quad (4)$$

and, most recently, J. Li and X. Xu [LX11] have obtained the bound

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} \leq \left(1 + \frac{\sinh(2Kt) - 2Kt}{2\sinh^2(Kt)}\right) \frac{\Delta P_t f}{P_t f} + \frac{nK}{2} (1 + \coth(Kt)). \quad (5)$$

All these inequalities are based on the maximum principle and are not comparable to each other.

There is a huge literature on extensions of the Li-Yau inequality (1), actually almost 400 citations on MathSciNet or Zentralblatt. Let us mention in particular the most recent [SZ06, Wan10, BG11, Qia12, Lee13, QZZ13, GM14, Qia14].

In this article we shall prove the following general Li-Yau inequality, *in negative and positive curvature, which improves all bounds known to us* (see Corollary 2.4). Assume that the Ricci curvature of the n -dimensional Riemannian manifold is (uniformly) bounded from below by $\rho \in \mathbb{R}^*$. Then

$$\frac{4}{n\rho} \frac{\Delta P_t f}{P_t f} < 1 + \frac{\pi^2}{\rho^2 t^2}$$

and

$$\frac{|\nabla P_t f|^2}{(P_t f)^2} < \frac{n}{2} \Phi_t \left(\frac{4}{n\rho} \frac{\Delta P_t f}{P_t f} \right)$$

for any positive function f and $t > 0$, where

$$\Phi_t(x) = \begin{cases} \frac{\rho}{2}(x - 2 + 2\sqrt{1-x} \coth(\rho t \sqrt{1-x})), & x \leq 1 \\ \frac{\rho}{2}(x - 2 + 2\sqrt{x-1} \cot(\rho t \sqrt{x-1})), & 1 \leq x < 1 + \frac{\pi^2}{\rho^2 t^2}. \end{cases}$$

This result will be obtained as an extension to any curvature lower bound of the equivalence between the following properties, which is due to [BL06] :

- i. The Ricci curvature of M is non-negative.
- ii. For any smooth positive function f and $t > 0$,

$$\exp\left(-\frac{2}{nP_t f}(\text{Ent}_{P_t}(f) + t\Delta P_t f)\right) \leq 1 + \frac{2t}{nP_t f}\left(\Delta P_t f - \frac{|\nabla P_t f|^2}{P_t f}\right) \quad (6)$$

where $\text{Ent}_{P_t}(f) = P_t(f \log f) - P_t f \log P_t f$.

- iii. For any smooth positive function f and $t > 0$,

$$\exp\left(\frac{2}{nP_t f}(\text{Ent}_{P_t}(f) - t\Delta P_t f)\right) \leq 1 + \frac{2t}{nP_t f}\left(P_t\left(\frac{|\nabla f|^2}{f}\right) - \Delta P_t f\right).$$

- iv. For any smooth positive function f and $t > 0$,

$$P_t f \Delta(\log P_t f) \geq P_t(f \Delta \log f) \left(1 + \frac{2t}{n} \Delta(\log P_t f)\right). \quad (7)$$

If we replace the left-hand side of (6) by 0, using the fact that an exponential is always positive, inequality (6) implies the Li-Yau inequality (1). Inequality (7) can be reformulated as a bound on the gradient of the heat kernel (commutation inequality), taking the dimension into account.

The present paper gives a generalisation of this equivalence with any lower bound of the Ricci curvature, positive or negative, instead of a non-negative Ricci curvature. As in [BL06], the result will be stated for general Markov semigroups, including diffusion semigroups on weighted Riemannian manifolds.

The paper is organised as follows. In the next section we state this generalisation, for a Markov diffusion semigroup under a $CD(\rho, n)$ curvature-dimension condition. We also derive first consequences, including our main result : a new Li-Yau inequality under this curvature-condition. The proof, which is very short and simple, is given in Section 3. Section 4 is devoted to applications : ultracontractive bounds in the positive curvature case, and new Harnack inequalities in the positive and negative cases, which are equivalent to our Li-Yau equality. Finally, in Section 5 we prove that our inequality implies all classical Li-Yau inequalities known to us, stressing in particular on the bounds (2) to (5).

2 Main result

2.1 Markov triple and curvature-dimension condition

A Markov diffusion triple (E, μ, Γ) , as defined in [BGL14, Chapter 3] consists in a nice state space E equipped with a Markov diffusion semigroup $(P_t)_{t \geq 0}$ with infinitesimal generator L ,

carré du champ Γ and invariant and reversible σ -finite measure μ . The carré du champ and Γ_2 operators are pointwise defined from the generator L by

$$\Gamma(f, g) = \frac{1}{2}(L(fg) - fLg - gLf) \quad \text{and} \quad \Gamma_2(f) = \frac{1}{2}(L\Gamma(f, f) - 2\Gamma(f, Lf))$$

for functions f and g in a suitable algebra \mathcal{A} of functions from E to \mathbb{R} . We let $\Gamma(f) = \Gamma(f, f)$. The generator L is assumed to satisfy the diffusion property, that is, for any smooth function φ and $f \in \mathcal{A}$,

$$L\varphi(f) = \varphi'(f)Lf + \varphi''(f)\Gamma(f).$$

In the Markov triple setting, the abstract curvature-dimension condition $CD(\rho, n)$, for $\rho \in \mathbb{R}$ and $n \geq 1$, is satisfied when

$$\Gamma_2(f) \geq \rho\Gamma(f) + \frac{1}{n}(Lf)^2$$

for any $f \in \mathcal{A}$.

Let us recall that under a $CD(\rho, n)$ condition with $\rho > 0$, then the semigroup is ergodic, that is, $P_t f \rightarrow \int f d\mu$ in $L^2(\mu)$.

The main example of a Markov diffusion triple is a smooth, connected and complete weighted Riemannian manifold (M, g) equipped with the generator $L = \Delta + \nabla V \cdot \nabla$, where Δ is the Laplace-Beltrami operator and V a smooth function on M , and the measure $d\mu = e^V dx$ where dx is the Riemannian measure. In this case, the carré du champ operator is $\Gamma(f) = |\nabla f|^2$ where $|\nabla f|$ is the length of the vector ∇f , and the algebra of functions \mathcal{A} consists in smooth and bounded functions on M .

For instance, when $V = 0$, the Bochner-Lichnerowicz inequality implies that the condition $CD(\rho, n)$ is satisfied when M is a n -dimensional Riemannian manifold with a Ricci curvature Ric (uniformly) bounded from below by ρ . For a general V , the condition holds on the m -dimensional manifold M as soon as $m < n$ and

$$Ric - \nabla^2 V \geq \rho + \frac{\nabla V \otimes \nabla V}{n - m}.$$

For a positive function f on E we let $\text{Ent}_{P_t}(f) = P_t(f \log f) - P_t f \log P_t f$ and $\text{Ent}_\mu(f) = \int f \log f d\mu - \int f d\mu \log \int f d\mu$.

Remark 2.1 *In this work we shall deal with a symmetric Markov semigroup for convenience, but all the results proved here can be stated in a non-symmetric case.*

2.2 Li-Yau inequality under the $CD(\rho, n)$ condition

Theorem 2.2 (Local logarithmic Sobolev inequalities) *Let (E, μ, Γ) be a Markov diffusion triple, $\rho \in \mathbb{R}^*$ and $n \geq 1$. Given a positive function f on E and $t > 0$, we let*

$$X = \frac{4}{n\rho} \frac{LP_t f}{P_t f}$$

and given $t > 0$ we define the functions Φ_t and $\tilde{\Phi}_t$ by

$$\Phi_t(x) = \begin{cases} \frac{\rho}{2}(x - 2 + 2\sqrt{1-x} \coth(\rho t \sqrt{1-x})), & x \leq 1 \\ \frac{\rho}{2}(x - 2 + 2\sqrt{x-1} \cot(\rho t \sqrt{x-1})), & 1 \leq x < 1 + \frac{\pi^2}{\rho^2 t^2} \end{cases} \quad (8)$$

and $\tilde{\Phi}_t(x) = \Phi_t(x) - \rho x + 2\rho$. Then the following properties are equivalent :

i. The Markov triple satisfies the $CD(\rho, n)$ condition.

ii. For any positive function $f \in \mathcal{A}$ and $t > 0$, then $X < 1 + \frac{\pi^2}{\rho^2 t^2}$ and

$$\exp\left(-\frac{2}{n} \frac{\text{Ent}_{P_t}(f)}{P_t f} + \frac{t\rho}{2} X - \rho t\right) \leq \begin{cases} \frac{\sinh(\rho t \sqrt{1-X})}{\rho \sqrt{1-X}} \left(\Phi_t(X) - \frac{2}{n} \frac{\Gamma(P_t f)}{(P_t f)^2}\right) & \text{if } X \leq 1 \\ \frac{\sin(\rho t \sqrt{X-1})}{\rho \sqrt{X-1}} \left(\Phi_t(X) - \frac{2}{n} \frac{\Gamma(P_t f)}{(P_t f)^2}\right) & \text{if } 1 \leq X < 1 + \frac{\pi^2}{\rho^2 t^2}. \end{cases} \quad (9)$$

iii. For any positive function $f \in \mathcal{A}$ and $t > 0$, then $X < 1 + \frac{\pi^2}{\rho^2 t^2}$ and

$$\exp\left(\frac{2}{n} \frac{\text{Ent}_{P_t}(f)}{P_t f} - \frac{t\rho}{2} X + \rho t\right) \leq \begin{cases} \frac{\sinh(\rho t \sqrt{1-X})}{\rho \sqrt{1-X}} \left(\tilde{\Phi}_t(X) + \frac{2}{n} \frac{P_t(\Gamma(f)/f)}{P_t f}\right) & \text{if } X \leq 1 \\ \frac{\sin(\rho t \sqrt{X-1})}{\rho \sqrt{X-1}} \left(\tilde{\Phi}_t(X) + \frac{2}{n} \frac{P_t(\Gamma(f)/f)}{P_t f}\right) & \text{if } 1 \leq X < 1 + \frac{\pi^2}{\rho^2 t^2}. \end{cases} \quad (10)$$

As is [BGL14], inequality (10) may be called a local logarithmic Sobolev inequality and (9) a local reverse logarithmic Sobolev inequality. We observe that the right-hand sides of (8), (9) and (10) are continuous in $X = 1$, justifying the way of writing.

Corollary 2.3 Under the $CD(\rho, n)$ condition, for any positive function $f \in \mathcal{A}$ and $t > 0$

$$\frac{4}{n\rho} \frac{LP_t f}{P_t f} < 1 + \frac{\pi^2}{\rho^2 t^2}. \quad (11)$$

The term in the right-hand side of (9) has to be positive, giving a general Li-Yau inequality:

Corollary 2.4 (General Li-Yau inequality) For any Markov diffusion triple satisfying a $CD(\rho, n)$ condition with $\rho \in \mathbb{R}^*$ and $n \geq 1$, then

$$\frac{\Gamma(P_t f)}{(P_t f)^2} < \frac{n}{2} \Phi_t \left(\frac{4}{n\rho} \frac{LP_t f}{P_t f} \right) \quad (12)$$

for any positive function $f \in \mathcal{A}$ and $t > 0$, where the function Φ_t is defined in (8).

Remark 2.5 i. When ρ tends to 0 in Theorem 2.2, we exactly recover the estimates ii. and iii. given in the introduction in the case $\rho = 0$. In particular the general Li-Yau inequality (12) converges to the classical Li-Yau inequality (1).

ii. Combining the two inequalities (9) and (10) leads to a commutation type inequality similar to (7), and which converges to (7) when ρ tends to 0. The inequality obtained in this general case where $\rho \neq 0$ is still equivalent to the $CD(\rho, n)$ condition, but less appealing than (7), which is why we omit it.

Remark 2.6 We shall see in Section 5 that for any t the bound (12) improves upon all classical bounds recalled in the introduction.

Remark 2.7 Our Li-Yau inequality (12) also improves on the following results obtained in [BQ99, Thms. 1 and 2] :

- Assume that $\rho > 0$. Then (12) holds for $X \leq X_0 := 1 + \pi^2/64$, where as above $X = 4LP_t f / (n\rho P_t f)$; if $X > X_0$, then the bound holds with $\Phi_t(X)$ replaced by the tangent $\Phi_t(X_0) + \Phi'_t(X_0)(X - X_0)$.

- Assume that $\rho < 0$. Then (12) holds for $X \leq 1$; moreover, for any X , then the bound holds with $\Phi_t(X)$ replaced by the tangent $\Phi_t(X_1) + \Phi'_t(X_1)(X - X_1)$ for any $X_1 \leq 1$.

But, by Lemma 2.8 below and for any given $t > 0$, the function Φ_t is a C^∞ and strictly concave function on the whole interval $(-\infty, 1 + \frac{\pi^2}{\rho^2 t^2})$; hence its graph is below its tangents, and (12) improves upon these bounds in [BQ99].

Lemma 2.8 (Properties of Φ_t) For any $t > 0$ and $\rho \in \mathbb{R}^*$, the function Φ_t is C^∞ and strictly concave on the interval $(-\infty, 1 + \frac{\pi^2}{\rho^2 t^2})$.

Proof

\triangleleft We have already observed that Φ_t is continuous in $X = 1$. Moreover Φ_t is C^∞ since in $X = 1$, the Taylor expansions (of any orders) are the same for $X = 1^-$ or $X = 1^+$.

Let us now prove that Φ_t is strictly concave, for instance in the case $\rho > 0$, the case $\rho < 0$ being similar. For fixed $u \in (0, 1)$, by direct computation, the map $y \mapsto \frac{\sin(yu)}{\sin(y)}$ is increasing and positive on $(0, \pi)$. This implies that the map

$$I : y \mapsto \int_0^1 \left(\frac{\sin(yu)}{\sin(y)} \right)^2 du = \frac{2y - \sin(2y)}{4y \sin^2(y)}$$

is increasing on $(0, \pi)$. Hence the function Φ_t has a decreasing derivative

$$\Phi'_t(x) = \rho/2 - \rho^2 t I(\rho t \sqrt{x-1})$$

on $(1, 1 + \frac{\pi^2}{\rho^2 t^2})$, so is strictly concave on this interval. The same argument can be performed on $X < 1$, starting from the function $y \mapsto \frac{\sinh(yu)}{\sinh(y)}$. This concludes the proof by differentiability of Φ_t in $X = 1$. \triangleright

On the figure 1 (resp. on the figure 2), we give the graphs of Φ_t , for $\rho = 1$ (resp. $\rho = -1$) and $t = 3/2, 2$ and $5/2$ (resp. $t = 1/4, 1/2$ and 1) from top to bottom. In both figures, the dashed lines correspond to the sine function, relevant for $X > 1$. Moreover, in the negative curvature case, the dotted (lowest) line corresponds to the limit case, obtained when t goes to infinity (with ρ fixed).

3 Proof of Theorem 2.2

We first assume that the Markov diffusion triple satisfies a $CD(\rho, n)$ condition and prove properties *i* and *ii*.

Let then $f \in \mathcal{A}$ be a fixed positive function, which we assume to be larger than an $\epsilon > 0$, without loss of generality. Let also $t > 0$ be fixed, and define $\Lambda(s) = P_s(P_{t-s} f \log P_{t-s} f)$ for




Figure 1: Graphs of Φ_t for positive curvatures




Figure 2: Graphs of Φ_t for negative curvatures

$s \in [0, t]$, so that $\text{Ent}_{P_t}(f) = \Lambda(t) - \Lambda(0)$. Classical properties of the derivative of Λ , see for instance [BGL14, Section 6.7.2], ensure that

$$\Lambda'(s) = P_s\left(\frac{\Gamma(g)}{g}\right) = P_s(g\Gamma(\log g)) \quad \text{and} \quad \Lambda''(s) = 2P_s(g\Gamma_2(\log g))$$

for all $s \in [0, t]$, where $g = P_{t-s}f$. Applying the $CD(\rho, n)$ condition, we obtain

$$\Lambda''(s) \geq 2\rho\Lambda'(s) + \frac{2}{n}P_s(gL(\log g)).$$

Since $L(\log g) = \frac{Lg}{g} - \frac{\Gamma(g)}{g^2}$ by diffusion property, the Cauchy-Schwarz inequality further gives

$$\Lambda''(s) \geq 2\rho\Lambda'(s) + \frac{2}{nP_t f}(LP_t f - \Lambda'(s))^2.$$

This computation has been performed for example in [BL06] and [BGL14, Section 6.7.2].

Let now $a = \frac{2}{nP_t f}$, $X = \frac{4}{n\rho} \frac{LP_t f}{P_t f}$, $B = \left(\frac{n\rho}{2} P_t f\right)^2 (X - 1)$ and $C = LP_t f - \frac{n\rho}{2} P_t f$. The above inequality can be written as

$$\Lambda''(s) \geq a\left((\Lambda'(s) - C)^2 + B\right),$$

or equivalently

$$H''(s) \leq -a^2 B H(s)$$

for all $s \in [0, t]$, where $H(s) = \exp(-a(\Lambda(s) - Cs))$.

Then Lemma 3.1 below has several consequences. First of all, since H is a positive function, then both cases *i* and *ii* in the lemma are impossible. This means that necessarily $a^2 B < \frac{\pi^2}{t^2}$, or in other words that for any $t > 0$

$$X < 1 + \frac{\pi^2}{\rho^2 t^2},$$

or equivalently :

$$\forall t > 0, \quad \frac{1}{4n\rho} \frac{LP_t f}{P_t f} < 1 + \frac{\pi^2}{\rho^2 t^2}.$$

Since we now know that $a^2 B < \frac{\pi^2}{t^2}$, Lemma 3.1 again ensures that for all $s \in [0, t]$

$$H(s) \geq \tau_{a^2 B}(t-s)H(0) + \tau_{a^2 B}(s)H(t). \quad (13)$$

The reverse local logarithmic Sobolev inequality (9) is now obtained by taking the derivative in (13) at $s = 0$, in the form

$$H'(0) \geq -\tau'_{a^2 B}(t)H(0) + \tau'_{a^2 B}(0)H(t).$$

In the first case where $a^2 B \in (0, \frac{\pi^2}{t^2})$, or equivalently $X \in (1, 1 + \frac{\pi^2}{t^2 \rho^2})$, then this inequality is exactly (9) with the sine function when written in terms of Λ , and then of $P_t f$. In the other case where $a^2 B < 0$, that is $X < 1$, then we obtain (9) with the hyperbolic sine function. The limit case $a^2 B = 0$, or equivalently $X = 0$, is the limit of any of the first two cases. Together, the obtained estimates are exactly inequality (9).

The local logarithmic Sobolev inequality (10) is obtained in the same way, by taking the derivative of (13) at $s = t$.

We now prove the converse implication, namely that *ii* implies *i*, the case *iii* being handled by the same method. Observing that inequality (9) is an equality when $f = 1$, the idea is to let $f = 1 + \varepsilon h$ with $h \in \mathcal{A}$ and to perform a second order Taylor expansion of (9) in the parameter ε tending to 0. The zeroth and first order terms in ε vanish, and recalling that $\text{Ent}_{P_t}(f) = \frac{\varepsilon^2}{2} [P_t(h^2) - (P_t h)^2] + o(\varepsilon^2)$, we obtain

$$P_t(h^2) - (P_t h)^2 \geq \frac{e^{2\rho t} - 1}{\rho} \Gamma(P_t h) + \frac{e^{2\rho t} - 1 - 2\rho t}{\rho^2} \frac{(LP_t h)^2}{n}$$

for all $t \geq 0$. Now, as in [BL06] for instance, a second order Taylor expansion in t tending to 0, with the zero and first order terms vanishing, gives the $CD(\rho, n)$ condition back.

In this proof we have used the following elementary but useful lemma, which is proved in [Vil09, Thm 14.28] for instance.

Lemma 3.1 *Let $t > 0$, $\lambda \in \mathbb{R}$ and h a C^2 non-negative function on $[0, t]$ such that $h'' \leq -\lambda h$ on $[0, t]$. Then :*

i. If $\lambda > \frac{\pi^2}{t^2}$ then for all $s \in [0, t]$, $h(s) = 0$.

ii. If $\lambda = \frac{\pi^2}{t^2}$ then for all $s \in [0, t]$, $h(s) = c \sin\left(\frac{s}{t}\right)$, for some $c \geq 0$.

iii. If $\lambda < \frac{\pi^2}{t^2}$ then for all $s \in [0, t]$, $h(s) \geq \tau_\lambda(t-s)h(0) + \tau_\lambda(s)h(t)$, where

$$\tau_\lambda(s) = \begin{cases} \frac{\sin(\sqrt{\lambda}s)}{\sin(\sqrt{\lambda}t)} & \text{if } \lambda > 0 \\ s/t & \text{if } \lambda = 0 \\ \frac{\sinh(\sqrt{-\lambda}s)}{\sinh(\sqrt{-\lambda}t)} & \text{if } \lambda < 0 \end{cases}$$

4 Applications to ultracontractive estimates and Harnack inequalities

4.1 Ultracontractive estimates in positive curvature

In this section we assume that $\rho > 0$. We will use the new Li-Yau inequality (12) in force to obtain uniform bounds on $L \log P_t f$, for positive f , and then ultracontractive estimate on the semigroup.

Under a curvature-dimension condition $CD(\rho, n)$ with $\rho > 0$ and $n \geq 1$, it is classical that a reversible Markov semigroup $(P_t)_{t \geq 0}$ is ultracontractive, that is, $P_t f$ is bounded for all $t > 0$ and integrable f .

A way of proving an ultracontractive bound is indeed the following: First, a $CD(\rho, n)$ condition with $\rho > 0$ implies a (Nash-type) logarithmic entropy-energy inequality :

$$\text{Ent}_\mu(f^2) \leq \frac{n}{2} \log \left(1 + \frac{4}{\rho n} \int \Gamma(f) d\mu \right) \quad (14)$$

for any $f \in \mathcal{A}$ such that $\int f^2 d\mu = 1$. Let us observe that, in this ergodic case where $P_t f$ converges to $\int f d\mu$ for large time, then (14) can be recovered by letting t go to infinity in our local inequality (10) (written for f^2 instead of f). Then (14) implies the following ultracontractive estimate: there exists a constant C such that

$$\|P_t f\|_\infty \leq \frac{C}{t^{n/2}} \int |f| d\mu, \quad t \in (0, 1] \quad (15)$$

for any $f \in \mathcal{A}$. These results can be found for instance in [BGL14, Chapter 6].

This bound is in fact included in the Li-Yau inequality (12) which also gives a quantitative estimate for large time. Observe indeed that $\Gamma(P_t f)/(P_t f)^2 \geq 0$ for any positive f and t , so (12) gives

$$\Phi_t \left(\frac{4}{n\rho} \frac{LP_t f}{P_t f} \right) > 0.$$

Let now $\rho, t > 0$ be fixed. By definition (8), it holds $\Phi_t(0) > 0$,

$$\lim_{x \rightarrow -\infty} \Phi_t(x) = -\infty \quad \text{and} \quad \lim_{x \rightarrow 1 + \frac{\pi^2}{\rho^2 t^2}} \Phi_t(x) = -\infty.$$

Hence the continuous and strictly concave function Φ_t admits exactly two roots $\xi_1^t < 0 < \xi_2^t$, and is positive in-between and negative outside its roots. Of course ξ_1^t and ξ_2^t depend on t and ρ but not on the dimension n . In particular

$$\xi_1^t < \frac{4}{n\rho} \frac{LP_t f}{P_t f} < \xi_2^t \quad (16)$$

for all positive f and t .

A first simple consequence is the following : if $t \geq 2/\rho$, then $\Phi_t(1) = 1/t - \rho/2$ is non-positive, so necessarily $\xi_2^t \leq 1$ (see also the graph of the second function in Figure 1). In other words :

Corollary 4.1 ([BQ99]) *Assume a $CD(\rho, n)$ condition with $\rho > 0$. Then for all positive function f in \mathcal{A}*

$$\frac{4}{n\rho} \frac{LP_t f}{P_t f} < 1, \quad t \geq \frac{2}{\rho}. \quad (17)$$

Let us remark that (17) provides additional information to the bound (11) in Corollary 2.3. Moreover, inequalities (11) and (17) are of course not optimal for large t since $LP_t f$ converges to 0 when t goes to $+\infty$. Next proposition gives an answer to this issue : it makes quantitative the fact that $\xi_1^t, \xi_2^t \rightarrow 0$ when t goes to $+\infty$ (and $\xi_1^t \rightarrow -\infty, \xi_2^t \rightarrow +\infty$ when t goes to 0), giving by (16) corresponding explicit upper and lower bounds for $LP_t f/P_t f$.

Lemma 4.2 *Assume that $\rho > 0$. Then the roots ξ_1^t and ξ_2^t of Φ_t are such that*

$$\begin{aligned} -4e^{-\rho t} - 4e^{-2\rho t} &\leq \xi_1^t \leq -4e^{-\rho t} + 8\rho t e^{-2\rho t}, & t \geq \frac{1}{2\rho} \\ 4e^{-\rho t} - 4e^{-2\rho t} &\leq \xi_2^t \leq 4e^{-\rho t} + 8\rho t e^{-2\rho t}, & t \geq \frac{6}{\rho} \\ \xi_1^t &= -\frac{2}{\rho t} + O(1) \quad \text{and} \quad \xi_2^t = \frac{\pi^2}{\rho^2 t^2} - \frac{4}{\rho t} + O(1), & t \rightarrow 0. \end{aligned}$$

Observe the compatibility of this last bound with Corollary 11.

Proof

\triangleleft We first consider the large time bounds on the negative root ξ_1^t . For $x < 0$, we observe that $\xi_1^t \geq x$ if and only if $\Phi_t(x) \leq 0$, or if and only if $u = \sqrt{1-x}$, which then is larger than 1, satisfies

$$(u-1)e^{\rho t u} \geq u+1.$$

For $u = 1 + 2e^{-\rho t}$, and by the elementary inequality $e^v \geq 1+v$, this property holds as soon as $t \geq 1/(2\rho)$. Therefore $\xi_1^t \geq 1 - u^2$ for these t . For $u = 1 + 2e^{-\rho t} - 4\rho t e^{-2\rho t}$, the reverse inequality holds also as soon as $t \geq 1/(2\rho)$. Therefore $\xi_1^t \leq 1 - u^2$ for these t . Together, this gives the large time lower and upper bounds on ξ_1^t .

Then we recall that the positive root ξ_2^t belongs to $(0, 1]$ as soon as $t \geq 2/\rho$, which we assume. Then, now, $\xi_2^t \leq x$ if and only if $\Phi_t(x) \leq 0$, or if and only if $(1-u)e^{\rho t u} \geq 1+u$. For $u = 1 - 2e^{-\rho t}$ the reverse inequality holds as soon as $t \geq 2/\rho$, so that $\xi_2^t \geq 1 - u^2$ for these t . For $u = 1 - 2e^{-\rho t} - 4\rho t e^{-2\rho t}$ the inequality holds as soon as $t \geq 6/\rho$, so that $\xi_2^t \leq 1 - u^2$ for these t . This gives the large time lower and upper bounds on ξ_2^t .

We proceed in the same manner to get the short time estimates. \triangleright

Let us now see how to turn these estimates on ξ_1^t, ξ_2^t into ultracontractivity estimates on the semigroup. Given $0 < t < s$, integrating the pointwise bound (16) over the interval $[t, s]$ gives

$$\exp\left(-\frac{n\rho}{4} \int_t^s \xi_2^u du\right) \leq \frac{P_t f}{P_s f} \leq \exp\left(-\frac{n\rho}{4} \int_t^s \xi_1^u du\right).$$

By ergodicity, letting s go to infinity implies :

$$\exp\left(-\frac{n\rho}{4} \int_t^\infty \xi_2^u du\right) \leq \frac{P_t f}{\int f d\mu} \leq \exp\left(-\frac{n\rho}{4} \int_t^\infty \xi_1^u du\right), \quad t > 0.$$

By Proposition 4.2, it follows that

$$\frac{P_t f}{\int f d\mu} \leq \exp\left[n\left(e^{-\rho t} + \frac{e^{-2\rho t}}{2}\right)\right], \quad t \geq \frac{1}{2\rho}$$

and

$$\frac{P_t f}{\int f d\mu} \geq \exp\left[-n\left(e^{-\rho t} + \frac{1+2\rho t}{2}e^{-2\rho t}\right)\right], \quad t \geq \frac{6}{\rho}.$$

Proceeding likewise for the short time estimates finally gives :

Corollary 4.3 *Assume a $CD(\rho, n)$ condition with $\rho > 0$ with $n \geq 1$. Then there exist constants C and D , depending on ρ and n , such that for any positive f*

$$\left|P_t f - \int f d\mu\right| \leq C e^{-\rho t} \int f d\mu, \quad t \geq 1, \quad (18)$$

and

$$\frac{1}{Dt^n} \exp\left(-\frac{n\pi^2}{\rho t}\right) \leq \frac{P_t f}{\int f d\mu} \leq \frac{D}{t^{n/2}}, \quad t \in (0, 1]. \quad (19)$$

By Proposition 4.2, the constant C in (18) can be made explicit in n and ρ . Inequality (18) does not give the asymptotic behavior of $P_t f$ in a satisfactory way. Indeed it is classical that the $CD(\rho, n)$ condition on a symmetric Markov semigroup with $\rho > 0$ and $n > 1$ implies a (Poincaré) spectral gap inequality with constant $(n-1)/(\rho n)$; it follows that $P_t f$ converges to its mean with an exponential speed with rate $\frac{2\rho n}{n-1}$:

$$\int \left(P_t f - \int f d\mu\right)^2 d\mu \leq e^{-\frac{2\rho n}{n-1}t} \int \left(f - \int f d\mu\right)^2 d\mu.$$

The right rate of convergence has been lost in (18). Let us in fact observe that the rate ρ in (18) cannot be improved into $\rho n/(n-1)$ by our method since neither ξ_1^t nor ξ_2^t depend on n .

On the other hand, since $|P_t f| \leq P_t |f|$, the upper bound in (19) can be extended to any f , recovering the classical ultracontractivity property (15), together with an explicit lower bound on $P_t f$ which is not included in (15). Moreover the method here is elementary and can also be written in the non-symmetric case.

One can also derive similar bounds on the gradients. For instance :

Corollary 4.4 *Assume a $CD(\rho, n)$ condition with $\rho > 0$ with $n \geq 1$. Then there exists a constant C , independent of ρ and n , such that for any positive f*

$$\Gamma(\log P_t f) = \frac{\Gamma(P_t f)}{(P_t f)^2} \leq C n \rho e^{-2\rho t}, \quad t \geq \frac{6}{\rho}.$$

Proof

◁ We use the above notation, together with $s = \rho t$, and assume that $s \geq 6$. Then, by Lemma 4.2, $|X| \leq 6e^{-2s} \leq 1$, so that in particular $\sqrt{1-X} \geq 1 - 6e^{-2s}$. Therefore, by this bound and the elementary $e^s \geq s^3/6$,

$$e^{-2s\sqrt{1-X}} \leq e^{12se^{-s}-2s} \leq e^{2-2s}.$$

In particular it is smaller than $1/3$, so

$$\coth(s\sqrt{1-X}) = \frac{1 + e^{-2s\sqrt{1-X}}}{1 - e^{-2s\sqrt{1-X}}} \leq 1 + 3e^{-2s\sqrt{1-X}}.$$

It follows that

$$\frac{2}{\rho} \Phi_t(X) \leq X - 2 + 2\sqrt{1-X}(1 + 3e^{-2s\sqrt{1-X}}) \leq X - 2 + 2\left(1 - \frac{X}{2}\right)(1 + 3e^{2-2s}) \leq Ce^{-2s}.$$

This concludes the argument by Corollary 2.3. ▷

4.2 Estimates in non-positive curvature

Let $\rho < 0$ be fixed. For any $t > 0$, by definition (8), it holds $\Phi_t(1) = 1/t - \rho/2 > 0$,

$$\lim_{x \rightarrow -\infty} \Phi_t(x) = +\infty \quad \text{and} \quad \lim_{x \rightarrow 1 + \frac{\pi^2}{\rho^2 t^2}} \Phi_t(x) = -\infty.$$

Hence the continuous and strictly concave function Φ_t admits exactly one root $1 < \xi^t < 1 + \frac{\pi^2}{\rho^2 t^2}$ (see for instance Figure 2). It is positive on $(-\infty, \xi^t)$ and in particular

$$\frac{4}{n\rho} \frac{LP_t f}{P_t f} \leq \xi^t$$

for all positive f and t , by the Li-Yau inequality (12). This recovers the bound (11). In fact (11) can not be improved by our method since $\xi^t \sim 1 + \frac{\pi^2}{\rho^2 t^2}$ for large time. Indeed, by direct computation as in the previous section,

$$1 + \frac{\pi^2}{\rho^2 t^2} \left(1 - \frac{2}{\rho t}\right) \leq \xi^t \leq 1 + \frac{\pi^2}{\rho^2 t^2}, \quad t \geq \frac{2}{|\rho|}.$$

4.3 Harnack inequalities

In this section we assume that the space E is a complete, connected and smooth Riemannian manifold (M, g) . This example has been described in Section 2.1. We shall let d denote the Riemannian distance on M .

The Li-Yau inequality (12) for $\rho \neq 0$ can be written as

$$-\frac{|\nabla P_t f|}{P_t f} \geq -\sqrt{\frac{n}{2} \Phi_t \left(\frac{4}{n\rho} \frac{LP_t f}{P_t f} \right)}$$

where Φ_t is defined in (8), or equivalently,

$$-\frac{|\nabla P_t f|}{P_t f} \geq \Psi_{t,\rho} \left(\frac{LP_t f}{P_t f} \right)$$

where

$$\Psi_{t,\rho}(x) = -\sqrt{\frac{n}{2}\Phi_t\left(\frac{4}{n\rho}x\right)}. \quad (20)$$

This extends to $\rho = 0$ by letting $\Psi_{t,0}(x) = -\sqrt{n/(2t)} + x$.

Basic properties of the function $\Psi_{t,\rho}$ are listed in the following remark. Their proofs are not complicated, and therefore are omitted.

Remark 4.5 *Let $t > 0$ be fixed.*

- *Let $\rho > 0$. The function $\Psi_{t,\rho}$ is defined on the interval $\mathcal{I}_{t,\rho} = [n\rho\xi_1^t/4, n\rho\xi_2^t/4]$, where the roots $\xi_1^t < 0 < \xi_2^t$ of Φ_t have been defined in Section 4.1 and depend only on ρ and t . Its derivative $\Psi'_{t,\rho}$ is an increasing one-to-one function from $(n\rho\xi_1^t/4, n\rho\xi_2^t/4)$ onto \mathbb{R} . The Legendre-Fenchel transform of $\Psi_{t,\rho} : \Psi_{t,\rho}^*(x) = \sup_{y \in \mathcal{I}_{t,\rho}} \{xy - \Psi_{t,\rho}(y)\}$ is defined and finite for every $x \in \mathbb{R}$.*
- *Let $\rho < 0$. The function Φ_t admits only one root $\xi^t > 0$ (see Section 4.2) and then $\Psi_{t,\rho}$ is defined on the interval $\mathcal{I}_{t,\rho} = [n\rho\xi^t/4, +\infty)$. Its derivative $\Psi'_{t,\rho}$ is an increasing one-to-one function from $(n\rho\xi^t/4, \infty)$ onto $(-\infty, 0)$. The Legendre transform of $\Psi_{t,\rho}$ is defined and finite on $(-\infty, 0)$.*
- *When $\rho = 0$, the function $\Psi_{t,0}(x) = -\sqrt{n/(2t)} + x$ is defined on $\mathcal{I}_{t,0} = [-n/(2t), +\infty)$. Its Legendre transform is also defined on $(-\infty, 0)$. The case $\rho = 0$ appears as limit case of the case $\rho < 0$, but not of the case $\rho > 0$.*
- *For any $\rho \in \mathbb{R}$ and $t > 0$, $\Psi_{t,\rho}$ is strictly convex on its interval of definition.*
- *For any $\rho \in \mathbb{R}$ and $t > 0$, $\Psi_{t,\rho}^*$ is non-negative.*

In the figure 3 (resp. figure 4) we have drawn the graph of $x \mapsto \Psi_{t,\rho}(x)$ with $\rho = 1$ (resp. $\rho = -1$), $t = 1$ and $n = 2$. The dashed line in the figure 4 corresponds to the graph of $\Psi_{t,0}$, again with $t = 1$ and $n = 2$.




Figure 3: Graph of $\Psi_{t,\rho}$ with a positive curvature




Figure 4: Graph of $\Psi_{t,\rho}$ with a negative curvature

It is standard since [LY86] that a Li-Yau inequality implies a parabolic Harnack inequality in the semigroup. Here we give the *equivalence* between these two types of bounds in our framework.

Theorem 4.6 (Harnack inequality) *Assume that L satisfies a $CD(\rho, n)$ condition on the manifold M , with $\rho \in \mathbb{R}$ and $n \geq 1$.*

i. Let us assume that $\rho > 0$.

For any $s, t > 0$, $x, y \in M$ and any positive function f in \mathcal{A} , we have

$$P_s f(x) \leq P_t f(y) \exp \left(\frac{d(x, y)}{t-s} \int_s^t \Psi_{u,\rho}^* \left(-\frac{t-s}{d(x, y)} \right) du \right). \quad (21)$$

Conversely, if (21) is satisfied for any positive function f in \mathcal{A} , $x, y \in M$ and $s, t > 0$ then the Li-Yau inequality (12) holds.

ii. Let us assume that $\rho \leq 0$.

For any $0 < s < t$, $x, y \in M$ and any positive function f in \mathcal{A} , then the inequality (21) holds.

Conversely, if (21) is satisfied for any positive function f in \mathcal{A} , $x, y \in M$ and $0 < s < t$, then the Li-Yau inequality (12) holds.

In both cases (21) also holds for any positive function f , in $L^1(\mu)$ for instance.

Proof

\triangleleft Let us prove the first part of the theorem, when $\rho > 0$. Let $s, t > 0$, $x, y \in M$ and f positive. Let for $u \in [0, 1]$,

$$\Lambda(u) = \log P_{b(u)} f(x_u),$$

where $b(u) = (1-u)s + ut$ (b is not necessarily increasing) and $(x_u)_{u \in [0,1]}$ is a constant speed geodesic between x and y . Then we get

$$\Lambda'(u) = (t-s) \frac{LP_{b(u)} f(x_u)}{P_{b(u)} f(x_u)} + \frac{\nabla P_{b(u)} f(x_u)}{P_{b(u)} f(x_u)} \cdot \dot{x}_u \geq (t-s) \frac{LP_{b(u)} f(x_u)}{P_{b(u)} f(x_u)} - d \frac{|\nabla P_{b(u)} f(x_u)|}{P_{b(u)} f(x_u)}$$

where $d = d(x, y)$. The Li-Yau inequality (12) ensures that

$$-\frac{|\nabla P_{b(u)} f|}{P_{b(u)} f} \geq \Psi_{b(u),\rho} \left(\frac{LP_{b(u)} f}{P_{b(u)} f} \right)$$

at the point x_u , where as in (20) $\Psi_{b(u),\rho}(x) = -\sqrt{\frac{n}{2}\Phi_{b(u)}(\frac{4x}{n\rho})}$. But now $\Psi_{b(u),\rho}$ is convex on $\mathcal{I}_{b(u),\rho}$, so for any $\alpha \in \mathcal{I}_{b(u),\rho}$

$$\Lambda'(u) \geq (t-s) \frac{LP_{b(u)}f}{P_{b(u)}} + d\Psi'_{b(u),\rho}(\alpha) \left(\frac{LP_{b(u)}f}{P_{b(u)}} - \alpha \right) + d\Psi_{b(u),\rho}(\alpha).$$

By Remark 4.5, for $\rho > 0$ there exists $\alpha \in \mathcal{I}_{b(u),\rho}$ such that $d\Psi'_{b(u),\rho}(\alpha) = -(t-s)$. Hence, for this α ,

$$\Lambda'(u) \geq (t-s)\alpha + d\Psi_{b(u),\rho}(\alpha).$$

It gives after integration over $u \in [0, 1]$,

$$P_s f(x) \leq P_t f(y) \exp \left(d \int_0^1 \left(-\frac{t-s}{d} \alpha - \Psi_{b(u),\rho}(\alpha) \right) du \right).$$

Now, by definition of the Legendre-Fenchel transform and of $b(u)$,

$$d \int_0^1 \left(-\frac{t-s}{d} \alpha - \Psi_{b(u),\rho}(\alpha) \right) ds \leq d \int_0^1 \Psi_{b(u),\rho}^* \left(-\frac{t-s}{d} \right) du = \frac{d}{t-s} \int_s^t \Psi_{u,\rho}^* \left(-\frac{t-s}{d} \right) du.$$

This concludes the argument.

Let us now prove the converse part. Let $s > 0$ and $y \in M$ be fixed. Let also x_ε (for $\varepsilon > 0$) be the exponential map starting from y and with initial tangent vector w . We apply (21) with $x = x_\varepsilon$ and $t = s + \varepsilon a$, $a \in \{-1, +1\}$. For f positive in \mathcal{A} , inequality (21) becomes

$$P_s f(x_\varepsilon) \leq P_{s+\varepsilon a} f(y) \exp \left(\frac{\varepsilon|w| + o(\varepsilon)}{\varepsilon a} \int_s^{s+\varepsilon a} \Psi_u^* \left(-\frac{\varepsilon a}{\varepsilon|w| + o(\varepsilon)} \right) du \right). \quad (22)$$

A first-order Taylor expansion of (22) in $\varepsilon > 0$ tending to 0 gives

$$\frac{\nabla P_s f}{P_s f} \cdot \frac{w}{|w|} \leq \frac{a}{|w|} \frac{LP_s f}{P_s f} + \Psi_{s,\rho}^* \left(-\frac{a}{|w|} \right)$$

at the point y . For $w = r \frac{\nabla P_s f}{|\nabla P_s f|}$ with $r > 0$, this can be written as

$$\frac{|\nabla P_s f|}{P_s f} \leq \frac{a}{r} \frac{LP_s f}{P_s f} + \Psi_{s,\rho}^* \left(-\frac{a}{r} \right).$$

Hence

$$\frac{|\nabla P_s f|}{P_s f} \leq - \left\{ \frac{LP_s f}{P_s f} z - \Psi_{t,\rho}^*(z) \right\}$$

for any $z \in \mathbb{R}$, since $r > 0$ and $a \in \{-1, +1\}$ are arbitrary. Since $\Psi_{s,\rho}^*$ is a convex function on \mathbb{R} , taking the infimum over z finally gives

$$\frac{|\nabla P_s f|}{P_s f} \leq -\Psi_{s,\rho}^* \left(\frac{LP_s f}{P_s f} \right)$$

at the point y . This is the Li-Yau inequality (12) at the arbitrary time $s > 0$ and point $y \in M$.

When $\rho \leq 0$, as explained in Remark 4.5, $\Psi'_{t,\rho}$ is a one-to-one function from $(n\rho\xi^t/4, \infty)$ onto $(-\infty, 0)$. We can use the same method as in the above case $\rho > 0$ but one can find such an α only if $0 < s < t$. In other words the argument works only for increasing functions b . The proof of the converse part is also the same in this case by considering only the $0 < s < t$. \triangleright

Remark 4.7 *Theorem 4.6 could be formally stated in the following general form : given a family Ψ_t of convex functions, an inequality of the form*

$$-\frac{|\nabla P_t f|}{P_t f} \geq \Psi_t \left(\frac{LP_t f}{P_t f} \right),$$

for any function $f > 0$ on M and $t > 0$ is equivalent to a Harnack inequality

$$P_s f(x) \leq P_t f(y) \exp \left(\frac{d(x, y)}{t-s} \int_s^t \Psi_u^* \left(-\frac{t-s}{d(x, y)} \right) du \right)$$

for any $s, t > 0$ and $x, y \in M$ such that $-(t-s)/d(x, y)$ is in the domain of the Legendre transforms Ψ_u^* .

Remark 4.8 *In the limit case $\rho = 0$, we have $\Psi_{t,0}(x) = -\sqrt{\frac{n}{2t} + x}$ for $x \geq -\frac{n}{2t}$. Then, $\Psi_{t,0}^*$ is only defined on $(-\infty, 0)$, with $\Psi_{t,0}^*(y) = -\frac{ny}{2t} - \frac{1}{4y}$ for $y < 0$. Therefore*

$$\frac{d}{t-s} \int_s^t \Psi_{u,0}^* \left(-\frac{t-s}{d} \right) du = \frac{d}{t-s} \int_s^t \left(\frac{n(t-s)}{2ud} + \frac{1}{4(t-s)} \right) du = \frac{n}{2} \log \left(\frac{t}{s} \right) + \frac{d^2}{4(t-s)}$$

for $0 < s < t$. Hence, under the $CD(0, n)$ condition, we recover the classical Harnack inequality

$$P_s f(x) \leq P_t f(y) \left(\frac{t}{s} \right)^{n/2} \exp \left(\frac{d(x, y)^2}{4(t-s)} \right), \quad 0 < s < t, \quad x, y \in M.$$

This Harnack inequality is equivalent to the classical Li-Yau inequality (1).

Let us now assume that the Markov semigroup admits a density kernel, that is, a function $p_t(x, y)$ such that for any function f , $P_t f(x) = \int f(y) p_t(x, y) dy$ where dy is the Riemannian measure. This is for instance the case if the semigroup is ultracontractive, so in particular if $\rho > 0$. Then a Harnack inequality classically implies a bound on the kernel. Here we obtain :

Corollary 4.9 (Heat kernel bound) *Under the $CD(\rho, n)$ condition, for any $x, y, z \in M$ it holds*

$$p_s(z, x) \leq p_t(z, y) \exp \left(\frac{d(x, y)}{t-s} \int_s^t \Psi_{u,\rho}^* \left(-\frac{t-s}{d(x, y)} \right) du \right) \quad (23)$$

for all $s, t > 0$ if $\rho > 0$, and all $0 < s < t$ if $\rho \leq 0$.

5 Comparison with earlier bounds

5.1 Linearisation of the Li-Yau inequality

For all given $t > 0$ and $\rho \neq 0$, the function Φ_t is concave (see Section 2). Hence the new Li-Yau inequality (12) admits a linearisation:

Proposition 5.1 (Linearisation of the Li-Yau inequality) *Under the $CD(\rho, n)$ condition with $\rho \in \mathbb{R}^*$ and $n \geq 1$*

i. for any $\alpha \geq 0$,

$$\frac{\Gamma(P_t f)}{(P_t f)^2} \leq A_1(\alpha) \frac{LP_t f}{P_t f} + \frac{n}{2} B_1(\alpha),$$

with

$$\begin{cases} A_1(\alpha) = 1 - \frac{\rho}{2\alpha \sinh^2(\alpha t)} (\sinh(2\alpha t) - 2\alpha t) \\ B_1(\alpha) = \frac{\alpha}{4 \sinh^2(\alpha t)} (\sinh(2\alpha t) + 2\alpha t) - \rho + \frac{\rho^2}{4\alpha \sinh^2(\alpha t)} (\sinh(2\alpha t) - 2\alpha t); \end{cases}$$

ii. for any $\beta \in (0, \pi/t)$,

$$\frac{\Gamma(P_t f)}{(P_t f)^2} \leq A_2(\beta) \frac{LP_t f}{P_t f} + \frac{n}{2} B_2(\beta),$$

with

$$\begin{cases} A_2(\beta) = 1 - \frac{\rho}{2\beta \sin^2(\beta t)} (2\beta t - \sin(2\beta t)) \\ B_2(\beta) = \frac{\beta}{4 \sin^2(\beta t)} (\sin(2\beta t) + 2\beta t) - \rho + \frac{\rho^2}{4\beta \sin^2(\beta t)} (2\beta t - \sin(2\beta t)). \end{cases}$$

Proof

◁ By Corollary 2.3 and concavity of the function Φ_t on $(-\infty, 1 + \frac{\pi^2}{\rho^2 t^2})$,

$$\frac{\Gamma(P_t f)}{(P_t f)^2} < \frac{n}{2} \Phi_t \left(\frac{4}{n\rho} \frac{LP_t f}{P_t f} \right) \leq \frac{2}{\rho} \Phi_t'(x_0) \frac{LP_t f}{P_t f} + \frac{n}{2} (\Phi_t(x_0) - x_0 \Phi_t'(x_0))$$

for any $x_0 < 1 + \pi^2/(\rho^2 t^2)$. By definition (8) of Φ_t , choosing $x_0 \leq 1$ gives case *i* with $\alpha = \rho\sqrt{1-x_0}$; observe then that the bound in *i* is the same for α and $-\alpha$. Choosing $1 < x_0 < 1 + \pi^2/(\rho^2 t^2)$ likewise gives case *ii* with $\beta = \rho\sqrt{x_0-1}$. ▷

Let us observe that the bounds in Proposition 5.1 can also be recovered by extending the method proposed in [BG11] (see also [Qia14]). In their Proposition 2.4, F. Baudoin and N. Garofalo use a close semigroup argument to prove that

$$\frac{\Gamma(P_t f)}{(P_t f)^2} \leq \left(1 - 2\rho \int_0^t V^2(s) ds \right) \frac{LP_t f}{P_t f} + \frac{n}{2} \left(\int_0^t V'(s) ds + \rho^2 \int_0^t V^2(s) ds - \rho \right)$$

for any positive function V on $[0, t]$ such that $V(0) = 1$ and $V(t) = 0$. They deduce diverse bounds for certain choices of V , which turn out to be suboptimal : in fact, it is a direct but long computation to see that optimising this inequality with respect to V , with $\int_0^t V^2(s) ds$ fixed, gives Proposition 5.1.

If we choose $\alpha = |\rho|$ in *i.* above we obtain the bound (5) :

Corollary 5.2 ([LX11]) *Under the $CD(-K, n)$ condition, then*

$$\frac{\Gamma(P_t f)}{(P_t f)^2} \leq \left(1 + \frac{\sinh(2Kt) - 2Kt}{2 \sinh^2(Kt)} \right) \frac{LP_t f}{P_t f} + \frac{n}{2} K(1 + \coth(Kt)) \quad (24)$$

for all positive t and f in \mathcal{A} .

By the same method as here, the authors in [LX11, Thm 1.3] deduced from (24) the following Harnack inequality : for every $0 < s < t$ and $f > 0$,

$$P_s f(x) \leq P_t f(y) \left(\frac{e^{2Kt} - 1 - 2Kt}{e^{2Ks} - 1 - 2Ks} \right)^{n/4} \exp \left(\frac{d(x, y)^2}{4(t-s)} \left(1 + \frac{t \coth(Kt) - s \coth(Ks)}{t-s} \right) \right). \quad (25)$$

Since our Harnack inequality (21) is equivalent to our Li-Yau inequality (12), which is stronger than (24), it follows that (21) is also stronger than (25).

5.2 Comparison with Davies' estimate (2) in negative curvature

Let us prove that the Li-Yau type inequality (12) given by Corollary 2.4 improves upon the bound (2) established for $\rho < 0$, in the notation $K = -\rho > 0$.

In our notation Φ_t and $X = 4LP_t f / (n\rho P_t f)$ we have to prove that

$$\Phi_t(X) \leq -\frac{\alpha K}{2} X + \frac{\alpha^2}{t} + \frac{K\alpha^2}{2(\alpha-1)}$$

for all $\alpha > 1, t > 0$ and X .

For $X < 1$, this can be written as

$$2ry \coth y \leq \frac{1}{\alpha-1} + 2r\alpha^2 + (\alpha-1)r^2 y^2$$

for all positive $r = (Kt)^{-1}$ and $y = Kt\sqrt{1-X}$. But $y \coth y \leq \sqrt{1+y^2}$ as can be seen by taking squares and using $y \leq \sinh(y)$ for $y \geq 0$. Hence

$$2ry \coth y \leq 2r\sqrt{1+y^2} \leq 2r + 2ry \leq 2r + \frac{1}{\alpha-1} + (\alpha-1)(ry)^2$$

by the Young inequality. This proves the claim for $X < 1$.

As regards the case where $X > 1$, we first recall the elementary bound : $y \cot y < 1 - y^2/4$ for all y in $(0, \pi)$. Letting indeed $u = y/2 \in (0, \pi/2)$, this is due to

$$y \cot y = u \frac{1 - \tan^2 u}{\tan u} < u \frac{1 - u^2}{u} = 1 - \frac{y^2}{4};$$

here we use that $\tan u > u$ with $x \mapsto \frac{1-x^2}{x} = \frac{1}{x} - x$ decreasing.

As a consequence, in the notation $r = (Kt)^{-1} > 0$ and $y = Kt\sqrt{X-1} \in (0, \pi)$,

$$\Phi_t(X) = K - \frac{K}{2} X + K r y \cot y \leq \frac{K}{2} [1 + 2r - r^2 y^2 - r y^2/2], \quad (26)$$

so arranging terms it is enough to prove that the second order polynomial

$$\frac{1}{\alpha-1} + 2r(\alpha^2 - 1) + y^2(r^2(1-\alpha) + r/2)$$

in y is non-negative.

We now observe that the left-hand side of (12) is non-negative, hence so is $\Phi_t(X)$, and finally the right-hand side of (26). In other words necessarily $y^2 \leq 2/r$ (that is, $X \leq 1 + 2/(Kt)$ for any $t > 0$). Now the above second order polynomial in y (with zero first order coefficient) is non-negative on the interval $[0, \sqrt{2/r}]$ if and only if it is so at 0 and $\sqrt{2/r}$, which is the case for any $\alpha > 1$ by direct computation. This proves the claim for $X > 1$.

The case $X = 1$ is covered as a limit case of any of these two cases, or can be directly considered. This concludes the argument, all cases being covered.

5.3 Comparison with estimate (3) in negative curvature

Let us now prove that the Li-Yau type inequality (12) also improves upon the bound (3) established for $\rho < 0$, in the notation $K = -\rho > 0$.

We first observe that, in the notation $r = (Kt)^{-1}$, $X = 4LP_t f / (n\rho)$ and $G = 4\Gamma(P_t f) / (nK)$, estimate (3) can be equivalently formulated as

$$X \leq -G + 2r + 2\sqrt{G + 1 + 2r} \quad (27)$$

In the case where $X < 1$, then estimate (12) can be written as

$$G \leq 2 - X + 2\sqrt{1 - X} \coth(r^{-1}\sqrt{1 - X}).$$

By the elementary inequality $y \coth y \leq \sqrt{1 + y^2} \leq 1 + y$, and the Young inequality, this implies

$$G \leq 2 - X + 2r + \frac{1}{\varepsilon} + \varepsilon(1 - X) \quad (28)$$

for all $\varepsilon > 0$, and then

$$X \leq -G + 2r + \delta G + \frac{1}{\delta}$$

for all $\delta = \varepsilon / (1 + \varepsilon) \in (0, 1)$. If $G > 1$ then we take $\delta = 1/\sqrt{G} < 1$, giving $X \leq -G + 2r + 2\sqrt{G}$; if $G \leq 1$ then we let δ tend to 1, giving $X \leq -G + 2r + G + 1 \leq -G + 2r + 2\sqrt{G + 1}$. In both cases this improves (27).

In the case where $X > 1$, then in the same notation the estimate (12) can be written as

$$G \leq 2 - X + 2\sqrt{X - 1} \cot(r^{-1}\sqrt{X - 1}).$$

Since moreover $y \cot y \leq 1$ for all $y \in (0, \pi)$, this implies the bound

$$X \leq -G + 2r + 2, \quad (29)$$

which in turn improves (27).

Since again the case $X = 1$ is a limit case or can be treated directly, all cases are covered.

5.4 Comparison with Hamilton's estimate (4) in negative curvature

Let us finally prove that (12) improves upon the bound (4) established for $\rho < 0$, in the notation $K = -\rho > 0$. Letting $s = Kt$, $X = 4LP_t f / (n\rho)$ and $G = 4\Gamma(P_t f) / (nK)$, estimate (4) can be written as

$$X \leq -e^{-2s} G + \frac{2}{s} e^{2s}. \quad (30)$$

In the case where $X < 1$, then choosing $\varepsilon = 2s$ in (28) ensures that (12) implies the bound

$$X \leq -\frac{1}{1 + 2s} G + \frac{5 + 4s + 4s^2}{2s(1 + 2s)}.$$

This in turn implies (30) by direct comparison of both coefficients on the right-hand sides. In the case where $X > 1$, then the bound (29), where $r = 1/s$, likewise implies (30). As above this concludes the argument.

Acknowledgements. This research was supported by the French ANR-12-BS01-0019 STAB project.

References

- [BG11] F. Baudoin and N. Garofalo. Perelman’s entropy and doubling property on Riemannian manifolds. *J. Geom. Anal.*, 21(4):1119–1131, 2011.
- [BGL14] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grund. Math. Wiss.* Springer, Cham, 2014.
- [BL06] D. Bakry and M. Ledoux. A logarithmic Sobolev form of the Li-Yau parabolic inequality. *Rev. Mat. Iberoam.*, 22(2):683–702, 2006.
- [BQ99] D. Bakry and Z. Qian. Harnack inequalities on a manifold with positive or negative Ricci curvature. *Rev. Mat. Iberoam.*, 15(1):143–179, 1999.
- [Dav89] E. B. Davies. *Heat kernels and spectral theory*, volume 92 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1989.
- [GM14] N. Garofalo and A. Mondino. Li-Yau and Harnack type inequalities in $RCD^*(K, N)$ metric measure spaces. *Nonlinear Anal.*, 95:721–734, 2014.
- [Ham93] R. Hamilton. A matrix Harnack estimate for the heat equation. *Comm. Anal. Geom.*, 1(1):88–99, 1993.
- [Lee13] P. W. Y. Lee. Generalized Li-Yau estimates and Huisken’s monotonicity formula. Preprint, 2013.
- [LX11] J. Li and X. Xu. Differential Harnack inequalities on Riemannian manifolds I: linear heat equation. *Adv. Math.*, 226(5):4456–4491, 2011.
- [LY86] P. Li and S.-T. Yau. On the parabolic kernel of the Schrödinger operator. *Acta Math.*, 156(3-4):153–201, 1986.
- [Qia12] B. Qian. A generalization of Hamilton’s gradient estimate. *J. Geom. Phys.*, 62(5):1064–1069, 2012.
- [Qia14] B. Qian. Remarks on differential Harnack inequalities. *J. Math. Anal. Appl.*, 409(1):556–566, 2014.
- [QZZ13] Z. Qian, H.-C. Zhang, and X.-P. Zhu. Sharp spectral gap and Li-Yau’s estimate on Alexandrov spaces. *Math. Z.*, 273(3-4):1175–1195, 2013.
- [SZ06] P. Souplet and Q. S. Zhang. Sharp gradient estimate and Yau’s Liouville theorem for the heat equation on noncompact manifolds. *Bull. London Math. Soc.*, 38(6):1045–1053, 2006.
- [Vil09] C. Villani. *Optimal transport, Old and new*, volume 338 of *Grund. Math. Wiss.* Springer, Berlin, 2009.
- [Wan10] F.-Y. Wang. Gradient and Harnack inequalities on noncompact manifolds with boundary. *Pacific J. Math.*, 245(1):185–200, 2010.
- [Yau95] S.-T. Yau. Harnack inequality for non-self-adjoint evolution equations. *Math. Res. Lett.*, 2(4):387–399, 1995.