



HAL
open science

A combinatorial move on the set of Jenkins-Strebel differentials

Corentin Boissy

► **To cite this version:**

| Corentin Boissy. A combinatorial move on the set of Jenkins-Strebel differentials. 2014. hal-01097028

HAL Id: hal-01097028

<https://hal.science/hal-01097028>

Preprint submitted on 18 Dec 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A COMBINATORIAL MOVE ON THE SET OF JENKINS-STREBEL DIFFERENTIALS

CORENTIN BOISSY

ABSTRACT. We describe an elementary combinatorial move on the set of quadratic differentials with a horizontal one cylinder decomposition. Computer experiment suggests that the corresponding equivalent classes are in one-to-one correspondence with the connected component of the strata.

1. INTRODUCTION

A nonzero quadratic differential with at most simple poles on a compact Riemann surface naturally defines a flat metric with conical singularities on this surface. Geometry and dynamics on such flat surfaces, in relation to geometry and dynamics on the corresponding moduli space of Abelian differentials is a very rich topic and has been widely studied in the last 30 years. It is related to interval exchange transformations, billiards in polygons, Teichmüller dynamics.

One important case is when the quadratic differential is the square of a holomorphic one-form (Abelian differential). It was first proven by Veech [6] that the strata of Abelian differentials are not connected, by using a combinatorial description, usually called *extended Rauzy classes*. The connected components of the moduli space of Abelian differentials were described by Kontsevich and Zorich in [2]. They showed that each stratum has up to three connected component, which are described by two invariants: hyperellipticity and parity of spin structure, that arise under some conditions on the set of zeroes. Later, Lanneau has described the connected components of the moduli space of quadratic differentials. In this case, the nonconnected strata either contain a hyperelliptic connected component (which is an infinite family of strata), or belong to a finite family of “exceptionnal strata”.

Date: December 18, 2014.

2000 Mathematics Subject Classification. Primary: 37E05. Secondary: 37D40.

Key words and phrases. Interval exchange maps, Rauzy induction, Abelian differentials, Moduli spaces, Teichmüller flow.

The author is partially supported by ANR “GeoDyM”.

The exceptional strata $\mathcal{Q}(-1, 9)$, $\mathcal{Q}(-1, 3, 6)$, $\mathcal{Q}(-1, 3, 3, 3)$ and $\mathcal{Q}(12)$ were initially proved non connected by the computation by Zorich of the corresponding extended Rauzy classes. More recently, Möller and Chen [4] discovered few more exceptional strata: $\mathcal{Q}(3, 9)$, $\mathcal{Q}(6, 6)$, $\mathcal{Q}(3, 3, 6)$ and $\mathcal{Q}(3, 3, 3, 3)$, and gave a proof of the non-connectedness of all exceptional strata, by using techniques of algebraic geometry.

2. CLASSES OF CYLINDRICAL PERMUTATIONS

2.1. Case of Abelian differentials. Consider a finite alphabet \mathcal{A} and a vector $\zeta \in \mathbb{R}^{\mathcal{A}}$, whose entries have positive real part. Choose two total ordering of \mathcal{A} , *i.e.* one-to-one maps $\pi_t, \pi_b : \mathcal{A} \rightarrow \{1, \dots, d\}$. The combinatorial data $\pi = (\pi_t, \pi_b)$ will be usually written as an array with two lines of symbols in \mathcal{A} :

$$\begin{pmatrix} \pi_t^{-1}(1) & \dots & \pi_t^{-1}(d) \\ \pi_b^{-1}(1) & \dots & \pi_b^{-1}(d) \end{pmatrix}$$

Such π will be referred to as a *labeled permutation*.

We construct a translation surface in the following way, that we will refer to as the *cylindrical construction* (note that *is not* the Veech construction): consider the broken line L_t on $\mathbb{C} = \mathbb{R}^2$ defined by concatenation of the vectors $\zeta_{\pi_t^{-1}(j)}$ (in this order) for $j = 1, \dots, d$ with starting point at $(0, l) \in \mathbb{R}^2$. Similarly, we consider the broken line L_b defined by concatenation of the vectors $\zeta_{\pi_b^{-1}(j)}$ (in this order) for $j = 1, \dots, d$ with starting point at the origin. Then join the two starting points of L_t, L_b a vertical segment and similarly with the two ending points. If l is large enough, so that L_t, L_b do not intersect, one gets a polygon whose sides come by pairs of a parallel segments of the same length. Gluing these segments by translations, one gets a translation surface $S(\pi, \zeta)$, with $\pi = (\pi_t, \pi_b)$ (see Figure 1).




FIGURE 1. Example of cylindrical construction

If ζ has zero imaginary part, $S(\pi, \zeta)$ is a translation surface with a one cylinder decomposition.

Now we choose any continuous data $\zeta \in \mathbb{C}^{\mathcal{A}}$ such that $Im(\zeta_\alpha) < 0$ we deform the S by changing the real part of ζ_α until we have $Re(\zeta_\alpha) < 0$ (see Figure 2). Keeping the other parameter unchanged and preserving the identifications, we still have a translation surface, that belongs to the same connected component as S (if $|Re(\zeta_\alpha)|$ is small enough). This surface is not any more presented as in the previous construction. However, after some suitable cutting and pasting, it is the case, with another combinatorial datum.

We assume that α is not the leftmost element of the top line component, nor the rightmost element of the bottom line. Let $\beta_1 \in \mathcal{A}$ be the index that correspond to the segment immediately on the left of α in L_t and $\beta_2 \in \mathcal{A}$ by the one immediately on the right of α in L_b . We now cut the triangle that contain the segments corresponding to α and β_1 on the top, and glue it along the segment corresponding to β_1 on the bottom. Similarly, we cut the triangle in the bottom corresponding to α, β_2 and glue it to the top along the segment corresponding to β_2 . So the flat surface is obtained by the same construction as before, where the combinatorial data

$$\begin{pmatrix} \dots \beta_1 \alpha \dots \beta_2 \dots \\ \dots \alpha \beta_2 \dots \beta_1 \dots \end{pmatrix}$$

as been replaced by

$$\begin{pmatrix} \dots \beta_1 \dots \alpha \beta_2 \dots \\ \dots \beta_2 \dots \beta_1 \alpha \dots \end{pmatrix}$$

and where ζ as been replaced by ζ' such that

$$\begin{aligned} \zeta'_\alpha &= -\zeta_\alpha \\ \zeta'_{\beta_1} &= \zeta_{\beta_1} + \zeta_\alpha \\ \zeta'_{\beta_2} &= \zeta_{\beta_2} + \zeta_\alpha \end{aligned}$$

Notation 2.1. If π is the initial combinatorial datum we will denote the new combinatorial datum by $T_\alpha(\pi)$.

One can also do an analogous procedure by starting from ζ_α with positive imaginary part. One gets the transformation

$$\begin{pmatrix} \dots \beta_1 \dots \alpha \beta_2 \dots \\ \dots \beta_2 \dots \beta_1 \alpha \dots \end{pmatrix} \rightarrow \begin{pmatrix} \dots \beta_1 \alpha \dots \beta_2 \dots \\ \dots \alpha \beta_2 \dots \beta_1 \dots \end{pmatrix}$$

i.e., one gets $T_\alpha^{-1}(\pi)$

The following proposition is clear from the definition of T_α .




FIGURE 2. Changing the combinatorial datum for the cylindrical construction

Proposition 2.1. π and $T_\alpha(\pi)$ corresponds to translation surfaces in the in the same connected component of the same stratum, when $T_\alpha(\pi)$ is well defined.

There are some other simple transformations that preserve the connected component.

$$U_t : \begin{pmatrix} \alpha_1 \alpha_2 \dots \alpha_n \\ \beta_1 \beta_2 \dots \beta_n \end{pmatrix} \mapsto \begin{pmatrix} \alpha_2 \dots \alpha_n \alpha_1 \\ \beta_1 \beta_2 \dots \beta_n \end{pmatrix}$$

$$U_b : \begin{pmatrix} \alpha_1 \alpha_2 \dots \alpha_n \\ \beta_1 \beta_2 \dots \beta_n \end{pmatrix} \mapsto \begin{pmatrix} \alpha_1 \alpha_2 \dots \alpha_n \\ \beta_2 \dots \beta_n \beta_1 \end{pmatrix}$$

and

$$S : \begin{pmatrix} \alpha_1 \alpha_2 \dots \alpha_n \\ \beta_1 \beta_2 \dots \beta_n \end{pmatrix} \mapsto \begin{pmatrix} \beta_n \dots \beta_2 \beta_1 \\ \alpha_n \dots \alpha_2 \alpha_1 \end{pmatrix}$$

A renumbering of a labeled permutation is obtained by precomposing the maps π_t, π_b with a permutation of the alphabet \mathcal{A} . For instance, the labeled permutation $\begin{pmatrix} b & c & d & a \\ a & d & c & b \end{pmatrix}$ is a renumbering of the labeled permutation $\begin{pmatrix} a & b & c & d \\ d & c & b & a \end{pmatrix}$. Of course, the set of labeled permutation on a alphabet \mathcal{A} up to renumbering is naturally identified with the set of permutations of \mathcal{A} .

Computing equivalent classes for these action, one gets the following experimental result, obtained by computer experiment¹.

Observation 2.2. *Two labeled permutations with up to 12 letters representing the same connected component for the cylindrical construction in the moduli space of Abelian differentials, can be joined by a succession of moves $T_\alpha, T_\alpha^{-1}, U_t, U_b$, up to renumbering.*

2.2. Case of quadratic differentials. One can define quadratic differentials in a similar way, by using *generalized permutation*. We will define a generalized permutation of type (l, m) as an array of two lines of symbols in a finite alphabet \mathcal{A} :

$$\pi = \begin{pmatrix} \alpha_1 & \dots & \alpha_l \\ \beta_1 & \dots & \beta_m \end{pmatrix}$$

with the condition that each letter appears exactly two times in the list $(\alpha_1, \dots, \alpha_l, \beta_1, \dots, \beta_m)$. We also assume that there exists $1 \leq i, j \leq l$ such that $\alpha_i = \alpha_j$ and $1 \leq i', j' \leq m$ such that $\beta_{i'} = \beta_{j'}$, *i.e.* there is at least a symbol that appear twice on the top line (and therefore, not on the bottom line) and there is at least a symbol that appear twice on the bottom line (and not on the top line).

Then, choosing a vector $\zeta \in \mathbb{C}^{\mathcal{A}}$ whose entries have positive real part, and with the extra condition that $\sum_{i=1}^l \zeta_{\alpha_i} = \sum_{j=1}^m \zeta_{\beta_j}$ one gets a flat surface be a similar construction as before. The resulting surface is a half-translation surface, *i.e.* a Riemann surface endowed with a

¹The SAGE programs can be found on the webpage of the author.

quadratic differential which is not the square of an Abelian differential. As before, if all the ζ_i have zero imaginary part, the corresponding flat surface has a one cylinder decomposition.

Definition 2.3. Let π be a generalized permutation. A letter $\alpha \in \mathcal{A}$ is said to be of type *top-bottom* if it appears both on the top and the bottom line. Similarly, it is of type *top-top* (resp. *bottom-bottom*) if it appears only on the top (resp. bottom).

We will define an analogous move as for the case of Abelian differential. However, the corresponding combinatorial operation depends on the type of the involved letters α, β_1, β_2 (and the definition of β_1, β_2 depends on the type of $\alpha \dots$). However, by writing π in a slightly different way, one gets a unified combinatorial operation.

Definition 2.4. Let $\pi = \left(\begin{smallmatrix} \alpha_1, \dots, \alpha_l \\ \beta_1, \dots, \beta_m \end{smallmatrix} \right)$. We define the *one line representation* of π to be

$$\alpha_1 \alpha_2 \dots \alpha_l | \beta_m \beta_{m-1} \dots \beta_1$$

Definition 2.5. Let π be a generalized permutation, and $\alpha \in \mathcal{A}$ with the following condition:

- α is either of type top-bottom, or is not the only top-top (resp. bottom-bottom) letter (such α is called *regular*).
- Each occurrence of α admits a letter on its left in the one line representation of π (*i.e.* no occurrence of α not the leftmost letter, or follows the symbol “|”).

Let π be written in a one line representation as (up to changing the order of the four blocks $\{\beta_1 \alpha, \beta_2 \alpha, \beta_1, \beta_2\}$):

$$\dots \beta_1 \alpha \dots \beta_2 \alpha \dots \beta_1 \dots \beta_2 \dots$$

Then we define $T_\alpha(\pi)$ to be the generalized permutation:

$$\dots \beta_1 \dots \beta_2 \dots \alpha \beta_1 \dots \alpha \beta_2 \dots$$

One has the following proposition:

Proposition 2.6. *Let α that satisfies the condition of Definition 2.5. Then, π and $T_\alpha(\pi)$ corresponds to the same connected component of the moduli space of quadratic differentials.*

Proof. The proof is made by doing similar surgeries as in the Abelian case. We assume that α is of type top-bottom, its top occurrence is not the leftmost element of the (top) line, and its bottom occurrence is not the rightmost of the (bottom) line. Let β_1 be the letter immediately on the left of α on the top line, and β_2 be the letter immediately on the right of α on the bottom line. Choose a continuous parameter ζ

such that $Im(\zeta_\alpha) < 0$. And change continuously the real part of ζ_α so that it becomes negative. Then, as before, we cut the triangle of sides corresponding to $\beta_1\alpha$, and glue it by along the side corresponding to the other occurrence of β_1 . If β_1 is on the bottom, the gluing is done by a translation, otherwise it is done by an half turn. We do a similar cutting and pasting for the triangle whose sides correspond to $\alpha\beta_2$.

If β_1, β_2 are both of type top-bottom, the combinatorial datum is changed in exactly the same way as before. Assume for instance that β_1 is top-top, and β_2 is top-bottom. Then, it is easy to see that π is changed in the following way:

$$\pi = \left(\begin{array}{c} \dots \beta_1 \alpha \dots \beta_1 \dots \beta_2 \dots \\ \dots \alpha \beta_2 \dots \dots \dots \end{array} \right) \rightarrow \pi' = \left(\begin{array}{c} \dots \beta_1 \dots \alpha \beta_1 \dots \alpha \beta_2 \dots \\ \dots \beta_2 \dots \dots \beta_1 \dots \dots \end{array} \right)$$

The condition on α and the generalized permutation π' , when written in a one line representation, corresponds precisely to Definition 2.5 and $T_\alpha(\pi)$.

When α is top-top (resp. bottom-bottom), we can define analogous transformation. We want to have ζ_α with negative real part and all other ζ_i with positive real part. Recall that there is a linear condition (1) on ζ that is needed in the definition of the surface. This condition forbid α to be the only top-top letter. \square

The maps U_t, U_b and S are defined analogously as in the case of Abelian differentials.

Definition 2.7. A generalized permutation is *mixed* if it contains top-bottom, top-top and bottom-bottom letter.

Lemma 2.8. *Any connected component contain a one cylinder surface defined by a mixed generalized permutation.*

Proof. We start from a flat surface S in a connected component \mathcal{C} with a one cylinder decomposition (such surface always exists, by [3], Theorem 3.6), and assume that it is not mixed. Each singularity appear on one of the horizontal side, but not both. Consider one horizontal side. If there are more than two singularities, and more than one saddle connection, then there exists a saddle connection joining a singularity P_1 , of order k_1 to a singularity P_2 of order k_2 , and we can shrink it to obtain a singularity of order $k_1 + k_2$ on a surface S' . We obtain a connected component with less singularities. Assume there exists a cylinder as in the lemma, then we can continuously change S' , until it is that cylinder. Breaking up continously the singularity of order $k_1 + k_2$ along this path (see [2] or [3] for the precise definition of breaking up

a singularity), we get a continuous path in \mathcal{C} , whose endpoint is the required cylinder.

The former procedure cannot be realized if each side contains only one singularity or is composed of only one saddle connection (*i.e.* contains exactly two simple poles). So we only need to check the result for such surfaces.

We use Rauzy induction (see [1], for the definition of the Rauzy moves $\mathcal{R}_t, \mathcal{R}_b$). Assume for instance that one side is a pair of poles. The generalized permutation for the cylinder decomposition is of the form

$$\begin{pmatrix} * & * & * \\ a & a & \end{pmatrix}$$

The corresponding generalized permutation for the Veech construction is

$$\begin{pmatrix} 0 & * & * & * \\ a & a & 0 & \end{pmatrix}$$

If “***” is not of the form $a_1 \dots a_n a_1 \dots a_n$, then using the Rauzy move \mathcal{R}_b , one gets

$$\begin{pmatrix} 0 \dots b \dots b \dots c \dots c \\ aa0 \end{pmatrix}$$

so that there is an element “ d ” between the two “ c ” whose other occurrence is not between the two “ c ”. Then, by applying suitably the Rauzy induction, one gets

$$\begin{aligned} & \begin{pmatrix} 0 \dots b \dots b \dots c \dots d \dots c \\ aa0 \end{pmatrix} \xrightarrow{\mathcal{R}_t} \begin{pmatrix} 0 \dots b \dots b \dots 0c \dots d \dots c \\ aa \end{pmatrix} \\ & \xrightarrow{\mathcal{R}_b^k} \begin{pmatrix} 0 \dots b \dots b \dots \\ c \dots d \dots c0aa \end{pmatrix} \xrightarrow{\mathcal{R}_t^2} \begin{pmatrix} 0 \dots b \dots b \dots \\ c \dots d \dots c0 \end{pmatrix} \end{aligned}$$

Then, the element “ d ”, is top-bottom. If the top line was initially $a_1 \dots a_n a_1 \dots a_n$, then the corresponding connected component was a hyperelliptic one. In such connected component, one can find mixed generalized permutations (for the cylindrical construction) of the kind

$$\begin{pmatrix} a_1 \dots a_n \alpha b_1 \dots b_m \alpha \\ \beta b_m \dots b_1 \beta a_n \dots a_1 \end{pmatrix}$$

The other cases are left to the reader. \square

We now can state the key result, which is proved by computer experiment².

²The SAGE programs can be found on the webpage of the author.

Observation 2.9. *Two mixed generalized permutations of at most 10 letter representing the same connected component can be joined by a succession of moves $T_\alpha, T_\alpha^{-1}, U_t, U_b$ and S , up to renumbering.*

Remark 2.10. In particular, we “see” all the connected components of the exceptional strata $\mathcal{Q}(-1, 9)$, $\mathcal{Q}(-1, 3, 6)$, $\mathcal{Q}(-1, 3, 3, 3)$, $\mathcal{Q}(12)$, $\mathcal{Q}(3, 9)$, $\mathcal{Q}(3, 3, 6)$, $\mathcal{Q}(6, 6)$ and $\mathcal{Q}(3, 3, 3, 3)$.

Remark 2.11. In fact, the operation S is not necessary, except the for strata $\mathcal{Q}(-1, 5)$, $\mathcal{Q}(-1, 1, 4)$ and $\mathcal{Q}(-1, 9)$, for which avoiding S leads to one extra class.

REFERENCES

- [1] C. Boissy and E. Lanneau. Dynamics and geometry of the raufy-veech induction for quadratic differentials. *Ergodic Theory and Dynamical Systems*, 29:767–816, 2009.
- [2] M. Kontsevich and A. Zorich. Connected components of the moduli spaces of Abelian differentials with prescribed singularities. *Invent. Math.*, 153(3):631–678, 2003.
- [3] Erwan Lanneau. Connected components of the strata of the moduli spaces of quadratic differentials. *Ann. Sci. Éc. Norm. Supér. (4)*, 41(1):1–56, 2008.
- [4] M. Möller and D. Chen. Quadratic differentials in low genus: exceptional and non-varying. arXiv:1204.1707, *Ann. Sci. Éc. Norm. Supér.* (to appear).
- [5] W. Veech. Gauss measures for transformations on the space of interval exchange maps. *Ann. of Math. (2)*, 115(1):201–242, 1982.
- [6] W. Veech. Moduli spaces of quadratic differentials. *J. Analyse Math.*, 55:117–170, 1990.

CORENTIN BOISSY, AIX MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 13453 MARSEILLE FRANCE
E-mail address: corentin.boissy@univ-amu.fr