



HAL
open science

A relevant glucose-insulin dynamics model for type 1 diabetes

Nicolas Magdelaine, Lucy Chaillous, Isabelle Guilhem, Jean-Yves Poirier,
Michel Krempf, Claude H. Moog, Eric Le Carpentier

► **To cite this version:**

Nicolas Magdelaine, Lucy Chaillous, Isabelle Guilhem, Jean-Yves Poirier, Michel Krempf, et al.. A relevant glucose-insulin dynamics model for type 1 diabetes. ATTD 2015; 8th International Conference on Advanced Technologies & Treatments of Diabetes, Feb 2015, Paris, France. hal-01097526

HAL Id: hal-01097526

<https://hal.science/hal-01097526v1>

Submitted on 18 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A relevant glucose-insulin dynamics model for type 1 diabetes

N. Magdelaine¹, L. Chaillous², I. Guilhem³, J.-Y. Poirier³, M. Krempf², C. Moog¹, E. Le Carpentier¹



1: IRCCyN, Nantes
2: CHU Nantes, Diabetes and Metabolism
3: CHU Rennes, Diabetes and Metabolism

Objective

A new model of glucose insulin dynamics for type 1 diabetes **without non desired equilibria**.

A standard identification algorithm provides a long term fit over two days with constant parameters.

As a byproduct, **the parameters** of the relevant model provide the computation of the tools for the **functional insulin therapy (FIT)**:

- the **basal infusion rate (BR)**,
- the **insulin sensitivity factor (ISF)** and
- the **insulin to carbo ratio (I:C)**.

This is a major outcome as they can be used for the education of type 1 diabetic patients.

Methods

Glucose dynamics

Increase of glycemia includes:

- the liver endogenous glucose production k_l
- the digestion of CHO $D(t)$.

The glucose absorption rate includes:


- an insulin-independent term k_b
- an insulin-dependent term $-k_{si} \cdot I(t)$

The dynamics of the blood glucose level (BG) $G(t)$ is:

$$\begin{cases} \dot{G}(t) = -k_{si} \cdot I(t) + k_l - k_b + D(t) \\ \dot{G}(t) = 0 \text{ if } G(t) = 0 \text{ for positivity} \end{cases}$$

Insulin dynamics

A second order from the pharmacokinetic profile :



$$I(s) = \frac{k_u/V_i}{(1+T_u \cdot s)^2} U(s)$$

Digestion dynamics

A second order with a single time constant

Six free parameters to be identified.

Insulin sensitivity k_{si} is one major parameter which has to be identified for each patient as it reflects the well-known variability of the metabolic effect of insulin.


Classical models

R.N. Bergman et al. "Quantitative estimation of insulin sensitivity," *Am. J. Physiol. Endocrinol. Metab.*, 236:667-677, 1979.

A. De Gaetano and O. Arino, "Mathematical modelling of the intravenous glucose tolerance test," *J. Math. Biol.*, vol. 40, no. 2, pp. 136-168, Feb. 2000.

Results

1. Equilibrium during fasting ($D(t)=0$) :



2. Computed tools for the functional insulin therapy :

$$U_b = \frac{k_l - k_b}{k_{si}} \frac{V_i}{k_u}$$

$$ISF = k_{si} \frac{k_u}{V_i}$$


$$I : C = \frac{V_i}{k_{si} k_u} \frac{k_r}{V_B}$$

3. Estimation algorithm

An off-line estimation, with the least square method on the output error, using standard clinical data (CGM, Injection, CHO) of a type 1 diabetic patient provides a fit over two days and the constant parameters allow to compute the tools for the FIT:

U_b 0.7 U/h	ISF 83 mg/dl/U	I:C 1: 17 U:g _{CHO}
------------------	-------------------	---------------------------------

The method is extended to a piecewise time varying model, improving the fit on an even longer period. k_{si} is assumed to be constant. The time varying parameter k_r reflects the glycemic index depending on the meal composition and the variations of the liver glucose production k_l displays a basal rate pattern as used by some patients.



Conclusion

This new model has realistic asymptotic properties. Its relevant parameters provide the tools for FIT. An adaptive estimation algorithm displays a long-term fit. This new model could impact the design of a control algorithm for the artificial pancreas.