



Concentration inequalities in the infinite urn scheme for occupancy counts and the missing mass, with applications

Anna Ben-Hamou, Stephane Boucheron, Mesrob I. Ohannessian

► To cite this version:

Anna Ben-Hamou, Stephane Boucheron, Mesrob I. Ohannessian. Concentration inequalities in the infinite urn scheme for occupancy counts and the missing mass, with applications. Bernoulli, 2017, 10.3150/15-BEJ743 . hal-01101671

HAL Id: hal-01101671

<https://hal.science/hal-01101671>

Submitted on 9 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Concentration inequalities in the infinite urn scheme for occupancy counts and the missing mass, with applications

Anna Ben-Hamou¹ Stéphane Boucheron^{1,2} and Mesrob I. Ohannessian³

¹*LPMA UMR 7599 CNRS & Université Paris-Diderot, 5, rue Thomas Mann 75013 Paris France. e-mail: anna.benhamou@univ-paris-diderot.fr*

²*DMA ENS Ulm, 45, rue d'Ulm 75005 Paris France. e-mail: stephane.boucheron@univ-paris-diderot.fr; url: stephane-v-boucheron.fr*

³*MSR-Inria Joint Centre, 1, rue Honoré d'Estienne d'Orves 91120 Palaiseau France e-mail: mesrob@gmail.com; url: sites.google.com/site/mesrob*

Abstract: An infinite urn scheme is defined by a probability mass function $(p_j)_{j \geq 1}$ over positive integers. A random allocation consists of a sample of N independent drawings according to this probability distribution where N may be deterministic or Poisson-distributed. This paper is concerned with occupancy counts, that is with the number of symbols with r or at least r occurrences in the sample, and with the missing mass that is the total probability of all symbols that do not occur in the sample. Without any further assumption on the sampling distribution, these random quantities are shown to satisfy Bernstein-type concentration inequalities. The variance factors in these concentration inequalities are shown to be tight if the sampling distribution satisfies a regular variation property. This regular variation property reads as follows. Let the number of symbols with probability larger than x be $\bar{\nu}(x) = |\{j: p_j \geq x\}|$. In a regularly varying urn scheme, $\bar{\nu}$ satisfies $\lim_{\tau \rightarrow 0} \bar{\nu}(\tau x) / \bar{\nu}(\tau) = x^{-\alpha}$ for $\alpha \in [0, 1]$ and the variance of the number of distinct symbols in a sample tends to infinity as the sample size tends to infinity. Among other applications, these concentration inequalities allow us to derive tight confidence intervals for the Good-Turing estimator of the missing mass.

Keywords and phrases: Concentration, regular variation, occupancy, missing mass, rare species.

1. Introduction

From the 20th century to the 21st, various disciplines have tried to infer something about scarcely observed events: zoologists about species, cryptologists about cyphers, linguists about vocabularies, and data scientists about almost everything. These problems are all about ‘small data’ within possibly much bigger data. Can we make such inference?

Problem setting

To move into a concrete setting, let U_1, U_2, \dots, U_n be i.i.d. observations from a fixed but unknown distribution $(p_j)_{j=1}^\infty$ over a discrete set of symbols $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$. We consider each j in \mathbb{N}^* as a discrete *symbol* devoid of numerical significance. The terminology of ‘infinite urn scheme’ comes from the analogy to n independent throws of balls over an infinity of urns, p_j being the probability of a ball falling into urn j , at any i -th throw. We alternatively adhere to the symbols or the urns perspective, based on which carries the intuition best. Species, cyphers, and vocabularies all being discrete, are well modeled as such. The sample size n may be fixed in advance; we call this the *binomial setting*. It may be randomly set by the duration of an experiment; this gives rise to the *Poisson setting*. More precisely, in the latter case we write it as N , a Poisson random variable independent of (U_i) and with expectation t . We index all Poisson-setting quantities by t and write them with functional notations, instead of subscripts used for the fixed- n scheme.

For each $j, n \in \mathbb{N}^*$, let $X_{n,j} = \sum_{i=1}^n \mathbb{I}_{\{U_i=j\}}$ be the number of times symbol j occurs in a sample of size n , and $X_j(t) = \sum_{i=1}^{N(t)} \mathbb{I}_{\{U_i=j\}}$ the Poisson version. In questions of underrepresented data, the central objects are sets of symbols that are repeated a small number r of times. The central quantities are the *occupancy counts* $K_{n,r}$ [respectively $K_r(t)$ for the Poisson setting], defined as the number of symbols that appear exactly r times in a sample of size n :

$$K_{n,r} = |\{j, X_{n,j} = r\}| = \sum_{j=1}^{\infty} \mathbb{I}_{\{X_{n,j}=r\}}.$$

The collection $(K_{n,r})_{r \geq 1}$ [resp. $(K_r(t))_{r \geq 1}$] has been given many names, such as the “profile” (in information theory [Orlitsky et al., 2004]) or the “fingerprint” (in theoretical computer science [Batu et al., 2001, Valiant and Valiant, 2011]) of the probability distribution $(p_j)_{j \in \mathbb{N}}$. Here we refer to them by *occupancy counts* individually, and *occupancy process* all together.

The occupancy counts then combine to yield the cumulated occupancy counts $K_{n,\bar{r}}$ [respectively $K_{\bar{r}}(t)$] and the total number of distinct symbols in the sample, or the total number of occupied urns, often called the *coverage* and denoted by K_n [respectively $K(t)$]:

$$K_{n,\bar{r}} = |\{j, X_{n,j} \geq r\}| = \sum_{j=1}^{\infty} \mathbb{I}_{\{X_{n,j} \geq r\}} = \sum_{s \geq r} K_{n,s},$$

and

$$K_n = |\{j, X_{n,j} > 0\}| = \sum_{j=1}^{\infty} \mathbb{I}_{\{X_{n,j} > 0\}} = \sum_{r \geq 1} K_{n,r}.$$

In addition to the occupancy numbers and the number of distinct symbols, we also address the *rare* (or small-count) *probabilities* $M_{n,r}$ [respectively $M_r(t)$],

defined as the probability mass corresponding to all symbols that appear exactly r times:

$$M_{n,r} = \mathbb{P}(\{j, X_{n,j} = r\}) = \sum_{j=1}^{\infty} p_j \mathbb{I}_{\{X_{n,j}=r\}}.$$

In particular, we focus on $M_{n,0} = \sum_{j=1}^{\infty} p_j \mathbb{I}_{\{X_{n,j}=0\}}$, which is called the missing mass, and which corresponds to the probability of all the unseen symbols.

Explicit formulas for the moments of the occupancy counts and masses can be derived in the binomial and Poisson settings. The occupancy counts' expectations are given by

$$\begin{aligned} \mathbb{E}K_n &= \sum_{j=1}^{\infty} (1 - (1 - p_j)^n) & \mathbb{E}K(t) &= \sum_{j=1}^{\infty} (1 - e^{-np_j}) \\ \mathbb{E}K_{n,r} &= \sum_{j=1}^{\infty} \binom{n}{r} p_j^r (1 - p_j)^{n-r} & \mathbb{E}K_r(t) &= \sum_{j=1}^{\infty} e^{-np_j} \frac{(np_j)^r}{r!} \\ \mathbb{E}M_{n,r} &= \sum_{j=1}^{\infty} \binom{n}{r} p_j^{r+1} (1 - p_j)^{n-r} & \mathbb{E}M_r(t) &= \sum_{j=1}^{\infty} e^{-np_j} p_j \frac{(np_j)^r}{r!}. \end{aligned}$$

Formulas for higher moments can also be computed explicitly but their expression, especially in the binomial setting where a lot of dependencies are involved, often has an impractical form.

This classical occupancy setting naturally models a host of different application areas, including computational linguistics, ecology, and biology. Urns may represent species, and we are interested in the number of distinct species observed in a sample (the ecological diversity) or in the probability of the unobserved species. In linguistics, urns may represent words. In both of these applications, the independence assumption of the random variables $\{U_i\}_{i=1\dots n}$ may seem unrealistic. For instance in a sentence, the probability of appearance of a word strongly depends on the previous words, both for grammatical and semantic reasons. Likewise, the nucleotides in a DNA sequence do not form an i.i.d. sample. In n -gram models, independence is only conditional and the observations are assumed to satisfy a Markovian hypothesis: the probability of occurrence of a word depends on the $n - 1$ previous words. But the i.i.d. case, although very simple, yields results that are interesting in themselves, and upon which a more sophisticated framework may be built.

Many practical questions may now be formulated in this setting. If we double the duration of a first experiment, how many yet unobserved specimens would we find (how does $K_{2n,r}$ compare to $K_{n,r}$ [resp. $K_r(2t)$ to $K_r(t)$]) [Fisher et al. \[1943\]](#)? If certain cypher keys have been observed, what is the probability for the next to be different (how does one estimate $M_{n,0}$)? For instance, [Good \[1953\]](#) and Turing observed that $(n + 1)\mathbb{E}M_{n,0} = \mathbb{E}K_{n+1,1}$ for all $n \geq 1$, and proposed to estimate the missing mass using the Jackknife estimator $G_{n,0} = K_{n,1}/n$ (the proportion of symbols seen just once).

Contributions

To study the Good-Turing estimator or other quantities that depend significantly on the small-count portion of the observations, we need to understand the occupancy counts well. Our contribution here is to give sharp concentration inequalities with explicit and reasonable constants, for K_n , $K_{n,r}$, and $M_{n,0}$ [resp. $K(t)$, $K_r(t)$, $M_0(t)$]. We give distribution-free results, and then exhibit a vast domain where these results are tight, namely the domain of distributions with a heavier tail than the geometric. In this domain, the non-asymptotic exponential concentration properties that we establish are sharp in the sense that the exponents are order-optimal, precisely capturing the scale of the variance. For this reason, we dedicate a portion of the paper to establishing bounds on various variances.

Organization

The paper is organized as follows. In Section 2, we present our terminology and give a concise summary of the results. In Section 3 we present our variance bounds and concentration results for the occupancy counts and the missing mass in great generality. In Section 4 we specialize these results to regularly varying distributions, the aforementioned domain of distributions where concentration can be characterized tightly. We then elaborate on some applications in Section 5, and conclude with a discussion of the results and possible extensions in Section 6. We group all proofs in the end, in Section 7.

2. Summary of results

Terminology

Our concentration results mostly take the form of bounds on the log-Laplace transform. Our terminology follows closely [Boucheron et al., 2013]. We say that the random variable Z is *sub-Gaussian* on the right tail (resp. on the left tail) with variance factor v if, for all $\lambda \geq 0$ (resp. $\lambda \leq 0$),

$$\log \mathbb{E} e^{\lambda(Z - \mathbb{E}Z)} \leq \frac{v\lambda^2}{2}. \quad (2.1)$$

We say that a random variable Z is *sub-Poisson* with variance factor v if, for all $\lambda \in \mathbb{R}$,

$$\log \mathbb{E} e^{\lambda(Z - \mathbb{E}Z)} \leq v\phi(\lambda), \quad (2.2)$$

with $\phi : \lambda \mapsto e^\lambda - \lambda - 1$.

We say that a random variable Z is *sub-gamma on the right tail* with variance factor v and scale parameter c if

$$\log \mathbb{E} e^{\lambda(X - \mathbb{E}X)} \leq \frac{\lambda^2 v}{2(1 - c\lambda)} \text{ for every } \lambda \text{ such that } 0 \leq \lambda \leq 1/c. \quad (2.3)$$

The random variable Z is *sub-gamma on the left tail* with variance factor v and scale parameter c , if $-Z$ is sub-gamma on the right tail with variance factor v and scale parameter c . If Z is sub-Poisson with variance factor v , then it is sub-Gaussian on the left tail with variance factor v , and sub-gamma on the right tail with variance factor v and scale parameter $1/3$.

These log-Laplace upper bounds then imply exponential tail bounds: Inequality (2.3) results in a Bernstein-type inequality for the right tail, that is, for $s > 0$ our inequalities have the form:

$$\mathbb{P}\{Z > \mathbb{E}[Z] + \sqrt{2vs} + cs\} \leq e^{-s},$$

while Inequality (2.1) for all $\lambda \leq 0$ entails

$$\mathbb{P}\{Z < \mathbb{E}[Z] - \sqrt{2vs}\} \leq e^{-s}.$$

We present such results first without making distributional assumptions, beyond the structure of those quantities themselves. These concentrations then specialize in various settings, such as that of regular variation.

Main results

We proceed by giving a coarse description of our main results. In the Poisson setting, for each $r \geq 1$, $(\mathbb{I}\{X_j(t) = r\})_{j \geq 1}$ are independent, hence $K_r(t)$ is a sum of independent Bernoulli random variables, and it is not too surprising that it satisfies sub-Poisson, also known as Bennett, inequalities. For $\lambda \in \mathbb{R}$, we have:

$$\log \mathbb{E} e^{\lambda(K_r(t) - \mathbb{E}K_r(t))} \leq \text{var}(K_r(t))\phi(\lambda) \leq \mathbb{E}[K_r(t)]\phi(\lambda).$$

The proofs are elementary and are based on the careful application of Efron-Stein-Steele inequalities and the entropy method [Boucheron et al., 2013].

As for the binomial setting, the summands are not independent but we can use negative association arguments [Dubhashi and Ranjan, 1998] (see Section 7) to obtain Bennett inequalities for the cumulated occupancy counts $K_{n,\bar{r}}$. These hold either with the Jackknife variance proxy given by the Efron-Stein inequality, $r\mathbb{E}K_{n,r}$ or with the variance proxy stemming from the negative correlation of the summands, $\mathbb{E}K_{n,\bar{r}}$. Letting $v_{n,\bar{r}} = \min(r\mathbb{E}K_{n,r}, \mathbb{E}K_{n,\bar{r}})$, we have, for all $\lambda \in \mathbb{R}$:

$$\log \mathbb{E} e^{\lambda(K_{n,\bar{r}} - \mathbb{E}K_{n,\bar{r}})} \leq v_{n,\bar{r}}\phi(\lambda).$$

This in turn implies a concentration inequality for $K_{n,r}$. Letting

$$v_{n,r} = 2 \min((r\mathbb{E}K_{n,r}) \vee ((r+1)\mathbb{E}K_{n,r+1}), \mathbb{E}K_{n,\bar{r}}),$$

we have, for all $s \geq 0$,

$$\mathbb{P}\left\{|K_{n,r} - \mathbb{E}K_{n,r}| \geq \sqrt{4v_{n,r}s} + 2s/3\right\} \leq 4e^{-s}.$$

We obtain distribution-free bounds on the log-Laplace transform of $M_{n,0}$, which result in sub-Gaussian concentration on the left tail, sub-gamma concentration on the right tail with scale proxy $1/n$. More precisely, letting $v_n^- = 2\mathbb{E}K_2(n)/n^2$ and $v_n^+ = 2\mathbb{E}K_{\bar{2}}(n)/n^2$, we show that, for all $\lambda \leq 0$,

$$\log \mathbb{E}e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \leq v_n^- \frac{\lambda^2}{2},$$

and, for all $\lambda \geq 0$,

$$\log \mathbb{E}e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \leq v_n^+ \frac{\lambda^2}{2(1 - \lambda/n)}.$$

Indeed, these results are distribution-free. But though the variance factor v_n^- is a sharp bound for the variance of the missing mass, v_n^+ may be much bigger. This leads us to look for distribution-specific conditions ensuring that v_n^+ captures the right order for the variance, such as by using a tail asymptotic stability condition as in extreme value theory.

Karlin [1967] pioneered such a condition by assuming that the function $\vec{v} : (0, 1] \rightarrow \mathbb{N}$, defined by $\vec{v}(x) = |\{j \in \mathbb{N}^*, p_j \geq x\}|$ satisfies a regular variation assumption, namely $\vec{v}(1/n) \sim n^\alpha \ell(n)$ near $+\infty$, with $\alpha \in (0, 1]$ [see also Gnedin et al., 2007, Ohannessian and Dahleh, 2012]. Here ℓ is a slowly varying function at ∞ , i.e. for all x , $\ell(\tau x)/\ell(\tau) \rightarrow 1$ as $\tau \rightarrow \infty$. This condition allows us to compare the asymptotics of the various occupancy scores. In particular, in this framework $\mathbb{E}K_2(n)$ and $\mathbb{E}K_{\bar{2}}(n)$ have the same order of growth, and, divided by n^2 they both are of the same order as the variance of the missing mass. Hence, the regular variation provides a framework in which our concentration inequalities are order-optimal.

To handle the case $\alpha = 0$, we move from *Karamata* to *de Haan* theory, and take \vec{v} to have an extended regular variation property, with the additional assumption that $\mathbb{E}K_1(n)$ tends to $+\infty$. This domain corresponds to light-tailed distributions which are still heavier than the geometric. In this case, we manage to show the sub-gamma concentration of the missing mass only for n large enough, that is, that there exists n_0 such that for all $n \geq n_0$, for $\lambda > 0$, we have $\log \mathbb{E}e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \leq (v_n \lambda^2)/2(1 - \lambda/n)$, with $v_n \asymp \text{Var}M_{n,0}$.

Back to our examples of applications, considerable insight may be gained from these concentration results. For instance, heavy tails lead to multiplicative concentration for $M_{n,0}$ and the consistency of the Good-Turing estimator: $\frac{G_{n,0}}{M_{n,0}} \xrightarrow{p} 1$. Generally, new estimators can be derived and shown to be consistent in a unified framework, once one is able to estimate α consistently. For instance, when $\vec{v}(1/\cdot)$ is regularly varying with index α , $\hat{\alpha} = K_{n,1}/K_n$ is a consistent estimator of α . Then, to estimate the number of new species in a sample twice the size of the original, we immediately get that $\hat{K}_{2n} = K_n + \frac{2^{\hat{\alpha}} - 1}{\hat{\alpha}} K_{n,1}$ is a consistent estimator of K_{2n} . This methodology is very similar to extreme value theory [Beirlant et al., 2006]: harnessing limiting expressions and tail parameter estimation. These results strengthen and extend the contribution of [Ohannessian and Dahleh, 2012], which is restricted to power-laws and implicit constants

in the inequalities. Beyond consistency results, we also obtain confidence intervals for the Good-Turing estimator in the Poisson setting, using empirical quantities.

Historical notes and related work

There exists a vast literature on the occupancy scheme, as formulated here and in many other variations. The most studied problems are the asymptotic behavior of K_n and $K_{n,r}$. This is done often in a finite context, or a scaling model where probabilities remain mostly uniform. Of particular relevance to this paper, we mention the work of Karlin [1967], who built on earlier work by Bahadur [1960], credited as one of the first to study the infinite occupancy scheme. Karlin's main results were to establish central limit theorems in an infinite setting, under a condition of regular variation. He also derived strong laws of large numbers. Gnedin et al. [2007] present a general review of these earlier results, as well as more contemporary work on this problem. The focus continues to be central limit theorems, or generally asymptotic results. For example the work of Hwang and Janson [2008] (effectively) provides a local limit theorem for K_n provided that the variance tends to infinity. Somewhat less asymptotic results have also been proposed, in the form of normal approximations, such as in the work of Barbour and Gnedin [2009].

Besides occupancy counts analysis, a distinct literature investigates the number of species and missing mass problems. These originated in the works of Fisher et al. [1943], Good [1953], and Good and Toulmin [1956], and generated a long line of research to this day [Bunge and Fitzpatrick, 1993]. Here, instead of characterizing the asymptotic behavior of these quantities, one is interested in estimating $K_{\lambda n} - K_n$ for a $\lambda > 1$, that is the number of discoveries when the sample size is multiplied by λ , or estimating $M_{n,0}$: estimators are proposed, and then their statistical properties are studied. One recently studied property by McAllester and Schapire [2000] and McAllester and Ortiz [2003], is that of concentration, which sets itself apart from the CLT-type results in that it is non-asymptotic in nature. Based on this, Ohannessian and Dahleh [2012] showed that in the regular variation setting of Karlin, one could show multiplicative concentration, and establish strong consistency results. Conversely, characterizing various aspects of concentration allows one to systematically design new estimators. For example, this was illustrated in Ohannessian and Dahleh [2012] for the estimation of rare probabilities, to both justify and extend Good's [Good, 1953] work that remains relevant in some of the aforementioned applications, especially computational linguistics.

3. Concentration

3.1. Occupancy counts

3.1.1. Variance bounds

In order to understand the fluctuations of occupancy counts K_n , $K(t)$, $K_{n,r}$, $K_r(t)$, we start by reviewing and stating variance bounds. We start with the Poisson setting where occupancy counts are sums of independent Bernoulli random variables with possibly different success probabilities, and thus variance computations are straightforward. Exact expressions may be derived [see for example [Gnedin et al., 2007](#), Equation (4)], but ingenuity may be used to derive more tractable and tight bounds. We start by stating a well-known connection between the variance of the number of occupied urns and the expected number of singletons [[Gnedin et al., 2007](#)][[Karlin, 1967](#)]. In the binomial setting, similar bounds can be derived using the Efron-Stein-Steele inequalities [see [Boucheron et al., 2013](#), Section 3.1].

Proposition 3.1. In the Poissonized setting,

$$\frac{\mathbb{E}K_1(2t)}{2} \leq \text{Var}(K(t)) \leq \mathbb{E}K_1(t).$$

In the binomial setting,

$$\text{Var}(K_n) \leq \mathbb{E}[K_{n,1}(1 - M_{n,0})] \leq \mathbb{E}K_{n,1}.$$

The upper bounds in these two propositions parallel each other, but in the binomial setting, we cannot hope to establish lower bounds like $\mathbb{E}K_{cn,1}/c \leq \text{Var}(K_n)$ for some constant $c > 0$ in full generality. To see this, consider the following example which shows that the variance of $K(t)$ and of K_n may differ significantly, and that the variance of K_n may be much smaller than the expected value of $K_{n,1}$.

Example 1. In the so-called *birthday paradox* scenario, n balls are thrown independently into n^2 urns with uniform probabilities $1/n^2$. In the Poisson setting for time $t = n$, since we have $\mathbb{E}K(n) = \sum_j (1 - e^{-np_j}) = n^2(1 - e^{-1/n})$, using an upper and lower Taylor expansion we can obtain the bounds:

$$n - \frac{1}{2} \leq \mathbb{E}K(n) \leq n - \frac{1}{2} + \frac{1}{6n}.$$

Since $\text{Var}(K(n)) = \sum_j e^{-np_j}(1 - e^{-np_j}) = \mathbb{E}K(2n) - \mathbb{E}K(n)$, it follows that:

$$n - \frac{1}{6n} \leq \text{Var}(K(n)) \leq n + \frac{1}{12n}.$$

Meanwhile, we have $\mathbb{E}[K_1(n)] = \sum_j np_j e^{-np_j} = ne^{-1/n}$, which can be bounded similarly:

$$n - 1 \leq \mathbb{E}[K_1(n)] \leq n - 1 + \frac{1}{2n}.$$

We can thus see that the Poisson birthday paradox satisfies the spirit of Proposition 3.1, if not its letter (because of being outside of our fixed- p setting). Namely, the Poisson quantities $\text{Var}(K(n))$ and $\mathbb{E}K_1(n)$ are of the same order of magnitude, roughly n .

On the other hand, in the binomial setting, since $1 - M_{n,0} = K_n/n^2$,

$$\text{Var}(K_n) \leq \mathbb{E}[K_{n,1}(1 - M_{n,0})] = \mathbb{E}\left[K_{n,1} \frac{K_n}{n^2}\right] \leq \frac{1}{n} \mathbb{E}K_{n,1} \leq 1,$$

where we have used the same variance bound as in the proof of Proposition 3.1 (Section 7.2.1). While this implies that the upper bound $\text{Var}(K_n) \leq \mathbb{E}K_{n,1}$ is satisfied, it also shows that a lower bound of the kind of $\mathbb{E}K_{cn,1}/c \leq \text{Var}(K_n)$ is not possible, since similarly to $\mathbb{E}K_1(n)$, we have $\mathbb{E}K_{n,1} = \sum_j np_j(1 - p_j)^n = n(1 - \frac{1}{n^2})^n \geq n - 1$. ■

Another straightforward bound on $\text{Var}(K_n)$ comes from the fact that the Bernoulli variables $(\mathbb{I}_{\{X_{n,j} > 0\}})_{j \geq 1}$ are negatively correlated. Thus, ignoring the covariance terms, we get

$$\text{Var}(K_n) \leq \sum_{j=1}^{\infty} \text{Var}(\mathbb{I}_{\{X_{n,j} > 0\}}) = \sum_{j=1}^{\infty} (1 - p_j)^n (1 - (1 - p_j)^n) = \mathbb{E}K_{2n} - \mathbb{E}K_n.$$

Let us denote this bound by $\text{Var}^{\text{ind}}(K_n) = \mathbb{E}K_{2n} - \mathbb{E}K_n$, as it is a variance proxy obtained by considering that the summands in K_n are independent. One can observe that the expression of $\text{Var}^{\text{ind}}(K_n)$ is very similar to the variance in the Poissonized setting, $\text{Var}(K(t)) = \mathbb{E}K(2t) - \mathbb{E}K(t)$. It is insightful to compare the true variance, the Poissonized proxy, and the negative correlation proxy, to quantify the price one pays by resorting to the latter two as an approximation for the first. We revisit this in more detail in Section 6.1.

We now investigate the fluctuations of the individual occupancy counts $K_{n,r}$ and $K_r(t)$, and that of the cumulative occupancy counts $K_{n,\bar{r}} = \sum_{j \geq r} K_{n,j}$ and $K_{\bar{r}}(t) = \sum_{j \geq r} K_j(t)$.

Proposition 3.2. In the Poisson setting, for $r \geq 1$, $t \geq 0$,

$$\text{Var}(K_{\bar{r}}(t)) \leq \min(r\mathbb{E}K_r(t), \mathbb{E}K_{\bar{r}}(t)).$$

In the binomial setting, for $r, n \geq 1$,

$$\text{Var}(K_{n,\bar{r}}) \leq \min(r\mathbb{E}K_{n,r}, \mathbb{E}K_{n,\bar{r}}).$$

For each setting, the first bound follows from Efron-Stein-Steele inequalities, the second from negative correlation. These techniques are presented briefly in Sections 7.1.1 and 7.1.2 respectively.

Remark 3.1. Except for $r = 1$, there is no clear-cut answer as to which of these two bounds is the tightest. In the regular variation scenario with index $\alpha \in]0, 1]$ as explored in [Gnedin et al., 2007], the two bounds are asymptotically of the same order, indeed,

$$\frac{r\mathbb{E}K_{n,r}}{\mathbb{E}K_{n,\bar{r}}} \underset{n \rightarrow +\infty}{\sim} \alpha,$$

see Section 4 for more on this. ■

Bounds on $\text{Var}(K_r(t))$ can be easily derived as $K_r(t)$ is a sum of independent Bernoulli random variables, however, noticing that $K_{n,r} = K_{n,\bar{r}} - K_{n,\bar{r}+1}$ and that $K_{n,\bar{r}}$ and $K_{n,\bar{r}+1}$ are positively correlated, the following bound is immediate.

Proposition 3.3. In the Poisson setting, for $r \geq 1$, $t \geq 0$,

$$\text{Var}(K_r(t)) \leq \mathbb{E}K_r(t).$$

In the binomial setting, for $r, n \geq 1$,

$$\begin{aligned} \text{Var}(K_{n,r}) &\leq \min(r\mathbb{E}K_{n,r} + (r+1)\mathbb{E}K_{n,r+1}, \mathbb{E}K_{n,\bar{r}} + \mathbb{E}K_{n,\bar{r}+1}) \\ &\leq 2 \min((r\mathbb{E}K_{n,r}) \vee ((r+1)\mathbb{E}K_{n,r+1}), \mathbb{E}K_{n,\bar{r}}). \end{aligned}$$

3.1.2. Concentration inequalities

Concentration inequalities refine variance bounds. These bounds on the logarithmic moment generating functions are indeed Bennett (sub-Poisson) inequalities with the variance upper bounds stated in the preceding section. For $K_{n,\bar{r}}$, the next proposition gives a Bernstein inequality where the variance factor is, up to a constant factor, the Efron-Stein upper bound on the variance.

Proposition 3.4. Let $r \geq 1$, and let $v_{n,\bar{r}} = \min(r\mathbb{E}K_{n,r}, \mathbb{E}K_{n,\bar{r}})$. Then, for all $\lambda \in \mathbb{R}$,

$$\log \mathbb{E}e^{\lambda(K_{n,\bar{r}} - \mathbb{E}K_{n,\bar{r}})} \leq v_{n,\bar{r}}\phi(\lambda),$$

with $\phi : \lambda \mapsto e^\lambda - \lambda - 1$.

It is worth noting that the variance bound $\mathbb{E}K_{n,\bar{r}}$ in this concentration inequality can also be obtained using a variant of Stein's method known as sized-bias coupling [Bartroff et al., 2014, Chen et al., 2010].

A critical element of the proof of Proposition 3.4 is to use the fact that each $K_{n,\bar{r}}$ is a sum of negatively associated random variables (Section 7.1.2). This is not the case for $K_{n,r}$, and thus negative association cannot be invoked directly. To deal with this, we simply use the observation of Ohannessian and Dahleh [2012] that since $K_{n,r} = K_{n,\bar{r}} - K_{n,\bar{r}+1}$, the concentration of $K_{n,r}$ follows from that of those two terms. We can show the following.

Proposition 3.5. Let

$$v_{n,r} = 2 \min((r\mathbb{E}K_{n,r}) \vee ((r+1)\mathbb{E}K_{n,r+1}), \mathbb{E}K_{n,\bar{r}}).$$

Then, for $s \geq 0$,

$$\mathbb{P}\left\{|K_{n,r} - \mathbb{E}K_{n,r}| \geq \sqrt{4v_{n,r}s} + 2s/3\right\} \leq 4e^{-s}.$$

3.2. Missing mass

3.2.1. Variance bound

Recall that $M_{n,0} = \sum_{j=1}^{\infty} p_j \mathbb{I}_{\{X_{n,j}=0\}} = \sum_{j=1}^{\infty} p_j Y_j$, and we can readily show that the summands are negatively associated weighted Bernoulli random variables (Section 7.1.2). This results in a handy upper bound for the variance of the missing mass.

Proposition 3.6. In the Poisson setting,

$$\text{Var}(M_0(t)) = 2\mathbb{E}K_2(t)/t^2 - \mathbb{E}K_2(2t)/2t^2 \leq 2\mathbb{E}K_2(t)/t^2,$$

while in the binomial setting,

$$\text{Var}(M_{n,0}) \leq \sum_{j=1}^{\infty} p_j^2 \text{Var}(Y_j) \leq \frac{2\mathbb{E}K_2(n)}{n^2}.$$

Note that whereas the expected value of the missing mass is connected to the number of singletons, its variance may be upper bounded using the number of *doubletons* (in the Poisson setting). This connection was already pointed out in [Good, 1953] and [Esty, 1982].

3.2.2. Concentration of the left tail

Moving on to the concentration properties of the missing mass, we first note that as a sum of weighted sub-Poisson random variables (following [Boucheron et al., 2013]), the missing mass is itself a sub-gamma random variable on both tails. It should not come as a surprise that the left tail of $M_{n,0}$ is sub-Gaussian with the variance factor derived from negative association. This had already been pointed out by McAllester and Schapire [2000] and McAllester and Ortiz [2003].

Proposition 3.7. [McAllester and Ortiz, 2003] In the Poisson setting, the missing mass $M_0(t)$ is sub-Gaussian on the left tail with variance factor the effective variance $\text{Var}(M_0(t)) = \sum_{j=1}^{\infty} p_j^2 e^{-tp_j} (1 - e^{-tp_j})$.

In the binomial setting, the missing mass $M_{n,0}$ is sub-Gaussian on the left tail with variance factor $v = \sum_{j=1}^{\infty} p_j^2 \text{Var}(Y_j)$ or $v_n^- = 2\mathbb{E}K_2(n)/n^2$.

For $\lambda \leq 0$,

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \frac{v\lambda^2}{2} \leq \frac{2v_n^-\lambda^2}{2}$$

3.2.3. Concentration of the right tail

The following concentration inequalities for the missing mass mostly rely on the observation that, for all $\lambda > 0$,

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \sum_{r=2}^{\infty} \left(\frac{\lambda}{n} \right)^r \mathbb{E}K_r(n). \quad (3.1)$$

This suggests that if we have a uniform control on the expected occupancy scores $(\mathbb{E}K_r(t))_{r \geq 2}$, then the missing mass has a sub-gamma right tail, with some more or less accurate variance proxy, and scale factor $1/n$.

The next theorem shows that the missing mass is sub-gamma on the right tail with variance proxy $2\mathbb{E}K_{\bar{2}}(n)/n^2$ and scale proxy $1/n$. Despite its simplicity and its generality, this bound exhibits an intuitively correct scale factor: if there exist symbols with probability of order $1/n$, they offer the major contribution to the fluctuations of the missing mass.

Theorem 3.8. *In the binomial as well as in the Poisson setting, the missing mass is sub-gamma on the right tail with variance factor $v_n^+ = 2\mathbb{E}K_{\bar{2}}(n)/n^2$ and scale factor $1/n$. For $\lambda \geq 0$,*

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \frac{v_n^+ \lambda^2}{2(1 - \lambda/n)}.$$

If the sequence $(\mathbb{E}K_r(n))_{r \geq 2}$ is non-increasing, the missing mass is sub-gamma on the right tail with variance factor $v_n^- = 2\mathbb{E}K_2(n)/n^2$ and scale factor $1/n$,

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \frac{v_n^- \lambda^2}{2(1 - \lambda/n)}.$$

Remark 3.2. McAllester and Ortiz [2003] and Berend and Kontorovich [2013] point out that for each Bernoulli random variable Y_j , for all $\lambda \in \mathbb{R}$

$$\log \mathbb{E} e^{\lambda(Y_j - \mathbb{E}Y_j)} \leq \frac{\lambda^2}{4C_{\text{LS}}(\mathbb{E}Y_j)},$$

where $C_{\text{LS}}(p) = \log(p/(1-p))/(1-2p)$ (or 2 if $p = 1/2$) is the optimal logarithmic Sobolev constant for Bernoulli random variables with success probability p (this sharp and non-trivial result has been proved independently by a number of people: Hoeffding [1963, Theorem 1, second inequality], Kearns and Saul [1998], Berend and Kontorovich [2013], Raginsky and Sason [2013]). From this observation, thanks to the negative association of the $(Y_j)_{j \geq 1}$, it follows that the missing mass is sub-Gaussian with variance factor

$$v = \sum_{j=1}^{\infty} \frac{p_j^2}{2C_{\text{LS}}((1-p_j)^n)} \leq \sum_{j=1}^{\infty} \frac{p_j^2}{2\log((1-p_j)^{-n})} \leq \sum_{j=1}^{\infty} \frac{p_j^2}{2np_j} \leq \frac{1}{2n}.$$

This variance factor is usually larger than $2\mathbb{E}K_{\bar{2}}(n)/n^2$. In the scenarios discussed in Section 4, $(2\mathbb{E}K_{\bar{2}}(n)/n^2)/(1/2n)$ even tends to 0 as n tends to infinity. ■

4. Regular variation

Motivation

Are the variance bounds in the concentration results of Section 3 tight? To see that in some pathological examples this may not be the case, consider the following example (which revisits Example 1).

Example 2. We may challenge the tail bounds offered by Proposition 3.7 and Theorem 3.8 in the simplest setting where all N symbols have equal probabilities $1/N$. Then the missing mass is $1 - K_n/N$, its variance is $\text{Var}(K_n)/N^2$. In the birthday paradox setting ($N = n^2$), $\text{Var}(M_{n,0}) \leq 1/n^4$, and the variance bound $2\mathbb{E}K_{\bar{2}}(n)/n^2$ is not tight. Indeed, one can verify that $\mathbb{E}K_{\bar{2}}(n) \geq \frac{1}{2}(1 - \frac{1}{n})$ so that $v_n^+ \geq \frac{1}{n^2} - \frac{1}{n^3}$. However, in what is called the *central domain* in [Kolchin et al., 1978], that is when $N \rightarrow \infty$ while $n/N \rightarrow t \in \mathbb{R}_+$, the tail bounds become relevant. The variance of K_n is equivalent to $Ne^{-t}(1 - e^{-t})$ while its expectation is equivalent to $N(1 - e^{-t})$. Note that in this setting all $\mathbb{E}K_r(n)$ and $\mathbb{E}K_{n,r}$ are of the same order of magnitude as $\mathbb{E}K_n$, indeed $\mathbb{E}K_r(n)/\mathbb{E}K(n) \rightarrow e^{-t}t^r/(r!(1 - e^{-t}))$.

These examples are illustrative although they do not fall in the fixed p regime we are considering in this paper. We use them because they have tractable expressions, and they provide informative diagnostics. To parallel the phenomenon of mismatched variance proxies in our setting, one can simply look at the geometric distribution for a concrete example. If $(p_k)_{k \geq 1}$ defines a geometric distribution $p_k = (1 - q)^{k-1}q$, then $\mathbb{E}K_2(n)$ remains bounded, while $\mathbb{E}K_{\bar{2}}(n)$ scales like $\log n$ as n tends to infinity. ■

In particular, we may conjecture that Theorem 3.8 is likely to be sharp when the first terms of the sequence $(\mathbb{E}K_r(n))_{r \geq 2}$ grow at the same rate as $\mathbb{E}K(n)$, or at least as $\mathbb{E}K_{\bar{2}}(n)$, which is not the case in the birthday paradox setting of Example 2. We see in what follows that the regular variation framework introduced by Karlin [1967] leads to such asymptotic equivalents. The most useful aspect of these equivalent growth rates is a simple characterization of the variance of various quantities, particularly relative to their expectation. We focus on the right tail of the missing mass, which exhibits the highest sensitivity to this asymptotic behavior, by trying to specialize Theorem 3.8 under regular variation.

Definition

To state the regular variation hypothesis, we first introduce some notation. Following [Karlin, 1967], [Gnedin et al., 2007] and [Ohannessian and Dahleh, 2012], we consider the counting measure on $[0, 1]$

$$\nu(dx) = \sum_{j=1}^{\infty} \delta_{p_j}(dx),$$

where δ_x is the Dirac mass at x .

This assumption is made on the *counting function* $\vec{\nu}$, defined for all $x > 0$ by

$$\vec{\nu}(x) = \nu[x, 1] = |\{j : j \geq 1, p_j \geq x\}|.$$

Following Karlin [1967], we say that frequencies p_j are regularly varying with index $\alpha \in [0, 1]$, if

$$\vec{\nu}(x) \underset{0+}{\sim} x^{-\alpha} \ell(1/x),$$

where ℓ is a slowly varying function, that is, for all $x > 0$, $\lim_{\tau \rightarrow +\infty} \frac{\ell(\tau x)}{\ell(\tau)} = 1$. In other words, this means that the function $\bar{\nu}(1/\cdot)$ is regularly varying with index $\alpha \in [0, 1]$, which is denoted by $\bar{\nu}(1/\cdot) \in \text{RV}_\alpha$.

In what follows, we treat three cases separately: the nominal regular variation case with $\alpha \in (0, 1)$ strictly, the *fast variation* case with $\alpha = 1$, and the *slow variation* case with $\alpha = 0$.

In the latter case, that is if frequencies p_j are regularly varying with index 0, we find that the mere regular variation hypothesis is not sufficient to obtain asymptotic formulas. For this reason, we introduce further control in the form of an *extended* regular variation hypothesis (given by Definition 1 of Section 4.3).

Remark 4.1. Before we proceed, as further motivation, we note that the regular variation hypothesis is very close to being a necessary condition for exponential concentration. For example, considering inequality (3.1), we see that if the sampling distribution is such that the ratio $\mathbb{E}K_{\bar{2}}(t)/\mathbb{E}K_2(t)$ remains bounded, then we are able to capture the right variance proxy. Now, defining the shorthand $\Phi_{\bar{2}}(t) = \mathbb{E}K_{\bar{2}}(t)$ and $\Phi_2(t) = \mathbb{E}K_2(t)$ following the notation of [Gnedin et al., 2007], we have

$$\Phi_{\bar{2}}'(t) = \frac{2\Phi_2(t)}{t}.$$

Hence, $\Phi_{\bar{2}}(t)/\Phi_2(t) = 2\Phi_{\bar{2}}(t)/t\Phi_{\bar{2}}'(t)$, and if instead of boundedness, we further require that this ratio converges to some finite limit, then, by the converse part of Karamata's Theorem [see de Haan and Ferreira, 2006, Theorem B.1.5], we find that $\Phi_{\bar{2}}$ (and then Φ_2) is regularly varying, which in turn implies that $\bar{\nu}(1/t)$ is regularly varying. ■

4.1. Case $\alpha \in (0, 1)$

We first consider the case $0 < \alpha < 1$. The next theorem states that when the sampling distribution is regularly varying with index $\alpha \in (0, 1)$, the variance factors in the Bernstein inequalities of Proposition 3.7 and Theorem 3.8 are of the same order as the variance of the missing mass.

Theorem 4.1. *Assume that the counting function $\bar{\nu}$ satisfies the regular variation condition with index $\alpha \in (0, 1)$, then the missing mass $M_{n,0}$ (or $M_0(n)$) is sub-Gaussian on the left tail with variance factor $v_n^- = 2\mathbb{E}K_2(n)/n^2$ and sub-gamma on the right tail with variance factor $v_n^+ = 2\mathbb{E}K_{\bar{2}}(n)/n^2$. The variance factors satisfy*

$$\lim_n \frac{v_n^-}{\text{Var}(M_{n,0})} = \frac{1}{1 - 2^{\alpha-2}},$$

$$\lim_n \frac{v_n^+}{\text{Var}(M_{n,0})} = \frac{2}{\alpha(1 - 2^{\alpha-2})},$$

and thus

$$\lim_n \frac{v_n^-}{v_n^+} = \frac{\alpha}{2}.$$

The second ratio deteriorates when α approaches 0, implying that the variance factor for the right tail gets worse when the distribution has a lighter tail. We do not prove Theorem 4.1 as the asymptotics are a consequence of Karlin's results surveyed in [Gnedin et al., 2007, Ohannessian and Dahleh, 2012]. In particular, we need Proposition 3.7, Theorem 3.8, and:

Theorem 4.2. [Karlin, 1967] *If the counting function \vec{v} is regularly varying with index $\alpha \in (0, 1)$, for all $r \geq 1$,*

$$\begin{aligned} - K_n &\stackrel{a.s.}{\sim}_{+\infty} \mathbb{E}K_n \sim \Gamma(1 - \alpha)n^\alpha \ell(n), \\ - K_{n,r} &\stackrel{a.s.}{\sim}_{+\infty} \mathbb{E}K_{n,r} \sim \frac{\alpha \Gamma(r - \alpha)}{r!} n^\alpha \ell(n), \\ - \text{Var}(M_{n,0}) &\sim \alpha \Gamma(2 - \alpha)(1 - 2^{\alpha-2})n^{\alpha-2} \ell(n) \end{aligned}$$

and the same hold for the corresponding Poissonized quantities.

Note that all expected occupancy counts are of the same order, and the asymptotic for $\mathbb{E}K_2(n)$ follows directly from the difference between $\mathbb{E}K(n)$ and $\mathbb{E}K_1(n)$.

4.2. Fast variation, $\alpha = 1$

We refer to the regular variation regime with $\alpha = 1$ as *fast variation*¹. From the perspective of concentration, this represents a relatively “easy” scenario. In a nutshell, this is because the variance of various quantities grows much slower than their expectation.

We first start by noting that Theorem 4.1 continues to hold as is for $\alpha = 1$. However, the rationale is slightly different. Moreover, the occupancy counts in this case present some singular behaviors which justify treating this case separately. We first note the following asymptotics that asserts that in this case, the number of occupied boxes K_n and the number of singletons $K_{n,1}$ are asymptotically equivalent, and that they dominate all the $K_{n,r}$, for $r \geq 2$. Intuitively, almost all the symbols of the sample are distinct and their probability lies around $1/n$, but a significant fraction of them has a probability smaller than $1/n$.

Theorem 4.3. [Karlin, 1967] *Assume $\vec{v}(1/x) = x^{-1}\ell(x)$ with $\ell \in \text{RV}_0$ (note that ℓ tends to 0 at ∞). Define $\ell_1 : [1, \infty) \rightarrow \mathbb{R}_+$ by*

$$\ell_1(y) = \int_y^\infty u^{-1}\ell(u)du.$$

Then $\ell_1 \in \text{RV}_0$ and $\lim_{t \rightarrow \infty} \ell_1(t)/\ell(t) = \infty$ and the following asymptotics hold:

¹This is also referred to sometimes as *rapid variation*, as in Gnedin et al. [2007], but we avoid that terminology as it also has a different connotation in the regular variation literature.

$$\begin{aligned}
& - K_n \stackrel{a.s.}{\underset{+\infty}{\sim}} \mathbb{E}K_n \underset{+\infty}{\sim} n\ell_1(n), \\
& - K_{n,1} \stackrel{a.s.}{\underset{+\infty}{\sim}} \mathbb{E}K_{n,1} \underset{+\infty}{\sim} \mathbb{E}K_n, \\
& - K_{n,r} \stackrel{a.s.}{\underset{+\infty}{\sim}} \mathbb{E}K_{n,r} \underset{+\infty}{\sim} \frac{1}{r(r-1)}n\ell(n) \quad , \quad r \geq 2,
\end{aligned}$$

and the same hold for the corresponding Poissonized quantities.

We can now see that the expected missing mass scales like $\mathbb{E}K_1(n)/n$ while its variance scales like $\mathbb{E}K_2(n)/n^2$. That said, it is not immediately clear that the variance factor $2\mathbb{E}K_{\bar{2}}(n)/n^2$ is of the same order as $\text{Var}(M_{n,0})$. As $\mathbb{E}K(n)$ and $\mathbb{E}K_1(n)$ are asymptotically equivalent, Theorem 4.3 is not sufficient to obtain the asymptotics for $\mathbb{E}K_{\bar{2}}(n)$. This is given by the next proposition, which is in fact sufficient to prove Theorem 4.1 for both $0 < \alpha < 1$ and $\alpha = 1$.

Proposition 4.4. Assume that the counting function \vec{v} satisfies the regular variation condition with index $\alpha \in (0, 1]$, then for all $r \geq 2$,

$$K_{\bar{r}}(n) \underset{+\infty}{\sim} \frac{\Gamma(r-\alpha)}{(r-1)!} \vec{v}(1/n) \quad \text{a.s.}$$

Thus, when $\alpha = 1$, the cumulated occupancy counts $\mathbb{E}K_{\bar{r}}(n)$, for $r \geq 2$, are of the order of $n\ell(n)$, and thus each is dominated by $\mathbb{E}K(n)$ and $\mathbb{E}K_1(n)$. In particular, $\mathbb{E}K_{\bar{2}}(n)$ is of order $n\ell(n)$ and our variance proxy still captures the right order for the variance of the missing mass.

Remark 4.2. Whereas in the case $0 < \alpha < 1$, $2\mathbb{E}K(n)/n^2$ would also have been a good variance proxy, here when $\alpha = 1$ we really need to remove the singletons and take $2\mathbb{E}K_{\bar{2}}(n)/n^2$ to get the right order.

We also note that when $\alpha = 1$, the missing mass is even more stable. If we let v_n denote $2\mathbb{E}K_2(n)/n^2$ or $2\mathbb{E}K_{\bar{2}}(n)/n^2$, then

$$\frac{\sqrt{v_n}}{\mathbb{E}M_{n,0}} \sim \begin{cases} c(\alpha) \cdot n^{-\alpha/2} \frac{\sqrt{\ell(n)}}{\ell(n)} & \text{for } 0 < \alpha < 1, \\ c \cdot \frac{n^{-1/2} \sqrt{\ell(n)}}{\ell_1(n)} & \text{for } \alpha = 1. \end{cases} \quad \blacksquare$$

4.3. Slow variation, $\alpha = 0$

The setting where the counting function \vec{v} satisfies the regular variation condition with index 0 represents a challenge. We refer to this regime simply as *slow variation*. Recall that this means that $\vec{v}(z/n)/\vec{v}(1/n)$ converges to 1 as n goes to infinity, yet to deal with this case we need to control the speed of this convergence, exemplified by the notion of extended regular variation that was introduced by de Haan [See Bingham et al., 1989, de Haan and Ferreira, 2006]. As we illustrate in the end of this section, one may face rather irregular behavior without such a hypothesis.

Definition 1. A measurable function $\ell : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ has the extended slow variation property, if there exists a non-negative measurable function $a : \mathbb{R}^+ \rightarrow$

\mathbb{R}^+ such that for all $x > 0$

$$\lim_{\tau \rightarrow \infty} \frac{\ell(\tau x) - \ell(\tau)}{a(\tau)} \xrightarrow{\tau \rightarrow \infty} \log(x).$$

The function $a(\cdot)$ is called an auxiliary function. When a function ℓ has the extended slow variation property with auxiliary function a , we denote it by $\ell \in \Pi_a$.

Note that the auxiliary function is always slowly varying and satisfies $\lim_{\tau \rightarrow \infty} \ell(\tau)/a(\tau) = \infty$, that is it grows slower than the original function. Furthermore, any two possible auxiliary functions are asymptotically equivalent, that is if a_1 and a_2 are both auxiliary functions for ℓ , then $\lim_{t \rightarrow \infty} a_1(t)/a_2(t) = 1$.

The notion of extended slow variation and the auxiliary function give us the aforementioned control needed to treat the $\alpha = 0$ case on the same footing as the $0 < \alpha < 1$ case. In particular, in what follows in this section we assume that $\bar{\nu}(1/\cdot) \in \Pi_a$, with the additional requirement that the auxiliary function a tends to $+\infty$.

Remark 4.3. This domain corresponds to light-tailed distributions just above the geometric distribution (the upper-exponential part of Gumbel's domain). For the geometric distribution with frequencies $p_j = (1 - q)q^{j-1}$, $j = 1, 2, \dots$, the counting function satisfies $\bar{\nu}(1/n) \sim_{\infty} \log_{1/q}(n) \in \text{RV}_0$, but the auxiliary function $a(n) = \log(1/q)$ does not tend to infinity. Frequencies of the form $p_j = cq^{\sqrt{j}}$ on the other hand do fit this framework. ■

Theorem 4.5. [*Gnedin et al., 2007*] Assume that ℓ defined by $\ell(t) = \bar{\nu}(1/t)$ is in Π_a where a is slowly varying and tends to infinity. The following asymptotics hold for each $r \geq n$

$$\begin{aligned} & - K_n \stackrel{\mathbb{P}}{\sim}_{+\infty} \mathbb{E}K_n \sim \ell(n) \\ & - K_{n,r} \stackrel{\mathbb{P}}{\sim}_{+\infty} \mathbb{E}K_{n,r} \sim \frac{a(n)}{r} \\ & - K_{n,\bar{r}} \stackrel{\mathbb{P}}{\sim}_{+\infty} \mathbb{E}K_{n,\bar{r}} \sim \ell(n) \\ & - M_{n,r} \stackrel{\mathbb{P}}{\sim}_{+\infty} \mathbb{E}M_{n,r} \sim \frac{a(n)}{n}. \end{aligned}$$

and the same equivalents hold for the corresponding Poissonized quantities.

Remark 4.4. In this case, the expectations $(\mathbb{E}K_{n,r})_{r \geq 1}$ are of the same order but are much smaller than $\mathbb{E}K_n$, and the variables K_n and $K_{n,\bar{r}}$ are all almost surely equivalent to $\ell(n)$. It is also remarkable that all the expected masses $(\mathbb{E}M_{n,r})_{r \geq 1}$ are equivalent. ■

As $\mathbb{E}K_{\bar{2}}(n) \sim \mathbb{E}K(n) \sim \ell(n)$, whereas the variance of the missing mass is of order $a(n)/n^2$, upper bounding $\mathbb{E}K_r(n)$, $r \geq 2$ by $\mathbb{E}K_{\bar{2}}(n)$, is no longer sufficient to capture the right order for the variance. However, exploiting more carefully the regular variation hypothesis, we manage to get a uniform control over the $\mathbb{E}K_r(n)$, which is valid for large enough n .

Theorem 4.6. Assume that ℓ defined by $\ell(x) = \vec{\nu}(1/x)$ is in Π_a where the slowly varying function a tends to infinity, then

1. letting $v_n = 12a(n)/n^2$,

$$v_n \asymp \text{Var}(M_{n,0}) \sim \frac{3a(n)}{4n^2}$$

2. there exists $n_0 \in \mathbb{N}$ that depends on $\vec{\nu}$ such that for all $n > n_0$, for all $\lambda > 0$,

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \frac{v_n \lambda^2}{2(1 - \lambda/n)}.$$

For all $s \geq 0$, $n \geq n_0$,

$$\mathbb{P} \left\{ M_{n,0} \geq \mathbb{E}M_{n,0} + \sqrt{2v_n s} + \frac{s}{n} \right\} \leq e^{-s}.$$

The same results hold for $M_0(t)$.

4.3.1. Too slow variation

We conclude this section by motivating why it is crucial to have a heavy-enough tail in order to obtain meaningful concentration. For example, even under regular variation when $\alpha = 0$, but $\vec{\nu}$ is not in a de Haan class Π_a with $a(n) \rightarrow \infty$, the behavior of the occupancy counts and their moments may be quite irregular. In this section, we collect some observations on those light-tailed distributions. We start with the geometric distribution which represents in many respects a borderline case.

The geometric case is an example of slow variation: $\vec{\nu}(1/\cdot) \in \text{RV}_0$. Indeed, with $p_k = (1 - q)^{k-1}q$, $0 < q < 1$, we have

$$\begin{aligned} \vec{\nu}(x) &= \sum_{k=1}^{+\infty} \mathbb{I}_{\{p_k \geq x\}} \\ &= |k \in \mathbb{N}^*, (1 - q)^{k-1}q \geq x| \\ &= 1 + \left\lfloor \frac{\log(x/q)}{\log(1 - q)} \right\rfloor, \end{aligned}$$

and thus $\vec{\nu}(x) \underset{x \rightarrow 0}{\sim} \ell(1/x)$, with ℓ slowly varying.

In this case, $\text{Var}(K(n)) = \mathbb{E}K(2n) - \mathbb{E}K(n) \rightarrow \frac{\log(2)}{\log(1/(1-q))}$.

Proposition 4.7. When the sampling distribution is geometric with parameter $q \in (0, 1)$, letting $M_n = \max(X_1, \dots, X_n)$,

$$\mathbb{E}M_n \geq \mathbb{E}K_n \geq \mathbb{E}M_n - \frac{1 - q}{q^2}.$$

In the case of geometric frequencies, the missing mass can fluctuate widely with respect to its expectation, and one cannot expect to obtain sub-gamma concentration with both the correct variance proxy and scale factor $1/n$. Indeed, intuitively, the symbol which primarily contributes to the missing mass' fluctuations, is the quantile of order $1 - 1/n$. With $F(k) = \sum_{j=1}^k p_j$, and F^\leftarrow the generalized inverse of F ,

$$\begin{aligned} j^* = F^\leftarrow(1 - 1/n) &= \inf\{j \geq 1, F(j) \geq 1 - 1/n\} \\ &= \inf\{j \geq 1, \sum_{k>j} p_k \leq 1/n\}. \end{aligned}$$

Omitting the slowly varying functions, when $\bar{\nu}(1/\cdot) \in \text{RV}_\alpha$, $0 < \alpha < 1$, j^* is of order $n^{\frac{\alpha}{1-\alpha}}$ and p_{j^*} is of order $n^{-\frac{1}{1-\alpha}}$. The closer to 1 is α , the smaller the probability of j^* . When α goes to 0, this probability becomes $1/n$. With geometric frequencies, j^* is $\frac{\log(n)}{\log(1/(1-q))}$ and p_{j^*} is $\frac{q}{n(1-q)}$. Hence, around the quantile of order $1 - 1/n$, there are symbols which may contribute significantly to the missing mass' fluctuations.

Another interesting case consists of distributions which are very light-tailed, in the sense that $\frac{p_{k+1}}{p_k} \rightarrow 0$ when $k \rightarrow \infty$. An example of these is the Poisson distribution $\mathcal{P}(\lambda)$, for which $\frac{p_{k+1}}{p_k} = \frac{\lambda}{k} \xrightarrow[k \rightarrow +\infty]{} 0$. The next proposition shows that for such concentrated distributions, the missing mass essentially concentrates on two points.

Proposition 4.8. In the infinite urns scheme with probability mass function $(p_k)_{k \in \mathbb{N}}$, if $p_k > 0$ for all k and $\lim_{k \rightarrow \infty} \frac{p_{k+1}}{p_k} = 0$, then there exists a sequence of integers $(u_n)_{n \in \mathbb{N}}$ such that

$$\lim_{n \rightarrow \infty} \mathbb{P}\{M_{n,0} \in \{\bar{F}(u_n), \bar{F}(u_n + 1)\}\} = 1,$$

where $\bar{F}(k) = \sum_{j>k} p_j$.

5. Applications

5.1. Estimating the regular variation index

When working in the regular variation setting, the most basic estimation task is to estimate the regular variation index α . We already mentioned in Section 2 the fact that, when $\bar{\nu} \in \text{RV}_\alpha$, $\alpha \in (0, 1)$, the ratio $K_{n,1}/K_n$ provides a consistent estimate of α . This is actually only one among a family of estimators of α that one may construct. The next result shows this, and is a direct consequence of Proposition 4.4.

Proposition 5.1. If $\bar{\nu} \in \text{RV}_\alpha$, $\alpha \in]0, 1]$, then for all $r \geq 1$

$$\frac{rK_{n,r}}{K_{n,\bar{r}}}$$

is a strongly consistent estimator of α .

Thus, writing $k_n = \max\{r, K_{n,r} > 0\}$, at time n , we can have up to k_n non-trivial estimators of α .

5.2. Estimating the missing mass

The Good-Turing estimation problem [Good, 1953] is that of estimating $M_{n,r}$ from the observation (X_1, X_2, \dots, X_n) . For large scores r , designing estimators for $M_{n,r}$ is straightforward, we assume that the empirical distribution mimics the sampling distribution, and that the empirical probabilities $\frac{rK_{n,r}}{n}$ are likely to be good estimators. The question is more delicate for rare events. In particular, for $r = 0$, it may be a bad idea to assume that there is no missing mass $M_{n,0} = 0$, that is to assign a zero probability to the symbols that do not appear in the sample. Various “smoothing” techniques were developed, in order to adjust the maximum likelihood estimator and obtain more accurate probabilities.

Good-Turing estimators attempt to estimate $(M_{n,r})_r$ from $(K_{n,r})_r$ for all r . They are defined as

$$G_{n,r} = \frac{(r+1)K_{n,r+1}}{n}.$$

The rationale for this choice comes from the following observations.

$$\mathbb{E}G_{n,0} = \frac{\mathbb{E}[K_{n,1}]}{n} = \mathbb{E}M_{n-1,0} = \mathbb{E}M_{n,0} + \frac{\mathbb{E}M_{n,1}}{n}, \quad (5.1)$$

and

$$\mathbb{E}G_{n,r} = \frac{(r+1)\mathbb{E}K_{n,r+1}}{n} = \mathbb{E}M_{n-1,r}. \quad (5.2)$$

In the Poisson setting, there is no bias: $\mathbb{E}G_r(t) = (r+1)\frac{\mathbb{E}K_{r+1}(t)}{t} = \mathbb{E}M_r(t)$.

Here, we primarily focus on the estimation of the missing masses $M_{n,0}$ and $M_0(t)$, though most of the methodology extends also to $r > 0$, with the appropriate concentration results. From (5.1) and (5.2), Good-Turing estimators look like slightly biased estimators of the relevant masses. In particular, the bias $\mathbb{E}G_{n,0} - \mathbb{E}M_{n,0}$ is always positive but smaller than $1/n$. It is however far from obvious to determine scenarios where these estimators are consistent and where meaningful confidence regions can be constructed.

When trying to estimate the missing mass $M_{n,0}$ or $\mathbb{E}M_{n,0}$, consistency needs to be redefined since the estimand is not a fixed parameter of interest but a random quantity whose expectation further depends on n . Additive consistency, that is bounds on $\widehat{M}_{n,0} - M_{n,0}$ is not a satisfactory notion, because, as $M_{n,0}$ tends to 0, the trivial constant estimator 0 would be universally asymptotically consistent. Relative consistency, that is control on $(\widehat{M}_{n,0} - M_{n,0})/M_{n,0}$ looks like a much more reasonable notion. It is however much harder to establish.

In order to establish relative consistency of a missing mass estimator, we have to check that $\mathbb{E}[\widehat{M}_{n,0} - M_{n,0}]$ is not too large with respect to $\mathbb{E}M_{n,0}$, and that both $\widehat{M}_{n,0}$ and $M_{n,0}$ are concentrated around their mean values.

As shown in [Ohannessian and Dahleh, 2012], the Good-Turing estimator of the missing mass is not universally consistent in this sense. This occurs principally in very light tails, such as those described in Section 4.3.1.

Proposition 5.2. [Ohanessian and Dahleh, 2012] When the sampling distribution is geometric with small enough $q \in (0, 1)$, there exists $\eta > 0$, and a subsequence n_i such that for i large enough, $G_{n_i,0}/M_{n_i,0} = 0$ with probability no less than η .

On the other hand, the concentration result of Corollary 4.1 entails a law of large numbers for $M_{n,0}$ (by a direct application of the Borel-Cantelli lemma), which in turns entails the strong multiplicative consistency of the Good-Turing estimate.

Corollary 5.3. We have the following two regimes of consistency for the Good-Turing estimator of the missing mass.

- (i) If the counting function $\vec{\nu}$ is such that $\mathbb{E}K_{n,2}/\mathbb{E}K_{n,1}$ remains bounded and $\mathbb{E}K_{n,1} \rightarrow +\infty$ (in particular, when $\vec{\nu}$ is regularly varying with index $\alpha \in (0, 1]$ or $\alpha = 0$ and $\vec{\nu} \in \Pi_a$ with $a \rightarrow \infty$),

$$\frac{M_{n,0}}{\mathbb{E}M_{n,0}} \xrightarrow{\mathbb{P}} 1,$$

and the Good-Turing estimator of $M_{n,0}$ defined by $G_{n,0} = K_{n,1}/n$, is multiplicatively consistent in probability:

$$\frac{G_{n,0}}{M_{n,0}} \xrightarrow{\mathbb{P}} 1.$$

- (ii) If furthermore $\mathbb{E}K_{n,\bar{2}}/\mathbb{E}K_{n,1}$ remains bounded and if, for all $c > 0$, $\sum_{n=0}^{\infty} \exp(-c\mathbb{E}K_{n,1}) < \infty$ (in particular, when $\vec{\nu}$ is regularly varying with index $\alpha \in (0, 1]$), then these two convergences occur almost surely.

Consistency is a desirable property, but the concentration inequalities provide us with more power, in particular in terms of giving confidence intervals that are asymptotically tight. For brevity, we focus here on the Poisson setting to derive concentration inequalities which in turn yield confidence intervals. A similar, but somewhat more tedious, methodology yields confidence intervals in the binomial setting as well.

5.2.1. Concentration inequalities for $G_0(t) - M_0(t)$

In the Poisson setting, the analysis of the Good-Turing estimator is illuminating. As noted in earlier, the first pleasant observation is that the Good-Turing estimator is an unbiased estimator of the missing mass. Second, the variance of $G_0(t) - M_0(t)$ is simply related to occupancy counts:

$$\text{Var}(G_0(t) - M_0(t)) = \frac{1}{t^2} (\mathbb{E}K_1(t) + 2\mathbb{E}K_2(t)). \quad (5.3)$$

Third, simple yet often tight concentration inequalities can be obtained for $G_0(t) - M_0(t)$.

Proposition 5.4. The random variable $G_0(t) - M_0(t)$ is sub-gamma on the right tail with variance factor $\text{Var}(G_0(t) - M_0(t))$ and scale factor $1/t$, and sub-gamma on the left tail with variance factor $3\mathbb{E}K(t)/t^2$ and scale factor $1/t$.

For all $\lambda \geq 0$,

- (i) $\log \mathbb{E}e^{\lambda(G_0(t) - M_0(t))} \leq \text{Var}(G_0(t) - M_0(t))t^2\phi\left(\frac{\lambda}{t}\right)$, and
- (ii) $\log \mathbb{E}e^{\lambda(M_0(t) - G_0(t))} \leq \frac{3\mathbb{E}K(t)}{2t^2} \frac{\lambda^2}{1 - \lambda/t}$.

We are now in a position to build confidence intervals for the missing mass.

Proposition 5.5. With probability larger than $1 - 4\delta$, the following hold

$$M_0(t) \leq G_0(t) + \frac{1}{t} \left(\sqrt{6K(t) \log \frac{1}{\delta}} + 5 \log \frac{1}{\delta} \right).$$

$$M_0(t) \geq G_0(t) - \frac{1}{t} \left(\sqrt{2(K_1(t) + 2K_2(t)) \log \frac{1}{\delta}} + 4 \log \frac{1}{\delta} \right).$$

To see that these confidence bounds are asymptotically tight, consider the following central limit theorem. A similar results can be paralleled in the binomial setting.

Proposition 5.6. If the counting function \vec{v} is regularly varying with index $\alpha \in (0, 1]$, the following central limit theorem holds for the ratio $G_0(t)/M_0(t)$:

$$\frac{\mathbb{E}K_1(t)}{\sqrt{\mathbb{E}K_1(t) + 2\mathbb{E}K_2(t)}} \left(\frac{G_0(t)}{M_0(t)} - 1 \right) \rightsquigarrow \mathcal{N}(0, 1).$$

Remark 5.1. Note that when $\alpha = 1$, this convergence occurs faster: the speed is of order $\sqrt{n\ell_1(n)}$ instead of $\sqrt{n^\alpha\ell(n)}$. ■

5.3. Estimating the number of species

Fisher's number of species problem [Fisher et al. \[1943\]](#) consists of estimating $K_{(1+\tau)n} - K_n$ for $\tau > 0$, the number of distinct new species one would observe if the data collection runs for an additional fraction τ of time. This was posed primarily within the Poisson model in the original paper [Fisher et al. \[1943\]](#) and later by [\[Efron and Thisted, 1976\]](#), but the same question may also be asked in the binomial model. The following estimates come from straightforward computations on the asymptotics given in Theorems 4.2, 4.3 and 4.5.

Proposition 5.7. If the counting function \vec{v} is regularly varying with index $\alpha \in (0, 1]$, letting $\hat{\alpha}$ be any of the estimates $rK_{n,r}/K_{n,\bar{r}}$ of α from Proposition 5.1, then any of the following quantities

$$(\tau^{\hat{\alpha}} - 1)K_n, \quad \frac{\tau^{\hat{\alpha}} - 1}{\hat{\alpha}} K_{n,1}, \quad \text{and} \quad \left(\prod_{k=2}^r \frac{k}{k - 1 - \hat{\alpha}} \right) \frac{\tau^{\hat{\alpha}} - 1}{\hat{\alpha}} K_{n,r}, \quad r \geq 2,$$

is a strongly consistent estimate of $K_{\tau n} - K_n$, the number of newly discovered species when the sample size is multiplied by τ .

If the counting function $\vec{\nu}$ is in Π_a , with $a(n) \rightarrow +\infty$, then, for each $r \geq 1$,

$$\log(\tau)rK_{n,r}$$

is an estimate of $K_{\tau n} - K_n$, consistent in probability.

6. Discussion

To conclude the paper, we review our results in a larger context, and propose some connections, extensions, and open problems.

6.1. The cost of Poissonization and negative correlation

Resorting to Poissonization or negative correlation may have a price. It may lead to variance overestimates. [Gnedin et al., 2007, Lemma 1] asserts that for some constant c

$$|\text{Var}(K(n)) - \text{Var}(K_n)| \leq \frac{c}{n} \max(1, \mathbb{E}K_1(n)^2)$$

This bound conveys a mixed message. As $\mathbb{E}K_1(n)/n$ tends to 0, it asserts that

$$|\text{Var}(K(n)) - \text{Var}(K_n)| / \mathbb{E}K_1(n)$$

tends to 0. But there exist scenarios where $\mathbb{E}K_1(n)^2/n$ tends to infinity. It is shown in [Gnedin et al., 2007] that $\mathbb{E}K_1(n)^2/(n\text{Var}(K(n)))$ tends to 0, so that, as soon as $n\text{Var}(K(n))$ tends to infinity (which might not always be the case), the two variances $\text{Var}(K_n)$ and $\text{Var}(K(n))$ are asymptotically equivalent.

It would be interesting to find the necessary and sufficient conditions under which there is equivalence. Though these aren't generally known, it is instructive to compare $\text{Var}(K(n))$, $\text{Var}(K_n)$ and $\text{Var}^{\text{ind}}(K_n)$ the variance upper bound obtained from negative correlation by bounding their differences. For instance, one can show that for any sampling distribution we have:

$$\frac{\mathbb{E}K_2(2n)}{n} \leq \text{Var}(K(n)) - \text{Var}^{\text{ind}}(K_n) \leq \frac{2\mathbb{E}K_2(n)}{n},$$

and

$$0 \leq \text{Var}^{\text{ind}}(K_n) - \text{Var}(K_n) \leq \frac{(\mathbb{E}K_{n,1})^2}{n} - \frac{\mathbb{E}K_{2n,2}}{2n-1}.$$

These bounds are insightful but, without any further assumptions on the sampling distribution, they are not sufficient to prove asymptotic equivalence.

6.2. Extensions of regular variation

The regular variation hypothesis is an elegant framework, allowing one to derive, thanks to Karamata and Tauberian Theorems, simple and comprehensible equivalents for various moments. As we have seen, it comes very close to being a necessary condition for exponential concentration. It may however seem too stringent. Without getting too specific about it, let us mention that other less demanding hypotheses also yield the asymptotic relative orders that work in favor of the concentration of the missing mass. For instance, referring back to Remark 4.1, one could ask instead that:

$$0 < \liminf_{t \rightarrow \infty} \frac{\Phi_2(t)}{\Phi_2(t)} \leq \limsup_{t \rightarrow \infty} \frac{\Phi_2(t)}{\Phi_2(t)} < \infty.$$

Recalling that $\Phi_2'(t) = \frac{2\Phi_2(t)}{t}$, and applying Corollary 2.6.2. of Bingham et al. [1989], one obtains that Φ_2 is in the class OR of O -regularly varying functions and Φ_2 is in the class ER of extended regularly varying functions, that is, for all $\lambda \geq 1$

$$0 < \liminf_{t \rightarrow \infty} \frac{\Phi_2(\lambda t)}{\Phi_2(t)} \leq \limsup_{t \rightarrow \infty} \frac{\Phi_2(\lambda t)}{\Phi_2(t)} < \infty,$$

and

$$\lambda^d \leq \liminf_{t \rightarrow \infty} \frac{\Phi_2(\lambda t)}{\Phi_2(t)} \leq \limsup_{t \rightarrow \infty} \frac{\Phi_2(\lambda t)}{\Phi_2(t)} \leq \lambda^c,$$

for some constants c and d . Observe that this result, which is the equivalent of Karamata's Theorem, differs from the regular variation setting, in the sense that the control on the derivative Φ_2 is looser than the one on Φ_2 , whereas, in Karamata Theorem, both the function and its derivative inherit the regular variation property.

We can in turn show that $\Phi(t) = \mathbb{E}K(t)$ is in the class OR and, by Theorem 2.10.2 of Bingham et al. [1989], this is equivalent to $\bar{\nu}(1/\cdot) \in OR$, as Φ is the Laplace-Stieltjes transform of $\bar{\nu}$.

6.3. Random measures

As noted by [Gnedin et al., 2007], the asymptotics for the moments of the occupancy counts in the regular variation setting is still valid when the frequencies $(p_j)_{j \geq 1}$ are random, in which case the measure ν is defined by

$$\mathbb{E} \left[\sum_{j=1}^{\infty} f(p_j) \right] = \int_0^1 f(x) \nu(dx),$$

for all functions $f \geq 0$. We can also define the measure ν_1 by

$$\mathbb{E} \left[\sum_{j=1}^{\infty} p_j f(p_j) \right] = \int_0^1 f(x) \nu_1(dx),$$

for all functions $f \geq 0$. This measure corresponds to the distribution of the frequency of the first discovered symbol.

For instance, when $(p_j)_{j \geq 1}$ are Poisson-Dirichlet($\alpha, 0$) with $0 < \alpha < 1$, the measure $\nu_1(dx)$ is the size-biased distribution of PD($\alpha, 0$), that is Beta $\mathcal{B}(1 - \alpha, \alpha)$ [see Pitman and Yor, 1997]. Thus we have:

$$\begin{aligned} \nu_1[0, x] &= \frac{1}{\mathcal{B}(1 - \alpha, \alpha)} \int_0^x t^{-\alpha} (1 - t)^{\alpha-1} dt \\ &\underset{x \rightarrow 0}{\sim} \frac{x^{1-\alpha}}{(1 - \alpha)\mathcal{B}(1 - \alpha, \alpha)} \end{aligned}$$

and, by [Proposition 13 Gneden et al., 2007], this is equivalent to

$$\vec{\nu}(x) \underset{x \rightarrow 0}{\sim} \frac{1}{\alpha \mathcal{B}(1 - \alpha, \alpha)} x^{-\alpha}.$$

Thus, denoting by $N(x)$ the random number of frequencies p_j which are larger than x , the expectation $\vec{\nu}(x) = \mathbb{E}N(x)$ is regularly varying. One can also show that the mass-partition mechanism of the distribution PD($\alpha, 0$) almost surely generates $N(x)$ to be regularly varying. To see this, refer to [Proposition 10 Pitman and Yor, 1997] or to [Proposition 2.6 Bertoin, 2006] which assert that the limit

$$L := \lim_{n \rightarrow \infty} np_n^\alpha$$

exists almost surely. This is equivalent to

$$N(x) \underset{x \rightarrow 0}{\sim} x^{-\alpha} L \quad \text{a.s.}$$

The PD($\alpha, 0$) distribution can be generated through a Poisson process with intensity measure $\nu([x, \infty]) = cx^{-\alpha}$. Without entering into further detail, let us mention that similar almost sure results hold even when the intensity measure ν is not a strict power, but satisfies the property

$$\nu([x, \infty]) \underset{x \rightarrow 0}{\sim} x^{-\alpha} \ell(x),$$

with ℓ slowly varying, [Section 6 Gneden, 2010]. Working with a regular variation hypothesis thus gives us more flexibility than assuming specific Bayesian priors.

7. Proofs

7.1. Fundamental techniques

7.1.1. Efron-Stein-Steele inequalities

Our variance bounds mostly follow from the Efron-Stein-Steele Inequality [Efron and Stein, 1981], which states that when a random variable is expressed as a function of many independent random variables, its variance can be controlled by the sum of the local fluctuations.

Theorem 7.1. *Let \mathcal{X} be some set, (X_1, X_2, \dots, X_n) be independent random variables taking values in \mathcal{X} , $f : \mathcal{X}^n \rightarrow \mathbb{R}$ be a measurable function of n variables, and $Z = f(X_1, X_2, \dots, X_n)$.*

For all $i \in \{1, \dots, n\}$, let $X^{(i)} = (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$ and $\mathbb{E}^{(i)}Z = \mathbb{E}[Z|X^{(i)}]$. Then, letting $v = \sum_{i=1}^n \mathbb{E}[(Z - \mathbb{E}^{(i)}Z)^2]$,

$$\text{Var}[Z] \leq v.$$

If X'_1, \dots, X'_n are independent copies of X_1, \dots, X_n , then letting $Z'_i = f(X_1, \dots, X_{i-1}, X'_i, X_{i+1}, \dots, X_n)$,

$$v = \sum_{i=1}^n \mathbb{E}[(Z - Z'_i)_+^2] \leq \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2],$$

where the random variables Z_i are arbitrary $X^{(i)}$ -measurable and square-integrable random variables.

7.1.2. Negative association

The random variables K_n , $K_{n,r}$, and $M_{n,r}$ are sums or weighted sums of Bernoulli random variables. These summands depends on the scores $(X_{n,j})_{j \geq 1}$ and therefore are not independent. Transforming the fixed- n binomial setting into a continuous time Poisson setting is one way to circumvent this problem. This is the Poissonization method. In this setting, the score variables $(X_j(n))_{j \geq 1}$ are independent Poisson variables with respective means np_j . Results valid for the Poisson setting can then be transferred to the fixed- n setting, up to approximation costs. For instance, [Gnedin et al., 2007] (Lemma 1) provide bounds on the discrepancy between expectations and variances in the two settings. (See also our discussion in Section 6.1).

Another approach to deal with the dependence is to invoke the notion of negative association, which provides a systematic comparison between moments of certain monotonic functions of the occupancy scores. In the bins and balls setting, this approach is mostly used to obtain bounds on the logarithmic moment generating function, and thus exponential concentration for the occupancy counts [see Dubhashi and Ranjan, 1998, Shao, 2000, McAllester and Schapire, 2000, Ohannessian and Dahleh, 2012]. We use this notion throughout the proofs, and therefore present it here formally.

Definition 2 (NEGATIVE ASSOCIATION). Real-valued random variables Z_1, \dots, Z_K are said to be negatively associated if, for any two disjoint subsets A and B of $\{1, \dots, K\}$, and any two real-valued functions $f : \mathbb{R}^{|A|} \mapsto \mathbb{R}$ and $g : \mathbb{R}^{|B|} \mapsto \mathbb{R}$ that are both either coordinate-wise non-increasing or coordinate-wise non-decreasing, we have:

$$\mathbb{E}[f(Z_A) \cdot g(Z_B)] \leq \mathbb{E}[f(Z_A)] \cdot \mathbb{E}[g(Z_B)].$$

In particular, as far as concentration properties are concerned, sums of negatively associated variables can only do better than sums of independent variables.

Theorem 7.2. [*Dubhashi and Ranjan, 1998*] For each $n \in \mathbb{N}$, the occupancy scores $(X_{n,j})_{j \geq 1}$ are negatively associated.

As any monotonic functions of negatively associated variables are also negatively associated, the variables $(\mathbb{I}_{\{X_{n,j} > 0\}})_{j \geq 1}$ (respectively $(\mathbb{I}_{\{X_{n,j} = 0\}})_{j \geq 1}$) are negatively associated as increasing (respectively decreasing) functions of $(X_{n,j})_{j \geq 1}$. This is of pivotal importance for our proofs of concentration results for K_n and $M_{n,0}$. For $r \geq 1$, the variables $(\mathbb{I}_{\{X_{n,j} = r\}})_{j \geq 1}$ appearing in $K_{n,r}$ are not negatively associated. However, following [*Ohannessian and Dahleh, 2012*], one way to deal with this problem is to observe that

$$K_{n,r} = K_{n,\bar{r}} - K_{n,\overline{r+1}},$$

recalling that $K_{n,\bar{r}} = \sum_{j=1}^{\infty} \mathbb{I}_{\{X_{n,j} \geq r\}}$ is the number of urns that contain at least r balls and that the Bernoulli variables appearing in $K_{n,\bar{r}}$ are negatively associated.

7.1.3. Potter's inequalities and its variants

One useful result from regular variation theory is provided by Potter's inequality [See *Bingham et al., 1989, de Haan and Ferreira, 2006*, for proofs and refinements].

Theorem 7.3. (POTTER-DREES INEQUALITIES.)

- (i) If $f \in \text{RV}_\gamma$, then for all $\delta > 0$, there exists $t_0 = t_0(\delta)$, such that for all $t, x: \min(t, tx) > t_0$,

$$(1 - \delta)x^\gamma \min(x^\delta, x^{-\delta}) \leq \frac{f(tx)}{f(t)} \leq (1 + \delta)x^\gamma \max(x^\delta, x^{-\delta})$$

- (ii) If $\ell \in \Pi_a$, then for all δ_1, δ_2 , there exists t_0 such that for all $t \geq t_0$, for all $x \geq 1$,

$$(1 - \delta_2) \frac{1 - x^{\delta_1}}{\delta_1} - \delta_2 < \frac{\ell(tx) - \ell(t)}{a(t)} < (1 + \delta_2) \frac{x^{\delta_1} - 1}{\delta_1} + \delta_2.$$

7.2. Occupancy counts

7.2.1. Variance bounds for occupancy counts

Proof of Proposition 3.1. Recall that in the Poisson setting,

$$\frac{d\mathbb{E}K(t)}{dt} = \frac{\mathbb{E}K_1(t)}{t} = \mathbb{E}M_0(t).$$

This entails

$$\begin{aligned}\mathrm{Var}(K(t)) &= \sum_{j=1}^{\infty} e^{-tp_j} (1 - e^{-tp_j}) \\ &= \mathbb{E}K(2t) - \mathbb{E}K(t) \\ &= \int_t^{2t} \mathbb{E}M_0(s) ds.\end{aligned}$$

Now, as $\mathbb{E}M_0(s)$ is non-increasing,

$$\frac{\mathbb{E}K_1(2t)}{2} = t\mathbb{E}M_0(2t) \leq \mathrm{Var}(K(t)) \leq t\mathbb{E}M_0(t) = \mathbb{E}K_1(t).$$

Moving on to the binomial setting, let K_n^i denote the number of occupied urns when the i^{th} ball is replaced by an independent copy. Then

$$\mathrm{Var}(K_n) \leq \mathbb{E} \left[\sum_{i=1}^n (K_n - K_n^i)_+^2 \right],$$

where $(K_n - K_n^i)_+$ denotes the positive part. Now, $K_n - K_n^i$ is positive if and only if ball i is moved from a singleton into a non-empty urn. Thus $\mathrm{Var}(K_n) \leq \mathbb{E}[K_{n,1}(1 - M_{n,0})]$. \square

Proof of Proposition 3.2. The bound $r\mathbb{E}K_{n,r}$ follows from the Efron-Stein inequality: denoting by $K_{n,\bar{r}}^{(i)}$ the number of cells with occupancy score larger than r when ball i is removed, then

$$K_{n,\bar{r}} - K_{n,\bar{r}}^{(i)} = \begin{cases} 1 & \text{if ball } i \text{ is in a } r\text{-ton} \\ 0 & \text{otherwise.} \end{cases}$$

And thus, we get $\sum_{i=1}^n (K_{n,\bar{r}} - K_{n,\bar{r}}^{(i)})^2 = rK_{n,r}$.

The second bound follows from the negative association of the variables $(\mathbb{I}_{\{X_{n,j} \geq r\}})_j$ (negative correlation is actually sufficient):

$$\mathrm{Var} \left(\sum_{j=1}^{\infty} \mathbb{I}_{\{X_{n,j} \geq r\}} \right) \leq \sum_{j=1}^{\infty} \mathrm{Var}(\mathbb{I}_{\{X_{n,j} \geq r\}}) \leq \mathbb{E}K_{n,\bar{r}}. \quad \square$$

7.2.2. Concentration inequalities for occupancy counts

Proof of Proposition 3.4. Let $X_{n,j}$ denote the occupancy score of cell $j, j \in \mathbb{N}$, then

$$K_{n,\bar{r}} = \sum_{j=1}^{\infty} \mathbb{I}_{\{X_{n,j} \geq r\}}.$$

As observed in [Ohanessian and Dahleh, 2012], $K_{n,\bar{r}}$ is a sum of negatively associated Bernoulli random variables. Moreover, the Efron-Stein inequality implies that for each $j \in \mathbb{N}$,

$$\text{Var}(\mathbb{I}_{\{X_{n,j} \geq r\}}) \leq r \mathbb{E} \mathbb{I}_{\{X_{n,j} = r\}}.$$

Thus we have

$$\begin{aligned} \log \mathbb{E} e^{\lambda(K_{n,\bar{r}} - \mathbb{E} K_{n,\bar{r}})} &\leq \sum_{j=1}^{\infty} \log \mathbb{E} e^{\lambda(\mathbb{I}_{\{X_{n,j} \geq r\}} - \mathbb{E} \mathbb{I}_{\{X_{n,j} \geq r\}})} \\ &\leq \sum_{j=1}^{\infty} \text{Var}(\mathbb{I}_{\{X_{n,j} \geq r\}}) \phi(\lambda) \\ &\leq \phi(\lambda) \sum_{j=1}^{\infty} r \mathbb{E} \mathbb{I}_{\{X_{n,j} = r\}} \\ &= \phi(\lambda) r \mathbb{E} K_{n,r}, \end{aligned}$$

where the first inequality comes from negative association, the second inequality is Bennett's inequality for Bernoulli random variables, and the last inequality comes from the Efron-Stein inequality. The other bound comes from the fact that $\text{Var}(\mathbb{I}_{\{X_{n,j} \geq r\}}) \leq \mathbb{E} \mathbb{I}_{\{X_{n,j} \geq r\}}$. \square

Proof of Proposition 3.5. As $K_{n,r} = K_{n,\bar{r}} - K_{n,\overline{r+1}}$,

$$\begin{aligned} \{K_{n,r} \geq \mathbb{E} K_{n,r} + x\} &\subseteq \left\{ K_{n,\bar{r}} \geq \mathbb{E} K_{n,\bar{r}} + \frac{x}{2} \right\} \cup \left\{ K_{n,\overline{r+1}} \leq \mathbb{E} K_{n,\overline{r+1}} - \frac{x}{2} \right\}. \end{aligned}$$

By Proposition 3.4, Bernstein inequalities hold for both $K_{n,\bar{r}}$ and $K_{n,\overline{r+1}}$, with variance proxies $\mathbb{E} r K_{n,r}$ (or $\mathbb{E} K_{n,\bar{r}}$) and $(r+1)K_{n,r+1}$ (or $\mathbb{E} K_{n,\overline{r+1}} \leq \mathbb{E} K_{n,\bar{r}}$) respectively. Hence,

$$\begin{aligned} \mathbb{P} \{K_{n,r} \geq \mathbb{E} K_{n,r} + x\} &\leq \exp \left(-\frac{x^2/4}{2(r \mathbb{E} K_{n,r} + x/6)} \right) + \exp \left(-\frac{x^2/4}{2((r+1) \mathbb{E} K_{n,r+1})} \right) \\ &\leq 2 \exp \left(-\frac{x^2/4}{2((r \mathbb{E} K_{n,r}) \vee ((r+1) K_{n,r+1}) + x/6)} \right). \end{aligned}$$

The same reasoning works for the alternative variance proxies and for the left tails. \square

7.3. Missing mass

7.3.1. Variance bounds for the missing mass

Proof of Proposition 3.6. In the Poisson setting,

$$\text{Var}(M_0(t)) = \sum_{j=1}^{\infty} p_j^2 e^{-tp_j} (1 - e^{-tp_j}) \leq \sum_{j=1}^{\infty} p_j^2 e^{-tp_j} = \frac{2}{t^2} \mathbb{E} K_2(t).$$

In the binomial setting, by negative correlation,

$$\text{Var}(M_{n,0}) \leq \sum_{j=1}^{\infty} p_j^2 (1 - (1 - p_j)^n) (1 - p_j)^n \leq \sum_{j=1}^{\infty} p_j^2 e^{-np_j} = \frac{2}{n^2} \mathbb{E}K_2(n).$$

□

7.3.2. Concentration inequalities for the missing mass

Proof of Proposition 3.7. For all $\lambda \in \mathbb{R}$,

$$\begin{aligned} \log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] &= \log \mathbb{E} \left[e^{\lambda \sum_{j=1}^{\infty} p_j (Y_j - \mathbb{E}Y_j)} \right] \\ &\leq \sum_{j=1}^{\infty} \log \mathbb{E} \left[e^{\lambda p_j (Y_j - \mathbb{E}Y_j)} \right] \\ &\leq \sum_{j=1}^{\infty} (1 - p_j)^n (1 - (1 - p_j)^n) \phi(\lambda p_j), \end{aligned}$$

where the first inequality comes from negative association, and the second is Bennett's inequality for Bernoulli random variables.

Noting that $\lim_{\lambda \rightarrow 0^-} \phi(\lambda)/\lambda^2 = \lim_{\lambda \rightarrow 0^+} \phi(\lambda)/\lambda^2 = 1/2$, the function $\lambda \mapsto \phi(\lambda)/\lambda^2$ has a continuous increasing extension on \mathbb{R} . Hence, for $\lambda \leq 0$, we have $\phi(\lambda) \leq \lambda^2/2$.

Thus, for $\lambda < 0$,

$$\begin{aligned} \log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] &\leq \sum_{j=1}^{\infty} p_j^2 (1 - p_j)^n (1 - (1 - p_j)^n) \frac{\lambda^2}{2} \\ &= \sum_{j=1}^{\infty} p_j^2 \text{Var}[Y_j] \frac{\lambda^2}{2}. \end{aligned}$$

Recall that $\sum_{j=1}^{\infty} p_j^2 \text{Var}[Y_j] \leq 2\mathbb{E}K_2(n)/n^2$ (Proposition 3.6). □

Proof of Theorem 3.8. From the beginning of the proof of Proposition 3.7, that is, thanks to negative association and to the fact that each Bernoulli random variable satisfies a Bennett inequality,

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \sum_{j=1}^{\infty} e^{-np_j} \phi(\lambda p_j).$$

Now, using the power series expansion of ϕ ,

$$\begin{aligned} \sum_{j=1}^{\infty} e^{-np_j} \phi(\lambda p_j) &= \sum_{j=1}^{\infty} e^{-np_j} \sum_{r=2}^{\infty} \frac{(\lambda p_j)^r}{r!} \\ &= \sum_{r=2}^{\infty} \left(\frac{\lambda}{n} \right)^r \sum_{j=1}^{\infty} e^{-np_j} \frac{(np_j)^r}{r!}. \end{aligned}$$

We recognize that for each r , $\sum_{j=1}^{\infty} e^{-np_j} \frac{(np_j)^r}{r!} = \mathbb{E}K_r(n)$ and that for each $r \geq 2$, $\mathbb{E}K_r(n) \leq \mathbb{E}K_2(n)$,

$$\begin{aligned} \sum_{j=1}^{\infty} e^{-np_j} \phi(\lambda p_j) &= \sum_{r=2}^{\infty} \left(\frac{\lambda}{n}\right)^r \mathbb{E}K_r(n) \\ &\leq \mathbb{E}K_2(n) \sum_{r=2}^{\infty} \left(\frac{\lambda}{n}\right)^r \\ &= \lambda^2 \frac{\mathbb{E}K_2(n)/n^2}{1 - \lambda/n}, \end{aligned}$$

which concludes the proof. \square

7.4. Regular variation

Proof of Proposition 4.4. By monotonicity of $K_{n,\bar{r}}$, we have the following strong law for any sampling distribution

$$K_{n,\bar{r}} = \sum_{s=r}^{\infty} K_{n,s} \underset{+\infty}{\sim} \sum_{s=r}^{\infty} \mathbb{E}K_s(n) \quad \text{a.s.},$$

[see [Gnedin et al., 2007](#), the discussion after Proposition 2]. Recall that $X_j(n) \sim \mathcal{P}(np_j)$ and that, if $Y \sim \mathcal{P}(\lambda)$, then $\mathbb{P}[Y \leq k] = \frac{\Gamma(k+1, \lambda)}{k!}$, where $\Gamma(z, x) = \int_x^{+\infty} e^{-t} t^{z-1} dt$ is the incomplete Gamma function. Hence

$$\begin{aligned} \sum_{s=r}^{\infty} \mathbb{E}K_s(n) &= \sum_{j=1}^{\infty} \mathbb{P}[X_j(n) \geq r] \\ &= \sum_{j=1}^{\infty} \frac{1}{(r-1)!} \int_0^{np_j} e^{-t} t^{r-1} dt \\ &= \frac{1}{(r-1)!} \int_0^1 \int_0^{nx} e^{-t} t^{r-1} dt \cdot \nu(dx) \\ &= \frac{1}{(r-1)!} \int_0^1 ne^{-nx} (nx)^{r-1} \vec{\nu}(x) dx \\ &= \frac{1}{(r-1)!} \int_0^{+\infty} e^{-z} z^{r-1} \vec{\nu}(z/n) dz \\ &\underset{+\infty}{\sim} \frac{\vec{\nu}(1/n)}{(r-1)!} \Gamma(r - \alpha). \end{aligned}$$

In particular,

$$K_{n,\bar{r}} \underset{+\infty}{\sim} \frac{rK_{n,r}}{\alpha} \quad \text{a.s.} \quad \square$$

Proof of Theorem 4.6. Recall the beginning of the proof of Theorem 3.8:

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \sum_{r=2}^{\infty} \left(\frac{\lambda}{n} \right)^r \mathbb{E}K_r(n).$$

Now, bounding each $\mathbb{E}K_r(n)$ by $\mathbb{E}K_{\bar{2}}(n)$ is not sufficient to get the right order for the variance: $\mathbb{E}K_{\bar{2}}(n)$ is of order $\ell(n)$ whereas $\text{Var}M_{n,0}$ is of order $a(n)/n^2$.

We explore more carefully the structure of $\mathbb{E}K_r(n)$ and show that these quantities are uniformly (in r) bounded by a function of order $a(n)$ for large enough n , that is, that there exists $n^* \in \mathbb{N}$ and $C \in \mathbb{R}_+$ such that for all $n \geq n^*$, for all $r \geq 1$, $\mathbb{E}K_r(n) \leq Ca(n)$.

Before going into the proof, we observe that for $r \geq n/a(n)$, the result is true. Indeed, from the identity $\sum_{r=1}^{\infty} r\mathbb{E}K_r(n) = n$, we deduce that $r\mathbb{E}K_{\bar{r}}(n) \leq n$, so that for $r \geq n/a(n)$, $\mathbb{E}K_{\bar{r}}(n) \leq a(n)$. Thus we assume that $r \leq n/a(n)$.

First, we easily deal with the contribution to $\mathbb{E}K_r(n)$ of the symbols with probability less than $1/n$. Indeed

$$I_1^r := \int_0^{1/n} e^{-nx} \frac{(nx)^r}{r!} \nu(dx) \leq \int_0^{1/n} e^{-nx} \frac{(nx)^2}{2!} \nu(dx) \leq \mathbb{E}K_2(n).$$

As $\mathbb{E}K_2(n) \sim a(n)/2$, for all δ_0 , there exists n_0 such that for all $n \geq n_0$, for all $r \geq 1$, $I_1^r \leq (1 + \delta_0)/2$.

For the the contribution of the symbols with probability larger than $1/n$, integration by part and change of variable yield:

$$\begin{aligned} I_2^r &:= \int_{1/n}^1 e^{-nx} \frac{(nx)^r}{r!} \nu(dx) \\ &= \left[e^{-nx} \frac{(nx)^r}{r!} (-\vec{\nu}(x)) \right]_{1/n}^1 + \int_{1/n}^1 e^{-nx} \frac{n^r}{r!} (rx^{r-1} - nx^r) \vec{\nu}(x) dx \\ &= \frac{\vec{\nu}(1/n)e^{-1}}{r!} + \int_1^{\infty} e^{-z} \left(\frac{z^{r-1}}{(r-1)!} - \frac{z^r}{r!} \right) \vec{\nu}(z/n) dz. \end{aligned}$$

As $\int_1^{\infty} e^{-z} \left(\frac{z^{r-1}}{(r-1)!} - \frac{z^r}{r!} \right) dz = -\mathbb{P}[\mathcal{P}(1) = r] = -e^{-1}/r!$, we can rearrange the previous expression:

$$I_2^r = a(n) \int_1^{\infty} e^{-z} \left(\frac{z^{r-1}}{(r-1)!} - \frac{z^r}{r!} \right) \frac{\vec{\nu}(z/n) - \vec{\nu}(1/n)}{a(n)} dz.$$

Notice that when $z \in [1, r]$, the integrand is negative, so we simply ignore this part of the integral and restrict ourselves to

$$I_3^r := \int_r^{\infty} e^{-z} \left(\frac{z^r}{r!} - \frac{z^{r-1}}{(r-1)!} \right) \frac{\vec{\nu}(1/n) - \vec{\nu}(z/n)}{a(n)} dz,$$

which we try to bound by a constant term for n greater than some integer that does not depend on r .

The main ingredient of our proof is the next version of the Potter-Drees Inequality (see Theorem 7.3 in Section 7.1.3 and [de Haan and Ferreira, 2006, point 4 of Corollary B.2.15]): for $\ell \in \Pi_a$, for arbitrary δ_1, δ_2 , there exists t_0 such that for all $t \geq t_0$, and for all $x \leq 1$ with $tx \geq t_0$,

$$(1 - \delta_2) \frac{1 - x^{-\delta_1}}{\delta_1} - \delta_2 < \frac{\ell(t) - \ell(tx)}{a(t)} < (1 + \delta_2) \frac{x^{-\delta_1} - 1}{\delta_1} + \delta_2.$$

Thus, for arbitrary δ_1, δ_2 , there exists n_1 such that, for all $n \geq n_1$, for all $z \in [1, n/n_1]$,

$$\frac{\bar{\nu}(1/n) - \bar{\nu}(z/n)}{a(n)} \leq (1 + \delta_2) \frac{z^{\delta_1} - 1}{\delta_1} + \delta_2.$$

As $r \leq n/a(n)$, taking, if necessary, n large enough so that $a(n) \geq n_1$, we have $r \leq n/n_1$ and

$$\begin{aligned} I_3^r &\leq \int_r^{n/n_1} e^{-z} \left(\frac{z^r}{r!} - \frac{z^{r-1}}{(r-1)!} \right) \left((1 + \delta_2) \frac{z^{\delta_1} - 1}{\delta_1} + \delta_2 \right) dz \\ &\quad + \int_{n/n_1}^{\infty} e^{-z} \left(\frac{z^r}{r!} - \frac{z^{r-1}}{(r-1)!} \right) \frac{\bar{\nu}(1/n) - \bar{\nu}(z/n)}{a(n)} dz \\ &=: I_4^r + I_5^r, \end{aligned}$$

with

$$\begin{aligned} I_4^r &\leq \delta_2 + \frac{1 + \delta_2}{\delta_1} \int_r^{\infty} e^{-z} \left(\frac{z^{r+\delta_1}}{r!} - \frac{z^r}{r!} + \frac{z^{r-1}}{(r-1)!} - \frac{z^{r-1+\delta_1}}{(r-1)!} \right) dz \\ &\leq \delta_2 + \frac{1 + \delta_2}{\delta_1} \int_r^{\infty} e^{-z} \left(\frac{z^{r+\delta_1}}{r!} - \frac{z^{r-1+\delta_1}}{(r-1)!} \right) dz \\ &= \delta_2 + \frac{1 + \delta_2}{\delta_1} \left(\frac{\Gamma(r+1+\delta_1, r)}{\Gamma(r+1)} - \frac{\Gamma(r+\delta_1, r)}{\Gamma(r)} \right), \end{aligned}$$

where $\Gamma(a, x) = \int_x^{\infty} e^{-t} t^{a-1} dt$ is the incomplete Gamma function. Using the fact that $\Gamma(a, x) = (a-1)\Gamma(a-1, x) + x^{a-1}e^{-x}$, we have

$$\begin{aligned} I_4^r &\leq \delta_2 + \frac{1 + \delta_2}{\delta_1 \Gamma(r+1)} (\Gamma(r+1+\delta_1, r) - (r+\delta_1)\Gamma(r+\delta_1, r) + \delta_1 \Gamma(r+\delta_1, r)) \\ &= \delta_2 + \frac{1 + \delta_2}{\delta_1} \left(\frac{r^{r+\delta_1} e^{-r}}{r!} + \delta_1 \frac{\Gamma(r+\delta_1, r)}{\Gamma(r+1)} \right). \end{aligned}$$

By Stirling's inequality, for all r ,

$$\frac{r^{r+\delta_1} e^{-r}}{r!} \leq r^{\delta_1} (2\pi r)^{-1/2}.$$

Thus, taking $\delta_1 = 1/4$, the right-hand term is uniformly bounded by 1. And $\frac{\Gamma(r+\delta_1, r)}{\Gamma(r+1)}$ is also bounded by 1. Thus

$$I_4^r \leq \delta_2 + \frac{1 + \delta_2}{\delta_1} (1 + \delta_1),$$

and

$$\begin{aligned}
I_5^r &\leq \frac{\vec{\nu}(1/n)}{a(n)} \int_{n/n_1}^{\infty} e^{-z} \left(\frac{z^r}{r!} - \frac{z^{r-1}}{(r-1)!} \right) dz \\
&= \frac{\vec{\nu}(1/n)}{a(n)} e^{-n/n_1} \frac{(n/n_1)^r}{r!} \\
&\leq \frac{\vec{\nu}(1/n)}{a(n)} e^{-n/n_1} \frac{(n/n_1)^{\lfloor n/n_1 \rfloor}}{\lfloor n/n_1 \rfloor!}.
\end{aligned}$$

By Stirling's inequality, this bound is smaller than $\frac{\vec{\nu}(1/n)}{a(n)} (2\pi(n/n_1))^{-1/2}$, which tends to 0 as $n \rightarrow \infty$. Thus there exists n_2 such that for all $n \geq n_2$, and all $r \leq n/n_1$, $I_5^r \leq \delta_2$.

In the end, we get that for all $\delta_0 \geq 0$, δ_1 with $0 \leq \delta_1 \leq 1/4$, and $\delta_2 \geq 0$, there exists $n^* = \max(n_0, n_1, n_2)$ such that for all $n \geq n^*$, for all $r \geq 1$,

$$\mathbb{E}K_r(n) \leq a(n) \left(\frac{1 + \delta_0}{2} + \delta_2 + \frac{1 + \delta_2}{\delta_1} (1 + \delta_1) + \delta_2 \right).$$

Taking for instance $\delta_1 = 1/4$ and $\delta_0 = \delta_2 = 1/15$, we have that for large enough n and for all $r \geq 1$,

$$\mathbb{E}K_r(n) \leq 6a(n),$$

and

$$\log \mathbb{E} \left[e^{\lambda(M_{n,0} - \mathbb{E}M_{n,0})} \right] \leq \frac{12a(n)}{n^2} \cdot \frac{\lambda^2}{2(1 - \lambda/n)}.$$

□

Proof of Proposition 4.8. Under the condition of the Proposition 4.8, from [Grübel and Hitczenko, 2009], with probability tending to 1, the sample is gap-free, hence the missing mass is $\bar{F}(\max(X_1, \dots, X_n))$.

The condition of the Proposition implies the condition described in [Anderson, 1970], i.e. $\lim_{n \rightarrow +\infty} \frac{\bar{F}(n+1)}{\bar{F}(n)} = 0$, to ensure the existence of a sequence of integers $(u_n)_{n \in \mathbb{N}}$ such that

$$\lim_{n \rightarrow \infty} \mathbb{P} \{ \max(X_1, \dots, X_n) \in \{u_n, u_n + 1\} \} = 1. \quad \square$$

7.5. Applications

Proof of Corollary 5.3. Let us assume that $\mathbb{E}K_{n,1} \rightarrow \infty$. Using the fact that $0 \leq \mathbb{E}G_{n,0} - \mathbb{E}M_{n,0} \leq 1/n$, we notice that as soon as $\mathbb{E}K_{n,1} \rightarrow \infty$, $\mathbb{E}G_{n,0} \underset{+\infty}{\sim} \mathbb{E}M_{n,0}$. Now by Chebyshev Inequality,

$$\begin{aligned}
\mathbb{P} \left[\left| \frac{M_{n,0}}{\mathbb{E}M_{n,0}} - 1 \right| > \epsilon \right] &\leq \frac{\text{Var}(M_{n,0})}{\epsilon^2 (\mathbb{E}M_{n,0})^2} \leq \frac{2\mathbb{E}K_2(n)}{\epsilon^2 n^2 (\mathbb{E}M_{n,0})^2} \\
&\sim \frac{2(\mathbb{E}K_{n,2} + o(1))}{\epsilon^2 (\mathbb{E}K_{n,1})^2},
\end{aligned}$$

where we used that $|\mathbb{E}K_2(n) - \mathbb{E}K_{n,2}| \rightarrow 0$ [see Lemma 1 [Gnedin et al., 2007](#)]. On the other hand,

$$\mathbb{P} \left[\left| \frac{K_{n,1}}{\mathbb{E}K_{n,1}} - 1 \right| > \epsilon \right] \leq \frac{\text{Var}(K_{n,1})}{\epsilon^2 (\mathbb{E}K_{n,1})^2} \leq \frac{\mathbb{E}K_{n,1} + 2\mathbb{E}K_{n,2}}{\epsilon^2 (\mathbb{E}K_{n,1})^2},$$

showing that if, furthermore, $\mathbb{E}K_{n,2}/\mathbb{E}K_{n,1}$ remains bounded, the ratios $M_{n,0}/\mathbb{E}M_{n,0}$, $G_{n,0}/\mathbb{E}G_{n,0}$ and thus $M_{n,0}/G_{n,0}$ converge to 1 in probability. To get almost sure convergence, we use Theorem 3.8 to get that when $\mathbb{E}K_{n,1} \rightarrow \infty$,

$$\begin{aligned} \mathbb{P} \left[\left| \frac{M_{n,0}}{\mathbb{E}M_{n,0}} - 1 \right| > \epsilon \right] &\leq 2 \exp \left(-\frac{\epsilon^2 (\mathbb{E}M_{n,0})^2}{2(2\mathbb{E}K_{n,2}(n)/n^2 + \mathbb{E}M_{n,0}/n)} \right) \\ &= 2 \exp \left(-\frac{\epsilon^2 (\mathbb{E}K_{n,1} + o(\mathbb{E}K_{n,1}))^2}{2(2\mathbb{E}K_{n,2} + \mathbb{E}K_{n,1} + o(\mathbb{E}K_{n,1}))} \right). \end{aligned}$$

If $\mathbb{E}K_{n,2}/\mathbb{E}K_{n,1}$ remains bounded, this becomes smaller than $c_1 \exp(-c_2 \epsilon^2 \mathbb{E}K_{n,1})$. Hence, if $\exp(-c \mathbb{E}K_{n,1})$ is summable for all $c > 0$, we can apply the Borel-Cantelli lemma and obtain the almost sure convergence of $M_{n,0}/\mathbb{E}M_{n,0}$ to 1. Moreover, by Proposition 3.5,

$$\mathbb{P} \left[\left| \frac{K_{n,1}}{\mathbb{E}K_{n,1}} - 1 \right| > \epsilon \right] \leq 4 \exp \left(-\frac{\epsilon^2 (\mathbb{E}K_{n,1})^2}{2(4(\mathbb{E}K_{n,1} \vee 2\mathbb{E}K_{n,2}) + 2/3)} \right),$$

which shows that under these assumptions $K_{n,1}/\mathbb{E}K_{n,1}$ also tends to 1 almost surely. \square

Proof of Proposition 5.4. The random variable $G_0(t) - M_0(t)$ is a sum of independent, centered and bounded random variables, namely

$$G_0(t) - M_0(t) = \frac{1}{t} \sum_{j=1}^{\infty} \mathbb{I}_{X_j(t)=1} - tp_j \mathbb{I}_{X_j(t)=0}.$$

Bound (i) follows immediately from the observation that each $\mathbb{I}_{X_j(t)=1} - tp_j \mathbb{I}_{X_j(t)=0}$ satisfies a Bennett inequality,

$$\begin{aligned} \log \mathbb{E} e^{\lambda(G_0(t) - M_0(t))} &\leq \sum_{j=1}^{\infty} \text{Var}(\mathbb{I}_{X_j(t)=1} - tp_j \mathbb{I}_{X_j(t)=0}) \phi \left(\frac{\lambda}{t} \right) \\ &= \text{Var}(G_0(t) - M_0(t)) t^2 \phi \left(\frac{\lambda}{t} \right). \end{aligned}$$

Bound (ii) follows from the observation that each $\mathbb{I}_{X_j(t)=0} - \frac{1}{tp_j} \mathbb{I}_{X_j(t)=1}$ sat-

ifies a Bennett inequality,

$$\begin{aligned}
\log \mathbb{E} e^{\lambda(M_0(t) - G_0(t))} &\leq \sum_{j=1}^{\infty} \text{Var} \left(\mathbb{I}_{X_j(t)=0} - \frac{1}{tp_j} \mathbb{I}_{X_j(t)=1} \right) \phi(\lambda p_j) \\
&= \sum_{j=1}^{\infty} \left(1 + \frac{1}{tp_j} \right) e^{-tp_j} \phi(\lambda p_j) \\
&= \sum_{r \geq 2} \left(\frac{\lambda}{t} \right)^r \sum_{j=1}^{\infty} \left(1 + \frac{1}{tp_j} \right) e^{-tp_j} \frac{(tp_j)^r}{r!} \\
&= \sum_{r \geq 2} \left(\frac{\lambda}{t} \right)^r \left(\mathbb{E} K_r(t) + \frac{1}{r} \mathbb{E} K_{r-1}(t) \right) \\
&\leq \sum_{r \geq 2} \left(\frac{\lambda}{t} \right)^r \frac{3 \mathbb{E} K(t)}{2},
\end{aligned}$$

which concludes the proof. \square

Proof of Proposition 5.5. With probability greater than $1 - 2\delta$, by Proposition 5.4,

$$G_0(t) - M_0(t) \leq \frac{1}{t} \sqrt{2(\mathbb{E} K_1(t) + 2\mathbb{E} K_2(t)) \log \frac{1}{\delta}} + \frac{\log \frac{1}{\delta}}{3t}$$

and

$$G_0(t) - M_0(t) \geq -\frac{1}{t} \sqrt{6\mathbb{E} K(t) \log \frac{1}{\delta}} - \frac{\log \frac{1}{\delta}}{t}.$$

We may now invoke concentration inequalities for $K_1(t) + 2K_2(t)$ and $K(t)$. Indeed, with probability greater than $1 - \delta$, $K(t) \geq \mathbb{E} K(t) - \sqrt{2\mathbb{E} K(t) \log \frac{1}{\delta}}$ which entails $\sqrt{\mathbb{E} K(t)} \leq \sqrt{K(t) + \frac{\log \frac{1}{\delta}}{2}} + \sqrt{\frac{\log \frac{1}{\delta}}{2}}$.

We have $2K_2(t) + K_1(t) \geq 2\mathbb{E} K_2(t) + \mathbb{E} K_1(t) - \sqrt{4(2\mathbb{E} K_2(t) + \mathbb{E} K_1(t)) \log \frac{1}{\delta}}$ with probability greater than $1 - \delta$, which entails

$$\sqrt{2\mathbb{E} K_2(t) + \mathbb{E} K_1(t)} \leq \sqrt{(2K_2(t) + K_1(t)) + \log \frac{1}{\delta}} + \sqrt{\log \frac{1}{\delta}},$$

which concludes the proof. \square

Proof of Proposition 5.6. The covariance matrix $\text{Cov}(t)$ of $(G_0(t), M_0(t))$ can be written in terms of the expected occupancy counts as

$$\text{Cov}(t) = \frac{1}{t^2} \begin{pmatrix} \mathbb{E} K_1(t) & 0 \\ 0 & 2\mathbb{E} K_2(t) \end{pmatrix} - \frac{\mathbb{E} K_2(2t)}{2t^2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

From [Karlin, 1967], we have

$$\text{Cov}(t)^{-1/2} \begin{pmatrix} G_0(t) - \mathbb{E} G_0(t) \\ M_0(t) - \mathbb{E} M_0(t) \end{pmatrix} \rightsquigarrow \mathcal{N}(0, I_2),$$

where I_2 is the identity matrix, which can be rewritten as

$$\Sigma(t)^{-1/2} \begin{pmatrix} \frac{G_0(t)}{\mathbb{E}G_0(t)} - 1 \\ \frac{M_0(t)}{\mathbb{E}M_0(t)} - 1 \end{pmatrix} \rightsquigarrow \mathcal{N}(0, I_2),$$

with $\Sigma(t) = (\mathbb{E}G_0(t))^{-2} \text{Cov}(t)$.

The delta method applied to the function $(x_1, x_2) \mapsto x_1/x_2$ yields

$$\left((1-1)\Sigma(t) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right)^{-1/2} \begin{pmatrix} \frac{G_0(t)}{M_0(t)} - 1 \end{pmatrix} \rightsquigarrow \mathcal{N}(0, 1),$$

and

$$\left((1-1)\Sigma(t) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right)^{-1/2} = \frac{\mathbb{E}K_1(t)}{\sqrt{\mathbb{E}K_1(t) + 2\mathbb{E}K_2(t)}},$$

which concludes the proof. \square

Remark 7.1. The proof for the binomial setting is very similar, the only difficulty being that $\mathbb{E}G_{n,0}$ and $\mathbb{E}M_{n,0}$ are no longer equal. However, the bias becomes negligible with respect to the fluctuations, that is, for $v_n = \sqrt{n^\alpha \ell(n)}$ or $\sqrt{n \ell_1(n)}$

$$v_n \left(\frac{\mathbb{E}G_{n,0}}{\mathbb{E}M_{n,0}} - 1 \right) \xrightarrow[n \rightarrow \infty]{} 0. \quad \blacksquare$$

References

- C. Anderson. Extreme value theory for a class of discrete distributions with applications to some stochastic processes. *J. Appl. Probability*, 7:99–113, 1970.
- R. Bahadur. On the number of distinct values in a large sample from an infinite discrete distribution. *Proc. Nat. Inst. Sci. India Part A*, 26A(Supp. II):67–75, 1960.
- A. D. Barbour and A. V. Gneden. Small counts in the infinite occupancy scheme. *Electron. J. Probab.*, 14:no. 13, 365–384, 2009.
- J. Bartroff, L. Goldstein, and Ü. Işlak. Bounded size biased couplings, log concave distributions and concentration of measure for occupancy models. *arXiv preprint arXiv:1402.6769*, 2014.
- T. Batu, L. Fortnow, E. Fischer, R. Kumar, R. Rubinfeld, and P. White. Testing Random Variables for Independence and Identity. In *Proceedings of FOCS*, pages 442–451, 2001.
- J. Beirlant, Y. Goegebeur, J. Segers, and J. Teugels. *Statistics of extremes: theory and applications*. John Wiley & Sons, 2006.
- D. Berend and A. Kontorovich. On the concentration of the missing mass. *Electron. Commun. Probab.*, 18:no. 3, 7, 2013.
- J. Bertoin. *Random fragmentation and coagulation processes*. Cambridge University Press Cambridge, 2006.
- N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27. Cambridge university press, 1989.

- L. Bogachev, A. Gnedin, and Y. Yakubovich. On the variance of the number of occupied boxes. *Adv. in Appl. Math.*, 40(4):401–432, 2008.
- S. Boucheron, G. Lugosi, and P. Massart. *Concentration inequalities*. Oxford University Press, 2013.
- J. Bunge and M. Fitzpatrick. Estimating the number of species: a review. *Journal of the American Statistical Association*, 88(421):364–373, 1993.
- L. H. Y. Chen, L. Goldstein, and Q.-M. Shao. *Normal approximation by Stein’s method*. Springer, 2010.
- L. de Haan and A. Ferreira. *Extreme value theory*. Springer-Verlag, 2006.
- D. Dubhashi and D. Ranjan. Balls and bins: A study in negative dependence. *Random Struct. & Algorithms*, 13(2):99–124, 1998.
- B. Efron and C. Stein. The jackknife estimate of variance. *Annals of Statistics*, 9(3):586–596, 1981.
- B. Efron and R. Tibshirani. Estimating the number of unseen species: How many words did Shakespeare know? *Biometrika*, 63(3):435–447, 1976.
- W. Ewens. Confidence intervals for the coverage of low coverage samples. *Ann. Statist.*, 10(1):190–196, 1982.
- R. A. Fisher, A. S. Corbet, and C. B. Williams. The relation between the number of species and the number of individuals in a random sample of an animal population. *Journal of Animal Ecology*, 12(1):42–58, May 1943.
- A. Gnedin, B. Hansen, J. Pitman, et al. Notes on the occupancy problem with infinitely many boxes: general asymptotics and power laws. *Probab. Surv.*, 4: 146–171, 2007.
- A. V. Gnedin. Regeneration in random combinatorial structures. *Probab. Surv.*, 7:105–156, 2010.
- I. J. Good. The population frequencies of species and the estimation of population parameters. *Biometrika*, 40:16–264, 1953.
- I. J. Good and G. H. Toulmin. The number of new species, and the increase in population coverage, when a sample is increased. *Biometrika*, 43(1/2):45–63, June 1956.
- R. Grübel and P. Hitczenko. Gaps in discrete random samples. *J. Appl. Probab.*, 46(4):1038–1051, 2009.
- W. Hoeffding. Probability inequalities for sums of bounded random variables. *J. Amer. Statist. Assoc.*, 58:13–30, 1963.
- H.-K. Hwang and S. Janson. Local limit theorems for finite and infinite urn models. *Ann. Probab.*, 36(3):992–1022, 2008.
- S. Karlin. Central limit theorems for certain infinite urn schemes. *J. Math. Mech.*, 17:373–401, 1967.
- M. Kearns and L. Saul. Large deviation methods for approximate probabilistic inference. In *Proceedings of the Fourteenth Conference on Uncertainty in Artificial Intelligence*, pages 311–319. Morgan Kaufmann Publishers Inc., 1998.
- V. Kolchin, B. Sevastyanov, and V. Chistyakov. *Random allocations*. V. H. Winston & Sons, Washington, D.C.; distributed by Halsted Press [John Wiley & Sons], New York-Toronto, Ont.-London, 1978. Translated from the Russian, Translation edited by A. V. Balakrishnan, Scripta Series in Mathematics.

- D. McAllester and L. Ortiz. Concentration inequalities for the missing mass and for histogram rule error. *Journal of Machine Learning Research*, 4:895–911, 2003.
- D. A. McAllester and R. E. Schapire. On the convergence rate of Good-Turing estimators. In *COLT 2000*, pages 1–6, 2000.
- M. I. Ohannessian and M. A. Dahleh. Rare probability estimation under regularly varying heavy tails. In *COLT 2012, Journal of Machine Learning Research-Proceedings Track*, volume 23, pages 21.1–21.24, 2012.
- A. Orlitsky, N. Santhanam, and J. Zhang. Universal compression of memoryless sources over unknown alphabets. *IEEE Trans. Inform. Theory*, 50(7):1469–1481, 2004. ISSN 0018-9448.
- J. Pitman and M. Yor. The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator. *Ann. Probab.*, pages 855–900, 1997.
- M. Raginsky and I. Sason. *Concentration of Measure Inequalities in Information Theory, Communications and Coding*, volume 10. NOW Publishers, 2013.
- Q.-M. Shao. A comparison theorem on moment inequalities between negatively associated and independent random variables. *Journal of Theoretical Probability*, 13(2):343–356, 2000.
- G. Valiant and P. Valiant. Estimating the unseen: an $N/\log(N)$ -sample estimator for entropy and support size, shown optimal via new CLTs. In *STOC '11*, pages 685–694, 2011.