

1 Supplemental Table S1:Values of water potentials for *Fagus sylvatica*, at predawn (ψ_{pd}) and at midday (ψ_{md}), reported in the literature; the most negative
2 value of both ψ_{pd} and ψ_{md} found in each study is given. Leaf positions within the crown or other particular conditions of measurements are also indicated. Tree
3 age and height, when available are given, as well as the date of measurements, stand characteristics and country (FR: France, GE: Germany, GR: Greece, IP:
4 Iberian Peninsula, PO: Poland, SW: Switzerland, UK: United kingdom).

5

Ψ_{pd} (MPa)	Ψ_{md} (MPa)	Leaf position within the crown or experimental condition	Age in years (height)	Date	Stand description / Country	Reference
-0.12 -0.11	-2.4 -1.3	Sun leaves Shade leaves	5 and 55	1999	Kranzberg Forest / GE	Wieser et al. (2003)
-0.2 -0.2	-2.8 -2.0	Sun leaves Shade leaves	29	1999	Same stand as in the present study / FR	Lemoine et al. (2002b)
-0.71	-2.19	Lower sunny crown	85-105	1990-1993	Beech-oak forest in Lower Saxony / GE	Backes and Leuschner (2000)
-0.8	-2.4	Upper canopy leaves	30 (10-12 m)	1993-1994	Beech-oak forest in the Sistema Central / IP	Aranda et al. (2000)
-0.81	-2.2	Leaves in uppermost canopy Upper third of the crown	(15 m)	2000	Beech-oak forest in the Sistema Central / IP	Aranda et al. (2005)
-1.15	-2.03	Lower sunny crown	90-110	1990-1993 and 1995	Beech-oak forest in Lower Saxony / GE	Leuschner et al. (2001)
-1.4 -1.3	-1.9 -2.0	Ambient ozone twice-ambient ozone regime	60 (>28 m)	2003	Mixed beech-spruce forest in the Kranzberg Forest / GE	Löw et al. (2006)
-1.44 -1.66	-1.77 -1.76	Ambient CO ₂ Elevated CO ₂	80-100 (30-35 m)	2003	Mixed forest / SW	Leuzinger et al. (2005)

-1.79	-2.6	Sunny crown	Maturetrees(31 m)	2006	Mixed broad-leaved forest in Thuringia / GE	Köcher et al (2009)
-1.99	-2.76	Lower canopy leaves	15 (7.7 m)	2003	Broad-leaved mixed stand in the state forest of Hesse /FR	Peiffer (2005)
-2.28 -2.04	-2.88 -2.24	Upper canopy leaves Lower canopy leaves	37 (17 m)	2003	90%-pure beech stand in the state forest of Hesse / FR	Present study Peiffer et al. (2014)
	-1.75		56 (21 m)	1989	Predominantly beech forest / UK	Roberts and Rosier (1994)
	-1.94		30 - 60 (17 m)	2003-2005	Pure beech stand, Paiko mountain / GR	Fotelli et al. (2009)
-2.37 -1.39	Full sunlight leaves Shade leaves	70	1993 and1997	Arboretum of Amance / FR		Cochard et al. (1999)
-4.0	South oriented middle crown	(9 m)	1998	Mixed open woodland, Mount Verticos / GR		Raftoyannis and Radoglou (2002)
-2.5		adult	2003	Arboretum of Amance / FR		Bréda et al. (2004)

6

7

8

9

10 REFERENCES

- 11 Aranda I, Gil L, Pardos J A (2000) Water relations and gas exchange in *Fagus sylvatica*L. and *Quercus petraea*(Mattuschka) Liebl. in a mixed stand at their
12 southern limit of distribution in Europe. Trees 14: 344-352.
- 13 Aranda I, Gil L, Pardos J A (2005) Seasonal changes in apparent hydraulic conductance and their implications for water use of European beech (*Fagus*
14 *sylvatica*L.) and sessile oak [*Quercus petraea*(Matt.) Liebl] in South Europe. Plant Ecology 179: 155-167.
- 15 Backes K, Leuschner C (2000) Leaf water relations of competitive *Fagus sylvatica*and *Quercus petraea*trees during 4 years differing in soil drought. Can J
16 For Res 30: 335-346.
- 17 Bréda N, Granier A, Aussénac G (2004) La sécheresse de 2003 dans le contexte climatique des 54 dernières années: analyse écophysiolgue et influence sur
18 les arbres forestiers. Rev For FrLVI : 109-131.
- 19 Cochard H, Lemoine D, Dreyer E (1999) The effects of acclimatation to sunlight on the xylem vulnerability to embolism in *Fagus sylvatica* L. Plant Cell
20 Environ 22: 101-108.
- 21 Fotelli MN. Nahm M, Radoglou K, Rennenberg H, Halyvopoulos G, Matzarakis A (2009) Seasonal and interannualecophysiological responses of beech
22 (*Fagus sylvatica*) at its south-eastern distribution limit in Europe. For Ecol Manage 257: 1157-1164.
- 23 Köcher P, Gebauer T, Horna V, Leuschner C (2009) Leaf water status and stem xylem flux in relation to soil drought in five temperate broad-leaved tree
24 species with contrasting water use strategies. Ann ForSci66: 101.<http://hal.archives-ouvertes.fr/hal-00883452>
- 25 Lemoine D, Jacquemin S, Granier A (2002b) Beech (*Fagus sylvatica*L.) branches show acclimation of xylem anatomy and hydraulic properties to increased
26 light after thinning. Ann ForSci59: 761-766.<http://hal.archives-ouvertes.fr/hal-00883666>

- 27 Leuschner Ch, Backes K, Hertel D, Schipka F, Schmitt U, Terborg O, Runge M (2001) Drought responses at leaf, stem and fine root levels of competitive
28 *Fagus sylvatica*L. and *Quercus petraea* (Matt.) Liebl. Trees in dry and wet years. *For Ecol Manage* 140: 33-46.
- 29 Leuzinger S, Zotz G, Asshoff R, Körner Ch (2005) Responses of deciduous forest trees to severe drought in central Europe. *TreePhysiol* 25: 641-650.
- 30 Löw M, Herbinger K, Nunn AJ, Häberle KH, Leuchner M, Heerdt C, Werner H, Wipfler P, Pretzsch H, Tausz M, Matyssek R (2006) Extraordinary drought
31 of 2003 overrules ozone impact on adult beech trees (*Fagus sylvatica*). *Trees* 20: 539–548.
- 32 Peiffer M (2005) Parameterisation of the water balance and establishment of water and nutrient fluxes in lowland beech forests. PhD Thesis, École Nationale
33 du Génie Rural, des Eaux et Forêts, Nancy, France, 135 p. (in French)
- 34 Peiffer M, Bréda N, Badeau V, Granier A (2014) Disturbances in European beech water relation during an extreme drought. *Ann ForSci*
- 35 Raftoyannis Y, Radoglou K (2002) Physiological responses of beech and sessile oak in a natural mixed stand during dry summer. *Annals of Botany* 89: 723-
36 730.
- 37 Roberts J, Rosier P T W (1994) Comparative estimates of transpiration of ash and beech forest at a chalk site in southern Britain. *J Hydrol* 162: 229-245.
- 38 Wieser G, Hecke K, Tausz M, Häberle KH, Grams TEE, Matyssek R (2003) The influence of microclimate and tree age on the defence capacity of European
39 beech (*Fagus sylvatica*L.) against oxidative stress. *Ann ForSci* 60: 131-135. <http://hal.archives-ouvertes.fr/hal-0088368>

