



**Melitea salexigens gen. nov., sp. nov., a
gammaproteobacterium from the Mediterranean Sea**

Laurent Urios, Hélène Agogué, Laurent Intertaglia, Françoise Lesongeur,
Philippe Lebaron

► **To cite this version:**

Laurent Urios, Hélène Agogué, Laurent Intertaglia, Françoise Lesongeur, Philippe Lebaron. Melitea salexigens gen. nov., sp. nov., a gammaproteobacterium from the Mediterranean Sea. International Journal of Systematic and Evolutionary Microbiology, 2008, 58, pp.2489-2483. 10.1099/ij.s.0.65685-0 . hal-01102906

HAL Id: hal-01102906

<https://hal.science/hal-01102906>

Submitted on 13 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Table 1. Characteristics that serve to distinguish strain 5IX/A01/131^T from the type strains of related species

Strains: 1, *M. maritimus* JCM 12187^T (data from Yoon *et al.*, 2004); 2, *M. hydrolyticus* DSM 11525^T (Gonzalez *et al.*, 1997); 3, *M. salipaludis* KCCM 41586^T (Yoon *et al.*, 2003a); 4, *M. elongatus* DSM 6810^T (Yoon *et al.*, 2003b); 5, *M. celer* KCTC 12973^T (Yoon *et al.*, 2007); 6, *H. salexigens* OOB 286^T (Urios *et al.*, 2008); 7, strain 5IX/A01/131^T. All strains were positive for the following characteristics: alkaline phosphatase, leucine arylamidase, acid phosphatase and naphthol-AS-BI-phosphohydrolase. All strains possessed Q-8 as the major quinone. +, Positive; –, negative; (+), weakly positive; ND, no data available.

Characteristic	1	2	3	4	5	6	7
Sampling environment	Marine sediment	Salt marsh	Salt marsh	Sand and seawater	Marine saltern	Seawater	Seawater
DNA G + C content (mol%)	59.9	57.7	59	58.2	57.7	61.4	57.2
Motility	–	–	–	+	–	+	+
Temperature for growth (°C)							
Range	15–48	10–41	10–45	25–30	10–48	10–37	15–37
Optimum	37	37	37	25–30	37	25	30
Salinity for growth (g NaCl l ^{–1})							
Range	2–10	6–60	2–10	ND	ND	7–70	7–70
Optimum	2–4	6–30	2–3	2–3	ND	42	28
pH for growth							
Range	5–7.5	6.5–8.5	5–8	ND	5–8	5–9	6–10
Optimum	6.5–7.5	7.5	7–8	ND	7–8	8	8
API ZYM tests							
Valine arylamidase	–	–	–	–	–	+	–
<i>N</i> -Acetyl- β -glucosaminidase	–	–	+	+	+	–	(+)
Esterase (C4)	+	+	+	+	+	–	–
Lipase (C8)	+	+	+	+	+	(+)	(+)
Substrates							
Arabinose	–	+	–	+	+	–	–
Cellobiose	+	+	+	+	–	–	(+)
Fructose	–	–	–	+	–	–	+
Galactose	–	–	–	+	–	–	–
Glucose	+	+	+	+	–	–	+
Lactose	–	–	–	+	+	–	(+)
Maltose	+	+	+	+	–	–	+
Mannose	–	–	–	+	–	(+)	+
Rhamnose	–	–	+	–	–	–	–
Raffinose	–	ND	–	–	–	–	+
Sucrose	–	–	–	+	–	–	+
Glycerol	ND	–	ND	–	ND	+	–
Acetate	+	+	ND	+	+	–	(+)
Aspartate	ND	–	ND	(+)	ND	+	–
Glutamate	–	+	ND	+	–	(+)	(+)
β -Hydroxybutyrate	ND	+	ND	ND	ND	+	–
Lactate	–	ND	ND	+	ND	–	–
Propionate	ND	+	ND	+	ND	–	–
Pyruvate	+	+	ND	ND	+	+	–
Succinate	+	+	ND	+	–	+	–
Alanine	ND	+	ND	(+)	ND	–	(+)
Leucine	ND	+	ND	+	ND	+	–
Proline	ND	+	ND	(+)	ND	+	–
Serine	+	+	ND	–	ND	–	–
Tween 80	ND	–	+	ND	+	+	–

used for the determination of growth parameters were as reported by Wery *et al.* (2001b). Growth was observed at 15–37 °C, with a clear optimum at 30 °C (Supplementary Fig. S2). The strain grew at NaCl concentrations ranging from 7 to 70 g l^{–1} and an optimum concentration could be defined as 42 g l^{–1} (Supplementary Fig. S2). Growth occurred at

pH 6.0–10.0, with an optimum at pH 8.0. Growth decreased by 49 % at pH 6.0 in comparison with the value obtained at pH 7.0. Growth rates were the same at pH 7.0 and pH 9.0; a small difference (14 %) was observed between pH 9.0 and 10.0 (Supplementary Fig. S2). To investigate the possibility of growth under anaerobic conditions, cultures were placed in




Fig. 1. Phylogenetic tree, based on 16S rRNA gene sequences, showing the position of strain 5IX/A01/131^T. The tree corresponds to an unrooted tree obtained using the neighbour-joining algorithm (with Kimura corrections). Accession numbers and type strains are indicated. Bootstrap percentages (based on 1000 resamplings) are shown at branch points. Bar, 0.02 substitutions per nucleotide position.

an anaerobic jar on MA plates at optimal conditions of temperature, pH and salinity (Supplementary Fig. S2). No growth was observed after 10 days and thus the strain should be considered as strictly aerobic.

The ability of isolate 5IX/A01/131^T to use various substrates was investigated using Biolog GN2 MicroPlates (Tang *et al.*, 1998) according to the manufacturer's instructions. Positive reactions are listed in the species description. A comparison between strain 5IX/A01/131^T and its closest relatives is presented in Table 1.

Enzyme activities were investigated using the API ZYM system (bioMérieux) according to the manufacturer's instructions. Positive reactions were obtained for alkaline phosphatase, leucine arylamidase, acid phosphatase and naphthol-AS-BI-phosphohydrolase. Weakly positive signals were obtained for esterase (C8) and *N*-acetylglucosaminidase (Table 1).

Analysis of the fatty acid methyl esters was performed by the Identification Service of the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ, Braunschweig, Germany). To allow comparisons between the fatty acid composition of strain 5IX/A01/131^T and those of its relatives, MA was used. Biomass was harvested from MA plates after cultivation at optimal conditions (Supplementary Fig. S2). The fatty acid composition (%) for isolate 5IX/A01/131^T was as follows: 17:1 ω 8c (34.1), 17:0 (13.0), 18:1 ω 7c (11.4), 16:1 ω 7c/iso-15:0 2-OH (9.0), 15:0 (5.3), 11:0 3-OH (4.6), iso-11:0 3-OH (4.3), 16:0 (3.9), iso-15:0 (2.0), 9:0 (1.9), 10:0 3-OH (1.7), iso-17:0 (1.3), iso-13:0 (1.0), iso-11:0 (0.9), 14:0 (0.8), 11:0 (0.7), 17:1 ω 6c (0.6), 18:0 (0.5), 12:1 3-OH (0.5), 13:0 (0.4), 12:0 (0.3) and 9:0 3-OH (0.3). Fatty acids 9:0, iso-13:0, 9:0 3-OH and 12:1 3-OH were only detected in strain 5IX/A01/131^T. Fatty acids 11:0, 12:0, 13:0, 17:1 ω 6c and 11:0 3-OH were found in strains 5IX/A01/131^T and *H. salexigens* DSM 19537^T but not in *Microbulbifer* species. Fatty acids iso-11:0, iso-15:0 and

Table 2. Cellular fatty acid compositions (%) of strain 5IX/A01/131^T and related type strains in the genus *Microbulbifer*

Strains: 1, *M. maritimus* JCM 12187^T (data from Yoon *et al.*, 2004); 2, *M. hydrolyticus* DSM 11525^T (Gonzalez *et al.*, 1997); 3, *M. salipaludis* KCCM 41586^T (Yoon *et al.*, 2003a); 4, *M. elongatus* DSM 6810^T (Yoon *et al.*, 2003b); 5, *M. celer* KCTC 12973^T (Yoon *et al.*, 2007); 6, *H. salexigens* OOB 286^T (Urios *et al.*, 2008); 7, strain 5IX/A01/131^T. –, Not detected/not reported.

Fatty acid	1	2	3	4	5	6	7
9:0	–	–	–	–	–	–	1.9
10:0	1.3	1.7	2.4	1.6	1.6	–	–
11:0	–	–	–	–	–	1.0	0.7
12:0	–	–	–	–	–	1.6	0.3
13:0	–	–	–	–	–	1.3	0.4
14:0	1.0	1.2	2.6	0.7	0.9	1.3	0.8
15:0	1.6	1.5	1.7	0.9	1.5	4.5	5.3
16:0	8.7	11.4	16.3	7.1	12.6	2.0	3.9
17:0	1.3	2.9	2.2	2.5	2.6	9.3	13.0
18:0	–	1.6	1.4	1.2	0.8	–	0.5
17:0 cyclo	2.3	5.7	–	–	6.3	–	–
19:0 ω 8c cyclo	1.4	1.0	–	–	5.3	–	–
iso-11:0	10.0	5.7	4.8	6.5	6.7	–	0.9
iso-13:0	–	–	–	–	–	–	1.0
iso-15:0	25.9	24.4	19.4	20.7	21.7	–	2.0
iso-15:1	0.8	1.0	0.7	1.0	0.4	–	–
iso-16:0	–	–	–	0.5	0.3	–	–
iso-17:0	6.9	10.4	5.5	9.9	10.0	–	1.3
anteiso-17:0	–	–	–	0.8	0.3	–	–
iso-17:1 ω 9c	12.6	10.1	9.5	11.3	8.6	–	–
15:1 ω 6c	–	–	–	–	–	5.8	–
17:1 ω 6c	–	–	–	–	–	2.7	0.6
17:1 ω 8c	–	0.5	1.0	1.8	0.5	23.9	34.1
18:1 ω 5c	–	–	0.7	–	–	–	–
18:1 ω 7c	5.6	8.9	11.8	16.3	6.5	17.5	11.4
9:0 3-OH	–	–	–	–	–	–	0.3
10:0 3-OH	1.7	1.0	1.2	1.6	1.4	1.8	1.7
11:0 3-OH	–	–	–	–	–	3.3	4.6
12:0 3-OH	–	–	–	–	–	1.1	–
12:1 3-OH	–	–	–	–	–	–	0.5
16:0 2-OH	–	–	0.9	–	–	–	–
iso-11:0 3-OH	14.2	6.2	5.7	7.7	8.9	3.3	4.3
iso-17:0 3-OH	–	–	0.9	–	–	–	–
16:1 ω 7c/iso-15:0 2-OH	2.2	2.7	7.1	6.0	2.4	21.2	9.0

iso-17:0 were found in strain 5IX/A01/131^T and in *Microbulbifer* species but not in *H. salexigens* DSM 19537^T. The major fatty acid found in all *Microbulbifer* strains, iso-15:0, was 10-fold less abundant in strain 5IX/A01/131^T and iso-17:1 ω 9c, one of the other major fatty acids of *Microbulbifer* strains, was not detected in the novel strain. Fatty acids 12:0 3-OH and 15:1 ω 6c, one of the most abundant fatty acids in *H. salexigens* DSM 19537^T, were not found in strain 5IX/A01/131^T.

Analyses of respiratory quinones and polar lipids were carried out by the Identification Service of the DSMZ.

Strain 5IX/A01/131^T contained Q-8 and the major polar lipids were diphosphatidylglycerol, phosphatidylglycerol and an undefined aminophospholipid.

Genomic DNA was extracted as described by Wery *et al.* (2001a). The G+C content was determined by means of thermal denaturation, using the method of Marmur & Doty (1962) and the conditions described by Raguénès *et al.* (1997). The G+C content of the genomic DNA of strain 5IX/A01/131^T was 57.2 ± 0.2 mol%. The 16S rRNA gene was amplified and sequenced as described by Agogué *et al.* (2005). The sequence was analysed as described by Urios *et al.* (2006). Strain 5IX/A01/131^T was shown to be phylogenetically affiliated to the family *Alteromonadaceae* in the class *Gammaproteobacteria* (Fig. 1). The closest relative was *H. salexigens* 3X/A02/235^T, with a sequence similarity of 91%.

Differences in several phenotypic properties, as shown in Tables 1 and 2, serve to distinguish strain 5IX/A01/131^T from its closest phylogenetic relatives. On the basis of these differences, we propose that strain 5IX/A01/131^T should be assigned to a novel species and genus. Because of the marine origin of strain 5IX/A01/131^T and its salinity requirement, the name *Melitea salexigens* gen. nov., sp. nov. is proposed.

Description of *Melitea* gen. nov.

Melitea (Me.li'te.a. N.L. fem. n. *Melitea* named after Melite, a nymph of the sea in Greek mythology, referring to the marine origin).

Cells are motile, Gram-negative rods. The major fatty acids are 17:1 ω 8c, 17:0, 18:1 ω 7c, 16:1 ω 7c, 15:0, 11:0 3-OH, iso-11:0 3-OH, 16:0, iso-15:0, 9:0, 10:0 3-OH, iso-17:0 and iso-13:0. The ubiquinone is Q-8 and the polar lipids are diphosphatidylglycerol and phosphatidylglycerol. Phylogenetically affiliated to the class *Gammaproteobacteria*. The type species is *Melitea salexigens*.

Description of *Melitea salexigens* sp. nov.

Melitea salexigens (sa.le'xi.gens. L. n. *sal*, *salis* salt, seawater; L. v. *exigo* to demand; N.L. part. adj. *salexigens* seawater-demanding).

Displays the following properties in addition to those described for the genus. Cream colonies are formed on MA medium. Cells are 1.8 ± 0.2 μ m long and 0.7 ± 0.1 μ m wide with single polar flagella. The DNA G+C content of the type strain is 57 mol%. Growth occurs at 15–37 °C (optimally at 30 °C), at pH 6.0–10.0 (optimally at pH 8.0) and at salinities in the range 7–70 g NaCl l⁻¹ (optimally at 42 g l⁻¹). Positive reactions with Biolog GN2 plates are obtained for D-fructose, D-glucose, maltose, D-mannose, D-psicose, raffinose, sucrose, trehalose, turanose, α -ketoglutaric acid, α -ketovaleric acid, succinamic acid, L-glutamic acid and L-serine. Positive API ZYM reactions for enzyme activities are obtained for alkaline phosphatase,

leucine arylamidase, acid phosphatase and naphthol-AS-BI-phosphohydrolase. Oxidase- and catalase-positive.

The type strain, 5IX/A01/131^T (=DSM 19753^T =CIP 109757^T =MOLA 225^T), was isolated from seawater sampled from the surface microlayer in the bay of Barcelona (41° 24' N 2° 16' E).

Acknowledgements

This work was supported by the Equipe Mixte de Recherche linking the Université Pierre et Marie Curie and the Centre National de la Recherche Scientifique to the Pierre Fabre Laboratories. The project was also carried out within the framework of the MarBEF Network of Excellence 'Marine Biodiversity and Ecosystem Functioning', which is funded within the Community's Sixth Framework Programme (contract no. GOCE-CT-2003-505446). This publication is contribution number MPS-07066 of MarBEF. It was also partly funded by the French programme 'Bio-diversité et Changement Global – project: development of a coastal microbial observatory' from the Institut Français de la Biodiversité (IFB-GICC, Paris, France).

References

- Agogué, H., Casamayor, E. O., Joux, F., Obernosterer, I., Dupuy, C., Lantoine, F., Catala, P., Weinbauer, M. G., Reinthaler, T. & other authors (2004). Comparison of samplers for the biological characterization of the sea surface microlayer. *Limnol Oceanogr Methods* 2, 213–225.
- Agogué, H., Casamayor, E. O., Bourrain, M., Obernosterer, I., Joux, F., Herndl, G. J. & Lebaron, P. (2005). A survey on bacteria inhabiting the sea surface microlayer of coastal ecosystems. *FEMS Microbiol Ecol* 54, 269–280.
- Gonzalez, J. M., Mayer, F., Moran, M. A., Hodson, R. E. & Whitman, W. B. (1997). *Microbulbifer hydrolyticus* gen. nov., sp. nov., and *Marinobacterium georgiense* gen. nov., sp. nov., two marine bacteria from a lignin-rich pulp mill waste enrichment community. *Int J Syst Bacteriol* 47, 369–376.
- Humm, H. J. (1946). Marine agar-digesting bacteria of the South Atlantic coast. *Duke Univ Mar Stn Bull* 3, 45–75.
- Marmur, J. & Doty, P. (1962). Determination of the base composition of deoxyribonucleic acid from its thermal denaturation temperature. *J Mol Biol* 5, 109–118.
- Palleroni, N. J. (1984). Genus *Pseudomonas* Migula 1894. In *Bergey's Manual of Systematic Bacteriology*, vol. 1, pp. 141–199. Edited by N. R. Krieg & J. G. Holt. Baltimore: Williams & Wilkins.
- Powers, E. M. (1995). Efficacy of the Ryu nonstaining KOH technique for rapidly determining Gram reactions of food-borne and waterborne bacteria and yeasts. *Appl Environ Microbiol* 61, 3756–3758.
- Raguénès, G., Christen, R., Guézennec, J., Pignet, P. & Barbier, G. (1997). *Vibrio diabolicus* sp. nov., a new polysaccharide-secreting organism isolated from a deep-sea hydrothermal vent polychete annelid, *Alvinella pompejana*. *Int J Syst Bacteriol* 47, 989–995.
- Tang, Y. W., Ellis, N. M., Hopkins, M. K., Smith, D. H., Dodge, D. E. & Persing, D. H. (1998). Comparison of phenotypic and genotypic techniques for identification of unusual aerobic pathogenic gram-negative bacilli. *J Clin Microbiol* 36, 3674–3679.
- Urios, L., Agogué, H., Lesongeur, F., Stackebrandt, E. & Lebaron, P. (2006). *Balneola vulgaris* gen. nov., sp. nov., a member of the phylum *Bacteroidetes* from the north-western Mediterranean Sea. *Int J Syst Evol Microbiol* 56, 1883–1887.

Urios, L., Intertaglia, L., Lesongeur, F. & Lebaron, P. (2008). *Haliea salexigens* gen. nov., sp. nov., a member of the *Gammaproteobacteria* from the Mediterranean Sea. *Int J Syst Evol Microbiol* 58, 1233–1237.

Wery, N., Lesongeur, F., Pignet, P., Derennes, V., Cambon-Bonavita, M.-A., Godfroy, A. & Barbier, G. (2001a). *Marinitoga camini* gen. nov., sp. nov., a rod-shaped bacterium belonging to the order *Thermotogales*, isolated from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* 51, 495–504.

Wery, N., Moricet, J., Cueff, V., Jean, J., Pignet, P., Lesongeur, F., Cambon-Bonavita, M.-A. & Barbier, G. (2001b). *Caloranaerobacter azorensis* gen. nov., sp. nov., an anaerobic thermophilic bacterium isolated from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* 51, 1789–1796.

Yoon, J. H., Kim, I. G., Shin, D. Y., Kang, K. H. & Park, Y. H. (2003a). *Microbulbifer salipaludis* sp. nov., a moderate halophile isolated from a Korean salt marsh. *Int J Syst Evol Microbiol* 53, 53–57.

Yoon, J. H., Kim, I. G., Kang, K. H., Oh, T. K. & Park, Y. H. (2003b). Transfer of *Pseudomonas elongata* Humm 1946 to the genus *Microbulbifer* as *Microbulbifer elongatus* comb. nov. *Int J Syst Evol Microbiol* 53, 1357–1361.

Yoon, J. H., Kim, I. G., Oh, T. K. & Park, Y. H. (2004). *Microbulbifer maritimus* sp. nov., isolated from an intertidal sediment from the Yellow Sea, Korea. *Int J Syst Evol Microbiol* 54, 1111–1116.

Yoon, J. H., Jung, S. Y., Kang, S. J. & Oh, T. K. (2007). *Microbulbifer celer* sp. nov., isolated from a marine solar saltern of the Yellow Sea in Korea. *Int J Syst Evol Microbiol* 57, 2365–2369.