



HAL
open science

Progress Towards a conjecture on the mean-value of titchmarsh series-II

K Ramachandra

► **To cite this version:**

K Ramachandra. Progress Towards a conjecture on the mean-value of titchmarsh series-II. Hardy-Ramanujan Journal, 1981, Volume 4 - 1981, pp.1 - 12. 10.46298/hrj.1981.91 . hal-01103883

HAL Id: hal-01103883

<https://hal.science/hal-01103883>

Submitted on 15 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PROGRESS TOWARDS A CONJECTURE ON THE MEAN-VALUE OF TITCHMARSH SERIES-II

By K. RAMACHANDRA

§ 1. Introduction

The result of this paper may be considered as complementary to that of my earlier paper [2], on Titchmarsh series. Although not as interesting as the earlier result, the result of the present paper finds a nice application, (See [1]). In [3] I defined a class of series called Titchmarsh series and I now start by recalling its definition.

Titchmarsh Series. (or briefly K D T series).

Let $A > 10$ be a constant.

Let $\frac{1}{A} < \mu_1 < \mu_2 < \dots$ where

$$\frac{1}{A} < \mu_{n+1} - \mu_n < A \text{ (for } n = 1, 2, 3, \dots \text{)}.$$

(In [2] the notation is slightly different and we have used there λ_n instead of μ_n and for simplicity assumed $\lambda_1 = 1$. Also we have used there a_1, a_2, \dots in place of our present b_1, b_2, \dots and assumed for simplicity $a_1 = 1$. We have written there $F(s)$ instead of $F_0(s)$). Let b_1, b_2, \dots be a sequence of complex numbers possibly depending on a parameter $H > 10$ such that

$$|b_n| < (\mu_n H)^A. \text{ Put } F_0(s) = \sum_{n=1}^{\infty} b_n \mu_n^{-s} \text{ where } s = \sigma + it.$$

$F_0(s)$ is called a KDT series if there exists a constant $A > 10$ and a system of rectangles $R(T, T+H)$ defined by $\{\sigma > 0, T \leq t \leq T+H\}$ where $10 < H \leq T$, and T (which may be related to H) tends to infinity, and $F_0(s)$ admits an analytic continuation into these rectangles and the maximum of $|F_0(s)|$ taken over $R(T, T+H)$ does not exceed $\text{Exp}(H^A)$.

I then made the following conjecture

Conjecture

For a KDT series $F_0(s)$, we have,

$$\frac{1}{H} \int_L |F_0(it)|^2 dt > C_A \mu_n^{\frac{1}{X}} |b_n|^2,$$

where $X = 2 + D_A H$, L denotes the side ($\sigma = 0, T \leq t \leq T+H$) of $R(T, T+H)$, and C_A and D_A are positive constants depending only on A , provided $\mu_1 = b_1 = 1$.

I proved the following theorem.

Theorem 1

Under the restrictions $\mu_1 = b_1 = 1$, we have,

$$\frac{1}{H} \int_L |F_0(it)|^2 dt > C_A \mu_n^{\frac{1}{X}} |b_n|^2 \left(1 - \frac{\log \mu_n}{\log H} + \frac{1}{\log \log H} \right),$$

where $X = 2 + D_A H$ and, C_A and D_A are effective positive constants depending only on A .

I now prove the following theorem.

Theorem 2

For some convenience let us assume in the definition of Titchmarsh series $F_0(s)$ the rectangles $R(T, T + H)$ to be $(\sigma > \beta, T < t < T + H)$ where β is a positive constant such that $0 < \beta < \alpha_1 < \frac{1}{2}$, where α_1 is another constant. Let

$k \geq 2$ be an integer and write $F(s) = (F_0(s))^k =$

$$\sum_{n=1}^{\infty} (a_n \lambda_n^{-s}),$$

a series which is surely convergent where $F_0(s)$

is absolutely convergent. Put $Y = (M + H)^{\lambda}$ where

$$\lambda = kA^{100} (\alpha_1 - \beta)^{-10}$$

and $M = \text{maximum of } |F_0(s)|$

taken over $R(T, T + H)$. Define the entire function $\phi(s)$ by

$$\phi(s) = \sum_{n=1}^{\infty} a_n \lambda_n^{-s} \Delta\left(\frac{Y}{\lambda_n}\right)$$

where for $X > 0$, $\Delta(X)$ is defined by

$$\Delta(X) = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} X^W \text{Exp}(W^{4a+2}) \frac{dW}{W},$$

a being a suitable positive integer constant at our choice. We now suppose that $\alpha, \beta, \alpha_1, \alpha_2, \alpha_3$ are constants satisfying

$$\alpha < \beta < \alpha_1 < \alpha_2 < \alpha_3 < \alpha + \frac{1}{2}.$$

Put $X = [2^{-A-100} H] + 2$ and

$$V(\sigma) = \frac{1}{H} \sum_{\mu_n < X} |b_n|^2 \left(\frac{H}{\mu_n}\right)^{2\sigma}.$$

Then we have,

$$\frac{1}{H} \int_L |\phi(\alpha + it)|^{2k-1} dt$$

$$\gg V(\alpha) H^{1-2\alpha} \left\{ \frac{V(\alpha_1)}{V(\alpha_3)} \right\}^{\frac{\alpha_2 - \alpha}{\alpha_3 - \alpha_2}},$$

provided only that $V(\alpha) \neq 0$ and $(\log V(\alpha)) (\log H)^{-1}$ is bounded below by a negative constant. Here the constant implied by the Vinogradov symbol \gg depends only on $A, k, \alpha, \beta, \alpha_1, \alpha_2, \alpha_3$ and the negative constant referred to just now. Further it is effective.

Remark: It will be clear from the proof that if

$$\mu_n = n \quad (n = 1, 2, 3, \dots)$$

then we can choose $X = \left[\frac{H}{100} \right] + 2$.

The object of this paper is to prove theorem 2. The proof of the theorem is fairly long. The proof depends upon a special case of a convexity theorem of R. M. Gabriel which we state below (in the notation of D. R. Heath-Brown's paper [2]; for the more general theorem of Gabriel see the reference in [2] or Titchmarsh's famous book [5] pages 203 and 337.)

Theorem 3

Let $f(z)$ be regular in the infinite strip $\alpha < \operatorname{Re} z < \beta$ and continuous for $\alpha < \operatorname{Re} z < \beta$. Suppose $f(z) \rightarrow 0$ as $|\operatorname{Im} z| \rightarrow \infty$, uniformly in $\alpha < \operatorname{Re} z < \beta$. Then for $\alpha \leq \gamma \leq \beta$, and any $q > 0$, we have

$$\int_{-\infty}^{\infty} |f(\gamma + it)|^q dt < \left(\int_{-\infty}^{\infty} |f(\alpha + it)|^q dt \right)^{\frac{\beta - \gamma}{\beta - \alpha}} \left(\int_{-\infty}^{\infty} |f(\beta + it)|^q dt \right)^{\frac{\gamma - \alpha}{\beta - \alpha}}$$

provided the right hand side is finite.

Apart from this we have to use a well-known theorem of Montgomery and Vaughan. For reference see for instance my paper [4], where I give a simple proof of a weaker result which is sufficient for the purposes of this paper.

We now split up the proof of theorem 2 into several steps and give a brief sketch of these steps.

§ 2. Proof of Theorem 2

Step 1. Let

$$I(\sigma) = \frac{1}{H} \int_T^{T+H} \left(\int_{-\infty}^{\infty} |\phi(s)|^{2/k} | \text{Exp}((s - it_0)^{4a+2}) | dt \right) dt_0,$$

and assume that $I(\alpha) < V(\alpha) H^{1-2\alpha}$. The constant a shall be a sufficiently large positive integer. As already stated we set

$$X = [2^{-A-100} H] + 2, \text{ and}$$

$$V(\sigma) = \frac{1}{H} \sum_{\mu_n < X} |b_n|^2 \left(\frac{H}{\mu_n} \right)^{2\sigma} \text{ and}$$

we impose $\alpha < \beta < \alpha_1 < \alpha_2 < \alpha_3 < \alpha + \frac{1}{2}$.

Step 2. Next we write

$$J(\sigma) = \frac{1}{H} \int_T^{T+H} \left(\int_{-\infty}^{\infty} \left| \phi(s) - P^k(s) \right| \left| \text{Exp}((s-it_0)^{4a+2}) \right| dt \right) dt_0$$

where $P(s) = \sum_{\mu_n < X} b_n \mu_n^{-s}$. It is easily seen that

$\Delta(X) = O(X^B)$ and also $1 + O(X^{-B})$ where $B > 0$ is an arbitrary constant and the O -constant depends only on B and a .

Again $a_n = \mu_{n_1} \dots \mu_{n_k} = \lambda_n (b_{n_1} b_{n_2} \dots b_{n_k})$ and for

all $N > 1$, we have $N < \mu_{n_1} \dots \mu_{n_k} \leq 2N = O(N^k)$. From

these remarks it is clear that $\phi(s) - P^k(s)$ decays fast enough to ensure $J(\sigma) < 1$ when σ is large enough. Now from an easy application of a theorem of Gabriel (Theorem 3 above) it follows that in $\sigma > \alpha$, $J(\sigma)$ is $\ll_{\epsilon} [J(\alpha)]^{1-\epsilon}$ for every positive constant ϵ uniformly in σ , and so in $(\sigma > \beta, T < t \leq T+H)$, $|\phi(s)|$ is bounded above by a constant power of H . (Here for getting the last bound we have to use the fact that for any analytic function $\phi(s)$, $|\phi(s)|^{2/k}$ is bounded by its mean value over a disc of (positive but sufficiently small) constant radius with s as centre).

Step 3. An easy application of a well-known Montgomery-Vaughan theorem (refer [4] for instance) shows that

$$\begin{aligned} \frac{1}{H} \int_T^{T+H} \left(\int_{-\infty}^{\infty} \left| P(s) \right|^2 \left| \text{Exp}[(s-it_0)^{4a+2}] \right| dt \right) dt_0 \\ = O(V(\sigma) H^{1-2\sigma}). \end{aligned}$$

From this and the estimate $J(\sigma) \ll_{\epsilon} (J(\alpha))^{1-\epsilon}$ it follows that

$$\int_{\alpha}^{A^3} \left(\frac{1}{H} \int_T^{T+H} \left(\int_{-\infty}^{\infty} (|\phi(s) - P^k(s)|^{2/k} + |P(s)|^2) |\text{Exp}((s-it_0)^{4a+2})| dt \right) dt_0 \right) d\sigma = O((J(\alpha))^{1-\epsilon} + V(\alpha) H^{1-2\alpha}).$$

Note that $V(\sigma)$ and $V^*(\sigma) H^{1-2\sigma}$ are respectively monotonic increasing and monotonic decreasing functions of σ , where $V^*(\sigma)$ is the same as $V(\sigma)$ with the terms $\mu_n \leq 1$ omitted. From now on we assume that $V(\alpha)$ is bounded below by a constant negative power of H . Under this assumption it follows that the integral just considered is

$$O_{\epsilon}(V(\alpha) H^{1-2\alpha+\epsilon}) \text{ for every positive constant } \epsilon.$$

Hence there exist intervals I_1 and I_2 contained in $(T, T + \frac{H}{10})$ and $(T + H - \frac{H}{10}, T + H)$ respectively, for which the lengths are $4H^{\delta}$ (δ being any fixed constant satisfying $0 < \delta < \frac{1}{100}$) each, and further $I(I_1, \alpha)$

$$= \int_{\alpha}^{A^3} \left(\frac{1}{H} \int_{I_1} \int_{-\infty}^{\infty} (|\phi(s) - P^k(s)|^{2/k} + |P(s)|^2) |\text{Exp}((s-it_0)^{4a+2})| dt dt_0 \right) d\sigma$$

and $I(I_2, \alpha)$ defined similarly (by replacing I_1 by I_2) satisfy

$$I(I_1, \alpha) + I(I_2, \alpha) = O(V(\alpha) H^{1-2\alpha+\epsilon}).$$

Hence by the principle for the mean value over discs referred to in the second step, we see that in $(\beta \leq \sigma \leq A^2, t$ in any of the intervals $I_1, I_2)$ we have

$$|\phi(s)|^{2/k} = O(V(\alpha) H^{-2\alpha + 2\epsilon + \delta}).$$

Step 4. Let H_1 and H_2 be the mid points of I_1 and I_2 respectively. We now obtain a lower bound for at least one of the mean-values $K(\alpha_1)$ or $K(\alpha_2)$ where $K(\sigma)$ is defined by

$$K(\sigma) = \frac{1}{H_2 - H_1} \int_{H_1}^{H_2} |F_0(s)|^2 dt, (\beta < \sigma < A^2).$$

Note that when we replace H_1 and H_2 by other points in I_1 and I_2 the mean value $K(\sigma)$ changes by an amount which is at most $O(E)$ where $E = V(\alpha) H^{-2\alpha + 2\epsilon + 2\delta}$. Hence if σ denotes any of α_2 or α_3 and $H_1 < t < H_2$, we see that if j is a large positive integer constant,

$$(1) \quad \frac{j!}{2\pi i} \int_{L_0} \frac{F_0(s+w) (2X)^w}{w(w+1)\dots(w+j)} dw \\ = \sum_{\mu_n < 2X} b_n \mu_n^{-s} \left(1 - \frac{\mu_n}{2X}\right)^j,$$

where L_0 is the line $\text{Re } w = A^2$. Deform the line L_0 to the contour described by the lines L_1, L_2, L_3, L_4, L_5 (in this order) defined as follows. Let $H_3 = H_1 - H^\delta$, $H_4 = H_2 + H^\delta$ where δ is a small positive constant. L_1 and

L_5 are the portions $\text{Im } w \leq -H_3$ and $\text{Im } w \geq H_4$ respectively of L_0 . L_2 is the line segment

$$(\text{Im } w = -H_3, \alpha_1 - \sigma \leq \text{Re } w \leq A^2)$$

and L_4 is the line segment

$$(\text{Im } w = H_4, \alpha_1 - \sigma \leq \text{Re } w \leq A^2).$$

L_3 is the line segment ($\text{Re } w = \alpha_1 - \sigma, -H_3 < \text{Im } w < H_4$).

Taking the mean square after deformation of L_0 we find from the equation (1), (Note that the only pole to be taken care of is $w = 0$),

$$(2) \begin{cases} K(\sigma) \ll V_1(\sigma) H^{1-2\sigma} + (K(\alpha_1) + E) H^{-2(\sigma-\alpha_1)} \\ \text{(where } V_1(\sigma) \text{ is defined below) and also} \\ V(\sigma) H^{1-2\sigma} \ll K(\sigma) + (K(\alpha_1) + E) H^{-2(\sigma-\alpha_1)}. \end{cases}$$

The reason for this is that the mean square of the RHS of (1) is \gg and $\ll V_1(\sigma) H^{1-2\sigma}$ where $V_1(\sigma)$ is defined by

$$V_1(\sigma) = \frac{1}{H} \sum_{\mu_n \leq X} |b_n|^2 \left(1 - \frac{\mu_n}{2X}\right)^{2j} \left(\frac{H}{\mu_n}\right)^{2\sigma}.$$

Since $V(\alpha) \ll V(\sigma)$ we may omit the term containing E in the second of the equations (2), provided $\alpha_1 - \alpha < \frac{1}{2}$ (which is true because of our assumptions). This gives us $V(\sigma) H^{1-2\sigma} \ll K(\sigma) + K(\alpha_1) H^{-2(\sigma-\alpha_1)}$; ($\sigma = \alpha_2, \alpha_3$)

If we put $\sigma = \alpha_2$ we get a lower bound for one at least of the quantities $K(\alpha_1)$ or $K(\alpha_2)$.

We now deduce from the last inequality

$$V(\sigma) H^{1-2\sigma} \ll I(\sigma) + I(\alpha_1) H^{-2(\sigma-\alpha_1)},$$

$$(\sigma = \alpha_2, \alpha_3).$$

This is possible since in the range $(\sigma > \alpha_1, H_3 < t \leq H_4)$,

$|F_0(s)|^2$ is a very good approximation (in the mean) to

$|\phi(s)|^{2/k}$ and we leave the details as an exercise. From the

last inequality it follows that $I(\sigma) \gg V(\sigma) H^{1-2\sigma}$, for one at least of the values $\sigma = \alpha_1$ or α_2 .

Next by the convexity theorem of Gabriel (Theorem 3), we find that with the value of σ (α_1 or α_2 determined),

$$(3) \quad (I(\sigma))^{\alpha_3 - \alpha} < (I(\alpha))^{\alpha_3 - \sigma} (I(\alpha_3))^{\sigma - \alpha}.$$

Moreover by the arguments used in the first of the inequalities in (2) we get (by taking X in place of $2X$ in (1)).

$$I(\alpha_3) \ll V(\alpha_3) H^{1-2\alpha_3} + (I(\sigma) + E) H^{-2(\alpha_3 - \sigma)} + E.$$

$$\text{Now } E = V(\alpha) H^{-2\alpha + 2\varepsilon + 2\delta} < V(\alpha_3) H^{1-2\alpha_3}$$

(since by our assumptions, $\alpha_3 - \alpha < \frac{1}{2}$) by a small choice of the positive constants ε, δ . Thus we get

$$I(\alpha_3) \ll V(\alpha_3) H^{1-2\alpha_3} + I(\sigma) H^{-2(\alpha_3 - \sigma)},$$

and so by (3)

$$(4) \quad (I(\sigma))^{\alpha_3 - \alpha} \ll (I(\alpha))^{\alpha_3 - \sigma} \left(V(\alpha_3) H^{1-2\alpha_3} + I(\sigma) H^{-2(\alpha_3 - \sigma)} \right)^{\sigma - \alpha}.$$

This holds for either $\sigma = \alpha_1$ or $\sigma = \alpha_2$ and gives us

$$(5) \left\{ \begin{array}{l} \text{Either } I(\alpha) \gg V(\sigma) H^{1-2\alpha} > V(\alpha) H^{1-2\alpha}, \\ \text{or } (I(\alpha))^{\alpha_3 - \sigma} \gg (V(\sigma) H^{1-2\sigma} \alpha_3^{-\alpha}) \\ \qquad \qquad \qquad (V(\alpha_3) H^{1-2\alpha_3} \alpha^{-\sigma}). \end{array} \right.$$

The second of these inequalities gives

$$I(\alpha) \gg H^{1-2\alpha} V(\sigma) \left\{ \frac{V(\sigma)}{V(\alpha_3)} \right\}^{\frac{\sigma - \alpha}{\alpha_3 - \sigma}}$$

Since $V(\alpha) < V(\alpha_1) < V(\sigma) < V(\alpha_3)$ and since $\frac{u - \alpha}{\alpha_3 - u}$

is an increasing function of u in $\beta < u < \alpha_3$, we get finally

$$I(\alpha) \gg V(\alpha) H^{1-2\alpha} \left\{ \frac{V(\alpha_1)}{V(\alpha_3)} \right\}^{\frac{\alpha_2 - \alpha}{\alpha_3 - \alpha_2}}$$

Step 5. Step 4 nearly completes the proof. For we could have started with a slight modification of $I(\sigma)$ by averaging over a slightly smaller interval contained in $(T, T + H)$ instead of $(T, T + H)$. For instance by cutting off bits of length H^δ on either side. The decaying factor $\text{Exp}((s - it_0)^{4a+2})$ enables us to replace the modified

$$I(\sigma) \text{ by } \frac{1}{H} \int_T^{T+H} |\phi(s)|^{\frac{2}{k}} dt \quad \sigma = \alpha$$

in the last lower bound.

Steps 1, 2, 3, 4 and 5 complete the proof of theorem 2.

Theorem 4

With the notation of theorem 2, we have,

$$\frac{1}{H} \int_T^{T+H} |\phi(\alpha + it)|^2 dt \gg \left(v(\alpha) H^1 - 2\alpha \left\{ \frac{v(\alpha_1)}{v(\alpha_3)} \right\} \frac{\alpha_2 - \alpha}{\alpha_3 - \alpha_2} \right)^k$$

where the constant implied by the Vinogradov symbol is effective

Remark. This theorem will be used in [1].

References

1. R. Balasubramanian And K. Ramachandra, Some problems of analytic number theory-III, *Hardy-Ramanujan Journal*, Vol. 4 (1981).
2. D. R. Heath-Brown, Fractional moments of the Riemann zeta-function, (to appear).
3. K. Ramachandra, Progress towards a conjecture on the mean-value of Titchmarsh Series, *Proceedings of the Durham Conference on Analytic number theory*, (July-August 1979), (to appear).
4. K. Ramachandra Some remarks on a theorem of Montgomery and Vaughan, *J. of Number theory*, Vol. 11 (1975), 465-471.
5. E. C. Titchmarsh, The theory of the Riemann zeta-function, *Oxford* (1951).

Manuscript Completed In The Final Form On 11-Oct-1980

School of Mathematics
Tata Institute of Fundamental Research
Homi Bhabha Road
Colaba, Bombay 400 005
(India).