



HAL
open science

On a question of Ramachandra

Hugh L Montgomery

► **To cite this version:**

Hugh L Montgomery. On a question of Ramachandra. Hardy-Ramanujan Journal, 1982, Volume 5 - 1982, pp.31 - 36. 10.46298/hrj.1982.95 . hal-01104228

HAL Id: hal-01104228

<https://hal.science/hal-01104228>

Submitted on 16 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON A QUESTION OF RAMACHANDRA

By Hugh L MONTGOMERY *

Let $a_k(n)$ be the Dirichlet series coefficients defined by the relation

$$(1) \quad \left(\sum_p \frac{1}{p^s} \right)^k = \sum_{n=1}^{\infty} \frac{a_k(n)}{n^s} \quad (\text{Re } s > 1).$$

Here k is a positive integer, and we see more explicitly that

if $n = p_1^{a_1} \dots p_t^{a_t}$ then

$$(2) \quad a_k(n) = \begin{cases} \frac{k!}{a_1! \dots a_t!} & \text{if } \Omega(n) = k, \\ 0 & \text{otherwise,} \end{cases}$$

With possible applications to the Riemann zeta function in mind, Ramachandra has wanted to know the asymptotic size of the quantity

$$\max_k \left(\sum_{n \leq N} \frac{a_k(n)^2}{n^{2\sigma}} \right)^{\frac{1}{2k}}$$

as a function of N and σ , where σ is fixed, $\frac{1}{2} < \sigma < 1$. We settle this question by demonstrating the following

Theorem: Let σ be fixed, $\frac{1}{2} < \sigma < 1$. With the $a_k(n)$ defined by (1), we have

$$(3) \quad \max_k \left(\sum_{n \leq N} \frac{a_k(n)^2}{n^{2\sigma}} \right)^{\frac{1}{2k}} \approx \frac{(\log N)^{1-\sigma}}{\log \log N}$$

* Research supported in part by NSF Grant MCS 80-02559.

In addition, for any integer $k > 1$

$$(4) \quad \left(\sum_n \frac{a_k(n)^2}{n^{2\sigma}} \right)^{\frac{1}{2k}} \approx \frac{k^{1-\sigma}}{(\log k)^\sigma}$$

Here the implicit constants may depend only on σ .

With a little more care one could show that the values of k for which the maximum in (3) is attained satisfy

$$k \approx (\log N) / \log \log N.$$

To establish that the right hand side majorizes the left above we shall require the following two lemmas.

Lemma 1: Let $Y > 2$, and put

$$R = R(Y) = \{r: p \mid r \Rightarrow p < Y\}.$$

Then for any $k > 1$, and fixed σ , $\frac{1}{2} < \sigma < 1$,

$$\left(\sum_{r \in R} \frac{a_k(r)^2}{r^{2\sigma}} \right)^{\frac{1}{2k}} \ll \frac{Y^{1-\sigma}}{\log Y}.$$

Proof: Let X_2, X_3, \dots, X_p be independent random variables, each uniformly distributed on the circle $|z| = 1$.

Then

$$\begin{aligned} \left(\sum_{r \in R} \frac{a_k(r)^2}{r^{2\sigma}} \right)^{\frac{1}{2k}} &= \left\| \sum_{p < y} \frac{X_p}{p^\sigma} \right\|_{2k}^{\frac{1}{2k}} \ll \left\| \sum_{p < y} \frac{X_p}{p^\sigma} \right\|_{\infty}^{\frac{1}{2k}} \\ &= \sum_{p < y} \frac{1}{p^\sigma} \ll \frac{Y^{1-\sigma}}{\log Y}. \end{aligned}$$

Lemma 2: Let $Y > 2$, and put

$$S = S(y) = \{s : p | s \Rightarrow p > y\}.$$

Then for any $k > 1$, and fixed σ , $\frac{1}{2} < \sigma < 1$.

$$\left(\sum_{s \in S} \frac{a_k(s)^2}{s^{2\sigma}} \right)^{\frac{1}{2k}} \ll \frac{k^{1/2} Y^{\frac{1}{2}-\sigma}}{(\log Y)^{1/2}}$$

Proof: From (2) we see that $a_k(s) < k!$ for all s .

Thus

$$\begin{aligned} \sum_{s \in S} \frac{a_k(s)^2}{s^{2\sigma}} &< k! \sum_{s \in S} \frac{a_k(s)}{s^{2\sigma}} = k! \left(\sum_{p > Y} \frac{1}{p^{2\sigma}} \right)^k \\ &< k! \left(\frac{c Y^{1-2\sigma}}{\log Y} \right)^k, \end{aligned}$$

and the stated bound follows on taking the $2k$ -th root.

We now prove the Theorem. We first show that the left hand side of (3) is at least as large as the right hand side. Let Y be the largest integer such that

$$P = \prod_{p < Y} p < N.$$

Then by the prime number theorem with remainder,

$$Y = \log N + O\left(\frac{\log N}{(\log \log N)^A}\right)$$

Take $k = \Omega(P) = \pi(Y)$. Then

$$\sum_{n < N} \frac{a_k(n)^2}{n^{2\sigma}} > \frac{a_k(P)^2}{P^{2\sigma}} = \frac{k!^2}{P^{2\sigma}} > \frac{k!^2}{N^{2\sigma}}$$

$$\text{so that } \left(\sum_{n < N} \frac{a_k(n)^2}{n^{2\sigma}} \right)^{\frac{1}{2k}} \gg \frac{k}{N^{\sigma/k}}$$

Clearly $k \sim \frac{\log N}{\log \log N}$, and by more careful use of the prime number theorem we see that $k > \frac{\log N}{\log \log N}$ for all large N . Thus $N^{1/k} < \log N$, and we have the desired lower bound. In (4) the value of k is prescribed; we choose Y so that $\pi(Y) = k$, we take $N = P$, and proceed as above. We obtain the desired lower bound since

$$\frac{(\log N)^{1-\sigma}}{\log \log N} \sim \frac{k^{1-\sigma}}{(\log k)^\sigma}$$

We now complete the proof of (3). Let R and S be as in Lemmas 1 and 2, where Y is a parameter to be chosen later. Any n is uniquely of the form $n = rs$ with $r \in R$, $s \in S$. If $\Omega(n) = k$ and $\Omega(r) = m$ then by (2)

$$a_k(n) = \binom{k}{m} a_m(r) a_{k-m}(s).$$

Hence

$$\sum_{n \leq N} \frac{a_k(n)^2}{n^{2\sigma}} = \sum_{m=0}^k \binom{k}{m}^2 \sum_{r \in R} \frac{a_m(r)^2}{r^{2\sigma}} \sum_{\substack{s \in S \\ rs \leq N}} \frac{a_{k-m}(s)^2}{s^{2\sigma}}$$

Since $\binom{k}{m} < 2^k$, this is

$$< 2^k \sum_{m=0}^k \binom{k}{m} \left(\sum_{r \in R} \frac{a_m(r)^2}{r^{2\sigma}} \right) \left(\sum_{\substack{s \in S \\ s \leq N}} \frac{a_{k-m}(s)^2}{s^{2\sigma}} \right)$$

$$(5) \quad < 2^k (R^2 + s^2)^k$$

where

$$R = \max_m \left(\sum_{r \in R} \frac{a_m(r)^2}{r^{2\sigma}} \right)^{\frac{1}{2m}}$$

and

$$S = \max_m \left(\sum_{\substack{s \in S \\ s < N}} \frac{a_m(s)^2}{s^{2\sigma}} \right)^{\frac{1}{2m}}$$

By Lemma 1

$$(6) \quad R \ll \frac{Y^{1-\sigma}}{\log Y}$$

To treat S we note that $\Omega(s) = m$, $s \in S$, then $s > Y^m$.

If $m > \frac{\log N}{\log Y}$ then it follows that $s > N$, and the sum

defining S will be empty. Thus we may suppose that $m < \frac{\log N}{\log Y}$.

From this bound and Lemma 2 we see that

$$(7) \quad S \ll \frac{(\log N)^{\frac{1}{2}} Y^{\frac{1}{2} - \sigma}}{\log Y}$$

On combining (5), (6), and (7) it follows that

$$\left(\sum_{n < N} \frac{a_k(n)^2}{n^{2\sigma}} \right)^{\frac{1}{2k}} \ll \frac{Y^{1-\sigma}}{\log Y} + \frac{(\log N)^{\frac{1}{2}} Y^{\frac{1}{2} - \sigma}}{\log Y},$$

and it suffices to take $Y = \log N$.

To complete the proof of (4) we may proceed more simply. With the X_p as in the proof of Lemma 1 we see that

$$\left(\sum_n \frac{a_k(n)^2}{n^{2\sigma}} \right)^{\frac{1}{2k}} = \left\| \sum_p \frac{X_p}{p^\sigma} \right\|_{2k}.$$

By the triangle inequality this is

$$\begin{aligned} &< \left\| \sum_{p < Y} \frac{X_p}{p^\sigma} \right\|_{2k} + \left\| \sum_{p > Y} \frac{X_p}{p^\sigma} \right\|_{2k} \\ &= \left(\sum_{r \in R} \frac{a_k(r)^2}{r^{2\sigma}} \right)^{\frac{1}{2k}} + \left(\sum_{s \in S} \frac{a_k(s)^2}{s^{2\sigma}} \right)^{\frac{1}{2k}} \end{aligned}$$

By Lemmas 1 and 2 this is

$$\ll \frac{Y^{1-\sigma}}{\log Y} + \frac{k^{\frac{1}{2}} Y^{\frac{1}{2}-\sigma}}{(\log Y)^{\frac{1}{2}}}$$

and it suffices to take $Y = k \log k$.

University of Michigan
Ann Arbor, Michigan 48109.