



HAL
open science

On the algebraic differential equations satisfied by some elliptic function I

P Chowla, S Chowla

► **To cite this version:**

P Chowla, S Chowla. On the algebraic differential equations satisfied by some elliptic function I . Hardy-Ramanujan Journal, 1984, Volume 7 - 1984, pp.11-12. 10.46298/hrj.1984.106 . hal-01104327

HAL Id: hal-01104327

<https://hal.science/hal-01104327>

Submitted on 16 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE ALGEBRAIC DIFFERENTIAL EQUATIONS SATISFIED BY SOME ELLIPTIC FUNCTIONS (I)

By

P. CHOWLA AND S. CHOWLA

Summary

When a is an odd positive integer it is implicit in the work of Jacobi that the functions

$$Y = \sum_{n=1}^{\infty} \sigma_a(n) X^n$$

where

$$\sigma_a(n) = \sum_{d|n} d^a$$

(the sum of the a th powers of the divisors of n) satisfy an algebraic differential equation i. e. there is a polynomial T , not identically 0, such that $T(X, Y, Y_1, \dots, Y_m) = 0$. When $a=1$ we give a new argument based on Ramanujan that we may take $m = 3$ here.

§ 1. With Ramanujan (Coll. Papers, pp 138-142) we write

$$P = 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) X^n$$

$$Q = 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) X^n$$

$$R = 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) X^n$$

Then we have

$$(\alpha) \quad X \frac{dP}{dX} = \frac{P^2 - Q}{12}$$

$$(\beta) \quad X \frac{dQ}{dX} = \frac{PQ - R}{3}$$

$$(\gamma) \quad X \frac{dR}{dX} = \frac{PR - Q^2}{2}$$

Now differentiate (α) twice w.r.t. "X". The first step gives us

$$(\delta) \quad \frac{d^2P}{dX^2} \text{ in terms of } P, Q, R, X.$$

Another differentiation gives us

$$(\epsilon) \quad \frac{d^3P}{dX^3} \text{ in terms of } P, Q, R, X.$$

Now eliminate Q,R from the 3 equations (α) , (δ) , (ϵ) .

This relation (which can be converted into a polynomial relation) gives us

$$T\left(X, P, \frac{dP}{dX}, \frac{d^2P}{dX^2}, \frac{d^3P}{dX^3}\right) = 0$$

Here T is a polynomial (not identically 0) in the 5 quantities

X, P, P_m ($1 \leq m \leq 3$) where P_m stands for $\frac{d^m P}{dX^m}$.

Thus we have a third order alg. diff. eqn. satisfied by P, and

therefore by $\sum_1^{\infty} \sigma_1(n) X^n$. We also risk a *conjecture*.

The function

$$Y = \sum_1^{\infty} \sigma_a(n) X^n$$

does *not* satisfy an alg. differential equation when a is an *even* positive integer.

We have been unable to prove anything in this direction.