



HAL
open science

On algebraic differential equations satisfied by some elliptic functions II

P Chowla, S Chowla

► **To cite this version:**

P Chowla, S Chowla. On algebraic differential equations satisfied by some elliptic functions II. Hardy-Ramanujan Journal, 1984, Volume 7 - 1984, pp.13 - 16. 10.46298/hrj.1984.107 . hal-01104334

HAL Id: hal-01104334

<https://hal.science/hal-01104334v1>

Submitted on 16 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON ALGEBRAIC DIFFERENTIAL EQUATIONS SATISFIED BY SOME ELLIPTIC FUNCTIONS (II)

By

P. CHOWLA and S. CHOWLA

Summary :

In (I) we obtained the "implicit" algebraic differential equation for the function defined by

$$Y = \sum_1^{\infty} \frac{n^a x^n}{1 - x^n}$$

where a is an odd positive integer, and conjectured that there are no algebraic differential equations for the case when a is an even integer. In this note we obtain a simple proof that (this has been known for almost 200 years)

$$Y = \sum_1^{\infty} x^{n^2} \quad (|x| < 1)$$

satisfies an algebraic differential equation, and conjecture that

$$Y = \sum_1^{\infty} x^{n^k} \quad (k \text{ a positive integer } > 2)$$

does not satisfy an algebraic differential equation.

§1 In the standard notation of elliptic functions we write

$$K = \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}$$

$$E = \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \phi} \, d\phi$$

Here $0 < k < 1$. We also define k' by

$$k^2 + k'^2 = 1$$

and the corresponding integrals

$$K' = \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - k'^2 \sin^2 \phi}}$$

$$E' = \int_0^{\pi/2} \sqrt{1 - k'^2 \sin^2 \phi} \, d\phi$$

We know that

$$KE' + K'E - KK' = \frac{\pi}{2}$$

which is often called Legendre's relation.

Our proof rests on several identities for K , K' , E , E' and their derivatives. Most important :

$$\sqrt{\frac{2k}{\pi}} = \sum_{-\infty}^{\infty} q^{n^2}$$

where

$$q = e^{-\frac{\pi K'}{k}}$$

§ 2. Next we need:

$$\frac{dK}{dk} = \frac{E - k'^2 K}{k k'^2},$$

$$\frac{dE}{dk} = \frac{E - K}{k}$$

Since

$$\frac{dk'}{dk} = -\frac{k}{k'}$$

we easily express

$$\frac{d E'}{d k} , \frac{d K'}{d k}$$

as easy functions of K, K', E, E' and k . From these

$$\frac{d}{d k} (E K' + K E' - K K') = 0$$

Hence

$$E K' + K E' - K K' = c$$

where c is a constant independent of k .

We easily see that $c = \frac{\pi}{2}$.

§ 3. Next

$$\frac{d q}{d k} = \frac{\pi^2 q}{2k k'^2 K^2}$$

(Ramanujan, Collected Papers, p. 32)

§ 4. Next find

$$\frac{d K}{d q} = \frac{d K}{d k} \frac{d k}{d q}$$

which is, from our previous relations a simple rational function of K, E, k, q .

Form

$$\frac{d^2 K}{d q^2} \text{ and } \frac{d^3 K}{d q^3}$$

which again are simple rational functions of K, E, k, q . Eliminate E and k from the relations obtained in this section.

We then get a "polynomial" relation between

$$K, q, \frac{dK}{dq}, \frac{d^2K}{dq^2}, \frac{d^3K}{dq^3}$$

which is a third order algebraic differential equation

$$T\left(K, q, \frac{dK}{dq}, \frac{d^2K}{dq^2}, \frac{d^3K}{dq^3}\right) = 0$$

where T is a polynomial in the five quantities following T .

§ 5. It would be nice to obtain "explicitly" the polynomial T of the last section, as well as those mentioned in the beginning of this paper i. e. algebraic differential equations satisfied by

$$Y = \sum_{n=1}^{\infty} \frac{n^a x^n}{1-x^n}$$

i. e. a relation with an explicit polynomial T with

$$T(Y, X, Y', Y'', Y''') = 0$$

As far we know this has never been done.

Conclusion.

Professor Dwork (Princeton University) pointed out to us that G. Valerion in a monograph (in French) on automorphic forms in analysis obtained the algebraic differential equation explicitly in the case of the function

$$Y = \sum_{n=1}^{\infty} \pi (1 - x^n).$$