



HAL
open science

A remark on goldbach's problem II

S Srinivasan

► **To cite this version:**

S Srinivasan. A remark on goldbach's problem II. Hardy-Ramanujan Journal, 1984, Volume 7 - 1984, pp.17 - 20. 10.46298/hrj.1984.110 . hal-01104342

HAL Id: hal-01104342

<https://hal.science/hal-01104342v1>

Submitted on 16 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A REMARK ON GOLDBACH'S PROBLEM II

By

S. SRINIVASAN

For $A > 0$, even integer $N (> 1)$, set

$$E_A(N) = \sum_{q \leq (\log N)^A} \frac{\mu^2(q) c_q(-N)}{\phi^2(q)}$$

Here, $c_q(n)$ is the Ramanujan's sum ; one evaluation is

$$(1) \quad c_q(-N) = \frac{\mu(q/(q, N)) \phi(q)}{\phi(q/(q, N))}$$

R. Balasubramanian and C. J. Mozzochi proved (cf. § 6 of [1]) the following theorem.

Theorem.

For any fixed $A (> 0)$, the relation

$$(2) \quad E_A(N) \sim E_\infty(N), N \rightarrow \infty,$$

is false.

Now we give an alternative proof of this theorem based on

$$(3) \quad E_\infty(N) - E_A(N) = O(E_\infty(N)/A), E_\infty(N) \asymp N/\phi(N),$$

proved in [2].

Remark.

In view of the above theorem, it is necessary to state the Theorem in [2] with the factor $N/\phi(N)$ in the hypothesis; i. e., the bound

$$\delta_B \frac{N}{(\log N)^2} \text{ replaced by } \delta_B \frac{N}{(\log N)^2} \frac{N}{\phi(N)}.$$

This can be done with no change in the proof there.

We require also the following

Lemma.

Let $\alpha > 0$. Then, as $x \rightarrow \infty$,

$$|\{n < x : \mu(n) \neq 0, p | n \Rightarrow p \leq x^\alpha\}| > c_\alpha x,$$

holds with some $c_\alpha > 0$.

Remark

Actually, the above number has a well-known asymptotic formula. However, we give here a direct simple proof of this Lemma, as it is sufficient for our present purpose.

Proof of the Lemma

Clearly, it suffices to prove the Lemma for a sequence $\alpha = \alpha_k \rightarrow 0$ (as $k \rightarrow \infty$). We choose $\alpha_k = 3/(5k+4)$; $k = 1, 2, \dots$. Consider $0 < \frac{2}{3}\alpha_k = \delta_0 < \dots < \delta_R = \alpha_k$ with $R = 2k+1$; θ defined through $\delta_j = \delta_0 + \theta j$ ($j = 0, 1, \dots, R$).

Note that

$$\begin{aligned} \delta_1 + \dots + \delta_{R-1} &= (R-1)\delta_0 + \frac{R-1}{2}R\theta \\ &= \frac{R-1}{2}(2\delta_0 + \delta_R - \delta_0), \end{aligned}$$

and on inserting the values of R and δ_R in terms of k we see that

$$(4) \quad 1 - 2\delta_0 = \delta_1 + \dots + \delta_{R-1} < 1 - (\delta_R + \delta)$$

with any fixed ($0 <$) $\delta < \frac{1}{2}\delta_0$.

Let q^* denote a typical product $\prod_{j=1}^R p_j$ with primes p_j

satisfying $x^{\delta_{j-1}} < p_j < x^{\delta_j}$. Now, from (4), $x^{1-\delta} > q^* > x^{1-\delta_0}$. Letting $q' < x/q^*$ run through square-free values, we see that $q^*q' < x$ are distinct square-free numbers, and their number is

$$\gg \sum \frac{x}{q^*} = x \prod_{j=1}^R \left(\sum \frac{1}{p_j} \right) > c'_k x$$

with some $c'_k > 0$. This completes the proof of the Lemma.

Proof of the Theorem.

In view of (3), it is sufficient to consider a lower bound for

$$(3') \quad E_{A,A'}(N) = E_{A',(N)} - E_A(N)$$

with a suitably large A' ($> 2A$, say). Now, we restrict N to the sequence

$$(5) \quad N_m = \prod_{p < \log m} p, \quad (\log N_m \sim \log m)$$

and note that if a square-free $q \times N$, then $q / (q, N) \gg \log N$. Thus, by (1), we see the contribution of $q \times N$ to $E_{A,A'}(N)$ is

$$O \left(\sum' \frac{\mu^2(q)}{\phi(q)} \frac{1}{\sqrt{\log N}} \right), \text{ say,}$$

where ' denotes the restriction on q in (3'). So this contribution is

$$(6) \quad O_{A,A'}(\log \log N / \sqrt{\log N}) = o(E_{\infty}(N)), \quad N \rightarrow \infty,$$

since, for N in (5), $E_{\infty}(N) \asymp \log \log N$, by (3). Next, the (remaining) contribution of $q \mid N$ to (3') is

$$\sum_{q|N} \frac{\mu^2(q)}{\phi(q)} > (\log N)^A < q \leq (\log N)^{A'}$$

$$\sum_{q|N} \frac{\mu^2(q)}{q} > (\log N)^A < q \leq (\log N)^{2A}$$

and this is easily seen (by (5) and Lemma) to be $> c_A^* \log \log N$. The proof is completed by a suitably large choice of A' (in terms of c_A^*) via (3) and (6).

References

- [1] R. Balasubramanian and C. J. Mozzochi, *Siegel zeros and the Goldbach Problem*, J. Number Theory, 16 (1983), 311-332.
- [2] S. Srinivasan, *A Remark on Goldbach Problem*, J. Number Theory, 12 (1980), 116-121.

School of Mathematics
Tata Institute of Fundamental Research
Homi Bhabha Road
Bombay 400 005 (India)