



Hybrid mean square of L -functions.

Mangala J Narlikar

► To cite this version:

Mangala J Narlikar. Hybrid mean square of L -functions.. Hardy-Ramanujan Journal, 1986, Volume 9 -1986, pp.11-16. 10.46298/hrj.1986.116 . hal-01104702

HAL Id: hal-01104702

<https://hal.science/hal-01104702v1>

Submitted on 19 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HYBRID MEAN SQUARE OF L-FUNCTIONS

By M. J. Narlikar

§1. Introduction. Let $k \geq 1$ be an integer and χ run through the set of characters mod k . Let $T \geq 0$ and $T_0 = T+2$. Then R. Balasubramanian and K. Ramachandra proved the following theorem. For the history of the theorem see [1].

Theorem 1. We have

$$\frac{1}{2\pi\phi(k)} \sum_{\chi \text{ mod } k} \int_0^T |L(\frac{1}{2}+it, \chi)|^2 dt = \frac{\phi(k)}{k^2} A(k, T) + O\left(\frac{\phi(k)}{k} T^{1/2} \log T_0\right)$$

where $A(k, T) = \frac{kT}{2\pi} \log \frac{kT}{2\pi} + \left[2\gamma - 1 + \sum_{p|k} \frac{\log p}{p-1} \right] \frac{kT}{2\pi}$, γ being the

Euler's constant.

In this paper I prove the following

Theorem 2. For $T \geq 2$ and $k \geq 3$,

$$\frac{1}{2\pi\phi(k)} \sum_{\chi \text{ mod } k} \int_0^T |L(\frac{1}{2}+it, \chi)|^2 dt = \frac{\phi(k)}{k^2} A(k, T) +$$

$$+ B(k) T^{1/2} + O\left(\frac{\phi(k)}{k} T^{5/12} (\log T)^2\right)$$

where $B(k)$ is a negative constant and depends only on k . Moreover

$$|B(k)| \gg \frac{\phi(k)}{k} \quad \text{and} \quad |B(k)| \ll \frac{\phi(k)}{k}.$$

§2. Proof of the Theorem 2.

Lemma 1. We have

$$\frac{1}{\phi(k)} \sum_{\chi \bmod k} \left| L\left(\frac{1}{2} + it, \chi\right) \right|^2 = \frac{1}{k} \sum_{\substack{(a,k)=1 \\ 1 \leq a \leq k}} \left| \zeta\left(\frac{1}{2} + it, \frac{a}{k}\right) \right|^2$$

where $\zeta(s, \alpha)$ for $0 < \alpha < 1$ is the Hurwitz Zeta-function defined by

$$\zeta(s, \alpha) = \sum_{n=0}^{\infty} (n+\alpha)^{-s} \quad \text{for } \operatorname{Re} s > 1.$$

Proof. Follows from the fact that if $F(a)$ is a complex number for every residue class $a \bmod k$ with $(a, k) = 1$, then

$$\frac{1}{\phi(k)} \sum_{\chi \bmod k} \left| \sum_{\substack{(a,k)=1 \\ 1 \leq a \leq k}} \chi(a) F(a) \right|^2 = \sum_{\substack{(a,k)=1 \\ 1 \leq a \leq k}} |F(a)|^2.$$

Lemma 2. Theorem 1 gives Theorem 2 for $1 \leq T \leq 2$ and hence it is sufficient to prove that for $T \geq 2$

$$\begin{aligned} \frac{1}{2\pi\phi(k)} \sum_{\chi \bmod k} \int_1^T \left| L\left(\frac{1}{2} + it, \chi\right) \right|^2 dt &= \frac{\phi(k)}{k^2} \{A(k, T) - A(k, 1)\} + \\ &\quad + B(k) T^{1/2} + O\left(\frac{\phi(k)}{k} T^{5/12} (\log T)^2\right). \end{aligned}$$

where $B(k)$ is as in Theorem 2.

Lemma 3. For $T \geq 2$, $0 < \alpha < 1$,

$$\begin{aligned} \frac{1}{2\pi} \int_1^T \left| \zeta\left(\frac{1}{2} + it, \alpha\right) \right|^2 dt &= \frac{T}{2\pi} \log \frac{T}{2\pi} + \frac{T}{2\pi} [C(\alpha) + \gamma - 1] - \frac{i}{2\pi\alpha} + A_1(\alpha) T^{1/2} + \\ &\quad + O(T^{5/12} (\log T)^2) + O\left(\frac{\log T}{(1/2\alpha)^0}\right). \end{aligned}$$

MEAN SQUARE OF L-FUNCTIONS

where $\|2\alpha\|$ denotes the distance of 2α from the nearest integer and

$$C(\alpha) = \lim_{N \rightarrow \infty} \left[\sum_{n=1}^N \frac{1}{(n+\alpha)} - \log N \right] \quad \text{and}$$

$$A_1(\alpha) = \frac{1}{\sqrt{2\pi}} \left[-2H(\alpha) + \int_{1-\alpha}^1 \frac{\cos\{\pi(u^2-u-1/4)\}}{\cos \pi u} du + 2 \sum_{r=1}^{\infty} \int_r^{\infty} \cos\{\pi(u^2-1/4)\} du \right],$$

$$H(\alpha) = \int_{\frac{1-\alpha}{2}}^{\frac{1+\alpha}{2}} G^2(u) du + \int_{\alpha/2}^{1-\alpha/2} G^2(u) du$$

$$\text{where } G(u) = \frac{\cos\{2\pi(u^2-u-1/16)\}}{\cos(2\pi u)}.$$

$$\text{Finally, } A_1(\alpha) < -\frac{1}{\sqrt{2\pi}} H(\alpha)$$

Proof. See [2].

Lemma 4. We have

$$B(k) = \frac{1}{\sqrt{2\pi} k} \sum_{\substack{(a,k)=1 \\ 1 \leq a \leq k}} \left[-2H\left(\frac{a}{k}\right) + \int_{1-\frac{a}{k}}^1 \frac{\cos\{\pi(u^2-u-1/4)\}}{\cos \pi u} du \right]$$

$$+ 2 \sum_{r=1}^{\infty} \int_r^{\infty} \cos\{\pi(u^2-1/4)\} du \Big]$$

$$\text{and } |B(k)| \ll \frac{\phi(k)}{k}.$$

Proof. Follows from the expression for $A_1(\alpha)$ and the last part of

Lemma 3.

Lemma 5. In Lemma 3, the error term $O\left(\frac{\log T}{\|2\alpha\|}\right)$ can be replaced by $O\left[\min\left\{\frac{\log T}{\|2\alpha\|}, T^{1/2} \log T\right\}\right]$.

Proof. The sums in Lemmas 11 to 14 of [2] which contribute to this O-term add up to

$$2 \operatorname{Re} \sum_{m=1}^{\lfloor \sqrt{T}/2\pi \rfloor} \sum_{0 \leq r \leq m} \frac{1}{1/4} \operatorname{Exp}\left[2\pi i(2m\alpha + 2r\alpha + \alpha^2)\right] \int_{r+\alpha}^{\infty} \operatorname{Exp}\left[i\pi\left(\theta^2 - \frac{1}{4}\right)\right] d\theta.$$

Now it is easy to see that this sum is $O(T^{1/2} \log T)$. We only have to integrate by parts and obtain an estimate $O(\frac{1}{r+\alpha})$ for $r \geq 1$ and $O(1)$ for $r=0$, for the expression inside the summation sign.

Lemma 6. We have

$$\frac{1}{k} \sum_{\substack{(a,k)=1 \\ 1 \leq a \leq k}} \min\left[\frac{\log T}{\|2a/k\|}, T^{1/2} \log T\right] \ll \epsilon \frac{\phi(k)}{k} T^\epsilon.$$

Proof. Put $S = \frac{1}{\phi(k)} \sum_{\substack{(a,k)=1 \\ 1 \leq a \leq k}} \min\left[\frac{\log T}{\|2a/k\|}, T^{1/2} \log T\right]$

$$\text{and } S_U = \frac{1}{\phi(k)} \sum_{\substack{1/2U \leq \|2a/k\| < 1/U \\ (a,k)=1, 1 \leq a \leq k}} \min\left[\frac{\log T}{\|2a/k\|}, T^{1/2} \log T\right]$$

where U runs through powers 2^r of 2, $2^r < k$. We divide the totality of S_U into 3 parts according as the integer nearest to $\frac{2a}{k}$ is 0, 1 or 2. In each case the number of integers a such that $\frac{1}{2U} \leq \|2a/k\| \leq \frac{1}{U}$ is $\ll \frac{\varphi(k)}{U} + 2^{\omega(k)}$, where $\omega(k)$ is the number of different prime factors of k . Hence, if $U \leq X$, taking the first term in the

MEAN SQUARE OF L-FUNCTIONS

minimum,

$$S_U \ll \left[1+U \frac{2^{\varphi(k)}}{\varphi(k)} \right] \log T.$$

For $U > X$, we take the second term in the minimum, so that

$$S_U \ll T^{1/2} \log T \left[\frac{1+2^{\omega(k)}}{U \varphi(k)} \right].$$

Now $2^{\omega(k)} \ll k^\epsilon$, hence

$$\frac{S}{\log T} \ll_\epsilon \log X + \frac{X}{X^{1-\epsilon}} + \frac{T^{1/2}}{k^{1-\epsilon}}.$$

If $k < T^{10}$ we take $X=k$ and we do not get the last two terms as the case $U > X$ does not arise. If $k \geq T^{10}$, take $X=T^{1/2}$ so that

$$\frac{S}{\log T} \ll T^\epsilon \text{ in either case.}$$

This proves the Lemma and Theorem 2 is now completely proved.

Acknowledgement. The author wishes to express her gratitude to Professor K. Ramachandra for encouragement.

References

- [1] R. Balasubramanian and K. Ramachandra, "A Hybrid version of a theorem of Ingham", Number Theory Proceedings, Edited by K. Alladi, Lecture notes in mathematics, 1122, Springer-Verlag (1984), 38-46.
- [2] M.J. Narlikar, "On the mean square value of Hurwitz zeta-function", Proc. Indian Acad. Sci. 90 (1981), 195-212.

M. J. NARLIKAR

701 Bhaskara
TIFR Housing Colony
Bombay 400 005

ADDED IN THE END. Since the submission of this paper the following paper has appeared :- Tom Meurman, A generalisation of Atkinson's formula to L-functions, Acta Arith., XLVII (1986), p.351-370.