



HAL
open science

Proof of some conjectures on the mean-value of titchmarsh series with applications to Titchmarsh's phenomenon

K Ramachandra

► **To cite this version:**

K Ramachandra. Proof of some conjectures on the mean-value of titchmarsh series with applications to Titchmarsh's phenomenon. Hardy-Ramanujan Journal, 1990, Volume 13 - 1990, pp.21 - 27. 10.46298/hrj.1990.119 . hal-01104713

HAL Id: hal-01104713

<https://hal.science/hal-01104713v1>

Submitted on 19 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hardy-Ramanujan Journal
Vol.13 (1990) 21-27

PROOF OF SOME CONJECTURES ON THE MEAN-VALUE
OF TITCHMARSH SERIES WITH APPLICATIONS TO
TITCHMARSH'S PHENOMENON

BY

K. RAMACHANDRA

§ 1. INTRODUCTION. This is a continuation of [R]₁ but no previous knowledge of this paper is necessary. In fact we improve these results a good deal. However for some applications of the main results of the present paper, a knowledge of the results of [BRS] is assumed. The main results of the present paper are the following two theorems on what I call weak Titchmarsh series. We begin with a definition.

WEAK TITCHMARSH SERIES. Let $0 \leq \epsilon < 1, D \geq 1, C \geq 1$ and $H \geq 10$. Put $R = H^\epsilon$. Let $a_1 = \lambda_1 = 1$ and $\{\lambda_n\}$ ($n = 1, 2, 3, \dots$) be any sequence of real numbers with $\frac{1}{C} \leq \lambda_{n+1} - \lambda_n \leq C$ ($n = 1, 2, 3, \dots$) and $\{a_n\}$ ($n = 1, 2, 3, \dots$) any sequence of complex numbers satisfying

$$\sum_{\lambda_n \leq X} |a_n| \leq D(\log X)^R$$

for all $X \geq 3C$. Then for complex $s = \sigma + it$ ($\sigma > 0$) we define the analytic function $F(s) = \sum_{n=1}^{\infty} a_n \lambda_n^{-s}$ as a weak Titchmarsh series associated with the parameters occurring in the definition.

THEOREM 1 (FOURTH MAIN THEOREM). For a weak Titchmarsh series $F(s)$ with $H \geq 36C^2 H^\epsilon$, we have

$$\liminf_{\sigma \rightarrow +0} \int_0^H |F(\sigma + it)| dt \geq H - 36C^2 H^\epsilon - 12CD.$$

THEOREM 2 (FIFTH MAIN THEOREM). For a weak Titchmarsh series $F(s)$ with $\log H \geq 4320 C^2(1 - \varepsilon)^{-5}$, we have,

$$\liminf_{\sigma \rightarrow +0} \int_0^H |F(\sigma + it)|^2 dt \geq \sum_{n \leq M} \left(H - \frac{H}{\log H} - 100C^2 n \right) |a_n|^2 - 2D^2$$

where $M = (36C^2)^{-1} H^{1-\varepsilon} (\log H)^{-4}$.

REMARKS. Theorems 1 and 2 have been referred to as the fourth and the fifth main theorems in $[R]_2$. Also we remark that it is not difficult to improve the conditions in the theorems slightly.

§ 2. **PROOF OF THEOREM 1.** We can argue with $\sigma > 0$ and then pass to the limit as $\sigma \rightarrow +0$. But formally the notation is simplified if we treat as though $F(s)$ is convergent absolutely if $\sigma = 0$ and there is no loss of generality. Let r be a positive integer and $0 < U \leq r^{-1}H$. Then since $|F(s)| \geq 1 + \operatorname{Re}(F(s))$, we have (with $\lambda = u_1 + \dots + u_r$),

$$\begin{aligned} \int_0^H |F(it)| dt &\geq U^{-r} \int_0^U du_r \dots \int_0^U du_1 \int_\lambda^{H-rU+\lambda} |F(it)| dt \\ &\geq U^{-r} \int_0^U du_r \dots \int_0^U du_1 \int_\lambda^{H-rU+\lambda} \{1 + \operatorname{Re}(F(it))\} dt \\ &\geq H - rU - 2^{r+1} U^{-r} J \end{aligned}$$

where $J = \sum_{n=2}^{\infty} |a_n| (\log \lambda_n)^{-r-1}$. Now $J = S_0 + \sum_{j=1}^{\infty} S_j$ where $S_0 = \sum_{\lambda_n \leq 3C} |a_n| (\log \lambda_n)^{-r-1}$ and $S_j = \sum_{3^j C \leq \lambda_n \leq 3^{j+1} C} |a_n| (\log \lambda_n)^{-r-1}$. In S_0 we use $\lambda_n \geq \lambda_2 \geq 1 + C^{-1}$ and so $(\log \lambda_n)^{-r-1} \leq (2C)^{r+1}$ and we obtain $S_0 \leq D(2C)^{r+1}(3C)^R$. Also, we have,

$$\begin{aligned} S_j &\leq D(\log(3^{j+1}C))^R (\log(3^j C))^{-r-1} \\ &\leq D2^R (\log(3^j C))^{R-r-1}, \text{ (since } 3^{j+1}C \leq (3^j C)^2\text{),} \\ &\leq D2^R j^{-2} \text{ by fixing } r = [3R]. \end{aligned}$$

Thus for $r = [3R]$ we have

$$\begin{aligned} J &\leq D(2C)^{r+1}(3C)^R + 2D2^R, \text{ (since } \sum_{j=1}^{\infty} j^{-2} < 2\text{),} \\ &\leq 3D(2C)^{r+1}(3C)^R. \end{aligned}$$

Collecting we have,

$$\begin{aligned} 2^{r+1}U^{-r}J &\leq 12CD(3C)^R\left(\frac{4C}{U}\right)^r \\ &\leq 12CD\left(\frac{12C^2}{U}\right)^R \text{ if } U \geq 4C \\ &\leq 12CD \text{ by fixing } U = 12C^2. \end{aligned}$$

The only condition which we have to satisfy is $rU \leq H$ which is secured by $H \geq 36C^2H^\epsilon$. This completes the proof of Theorem 1.

§ 3. PROOF OF THEOREM 2. We write $\lambda = u_1 + \dots + u_r$, where $0 \leq u_i \leq U$ and $0 < U \leq r^{-1}H$. We put $M_1 = [M]$, $A(s) = \sum_{m \leq M_1} a_m \lambda_m^{-s}$ and $B(s) = \sum_{n \geq M_1+1} a_n \lambda_n^{-s}$ so that $F(s) = A(s) + B(s)$. For the moment we suppose M to be a free parameter with the restriction $3 \leq M \leq H$. We use

$$|F(it)|^2 \geq |A(it)|^2 + 2 \operatorname{Re}(A(it)\overline{B(it)}).$$

Now by a well-known theorem of H.L. Montgomery and R.C. Vaughan we have

$$\int_{\lambda}^{H-rU+\lambda} |A(it)|^2 dt \geq \sum_{n \leq M} (H - rU - 100C^2n) |a_n|^2$$

Next the absolute value of

$$2U^{-r} \int_0^U du_r \dots \int_0^U du_1 \int_{\lambda}^{H-rU+\lambda} (A(it)\overline{B(it)}) dt \quad (3.1)$$

does not exceed

$$\begin{aligned} &2^{r+2}U^{-r} \sum_{m \leq M_1, n \geq M_1+1} |a_m \bar{a}_n| \left(\log \frac{\lambda_n}{\lambda_m}\right)^{-r-1} \\ &\leq 2^{r+2}U^{-r} \left(\sum_{m \leq M_1} |a_m| \right) \left(\sum_{n \geq M_1+1} |a_n| \left(\log \frac{\lambda_n}{\lambda_{M_1}}\right)^{r+1} \right) \end{aligned}$$

Here the m -sum is $\leq D(\log \lambda_{M_1})^R \leq D(\log(3MC))^R$, since $\lambda_{M_1} \leq M_1C \leq MC$. It is enough to choose $M \geq 1$ for the bound for the m -sum. The n -sum can be broken up into $\lambda_n \leq 3\lambda_{M_1}$ and $3^j\lambda_{M_1} < \lambda_n \leq 3^{j+1}\lambda_{M_1}$ ($j =$

1, 2, 3, ...). Let us denote these sums by S_0 and S_j . Now since $\left(\log \frac{\lambda_n}{\lambda_{M_1}}\right) \geq \left(\log \frac{\lambda_{M_1+1}}{\lambda_{M_1}}\right) \geq \log \left(1 + \frac{1}{C\lambda_{M_1}}\right) \geq (2C\lambda_{M_1})^{-1} \geq (2C^2M)^{-1}$, we obtain

$$S_0 \leq D(\log(3\lambda_{M_1}))^R (2C^2M)^{r+1} \leq D(\log(3MC))^R (2C^2M)^{r+1}.$$

Also

$$\begin{aligned} S_j &\leq D(\log(3^{j+1}\lambda_{M_1}))^R (j \log 3)^{-r-1} \\ &\leq D(j \log 3 + \log(3MC))^R j^{-r-1} \\ &\leq 2^R D(j \log 3)^R (\log(3MC))^R j^{-r-1} \\ &\leq 4^R D(\log(3MC))^R j^{-2}, \text{ if } r \geq R+1, \end{aligned}$$

and so (since $\sum_{j=1}^{\infty} j^{-2} < 2$),

$$\left(\sum_m \dots\right) \left(\sum_n \dots\right) \leq D^2(\log(3MC))^R (\log(3MC))^R Y$$

(where $Y = (2C^2M)^{r+1} + 2(4^R)$)

$$\leq D^2(\log(3MC))^{2R} ((2C^2M)^{r+1} + 2(4^R)).$$

Hence the absolute value of the expression (3.1) does not exceed

$$\begin{aligned} &D^2(\log(3MC))^{2R} \left((8C^2M) \left(\frac{4C^2M}{U}\right)^r + 2 \left(\frac{4^R}{U^r}\right) \right) \quad (3.2) \\ &\leq D^2 \left\{ 8C^2M \left(\frac{4C^2M(\log(3MC))^2}{U}\right)^{R+\log(8C^2M)} + 2 \left(\frac{4}{U}\right)^R \right\} \end{aligned}$$

if $U \geq 4C^2M$ and $r \geq R + \log(8C^2M)$. We put $U = 12C^2M(\log(3MC))^2$ and obtain for (3.2) the bound $D^2\{1+1\} \leq 2D^2$. The conditions to be satisfied are $M \geq 1$ and

$$12C^2M(\log(3MC))^2(R + \log(8C^2M) + 1) \leq H.$$

In fact we can satisfy $Ur \leq \frac{H}{\log H}$ by requiring

$$12C^2M(\log(3MC))^2(R + \log(8C^2M) + 1) \leq \frac{H}{\log H}.$$

This is satisfied if

$$36C^2M(\log(8C^2M))^3R \leq H(\log H)^{-1}$$

Let $8C^2M \leq H$. Then $36C^2MR \leq H(\log H)^{-4}$ gives what we want. We choose $M = (36C^2)^{-1}H^{1-\varepsilon}(\log H)^{-4}$. Clearly this satisfies $8C^2M \leq H$. In order to satisfy $M \geq 1$ we have to secure that

$$(36C^2)^{-1} \frac{((1-\varepsilon)(\log H))^5}{120} (\log H)^{-4} \geq 1$$

i.e. $\log H \geq 4320C^2(1-\varepsilon)^{-5}$.

This completes the proof of Theorem 2.

§ 4. APPLICATIONS OF THEOREMS 1 AND 2. An immediate application of Theorem 1 is

THEOREM 3. Let $\zeta(s, a) = \sum_{n=0}^{\infty} (n+a)^{-s}$ (where $0 < a \leq 1$) be the Hurwitz zeta-function in $\sigma > 1$ and consider its analytic continuation in $\sigma \geq 1$. Then

$$\min_{T \geq 1} \int_T^{T+H} |\zeta(1+it, a)| dt \geq \frac{1}{a}H + o(H).$$

Let $(\zeta(s))^u = \sum_{n=1}^{\infty} d_u(n)n^{-s}$ where u is any complex constant. Consider the analytic continuation of $(\zeta(s))^u$ in $\sigma \geq 1, t \geq 1$. An immediate corollary to Theorem 2 is

THEOREM 4. We have,

$$\min_{T \geq 1} \left(\frac{1}{H} \int_T^{T+H} |\zeta(1+it)^u|^2 dt \right) \geq \sum_{n=1}^{\infty} \frac{|d_u(n)|^2}{n^2} + o(1),$$

and in particular for $u = 1$, we have,

$$\min_{T \geq 1} \left(\frac{1}{H} \int_T^{T+H} |\zeta(1+it)|^2 dt \right) \geq \frac{\pi^2}{6} + o(1).$$

It is possible to prove by using Theorem 2 a very nice Ω theorem for $|\zeta(1+it)^z|$, where $z = e^{i\theta}$ ($0 \leq \theta < 2\pi, \theta$ fixed) as $t \rightarrow \infty$. It is

THEOREM 5. *We have,*

$$\min_{T \geq 1} \max_{T \leq t \leq T+H} |(\zeta(1+it))^z| \geq e^{\gamma} \lambda(\theta) (\log \log H - \log \log \log H) + O(1), \quad (4.1)$$

where

$$\lambda(\theta) = \prod_p \left\{ \left(1 - \frac{1}{p}\right) \left(\frac{\sqrt{p^2 - \sin^2 \theta} + \cos \theta}{p - p^{-1}} \right)^{\cos \theta} \operatorname{Exp} \left(\sin \theta \sin^{-1} \left(\frac{\sin \theta}{p} \right) \right) \right\}$$

PROOF. By Theorem 2 we have with $\varepsilon = \frac{1}{3}$, $u = kz$,

$$\frac{1}{H} \int_T^{T+H} |(\zeta(1+it))^u|^2 dt \geq \frac{1}{2} \sum_{n \leq H^{\frac{1}{2}}} \frac{|d_u(n)|^2}{n^2} \quad (4.2)$$

uniformly in $T \geq 1$, and k any positive integer satisfying $1 \leq k \leq \log H$, provided H exceeds an absolute constant. Denote by S the RHS in (4.2). Then $S^{\frac{1}{2k}}$ has been studied in [BRS] as a function of H as k runs over $1 \leq k \leq \log H$. It has been proved (by considering the maximum term of the sum in S) that

$$\max_{1 \leq k \leq \log H} \left(S^{\frac{1}{2k}} \right) \geq e^{\gamma} \lambda(\theta) (\log \log H - \log \log \log H) + O(1).$$

This completes the proof of Theorem 5.

These ideas are quite general (applicable to zeta and L -functions of algebraic number fields). For example for ordinary L -series $L(s) = L(s, \chi)$ where χ is a non-principal character *mod* q , we can prove

THEOREM 6. *We have,*

$$\min_T \max_{T \leq t \leq T+H} |(L(1+it))^z| \geq e^{\gamma} \lambda(\theta) \frac{\varphi(q)}{q} \{(\log \log H - \log \log \log H) + O(1)\}$$

uniformly in $q \geq 3$.

ACKNOWLEDGEMENT. The author is very much thankful to Professor R. Balasubramanian for encouragement.

REFERENCES

- [BRS] R. BALASUBRAMANIAN, K. RAMACHANDRA AND A. SANKARA-NARAYANAN, *On the frequency of Titchmarsh's phenomenon for $\zeta(s)$ -VIII* (to appear).
- [R₁] K. RAMACHANDRA, *On the frequency of Titchmarsh's phenomenon for $\zeta(s)$ -VII*, Annales Acad. Sci. Fenn. Ser AI, 14 (1989), 27-40.
- [R₂] K. RAMACHANDRA, *A short monograph on the Riemann zeta function* (a book to appear).

P.S. The theorems of § 4 can be stated for σ satisfying $|\sigma - 1| \leq \psi(t)$ for suitable $\psi(t) \rightarrow 0$ as $t \rightarrow \infty$.

ADDRESS OF THE AUTHOR

PROFESSOR K. RAMACHANDRA
SCHOOL OF MATHEMATICS
TATA INSTITUTE OF FUNDAMENTAL RESEARCH
HOMI BHABHA ROAD
BOMBAY 400 005
INDIA

MANUSCRIPT COMPLETED ON 21 OCTOBER 1990.