



HAL
open science

On the frequency of Titchmarsh's phenomenon for $\zeta(s)$ IX.

K Ramachandra

► **To cite this version:**

| K Ramachandra. On the frequency of Titchmarsh's phenomenon for $\zeta(s)$ IX.. Hardy-Ramanujan Journal, 1990, Volume 13 - 1990, pp.28 - 33. 10.46298/hrj.1990.120 . hal-01104718

HAL Id: hal-01104718

<https://hal.science/hal-01104718>

Submitted on 19 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hardy-Ramanujan Journal

Vol. 13 (1990), 28-33

ON THE FREQUENCY OF TITCHMARSH'S PHENOMENON
FOR $\zeta(s)$ -IX

BY

K. RAMACHANDRA

§ 1. INTRODUCTION. In a previous paper [R], I proved the following result on $\zeta(1+it)$.

THEOREM 1. Let $0 \leq \theta < 2\pi$, $z = e^{i\theta}$ and

$$f(H) = \min_{T \geq 1} \max_{T \leq t \leq T+H} |(\zeta(1+it))^z| \quad (1)$$

Then

$$f(H) \geq e^\gamma \lambda(\theta) (\log \log H - \log \log \log H) + O(1) \quad (2)$$

where $H \geq \text{Exp}(e^e)$,

$$\lambda(\theta) = \prod_p \lambda_p(\theta),$$

$$\lambda_p(\theta) = \left\{ \left(1 - \frac{1}{p}\right) \left(\sqrt{1 - \frac{\text{Sin}^2 \theta}{p^2}} - \frac{\text{Cos } \theta}{p} \right)^{-\text{Cos } \theta} \text{Exp} \left(\text{Sin } \theta \text{Sin}^{-1} \left(\frac{\text{Sin } \theta}{p} \right) \right) \right\}. \quad (3)$$

In the present paper I prove that

$$f(H) \leq e^\gamma \lambda(\theta) (\log \log H + \log \log \log H) + O(1). \quad (4)$$

This result together with Theorem 1 gives the following Theorem.

THEOREM 2. We have

$$|f(H)e^{-\gamma(\lambda(\theta))^{-1}} - \log \log H| \leq \log \log \log H + O(1), \quad (5)$$

where $H \geq \text{Exp}(e^e)$.

REMARK. It is interesting to prove (or disprove!) $f(H)e^{-\gamma(\lambda(\theta))^{-1}} =$

$\log \log H + O(1)$.

§ 2. PROOF OF (4). We begin by

LEMMA 1. *Let $T = \text{Exp}((\log H)^2)$ where H exceeds an absolute constant. Then there exists a sub-interval I of $[T, 2T]$ of length $H + 2(\log H)^{10}$, such that the rectangle $(\sigma \geq \frac{3}{4}, t \in I)$ does not contain any zero of $\zeta(s)$ and moreover*

$$\max |\log \zeta(\sigma + it)| = O((\log H)^{\frac{1}{4}} (\log \log H)^{-\frac{3}{4}}) \quad (6)$$

the maximum being taken over the rectangle referred to.

PROOF. Follows from [BR] and the result (due to A.E. Ingham [I], see also [T] page 236 and p. 293-295 [AI]) that the number of zeros of $\zeta(s)$ in $(\sigma \geq \frac{3}{4}, T \leq t \leq 2T)$ is $O(T^{\frac{1}{4}})$.

LEMMA 2. *Let J be the interval obtained by removing from I intervals of length $(\log H)^{10}$ from both ends. Then for $t \in J$, we have,*

$$\log \zeta(1 + it) = \sum_{m \geq 1} \sum_p (mp^{ms})^{-1} \text{Exp} \left(-\frac{p^m}{X} \right) + O((\log \log H)^{-1}) \quad (7)$$

where $X = \log H \log \log H$ and $s = 1 + it$.

PROOF. The lemma follows from the fact that the double sum on the right is

$$\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \log \zeta(s+w) X^w \Gamma(w) dw \quad (8)$$

where $w = u + iv$ is a complex variable. Here we break off the portion $|v| \geq (\log H)^9$ with an error $O((\log \log H)^{-1})$ and move the line of integration to $u = -\frac{1}{4}$. Using Lemma 1 it is easily seen that the horizontal portions and the main integral contribute together $O((\log \log H)^{-1})$.

LEMMA 3. *Denote the double sum in (7) by S . Then*

$$S = \log \prod_{p \leq X} (1 - p^{-s})^{-1} + O((\log \log H)^{-1}). \quad (9)$$

PROOF. We use the fact that $\text{Exp}(-p^m X^{-1}) = 1 + O(p^m X^{-1})$ if $p^m \leq X$ and $= O(X p^{-m})$ if $p^m \geq X$. Using this it is easy to see that

$$\begin{aligned} S &= \sum_{p^m \leq X} \sum_{p^m \leq X} (m p^{ms})^{-1} + O\left(\sum_{p^m \leq X} \sum_{p^m \leq X} X^{-1}\right) + O\left(\sum_{p^m \geq X} \sum_{p^m \geq X} X(m p^{2m})^{-1}\right) \\ &= \sum_{p^m \leq X} \sum_{p^m \leq X} (m p^{ms})^{-1} + O((\log \log H)^{-1}). \end{aligned}$$

Denoting the last double sum by S_0 , we have,

$$S_0 - \sum_{p \leq X} \log(1 - p^{-s})^{-1} = O\left(\sum_{p^m \geq X, m \geq 2} \sum_{p^m \geq X, m \geq 2} (m p^m)^{-1}\right) = O((\log \log H)^{-1}).$$

LEMMA 4. We have, for $t \in J$,

$$\log \zeta(1 + it) = \sum_{p \leq X} \log(1 - p^{-s})^{-1} + O((\log \log H)^{-1}), \quad (10)$$

where $s = 1 + it$.

PROOF. Follows from Lemmas 1, 2 and 3.

LEMMA 5. Let $0 \leq r < 1, 0 \leq \phi < 2\pi$. Then, we have,

$$\log |(1 - r e^{i\phi})^{-z}| \leq -\text{Cos } \theta \log \left(\sqrt{1 - r^2 \text{Sin}^2 \theta} - r \text{Cos } \theta\right) + \text{Sin } \theta \text{Sin}^{-1}(r \text{Sin } \theta). \quad (11)$$

REMARK. Put

$$\lambda_p(\theta) = (1 - p^{-1}) \left(\sqrt{1 - p^{-2} \text{Sin}^2 \theta} - p^{-1} \text{Cos } \theta\right)^{-\text{Cos } \theta} \text{Exp} \left(\text{Sin } \theta \text{Sin}^{-1} \left(\frac{\text{Sin } \theta}{p}\right)\right). \quad (12)$$

In the lemma replace $r e^{i\phi}$ by p^{-s} . Lemmas 4 and 5 complete the proof of (4) and hence that of Theorem 2 since $\sum_{p \geq X} \log \lambda_p(\theta) = O(X^{-1})$ and

$$\prod_{p \leq X} (1 - p^{-1})^{-1} = e^\gamma \log X + O(1). \quad (\text{See [P] page 81}).$$

PROOF OF LEMMA 5. Denote the LHS of (11) by $g(\phi)$. Then

$$\begin{aligned} g(\phi) &= \sum_{n=1}^{\infty} n^{-1} r^n \text{Cos}(n\phi + \theta) \\ g'(\phi) &= -\sum_{n=1}^{\infty} r^n \text{Sin}(n\phi + \theta) \\ &= \text{Im} \left\{ \frac{-r e^{i(\phi + \theta)} (1 - r e^{-i\phi})}{(1 - r e^{i\phi})(1 - r e^{-i\phi})} \right\}. \end{aligned}$$

Hence $g'(\phi) = 0$ if $\text{Sin}(\phi + \theta) = r \text{Sin } \theta$, i.e. if

$$\phi = -\theta + \text{Sin}^{-1}(r \text{Sin } \theta). \quad (13)$$

At this point $g(\phi)$ attains the maximum as we shall show in the end. Now

$$\begin{aligned} g(\phi) &= \text{Re} \left\{ -e^{i\theta} \left(\log \sqrt{1 - 2r \text{Cos } \phi + r^2} - i \text{Sin}^{-1} \frac{r \text{Sin } \phi}{\sqrt{1 - 2r \text{Cos } \phi + r^2}} \right) \right\} \\ &= -\text{Cos } \theta \log \sqrt{1 - 2r \text{Cos } \phi + r^2} - \text{Sin } \theta \text{Sin}^{-1} \left(\frac{r \text{Sin } \phi}{\sqrt{1 - 2r \text{Cos } \phi + r^2}} \right) \end{aligned} \quad (14)$$

From (13) we have

$$\begin{aligned} \text{Sin } \phi &= r \text{Sin } \theta \text{Cos } \theta - \sqrt{1 - r^2 \text{Sin}^2 \theta} \text{Sin } \theta \\ &= -\text{Sin } \theta \left(\sqrt{1 - r^2 \text{Sin}^2 \theta} - r \text{Cos } \theta \right), \\ \text{Cos } \phi &= \sqrt{1 - r^2 \text{Sin}^2 \theta} \text{Cos } \theta + r \text{Sin}^2 \theta, \\ 1 - 2r \text{Cos } \phi + r^2 &= 1 - 2r \text{Cos } \theta \sqrt{1 - r^2 \text{Sin}^2 \theta} - 2r^2 \text{Sin}^2 \theta + r^2 \\ &= \left(\sqrt{1 - r^2 \text{Sin}^2 \theta} - r \text{Cos } \theta \right)^2, \end{aligned}$$

since $-r^2 \text{Sin}^2 \theta + r^2 \text{Cos}^2 \theta = -2r^2 \text{Sin}^2 \theta + r^2$. Hence

$$g(\phi) \leq h(\theta) \quad (15)$$

where $h(\theta)$ is the RHS of (11), provided $g(\phi)$ attains its maximum for the value ϕ given by (13). We now show that

(a) If $\text{Cos } \theta \geq 0$ then $g(\pi) < h(\theta)$

and

(b) If $\text{Cos } \theta < 0$ then $g(0) < h(\theta)$.

Note that $\text{Sin } \theta \text{Sin}^{-1}(r \text{Sin } \theta) \geq 0$. Hence it suffices to prove (in case (a))

$$\begin{aligned} g(\pi) &= \text{Re} \log \left\{ (1 - re^{i\phi})^{-z} \right\}_{\phi=\pi} \\ &= -\text{Cos } \theta \log(1 + r) < -\text{Cos } \theta \log(\sqrt{1 - r^2 \text{Sin}^2 \theta} - r \text{Cos } \theta) \end{aligned}$$

$$\text{i.e. } \log(1+r) > \log(\sqrt{1-r^2 \sin^2 \theta} - r \cos \theta)$$

$$\text{i.e. } (1+r+r \cos \theta)^2 > 1-r^2 \sin^2 \theta$$

$$\text{i.e. } (1+r)^2 + 2r(1+r) \cos \theta > 1-r^2$$

$$\text{i.e. } 1+r+2r \cos \theta > 1-r \text{ (true since } \cos \theta \geq 0)$$

In case (b) it suffices to prove

$$\begin{aligned} g(0) &= \operatorname{Re} \log\{(1-re^{i\phi})^{-z}\}_{\phi=0} \\ &= -\cos \theta \log(1-r) < -\cos \theta \log(\sqrt{1-r^2 \sin^2 \theta} - r \cos \theta) \end{aligned}$$

$$\text{i.e. } \log(1-r) < \log(\sqrt{1-r^2 \sin^2 \theta} - r \cos \theta)$$

$$\text{i.e. } 1-r < \sqrt{1-r^2 \sin^2 \theta} - r \cos \theta$$

$$\text{i.e. } (1-r+r \cos \theta)^2 < 1-r^2 \sin^2 \theta$$

$$\text{i.e. } (1-r)^2 + 2r(1-r) \cos \theta < 1-r^2$$

$$\text{i.e. } 1-r+2r \cos \theta < 1+r \text{ (which is true).}$$

Thus Lemma 5 is completely proved and hence (4) and Theorem 2 are completely proved.

REMARK 1. In the notation of [BR] our method gives $\gg TK^{-E}$ disjoint sub-intervals I (of length K) of $[T, 2T]$ for which $\max_{t \in I} |(\zeta(1+it))^z|$ lies between $e^{\gamma\lambda(\theta)}(\log \log K - \log \log \log K) + O(1)$ and $e^{\gamma\lambda(\theta)}(\log \log K + \log \log \log K) + O(1)$.

REMARK 2. By our method we can show that if $\zeta(s) \neq 0$ in the open half plane $\sigma > \frac{1}{2}$ then for $t \geq 100$ we have

$$|(\zeta(1+it))^z| \leq 2e^{\gamma\lambda(\theta)} \log \log t + O(1).$$

REFERENCES

- [BR] R. BALASUBRAMANIAN AND K. RAMACHANDRA, *On the frequency of Titchmarsh's phenomenon for $\zeta(s)$ -V*, Arkiv för Math., Vol. 26 No.1 (1988), 13-20.
- [I] A.E. INGHAM, *On the estimation of $N(\sigma, T)$* , Quart J. Oxford, 11 (1940), 291-292.
- [AI] A. IVIĆ, *The Riemann zeta-function*, John Wiley and Sons, New York (1985).
- [P] K. PRACHAR, *Primzahlverteilung*, Springer-Verlag (1957).
- [R] K. RAMACHANDRA, *Proof of some conjectures on the mean-value of Titchmarsh series with applications to Titchmarsh's phenomenon*, Hardy-Ramanujan J., vol. 13 (1990), 21-27.
- [T] E.C. TITCHMARSH, *The theory of the Riemann zeta-function*, Second edition, Revised by D.R. Heath-Brown, Oxford University Press, (1986).

ADDRESS OF THE AUTHOR

PROFESSOR K. RAMACHANDRA
SCHOOL OF MATHEMATICS
TATA INSTITUTE OF FUNDAMENTAL RESEARCH
HOMI BHABHA ROAD
BOMBAY 400 005
INDIA

MANUSCRIPT COMPLETED ON 19 NOVEMBER 1990.