



HAL
open science

On the zeros of a class of generalised Dirichlet series-VIII

R Balasubramanian, K Ramachandra

► **To cite this version:**

R Balasubramanian, K Ramachandra. On the zeros of a class of generalised Dirichlet series-VIII. Hardy-Ramanujan Journal, 1991, Volume 14 - 1991, pp.21 - 33. 10.46298/hrj.1991.122. hal-01104792

HAL Id: hal-01104792

<https://hal.science/hal-01104792v1>

Submitted on 22 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hardy-Ramanujan Journal

Vol.14 (1991) 21-33

ON THE ZEROS OF A CLASS OF GENERALISED
DIRICHLET SERIES-VIII

BY

R. BALASUBRAMANIAN AND K. RAMACHANDRA

§ 1. INTRODUCTION AND NOTATION. In the paper VII^[4] of this series (for the earlier papers of the series see the list of references in the paper VII^[4]) K. Ramachandra started a new problem "Let $s = \sigma + it, T \geq T_0$. For what values $\alpha = \alpha(T)$ the rectangle $(\sigma \geq \alpha(T), T \leq t \leq 2T)$ contains infinity of zeros of a generalised Dirichlet series of a certain type?" (In the earlier papers of this series he and R. Baslasubramanian, sometimes individually and sometimes jointly, considered the problem where $\alpha = \alpha(T)$ is independent of T). Since the series considered in that paper were too general the answer $(\alpha(T) = \frac{1}{2} - \frac{D}{\log \log T})$ was perhaps too weak. In the present paper we consider some of the Dirichlet series of the form $F(s) = \sum_{n=1}^{\infty} (a_n b_n \lambda_n^{-s})$ which were considered in the paper V^[3] of this series. (The method of the present paper does not succeed for *all* the series considered in V^[3] let alone those considered in VI^[2]). Before we recall the general series of V^[3], we record two neat results (the second being deeper than the first) as two theorems. In what follows T is the only variable and we assume that T exceeds a large positive constant.

THEOREM 1. *Let $\{\chi(n)\}(n = 1, 2, 3, \dots)$ be any sequence of complex*

numbers with $\sum_{n \leq x} \chi(n) = O(1)$. Let, as usual, $s = \sigma + it$. Then the number of zeros of $\zeta(s) + \sum_{n=1}^{\infty} (\chi(n)n^{-s})$ in the rectangle

$$\left\{ \sigma \geq \frac{1}{2} - C_0(\log \log T)^{\frac{3}{2}}(\log T)^{-\frac{1}{2}}, T \leq t \leq 2T \right\}$$

is $\gg T(\log \log T)^{-1}$ for a suitable positive constant C_0 .

THEOREM 2. Let $1 = \lambda_1 < \lambda_2 < \lambda_3 < \dots$ be an infinite sequence of real numbers such that for $n \geq n_0$ (n_0 , a constant), λ_n is the restriction to integers of a twice continuously differentiable function $g(x)$ of a real variable x with the following properties.

- (1) As $x \rightarrow \infty$, $x^{-1}g(x)$ tends to a positive limit.
- (2) There exist positive constants a and b such that for all $x \geq n_0$, we have,

$$a \leq g'(x) \leq b$$

and

$$a \leq (g'(x))^2 - g(x)g''(x) \leq b.$$

Then the number of zeros of $F(s) = \sum_{n=1}^{\infty} ((-1)^n \lambda_n^{-s})$ in the rectangle

$$\left\{ \sigma \geq \frac{1}{2} - C_0(\log \log T)^{\frac{3}{2}}(\log T)^{-\frac{1}{2}}, T \leq t \leq 2T \right\}$$

is $\gg T(\log \log T)^{-1}$ for a suitable positive constant C_0 .

REMARK. For $n = 1, 2, 3, \dots$, let $\beta_n = \beta_n^{(1)} + \beta_n^{(2)}$ where $\beta_n^{(1)}$ and $\beta_n^{(2)}$ are two bounded monotonic sequences of real numbers. Then for $n \geq n_0$ we can replace λ_n by $\lambda_n + \beta_n$ and the result is practically unchanged (i.e. except for a change of C_0).

The general theorem is too lengthy to state. We now proceed to state it. We consider series of the form $F(s) = \sum_{n=1}^{\infty} (a_n b_n \lambda_n^{-s})$ where λ_n has been introduced already (the change of λ_n to $\lambda_n + \beta_n$ mentioned in the remark

below Theorem 2 is certainly permissible in what follows). Let $f(x)$ be a positive real valued function with the following properties.

(1) $f(x)x^\eta$ is increasing and $f(x)x^{-\eta}$ is decreasing for every $\eta > 0$ and all $x \geq x_0(\eta)$.

(2) For $n \geq n_0$, $a \leq |b_n| (f(n))^{-1} \leq b$.

(3) For all $x \geq 1$, $\sum_{x \leq n \leq 2x} |b_{n+1} - b_n| \leq bf(x)$. We next assume that $\{a_n\}$ and $\{b_n\}$ satisfy one at least of the following two conditions.

(4) **Monotonicity condition.** Let $a_n (n = 1, 2, 3, \dots)$ be a bounded sequence of complex numbers such that $x^{-1} \sum_{n \leq x} a_n$ tends to a non-zero limit (which may be complex) and further $|b_n| \lambda_n^{-\eta}$ is monotonic decreasing for every $\eta > 0$ and all $n \geq n_0(\eta)$.

(5) **Real part condition.** There exists an infinite arithmetic progression J of positive integers such that

$$\liminf_{x \rightarrow \infty} \left(\frac{1}{x} \sum_{\substack{x \leq \lambda_n \leq 2x, \\ n \in J}} \operatorname{Re} a_n \right) > 0$$

and

$$\lim_{x \rightarrow \infty} \left(\frac{1}{x} \sum_{\substack{x \leq \lambda_n \leq 2x, \\ n \in J}} a_n \right) = 0.$$

We are now in a position to state our general theorem.

THEOREM 3. Let $F(s) = \sum_{n=1}^{\infty} (a_n b_n \lambda_n^{-s})$ be as described above. Let $\operatorname{Exp}(-\sqrt{\log x}) \leq f(x)$ for $x \geq x_0$. Let β be a positive constant $< \frac{1}{2}$ and that $F(s)$ can be continued analytically in $(\sigma \geq \beta, \frac{1}{2}T \leq t \leq \frac{5}{2}T)$ and here $\max |F(s)| \leq T^{A_1}$ where $A_1 \geq 2$ is a positive constant. Finally let

$$\frac{1}{T} \int_{\frac{1}{2}T}^{\frac{5}{2}T} |F(\frac{1}{2} + it)|^2 dt \leq (\log T)^{A_2}$$

where $A_2 \geq 2$ is a constant. Then the number of zeros of $F(s)$ in the rectangle

$$\left\{ \sigma \geq \frac{1}{2} - C_0(\log \log T)^{\frac{3}{2}}(\log T)^{-\frac{1}{2}}, T \leq t \leq 2T \right\}$$

is $\gg T(\log \log T)^{-1}$ where $C_0 \geq 0$ is a certain constant.

REMARK 1. The restriction of the theorem regarding the upper bound for the mean square of $|F(\frac{1}{2} + it)|$ is very strong. Practically (since the mean square can be proved to be $\gg (f(T))^2$) it forces us to consider the series of $V^{[3]}$, with the extra restriction $f(x) \leq (\log x)^A$ for some constant $A \geq 2$ and all $x \geq x_0(A)$. Further the restriction $f(x) \geq \text{Exp}(-\sqrt{\log x})$ forces us to consider only a sub-class of functions considered in $V^{[3]}$. It may be remarked that the mean square hypothesis is satisfied for all functions considered in $V^{[3]}$ by imposing $f(x) \leq (\log x)^A$.

REMARK 2. A nice example of the functions covered by Theorem 3 is $\sum_{n=1}^{\infty} ((-1)^n \text{Exp}(-\sqrt{\log n}) n^{-s})$. It may be noted (as a special case of a very general Theorem [1]) that this is an entire function.

REMARK 3. In the theorem it is not difficult to relax the rectangle of analytic continuation to $(\sigma \geq \beta, T \leq t \leq 2T)$ and replace the mean-value condition by

$$\frac{1}{T} \int_T^{2T} |F(\frac{1}{2} + it)|^2 \leq (\log T)^{A_2}$$

where $A_2 \geq 2$ is a constant.

REMARK 4. It is possible to generalise our results further. As a simple example we can in Theorem 1 replace $\zeta(s) + \sum_{n=1}^{\infty} (\chi(n)n^{-s})$ by

$$K^{-s}(\zeta(s) + \sum_{n=1}^{\infty} (\chi(n)n^{-s})) + \sum_{n=1}^{\infty} d_n \lambda_n^{-s}$$

where $\sum_{n \leq x} d_n = O(1)$, K is a positive constant, $|\lambda_m - Kn| \geq (100)^{-1}$ for all m, n , $1 \ll \lambda_{n+1} - \lambda_n$ and finally $\lambda_n = O(n)$.

REMARK 5. We have imposed the restriction $f(x) \geq \text{Exp}(-\sqrt{\log x})$ for

$x \geq x_0$ to obtain some worthwhile results, but it is possible to obtain weaker results by relaxing this condition.

NOTATION. The letter A with or without subscripts will denote constants ≥ 2 . The letter C with or without subscripts will denote positive constants.

§ 2. **A GENERAL LEMMA.** Let $1 = \lambda_1 < \lambda_2 < \lambda_3 < \dots$ be an infinite sequence of real numbers with $1 \gg \lambda_{n+1} - \lambda_n \gg 1$ and $\{k_n\} (n = 1, 2, 3, \dots)$ be any sequence of complex numbers such that $k_1 = 1$ and the series $\phi(s) = \sum_{n=1}^{\infty} (k_n \lambda_n^{-s})$ is convergent in $\sigma \geq A_1$ and is continuable analytically in $(\sigma \geq \beta, T - (\log T)^2 \leq t \leq T + (\log T)^2)$ and there $\max |\phi(s)| \leq T^{A_2}$, where $\beta < \frac{1}{2}$ is a positive constant. Let

$$\frac{1}{T} \int_{T - (\log T)^2}^{2T + (\log T)^2} \left| \phi\left(\frac{1}{2} + it\right) \right|^2 dt \leq (\log T)^{A_3}.$$

Then, we have,

$$\frac{1}{T} \int_{\frac{1}{2} - (\log T)^{-1}}^{A_1 + 2} \int_{T-1}^{2T+1} \left| \phi(\sigma + it) \right|^2 dt d\sigma \leq (\log T)^{A_4}.$$

REMARK. This lemma is well-known to experts in the subject and so its proof will be postponed to the last section. Also it is possible to replace $(\log T)^2$ by a constant multiple of $\log \log T$.

§ 3. **THE FUNCTION $F_2(s)$.** As in VI^[2] we introduce the function (in VI^[2] we have used the kernel $\text{Exp}(W^{4a+2})$ but we now use the kernel $\text{Exp}((\text{Sin } W)^2)$)

$$F_2(s) = \sum_{n=1}^{\infty} a_n b_n (\Delta(T) - \Delta(TD^{-1})) \lambda_n^{-s}$$

where D is a large positive constant and $\Delta(x)$ for $x > 0$ is defined by

$$\Delta(x) = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} F(W) x^W \text{Exp}\left(\left(\text{Sin } \frac{W}{1000}\right)^2\right) \frac{dW}{W}.$$

As in VI^[2] we have

LEMMA 1. *Let q be any real constant satisfying $\beta < q < \frac{1}{2}$. Then we have the inequalities*

$$(1) \quad \frac{1}{T} \int_T^{2T} |F_2(q+it)|^2 dt \ll T^{1-2q} (f(T))^2,$$

and

$$(2) \quad \frac{1}{T} \int_T^{2T} |F_2(q+it)| \gg T^{\frac{1}{2}-q} f(T).$$

PROOF. Similar to the proof of Lemma 10 of VI^[2].

LEMMA 2. *Let T be an integer. Then the number of integers M in the range $T \leq M \leq 2T - 1$ for which*

$$\int_M^{M+1} |F_2(q+it)| dt > C_1 T^{\frac{1}{2}-q} f(T)$$

exceeds $C_2 T$.

PROOF. Similar to that of Lemma 4 of VI^[2].

LEMMA 3. *There exist at least $C_3 T (\log \log T)^{-1}$ points t_j with*

$$|F_2(q+it_j)| > c_1 T^{\frac{1}{2}-q} f(T)$$

and such that any two points t_j and $t_{j'}$ with $j \neq j'$ differ by at least $C_4 \log \log T$

REMARK. Here C_4 is arbitrary and C_3 depends on it.

PROOF. Follows from Lemma 2.

LEMMA 4. *Let r be a constant satisfying $\beta < r < q < \frac{1}{2}$. Put $C_5 = \frac{1}{100} C_4$ and $H = C_5 \log \log T$. Then*

$$\int_{t_j-H}^{t_j+H} |F_2(r+it)| \geq C_6 V \log \log T$$

where $V = T^{\frac{1}{2}-r} f(T)$ for at most $C_7 C_6^{-1} T (\log \log T)^{-1}$ points t_j .

REMARK. Here C_6 is arbitrary and C_7 is independent of C_6 .

PROOF. By (1) of Lemma 1, the sum over j of the quantity on the LHS does not exceed C_7VT and this gives Lemma 4.

LEMMA 5. *There are at least $\frac{1}{2}C_3T(\log\log T)^{-1}$ points t_j separated by (distances) at least $C_4\log\log T$ such that if $H = \frac{1}{100}C_4\log\log T$ then with $V = T^{\frac{1}{2}-r}f(T)$, we have,*

$$\int_{t_j-H}^{t_j+H} |F_2(r+it)| dt \leq C_6V \log\log T.$$

REMARK. Here C_4 is arbitrary and C_3 depends on it.

PROOF. The lemma follows by choosing a large C_6 in Lemma 4.

LEMMA 6. *Uniformly in σ with $q < \sigma_0 \leq \sigma < \frac{1}{2}$, we have, for the points t_j of Lemma 5,*

$$\int_{t_j-2H}^{t_j+2H} |F_2(\sigma+iv) \text{Exp}((\text{Sin} \frac{W}{1000})^2) \frac{dW}{W}| > C_8T^{\frac{1}{2}-\sigma} f(T)(\log\log T)^{-\theta}$$

where σ_0 is a constant $W = \sigma - q + iv$, and $\theta = \frac{1}{2(q-r)}$.

PROOF. Put $s_0 = q + it_j$, we have

$$F_2(s_0) = \frac{1}{2\pi i} \int F_2(s_0+W) X^W \text{Exp}((\text{Sin} \frac{W}{1000})^2) \frac{dW}{W}$$

where the integral is taken over the (anticlockwise) boundary of the rectangle bounded by the lines $\text{Re } W = r - q$, $\text{Re } W = \sigma - q$, $\text{Im } W = \pm H$. We take the absolute values (using Lemma 3) of the integrand on the RHS and choose $X = C_8T(\log\log T)^{(q-r)^{-1}}$, where C_8 is a large positive constant. This leads to Lemma 6.

LEMMA 7. *Given any σ in $\sigma_0 \leq \sigma < \frac{1}{2}$, there exist points v_j satisfying $t_j - 2H \leq v_j \leq t_j + 2H$, such that uniformly in σ there holds*

$$|F_2(\sigma+iv_j)| > C_9T^{\frac{1}{2}-\sigma} f(T)(\log\log T)^{-\theta}$$

where $\theta = (2(q-r))^{-1}$.

REMARK. Note that v_j are separated by (distances) at least $\frac{24}{25}C_4\log\log T$

where C_4 is at our disposal.

PROOF. Follows from Lemma 6.

LEMMA 8. *Given any σ in $\sigma_0 \leq \sigma < \frac{1}{2}$ there exist points p_j satisfying $v_j - H \leq p_j \leq v_j + H$ such that uniformly in σ , there holds,*

$$|F(\sigma + ip_j)| > C_{10} T^{\frac{1}{2} - \sigma} f(T) (\log \log T)^{-\theta}$$

where θ is the constant defined before.

REMARK 1. Note that p_j are separated by (distances) at least $\frac{1}{2} C_4 \log \log T$. Also the number of points p_j is at least $\frac{1}{2} C_3 T (\log \log T)^{-1}$. Here C_4 is arbitrary and C_3 depends on it. (Both are independent of σ).

REMARK 2. We can refine the lower bound for $|F(\sigma + ip_j)|$ but we do not do it since it does not have an application.

PROOF. We start with

$$F_2(\sigma + iv_j) = \frac{1}{2\pi i} \int F(\sigma + iv_j + W) T^W (1 - D^{-W}) \text{Exp}((\text{Sin} \frac{W}{1000})^2) \frac{dW}{W}$$

where the integration is over $\text{Re } W = 2$. We break off the portion $|v| \geq C_{11} \log \log T$ with a small error and move the line of integration in the rest to $\text{Re } W = 0$. Here C_{11} is a specific constant and not arbitrary. We now use Lemma 7 and majorise the integrand. This leads to the lemma.

The rest of the proof consists in proving that at least $\frac{1}{3} C_3 T (\log \log T)^{-1}$ of the rectangles

$$\left\{ \sigma \geq \frac{1}{2} - C_0 (\log \log T)^{\frac{3}{2}} (\log T)^{-\frac{1}{2}}, p_j - H \leq t \leq p_j + H \right\}$$

contain a zero of $F(s)$ if C_0 is a large positive constant. This would complete the proof of Theorem 3.

§ 4. TWO APPLICATIONS OF BOREL-CARATHÉODORY THEOREM. Suppose that the rectangle

$$\left\{ \sigma \geq \frac{1}{2} - K\delta, p_j - H \leq t \leq p_j + H \right\}$$

is zero free for $F(s)$, where δ and K are positive quantities to be chosen in the next section. (The quantity δ will be chosen to be small and K to be large).

LEMMA 1. (Borel-Carathéodory Theorem. See [5] page 174). *Suppose $G(z)$ is analytic in $|z - z_0| \leq R$ and on $|z - z_0| = R$ we have $\operatorname{Re} G(z) \leq U$. Then in $|z - z_0| \leq r < R$, we have,*

$$|G(z)| \leq \frac{2rU}{R-r} + \frac{R+r}{R-r} |G(z_0)|.$$

REMARK. The r of this lemma is not to be confused with that of the preceding section.

LEMMA 2. *In the rectangle*

$$\left\{ \sigma \geq \frac{1}{2} - (K-1)\delta, p_j - H + C_{12} \leq t \leq p_j + H - C_{12} \right\}$$

we have,

$$|\log F(s)| \leq C_{13} \delta^{-1} \log T.$$

PROOF. Choose z_0 to be a point in

$$\left\{ \sigma \geq 2, p_j - H + C_{12} \leq t \leq p_j + H - C_{12} \right\}$$

where $\log F(s)$ is bounded and then take R to be such that the circle with centre z_0 and radius R touches $\sigma = \frac{1}{2} - K\delta$ and lies within the rectangle $\left\{ \sigma \geq \frac{1}{2} - K\delta, p_j - H \leq t \leq p_j + H \right\}$. Next choose $r = R - \delta$. This proves Lemma 2.

LEMMA 3. *Let M_j denote the maximum of $|F(s)|$ in $\left\{ \sigma \geq \frac{1}{2}, p_j - H \leq t \leq p_j + H \right\}$. Then, we have,*

$$\sum_j M_j^2 \leq T(\log T)^{A_5}.$$

PROOF. Let M_j be attained at s_j say. Then M_j^2 is majorised by the mean of $|F(s)|^2$ over a disc of radius $(\log T)^{-1}$ with centre s_j . The lemma now follows from the general result of § 2.

LEMMA 4. *We have,*

$$M_j^2 \geq (\log T)^{11A_5}$$

for at most $T(\log T)^{-10}$ values of j . Hence we are still left with at least $\frac{1}{3} C_3 T (\log \log T)^{-1}$ values of j for which

$$M_j^2 \leq (\log T)^{11A_5}.$$

REMARK. From now on we restrict j only to these values.

PROOF. Follows from Lemma 3.

LEMMA 5. *In the rectangle*

$$\left\{ \sigma \geq \frac{1}{2} - \delta, p_j - H + C_{12} \leq t \leq p_j + H - C_{12} \right\}$$

we have,

$$|\log F(s)| \leq C_{14} \delta^{-1} \log \log T.$$

PROOF. Choose z_0 to be a point in

$$\left\{ \sigma \geq 2, p_j - H + C_{12} \leq t \leq p_j + H - C_{12} \right\}$$

and then take R to be such that the circle with centre z_0 and radius R touches $\sigma = \frac{1}{2}$ and lies within the rectangle $\left\{ \sigma \geq \frac{1}{2}, p_j - H \leq t \leq p_j + H \right\}$. Next choose $r = R - \delta$. The lemma now follows from Lemma 4.

§ 5. COMPLETION OF THE PROOF. Suppose that for a certain j , the rectangle $\left\{ \sigma \geq \frac{1}{2} - K\delta, |p_j - t| \leq H \right\}$ does not contain a zero of $F(s)$. We obtain a contradiction in the following way. Put $s_0 = \sigma + ip_j$ where $\sigma = \frac{1}{2} - \delta$, and also let $\sigma_1 = \frac{1}{2} - (K-1)\delta$, $\sigma_2 = \sigma$, and $\sigma_3 = \frac{1}{2} + \delta$. We apply maximum modulus principle to

$$\psi(W) = \log F(s_0 + W) X^W \text{Exp}\left(\left(\text{Sin} \frac{W}{1000}\right)^2\right)$$

according to which

$$|\psi(0)| \leq \max |\psi(W)|$$

maximum being taken over the boundary of the rectangle bounded by $Re W = -(K - 2)\delta, Re W = 2\delta, Im W = \pm \frac{1}{2}H$. If $\delta \geq 6(\log T)^{-\frac{1}{2}}$ we have (by a suitable choice of X and C_4)

$$\begin{aligned} \frac{\delta}{2} \log T &\leq \delta \log T - 3\sqrt{\log T} \leq |\psi(0)| \\ &\leq C_{15}(\delta^{-1} \log T)^{\frac{2}{K}} (\delta^{-1} \log \log T)^{\frac{K-2}{K}}. \end{aligned}$$

We now choose $K = \log \log T$ and obtain

$$\frac{\delta}{2} \log T \leq C_{16} \delta^{-1} \log \log T.$$

This is a contradiction if we choose $\delta = C_{17}(\log \log T)^{\frac{1}{2}}(\log T)^{-\frac{1}{2}}$ and $C_{17}^2 > 2C_{16}$. This proves Theorem 3 provided we prove the general lemma of § 2.

§ 6. PROOF OF THE GENERAL LEMMA. Let $\varepsilon > 0$ be arbitrary but fixed. Then in $\{\sigma \geq \beta + \varepsilon, T \leq t \leq 2T\}$, we have, by Cauchy's theorem $|\phi'(s)| \leq T^{A_2+1}$ and so in $\{|\sigma - \frac{1}{2}| \leq T^{-4A_2}, T \leq t \leq 2T\}$ we have

$$|\phi^2(\frac{1}{2} + it) - \phi^2(\sigma + it)| \leq 1.$$

Hence it suffices to consider in this rectangle the portion $|\sigma - \frac{1}{2}| \geq T^{-4A_2}$. If now $\frac{1}{2} - (\log T)^{-1} \leq \sigma \leq \frac{1}{2} - T^{-4A_2}$ we have

$$|\phi^2(s) = \frac{1}{2\pi i} \int \phi^2(s + W) X^W \text{Exp}(W^2) \frac{dW}{W}$$

the contour being the (anticlockwise) boundary of the rectangle bounded by $Re W = \beta - \sigma, Re W = \frac{1}{2} - \sigma, Im W = \pm \log T$. We choose X to be a large power of T so that the integral over the left boundary is negligible. Clearly the integrals over the horizontal boundaries are together negligible. We take absolute values and integrate with respect to t from $t = T$ to $t = 2T$. This leads to the result since on the right boundary $|X^W| \leq 1$ and $\int | \frac{dW}{W} | \ll \log T$.

If now $\sigma \geq \frac{1}{2} + T^{-4A_2}$ we start with

$$\phi^2(s) = \frac{1}{2\pi i} \int \phi^2(s + W) \text{Exp}(W^2) \frac{dW}{W}$$

the contour being the (anticlockwise) boundary of the rectangle bounded by $Re W = \frac{1}{2} - \sigma$, $Re W = 3A_1 - \sigma$, $Im W = \pm \log T$. The proof proceeds as before using $\phi(s + W) = O(1)$ on the right boundary and negligible on the horizontal boundaries and the fact $\int \left| \frac{dW}{W} \right| \ll \log T$ on the left boundary. This completes the proof of the general lemma.

Theorem 3 is now completely proved.

ACKNOWLEDGEMENT. The authors are grateful to Professor M. JUTILA for encouragement.

REFERENCES

- [1] R. BALASUBRAMANIAN AND K. RAMACHANDRA, *The proof that certain functions are entire*, Maths. Teacher (India) (1985), p.7.
- [2] R. BALASUBRAMANIAN AND K. RAMACHANDRA, *On the zeros of a class of generalised Dirichlet series-VI*, Arkiv för Matematik, no. 2, 19 (1981), p. 239-250.
- [3] K. RAMACHANDRA, *On the zeros of a class of generalised Dirichlet series-V*, Reine u. Angew. Math., 303/304 (1978), p. 295-313.
- [4] K. RAMACHANDRA, *On the zeros of a class of generalised Dirichlet series-VII*, (to appear).
- [5] E.C. TITCHMARSH, *Theory of functions*, Oxford University Press (1939).

ADDRESS OF THE AUTHORS

- 1) PROFESSOR R. BALASUBRAMANIAN
MATSCIENCE
THARAMANI P.O.
MADRAS 600 113
INDIA

- 2) PROFESSOR K. RAMACHANDRA
SCHOOL OF MATHEMATICS
TATA INSTITUTE OF FUNDAMENTAL RESEARCH
HOMI BHABHA ROAD
COLABA
BOMBAY 400 005
INDIA

MANUSCRIPT COMPLETED ON 17 FEBRUARY 1991