



HAL
open science

On the numbers e^e , e^{e^2} and e^{π^2} .

Michel Waldschmidt

► **To cite this version:**

Michel Waldschmidt. On the numbers e^e , e^{e^2} and e^{π^2} .. Hardy-Ramanujan Journal, 1998, pp.27 - 34.
hal-01109388

HAL Id: hal-01109388

<https://hal.science/hal-01109388>

Submitted on 26 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hardy-Ramanujan Journal
Vol.21 (1998) 27-34

On the numbers e^e , e^{e^2} and e^{π^2}

by

Michel WALDSCHMIDT

Abstract. We give two measures of simultaneous approximation by algebraic numbers, the first one for the triple (e, e^e, e^{e^2}) and the second one for (π, e, e^{π^2}) . We deduce from these measures two transcendence results which had been proved in the early 70's by W.D. Brownawell and the author.

Introduction

In 1949, A.O. Gel'fond introduced a new method for algebraic independence, which enabled him to prove that the two numbers $2^{\sqrt[3]{2}}$ and $2^{\sqrt[3]{4}}$ are algebraically independent. At the same time, he proved that one at least of the three numbers e^e , e^{e^2} , e^{e^3} is transcendental (see[G] Chap. III, §4).

At the end of his book [S] on transcendental numbers, Th. Schneider suggested that one at least of the two numbers e^e , e^{e^2} is transcendental; this was the last of a list of eight problems, and the first to be solved, in 1973, by W.D. Brownawell [B] and M. Waldschmidt [W 1], independently and simultaneously. For this result they shared the Distinguished Award of the Hardy-Ramanujan Society in 1986. Another consequence of their main result is that one at least of the two following statements holds true:

- (i) The numbers e and π are algebraically independent
- (ii) The number e^{π^2} is transcendental

Our goal is to shed a new light on these results. It is hoped that our approach will yield further progress towards a solution of the following open problems:

(?) Two at least of the three numbers e , e^e , e^{e^2} are algebraically independent.

(?) Two at least of the three numbers π , e , e^{π^2} are algebraically independent.

Further conjectures are as follows:

(?) Each of the numbers e^e , e^{e^2} , e^{π^2} is transcendental

(?) The numbers e and π are algebraically independent.

We conclude this note by showing how stronger statements are consequences of Schanuel's conjecture.

1. Heights

Let γ be a complex algebraic number. The *minimal polynomial* of γ over \mathbb{Z} is the unique polynomial

$$f(X) = a_0X^d + a_1X^{d-1} + \dots + a_d \in \mathbb{Z}[X]$$

which vanishes at the point γ , is irreducible in the factorial ring $\mathbb{Z}[X]$ and has leading coefficient $a_0 > 0$. The integer $d = \deg f$ is the *degree* of γ , denoted by $[Q(\gamma) : \mathbb{Q}]$. The *usual height* $H(\gamma)$ of γ is defined by

$$H(\gamma) = \max\{|a_0|, \dots, |a_d|\}.$$

It will be convenient to use also the so-called *Mahler's measure* of γ , which can be defined in three equivalent ways. The first one is

$$M(\gamma) = \exp \left(\int_0^1 \log |f(e^{2i\pi t})| dt \right).$$

For the second one, let $\gamma_1, \dots, \gamma_d$ denote the complex roots of f , so that

$$f(X) = a_0 \prod_{i=1}^d (X - \gamma_i).$$

Then, according to Jensen's formula, we have

$$M(\gamma) = |a_0| \prod_{i=1}^d \max\{1, |\gamma_i|\}.$$

For the third one, let K be a number field (that is a subfield of \mathbb{C} which is a \mathbb{Q} -vector space of finite dimension $[K:\mathbb{Q}]$) containing γ , and let M_K be the set of (normalized) absolute

values of K . Then

$$M(\gamma) = \prod_{v \in M_K} \max\{1, |\gamma|_v\}^{[K_v:\mathbb{Q}]}$$

where K_v is the completion of K for the absolute value v and Q_v the topological closure of Q in K_v and $[K_v : Q_v]$ the local degree.

Mahler's measure is related to the usual height by

$$2^{-d}H(\gamma) \leq M(\gamma) \leq \sqrt{d+1}.H(\gamma).$$

From this point of view it does not make too much difference to use H or M , but one should be careful that d denotes the exact degree of γ , not an upper bound. We shall deal below with algebraic numbers of degree d bounded by some parameter D .

Definition. For an algebraic number γ of degree d and Mahler's measure $M(\gamma)$, we define the *absolute logarithmic height* $h(\gamma)$ by

$$h(\gamma) = \frac{1}{d} \log M(\gamma).$$

2. Simultaneous Approximation

We state two results dealing with simultaneous Diophantine approximation. Both of them are consequences of the main result in [W 2]. Details of the proof will appear in the forthcoming book [W 3].

2.1. Simultaneous Approximation to e, e^e and e^{e^2}

Theorem 1. *There exists a positive absolute constant c_1 such that, if $\gamma_0, \gamma_1, \gamma_2$ are algebraic numbers in a field of degree D , then*

$$|e - \gamma_0| + |e^e - \gamma_1| + |e^{e^2} - \gamma_2| > \exp\{-c_1 D^2 (h_0 + h_1 + h_2)^{1/2} (h_1 + h_2)^{1/2} (h_0 + \log D) (\log D)^{-1}\}$$

where $h_i = \max\{e, h(\gamma_i)\}$ ($i = 0, 1, 2$).

2.2. Simultaneous Approximation to π, e and e^{π^2}

Theorem 2. *There exists a positive absolute constant c_2 such that, if $\gamma_0, \gamma_1, \gamma_2$ are algebraic numbers in a field of degree D , then*

$$|\pi - \gamma_0| + |e^e - \gamma_1| + |e^{e^2} - \gamma_2| > \exp\{-c_2 D^2 (h_0 + \log(Dh_1 h_2))^{1/2} h_1^{1/2} h_2^{1/2} (\log D)^{-1}\}$$

where $h_i = \max\{e, h(\gamma_i)\}$ ($i = 0, 1, 2$).

3. Transcendence Criterion

3.1 Algebraic Approximations to a Given Transcendental Number

The following result is Théorème 3.2 of [R-W 1]; see also Theorem 1.1 of [R-W 2].

Theorem 3. *Let $\theta \in \mathbb{C}$ be a complex number. The two following conditions are equivalent:*

(i) *the number θ is transcendental.*

(ii) *For any real number $h \geq 10^7$, there are infinitely many integers $d \geq 1$ for which there exists an algebraic number γ of degree d and absolute logarithmic height $h(\gamma) \leq h$ which satisfies*

$$0 < |\theta - \gamma| \leq \exp(-10^{-7} h d^2).$$

Notice that the proof of (ii) \Rightarrow (i) is an easy consequence of Liouville's inequality.

3.2. Application to e^e and e^{e^2}

Corollary to Theorem 1. *One at least of the two numbers e^e, e^{e^2} is transcendental.*

Proof of the corollary. Assume that the two numbers e^e, e^{e^2} are algebraic, say γ_1 and γ_2 . Then, according to Theorem 1, there exists a constant $c_3 > 1$ such that, for any algebraic number γ of degree $\leq D$ and height $h(\gamma) \leq h$ with $h \geq e$,

$$|e - \gamma| > \exp\{-c_3 D^2 h^{1/2} (h + \log D) (\log D)^{-1}\}.$$

We now use Theorem 3 for $\theta = e$ with $h = 10^{15} c_3^2$ and derive a contradiction.

4. Algebraic Independence

4.1 Simultaneous Approximation

The proof of the following result is given in [R-W 1], Théorème 3.1, as a consequence of Theorem 3 (see also [R-W 2] Corollary 1.2).

Corollary to Theorem 3. *Let $\theta_1, \dots, \theta_m$ be complex numbers such that the field $Q(\theta_1, \dots, \theta_m)$ has transcendence degree 1 over Q . There exists a constant $c > 0$ such that, for any real number $h \geq c$, there are infinitely many integers D for which there exists a tuple $(\gamma_1, \dots, \gamma_m)$ of algebraic numbers satisfying*

$$[Q(\gamma_1, \dots, \gamma_m) : Q] \leq D, \quad \max_{1 \leq i \leq m} h(\gamma_i) \leq h$$

and

$$\max_{1 \leq i \leq m} \{|\theta_i - \gamma_i|\} \leq \exp(-c^{-1}hD^2).$$

4.2. Application to π, e and e^{π^2}

Corollary to Theorem 2. *One at least of the two following statements is true:*

- (i) *The numbers e and π are algebraically independent.*
- (ii) *The number e^{π^2} is transcendental.*

Remark. This corollary can be stated in an equivalent way as follows:

For any non constant polynomial $P \in Z[X]$, the complex number

$$e^{\pi^2} + iP(e, \pi)$$

is transcendental.

The idea behind this remark originates in [R].

Proof of the Corollary. Assume that the number e^{π^2} is algebraic. Theorem 2 with $\gamma_2 = e^{\pi^2}$ shows that there exists a constant $c_4 > 0$ such that, for any pair (γ_0, γ_1) of algebraic numbers, if we set

$$D = [Q(\gamma_0, \gamma_1) : Q] \quad \text{and} \quad h = \max\{e, h(\gamma_0), h(\gamma_1)\},$$

then

$$|\pi - \gamma_0| + |e - \gamma_1| > \exp\{-c_4 D^2 (h + \log D)^{1/2} h^{1/2} (\log D)^{-1}\}$$

Therefore we deduce from the Corollary to Theorem 3 that the field $Q(\pi, e)$ has transcendence degree 2.

5. Schanuel's Conjecture

The following conjecture is stated in [L] Chap. III p. 30: (The results of this section are based on the conjecture to be stated).

Schanuel's Conjecture. *Let x_1, \dots, x_n be Q -linearly independent complex numbers. Then, among the $2n$ numbers*

$$x_1, \dots, x_n, e^{x_1}, \dots, e^{x_n},$$

at least n are algebraically independent.

Let us deduce from Schanuel's Conjecture the following statement (which is an open problem):

(?) *The 7 numbers*

$$e, \pi, e^e, e^{e^2}, e^{\pi^2}, 2^{\sqrt[3]{2}}, 2^{\sqrt[3]{2}}$$

are algebraically independent.

We shall use Schanuel's conjecture twice. We start with the numbers $1, \log 2$, and $i\pi$ which are linearly independent over Q because $\log 2$ is irrational. Therefore, according to Schanuel's conjecture, three at least of the numbers

$$1, \log 2, i\pi, e, 2, -1$$

are algebraically independent. This means that the three numbers $\log 2, \pi$ and e are algebraically independent. Therefore the 8 numbers

$$1, i\pi, \pi^2, e, e^2, \log 2, 2^{\frac{1}{3}} \log 2, 4^{\frac{1}{3}} \log 2$$

are Q -linearly independent. Again, Schanuel's conjecture implies that 8 at least of the numbers

$$1, i\pi, \pi^2, e, e^2, \log^2, 2^{\frac{1}{3}} \log 2, 4^{\frac{1}{3}} \log 2, e, -1, e^{\pi^2}, e^e, e^{e^2}, 2, 2^{\sqrt[3]{2}}, 2^{\sqrt[3]{4}}$$

are algebraically independent, and this means that the 8 numbers

$$e, \pi, e^e, e^{e^2}, e^{\pi^2}, 2^{\sqrt[3]{2}}, 2^{\sqrt[3]{4}}, \log 2$$

are algebraically independent.

ACKNOWLEDGEMENTS.

This research was performed in March/April 1998 while the author visited the Tata Institute of Fundamental Research under a cooperation program from the Indo-French Center IFCEPAR/CEFIPRA Research project 1601-2 << Geometry >>, The author wishes also to thank Professor K.Ramachandra for his invitation to the National Institute of Advanced Studies at Bangalore.

REFERENCES

- [B] Brownawell, W.Dale. - *The algebraic independence of certain numbers related to the exponential function.* J.Number Th. 6 (1974), 22-31.
- [G] Gel'fond, Aleksandr O, - *Transcendental and algebraic numbers.* Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow, 1952. 224 pp. *Transcendental and algebraic numbers.* Translated from the first Russian edition by Leo F. boron Dover Publications, Inc., New York 1960 vii+190 pp.
- [L] Lang, Serge. - *Introduction to transcendental numbers.* Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont. 1966 vi+105 pp.
- [R] Ramachandra, K. - *Contributions to the theory of transcendental numbers.*I, II. Acta Arith. 14 (1967/68), 65-72; *ibid.*, 14 (1967/1968), 73-88.
- [R-W 1] Roy, Damien; Waldschmidt, Michel. - *Approximation diophantienne et indépendance algébrique de logarithmes.* Ann. Scient. Ec. Norm. Sup., 30 (1997), no 6, 753-796.
- [R-W 2] Roy, Damien; Waldschmidt, Michel. - *Simultaneous approximation and algebraic independence.* The Ramanujan Journal, 1 Fasc. 4(1997), 379-430.

- [S] Schneider, Theodor,- *Einführung in die transzendenten Zahlen*. Springer-Verlag, berlin-Göttingen-Heidelberg, 1957, v+150 pp. *Introduction aux nombres transcendants*. Traduit de l'allemand par P. Eymard. Gauthier-Villars, paris 1959 viii+151 pp.
- [W 1] Waldschmidt, Michel. - *Solution du huitième problème de Schneider*. J.Number Theory, **5** (1973), 191-202.
- [W 2] Waldschmidt, Michel. - *Approximation diophantienne dans les groupes algébriques commutatifs - (I) : Une version effective du theoreme du sous-groupe algébrique*. J. reine angew. Math. **493** (1997), 61-113.
- [W 3] Waldschmidt, Michel. - *Diophantine Approximation on Linear Algebraic Groups. Transcendence Properties of the Exponential Function in Several Variables*. In preparation.

Michel Waldschmidt
Université P.et M. Curie (Paris VI)
Institut Mathématique de Jussieu
Problèmes Diophantiens, Case 247
4, Place Jussieu
F-75252, Paris CEDEX05
FRANCE.

E-mail: miw@math.jussieu.fr

Received 28th March 1998

Accepted 30th April 1998.