



HAL
open science

Mean-square upper bound of Hecke L -functions on the critical line.

A Sankaranarayanan

► **To cite this version:**

A Sankaranarayanan. Mean-square upper bound of Hecke L -functions on the critical line.. Hardy-Ramanujan Journal, 2003, Volume 26 - 2003, pp.2-17. 10.46298/hrj.2003.147 . hal-01109810

HAL Id: hal-01109810

<https://hal.science/hal-01109810>

Submitted on 27 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mean-Square Upper Bound of Hecke L - Functions On The Critical Line

A. Sankaranarayanan

* †

ABSTRACT. We prove the upper bound for the mean-square of the absolute value of the Hecke L -functions (attached to a holomorphic cusp form) defined for the congruence subgroup $\Gamma_0(N)$ on the critical line uniformly with respect to its conductor N .

1. Introduction

Let χ denote a character mod q and $N_\chi(\alpha, T)$, the number of zeros of $L(s, \chi)$ in the rectangle $\alpha < \sigma < 1, |t| < T$. Gallagher (see [7]) proved that

$$(1.1) \quad \sum_{\chi} \left| L\left(\frac{1}{2} + it, \chi\right) \right|^2 \ll (q + |t|) \log(q|t| + 2),$$

$$(1.2) \quad \int_{-T}^T \left| L\left(\frac{1}{2} + it, \chi\right) \right|^2 \ll (q + T) \log qT,$$

and

$$(1.3) \quad N_\chi(\alpha, T) \ll T^{3(1-\alpha)} (\log T)^C$$

uniformly for all $q \leq T$. He also established that

$$(1.4) \quad \sum_{\chi} (N_\chi(\alpha, T+1) - N_\chi(\alpha, T)) \ll q^{3(1-\alpha)} (\log q)^C$$

for $T \leq q$.

Meurman (see [24]) proved that for any $q \geq 1$,

$$(1.5) \quad \sum_{\chi(\text{mod } q)} \int_T^{T+H} \left| L\left(\frac{1}{2} + it, \chi\right) \right|^2 \ll_{\epsilon} \left(qH + (qT)^{\frac{1}{3}} \right) (q(T+H))^{\epsilon}$$

*2000 *Mathematics subject classification* : primary 11M , secondary 11M06 .

†*Key Words* : Hecke L -functions, Holomorphic cusp forms, Mean-value theorems, Montgomery-Vaughan Theorem.

for any $\epsilon > 0$ and $3 \leq H \leq T$.

Balasubramanian and Ramachandra have given a simple proof of the above result (1.5) (see [2]) and they could replace $(qT)^\epsilon$ by $\log qT$ times a power of $d(q)$ for primitive characters mod q . Jutila has proved ([20] and [21]) the fourth power mean for $\zeta(s)$ and mean-square for L -functions attached to cusp forms (holomorphic as well Maass wave forms) via Laplace transformation method. In fact he studies the error term mean values in a greater generality with an emphasis as a function of T .

It should be mentioned here that mean values of derivatives of modular L -series had been studied by Ram Murty and Kumar Murty in [34]. For several interesting mean value results which were obtained by many mathematicians, see for example [1], [11], [16], [19], [22], [23], [28], [37], [38].

Let $f(z) = \sum_{n=1}^{\infty} a_f(n)e^{2\pi inz}$, $\Im z > 0$ be a cusp form of even integral weight $k > 2$ for the full modular group $SL(2, Z)$. Hecke L -function attached to f is defined as

$$(1.6) \quad L_f(s) = \sum_{n=1}^{\infty} a_f(n)n^{-s}$$

which is absolutely convergent in $\Re s > \frac{k+1}{2}$, and it satisfies the functional equation

$$(1.7) \quad (2\pi)^{-s}\Gamma(s)L_f(s) = (-1)^{\frac{k}{2}}(2\pi)^{s-k}\Gamma(k-s)L_f(k-s).$$

In [10], A. Good proved the following

THEOREM A. *If C_{-1} denotes the residue and $C_{-1}C_0$ the constant term in the Laurent expansion of*

$$D(s) = \sum_{n=1}^{\infty} |a_n|^2 n^{-s} \quad \text{at } s = k,$$

then

$$\int_0^T \left| L_f \left(\frac{k}{2} + it \right) \right|^2 dt = 2C_{-1}T \left(\log \left(\frac{T}{2\pi e} \right) + C_0 \right) + O \left(T(\log T)^{\frac{2}{3}} \right)$$

as $T \rightarrow \infty$.

As a corollary, he deduced that

$$L_f \left(\frac{k}{2} + it \right) \ll |t|^{\frac{1}{3}} (\log |t|)^{\frac{5}{6}}.$$

Our main goal in this paper is the following. Let $f(z) = \sum_{n=1}^{\infty} a_f(n)e^{2\pi inz}$, $\Im z > 0$ be a holomorphic cusp form of even integral weight $k > 2$ with level N (i.e in $\Gamma_0(N)$) and

$a_f(1) = 1$. We also assume that f is a Hecke eigenform so that $a_f(n)$ are eigenvalues of all the Hecke operators. We are interested to estimate an upper bound for the average integral namely

$$I = \sum_f' \frac{1}{\langle f, f \rangle} \int_T^{2T} \left| L_f \left(\frac{k}{2} + it \right) \right|^2 dt$$

where \prime indicates that f runs over an orthonormal basis set, uniformly with respect to the conductor N . We prove

MAIN THEOREM. *We have*

$$\begin{aligned} I &= \sum_f' \frac{1}{\langle f, f \rangle} \int_T^{2T} \left| L_f \left(\frac{k}{2} + it \right) \right|^2 dt \\ &\ll_{k,\epsilon} NT(\log NT)^3(\log \log T)^2 + N^{5+10\epsilon} e^{-C(\log T)^2} \end{aligned}$$

uniformly for all levels N of f .

Remark 1. This main theorem may be compared with the result of A. Good. As a corollary to this main theorem, we obtain uniformly for all $N \ll T^\epsilon$, the inequality

$$I = \sum_f' \frac{1}{\langle f, f \rangle} \int_T^{2T} \left| L_f \left(\frac{k}{2} + it \right) \right|^2 dt \ll_{k,\epsilon} T^{1+\epsilon}$$

Remark 2. The main feature of the proof here is that we avoid the approximate functional equation (whereas Good's proof depends upon the approximate functional equation). Whenever a Dirichlet series has a functional equation, in [4] K. Chandrasekharan and Raghavan Narasimhan established that a form of the approximate functional equation can be proved and it has a nice form whenever the coefficients of the Dirichlet series under consideration are positive which can be utilised to study mean value theorems. Of course, in some special circumstances even if we do not have the coefficients of the Dirichlet series to be positive it is possible to use that form to study mean-value theorems. However K. Ramachandra observed that just with the functional equation, one can prove reasonable upper bounds for mean-value questions on certain lines and he has used this idea in many of his research work (see for example [28] and [29]). This crucial idea what we are going to use in proving our main theorem.

Remark 3. It should be mentioned here that a general mean value theorem (see theorem 1 of [22]) is available. It is not difficult to see that (choosing the parameters appropriately and

combining lemma 3.1 of section 3)

$$I = \sum_f' \frac{1}{\langle f, f \rangle} \int_T^{2T} \left| L_f \left(\frac{k}{2} + it \right) \right|^2 dt \ll_{k,\epsilon} T^{1+\epsilon}.$$

However, this estimate holds only when the conductor N of f satisfies the condition $N \ll 1$.

Acknowledgement : The author has great pleasure in thanking Professor Ram Murty for posing this problem some times back and for having fruitful and stimulating discussions related to the Poincare series topic. The author is highly indebted to Instituto de Matemáticas, UNAM, Morelia, Mexico for it's warm hospitality and grand financial support during his sabbatical stay there. He also wishes to express his gratitude to the referee for valuable comments.

2. Notation and preliminaries

The letters C and A (with or without suffixes) denote effective positive constants unless it is specified. It need not be the same at every occurrence. Throughout the paper we assume $T \geq T_0$ where T_0 is a large positive constant. We write $f(x) \ll g(x)$ to mean $|f(x)| < C_1 g(x)$ (sometimes we denote this by the O notation also). Let $s = \sigma + it$, and $w = u + iv$. The implied constants are all effective.

Let us recall some basic facts concerning the Poincaré series (see for example [33]).

Let Γ_∞ be the stabilizer of $i\infty$ in $\Gamma_0(N)$. The space $S_k(N)$ of cusp forms of weight k and level N is a finite dimensional vector space over C , spanned by the Poincaré series : for $m \geq 1$,

$$P_m(z, k, N) = \sum_{\gamma \in \Gamma_\infty \backslash \Gamma_0(N)} j(\gamma, z)^{-k} e(m\gamma z)$$

where

$$e(z) = e^{2\pi iz}, \quad j(\gamma, z) = (\det \gamma)^{-\frac{1}{2}} (cz + d)$$

and

$$\gamma = \begin{pmatrix} * & * \\ c & d \end{pmatrix}$$

In the case of $k = 2$, we do not have absolute convergence of the Poincaré series. However, the discussion below holds in this case as well. If $f \in S_k(N)$, we write the Fourier expansion of f as :

$$f(z) = \sum_{n=1}^{\infty} a_f(n) e(nz)$$

at $i\infty$. The space $S_k(N)$ has an inner product (Petersson inner product) :

$$\langle f, g \rangle = \int_{\Gamma_0(N)\backslash h} f(z)\overline{g(z)}y^k \frac{dx dy}{y^2}$$

where h denotes the upper half plane. Petersson proved that

$$a_f(n) = \frac{(4\pi n)^{k-1}}{\Gamma(k-1)} \langle f, P_n(\cdot, k, N) \rangle$$

so that if f_1, \dots, f_r is an orthonormal basis for $S_k(N)$, and

$$P_n(\cdot, k, N) = \sum_i c_i f_i$$

we obtain

$$c_i = \langle f_i, P_n(\cdot, k, N) \rangle = \frac{\Gamma(k-1)}{(4\pi n)^{k-1}} a_{f_i}(n).$$

Therefore, we have

$$\frac{(4\pi n)^{k-1}}{\Gamma(k-1)} P_n(\cdot, k, N) = \sum_i a_{f_i}(n) f_i.$$

If we compare the m -th coefficients on both sides, then we obtain

$$\frac{(4\pi n)^{k-1}}{\Gamma(k-1)} a_{P_n(m, k, N)} = \sum_i a_{f_i}(n) a_{f_i}(m).$$

The m -th Fourier coefficient of the n -th Poincaré series $P_n(z, k, N)$ can be computed as :

$$a_{P_n(m, k, N)} = \left(\frac{m}{n}\right)^{\frac{k-1}{2}} \left\{ \delta_{mn} + 2\pi i^{-k} \sum_{c \equiv 0 \pmod{N}} c^{-1} J_{k-1} \left(\frac{4\pi \sqrt{mn}}{c} \right) S_{\infty\infty}(m, n, c) \right\}$$

where $\delta_{mn} = 0$ unless $m = n$ in which case it is 1.

It is well known that (see [12], [13]), $L_f(s)$ (f with level N) satisfies the functional equation

$$(2.1) \quad L_f(s) = \chi_f(s) L_f(k-s),$$

where

$$(2.2) \quad \chi_f(s) = C \left(\frac{2\pi}{N}\right)^{2s-k} \frac{\Gamma(k-s)}{\Gamma(s)},$$

with $|C| = 1$. We also know that, in any fixed strip $a \leq \sigma \leq b$, as $t \rightarrow \infty$, we have

$$(2.3) \quad \Gamma(\sigma + it) = t^{\sigma+it-1/2} e^{-\pi/2-it+(i\pi/2)(\sigma-1/2)} \sqrt{2\pi} \left(1 + O\left(\frac{1}{t}\right)\right).$$

We observe that

$$(2.4) \quad |\chi_f(s)| = \left(\frac{2\pi}{N}\right)^{2\sigma-k} |t|^{k-2\sigma} \left(1 + O\left(\frac{1}{|t|}\right)\right)$$

as $t \rightarrow \infty$. Since, $L_f(s)$ is absolutely convergent in $\sigma > \frac{k+1}{2}$, we have

$$(2.5) \quad L_f\left(\frac{k+1}{2} + \epsilon + it\right) \ll_{\epsilon} 1.$$

From the functional equation, we have

$$(2.6) \quad \begin{aligned} \left|L_f\left(\frac{k}{2} - 1 + it\right)\right| &= \left|\chi_f\left(\frac{k}{2} - 1 + it\right) L_f\left(\frac{k}{2} + 1 + it\right)\right| \\ &\ll \left(\frac{2\pi}{N}\right)^{2\left(\frac{k}{2}-1\right)-k} |t|^{k-2\left(\frac{k}{2}-1\right)} \\ &\ll \left(\frac{2\pi}{N}\right)^{-2} |t|^2. \end{aligned}$$

Let H be the set of all holomorphic cusp forms (Hecke eigenforms) of even integral weight $k(\geq 2)$ of level N . We assume that $a_f(1) = 1 \forall f \in H$. We also observe that the set

$$\left\{ \frac{f}{\langle f, f \rangle} : f \in H \right\}$$

forms an orthonormal basis for H . From the Poincare series discussion above (see for example [33] also), we have (for $k \geq 2$)

$$(2.7) \quad \begin{aligned} E &= : \frac{\Gamma(k-1)}{(4\pi)^{k-1}} \sum'_f \frac{a_f(n)a_{\bar{f}}(m)}{\langle f, f \rangle (mn)^{\frac{k-1}{2}}} \\ &= \delta_{mn} + 2\pi i^{-k} \sum_{c \equiv 0 \pmod{N}} c^{-1} J_{k-1}\left(\frac{4\pi\sqrt{mn}}{c}\right) S_{\infty\infty}(m, n, c), \end{aligned}$$

where $\delta_{mn} = 0$ unless $m = n$ in which case it is 1. $J_{k-1}(x)$ is the Bessel function of order $k-1$, defined as

$$(2.8) \quad J_k(z) = \frac{\left(\frac{z}{2}\right)^k}{\sqrt{\pi}\Gamma\left(k + \frac{1}{2}\right)} \int_0^\pi (\sin \theta)^{2k} \cos(z \cos \theta) d\theta$$

and $S_{\infty\infty}(m, n, c)$ is the Kloosterman sum

$$(2.9) \quad S_{\infty\infty}(m, n, c) = S(m, n, Nl) = \sum_{\substack{d \pmod{Nl} \\ d\bar{d} \equiv 1 \pmod{Nl}}} e\left(\frac{md + n\bar{d}}{Nl}\right).$$

Here $e(z) = e^{2\pi iz}$. We have (from Deligne's estimate)

$$(2.10) \quad a_f(n) \ll d(n)n^{\frac{k-1}{2}},$$

and (from Weil's estimate)

$$(2.11) \quad S(m, n, c) \leq (m, n, c)^{\frac{1}{2}} d(c) c^{\frac{1}{2}}.$$

It is known that

$$(2.12) \quad J_k(x) \ll \frac{x^k}{2^k \Gamma\left(k + \frac{1}{2}\right)}$$

3. Some lemmas

LEMMA 3.1 (i) For $k = 2$

$$\sum_f' \frac{1}{\langle f, f \rangle} = \frac{(4\pi)^{k-1}}{\Gamma(k-1)} + O\left(N^{-\frac{3}{2}+\epsilon}\right)$$

and

(ii) for $k > 2$,

$$\sum_f' \frac{1}{\langle f, f \rangle} = \frac{(4\pi)^{k-1}}{\Gamma(k-1)} + O\left(N^{-(k-1)+\epsilon}\right).$$

Proof. From (2.7), (2.11) and (2.12), we get (for $k = 2$)

$$(3.1) \quad \begin{aligned} \sum_{c \equiv 0 \pmod{N}} c^{-1} J_{k-1}\left(\frac{4\pi\sqrt{mn}}{c}\right) S(m, n, c) &\ll (mn)^{\frac{1}{2}} (m, n)^{\frac{1}{2}} \left(\sum_{j=1}^{\infty} \frac{d(Nj)}{(Nj)^{\frac{3}{2}}}\right) \\ &\ll N^{-\frac{3}{2}+\epsilon} (mn)^{\frac{1}{2}} (m, n)^{\frac{1}{2}}, \end{aligned}$$

and for $k > 2$,

$$(3.2) \quad \sum_{c \equiv 0 \pmod{N}} c^{-1} J_{k-1} \left(\frac{4\pi \sqrt{mn}}{c} \right) S(m, n, c) \ll_k \sum_{c \equiv 0 \pmod{N}} \frac{(mn)^{\frac{k-1}{2}} d(c)}{c^{k-1}} \\ \ll_k N^{-(k-1)+\epsilon} (mn)^{\frac{k-1}{2}}.$$

From (3.1.1) and (3.1.2), we get

$$(3.3) \quad E =: \frac{\Gamma(k-1)}{(4\pi)^{k-1}} \sum'_{f < f, f >} \frac{a_f(n) a_f(\bar{m})}{(mn)^{\frac{k-1}{2}}} \\ = \delta_{mn} + O \left(\max \left(N^{-\frac{3}{2}+\epsilon} (mn)^{\frac{1}{2}} (m, n)^{\frac{1}{2}}, N^{-(k-1)+\epsilon} (mn)^{\frac{k-1}{2}} \right) \right).$$

Taking $m = n = 1$ in (3.1.3) and noticing that $a_f(1) = 1$, we obtain

$$(3.4) \quad \frac{\Gamma(k-1)}{(4\pi)^{k-1}} \sum'_{f < f, f >} \frac{1}{f} = 1 + O \left(\max \left(N^{-\frac{3}{2}+\epsilon}, N^{-(k-1)+\epsilon} \right) \right).$$

This proves the lemma.

LEMMA 3.2 For σ in the range $\frac{k}{2} - 1 \leq \sigma \leq \frac{k+1}{2} + \epsilon$, uniformly, we have the estimate

$$L_f(\sigma + it) \ll \left(\frac{|t|N}{2\pi} \right)^{\frac{4}{3} \left(\frac{k+1}{2} - \sigma \right) + \epsilon}.$$

Proof. Follows by applying maximum-modulus principle to a suitable function, namely

$$F(w) = L_f(w) e^{(w-s)^2} X^{w-s}$$

over a suitable rectangle and we can choose $X = \left(\frac{|t|N}{2\pi} \right)^{\frac{4}{3} + \epsilon}$.

LEMMA 3.3. (*Montgomery-Vaughan*) If b_n is an infinite sequence of complex numbers such that $\sum_{n=1}^{\infty} n |b_n|^2$ is convergent, then

$$\int_T^{T+H} \left| \sum_{n=1}^{\infty} b_n n^{-it} \right|^2 dt = \sum_{n=1}^{\infty} |b_n|^2 (H + O(n)).$$

Proof. See for example lemma 3.3 of [26] or [31].

4. Proof of the theorem

Let $s_1 = \frac{k}{2} + it$ and let Y and Y_1 be two parameters satisfying $T^{\frac{1}{2}} \leq Y, Y_1 \leq (NT)^A$ to be chosen appropriately later. Let $\epsilon_1 = (\log T)^{-1}$. By Mellin's transformation,

$$\begin{aligned}
S &=: \sum_{n=1}^{\infty} \frac{a_f(n)}{n^{s_1}} e^{-\frac{n}{Y}} \\
&= \frac{1}{2\pi i} \int_{\substack{\Re w = 1/2 + \epsilon, \\ |v| \leq (\log T)^2}} L_f(s_1 + w) Y^w \Gamma(w) dw + O\left(Y^{\frac{1}{2} + \epsilon} e^{-C(\log T)^2}\right) \\
&= L_f(s_1) + \frac{1}{2\pi i} \int_{\substack{\Re w = -1 + 2\epsilon_1, \\ |v| \leq (\log T)^2}} L_f(s_1 + w) Y^w \Gamma(w) dw + O\left(Y^{\frac{1}{2} + \epsilon} e^{-C(\log T)^2}\right) \\
&+ O\left(Y^{\frac{1}{2} + \epsilon} \left(\frac{TN}{2\pi}\right)^{2+\epsilon} e^{-C(\log T)^2}\right) \\
(4.1) \quad &= L_f(s_1) + I + O\left(Y^{\frac{1}{2} + \epsilon} N^{2+\epsilon} e^{-C(\log T)^2}\right) \text{ (say)}
\end{aligned}$$

by moving the line of integration to $\Re w = -1 + 2\epsilon_1$. Now

$$\begin{aligned}
I &= \frac{1}{2\pi i} \int_{\substack{\Re w = -1 + 2\epsilon_1, \\ |v| \leq (\log T)^2}} L_f(s_1 + w) Y^w \Gamma(w) dw \\
&= \frac{1}{2\pi i} \int_{\substack{\Re w = -1 + 2\epsilon_1, \\ |v| \leq (\log T)^2}} \chi(s_1 + w) \{Q_1 + Q_2\} Y^w \Gamma(w) dw \\
(4.2) \quad &= I_1 + I_2,
\end{aligned}$$

where

$$Q_1 = \sum_{n \leq Y_1} a_f(n) n^{s_1 + w - k} ; \quad Q_2 = \sum_{n > Y_1} a_f(n) n^{s_1 + w - k}.$$

We note that in I_2 , $\Re(s_1 + w) = \frac{k}{2} - 1 + 2\epsilon_1$. We have

$$\begin{aligned}
\sum_{n > Y_1} a_f(n) n^{s_1 + w - k} &= \sum_{Y_1 < n \leq Y_1^{10}} a_f(n) n^{s_1 + w - k} \\
&+ O\left(\sum_{n > Y_1^{10}} d(n) n^{\frac{k-1}{2} + \frac{k}{2} - 1 + 2\epsilon_1 - k}\right) \\
(4.3) \quad &= \sum_{Y_1 < n \leq Y_1^{10}} a_f(n) n^{s_1 + w - k} + O\left(Y_1^{-5 + 20\epsilon_1}\right).
\end{aligned}$$

Note that we have used the inequality (2.10). Using Hölder's inequality and a theorem of Montgomery and Vaughan (see [2]), we get

$$\int_T^{2T} |I_2|^2 dt \ll \int_T^{2T} \left| \int_{\substack{\Re w = -1 + 2\epsilon_1, \\ |v| \leq (\log T)^2}} \chi(s_1 + w) \left\{ \sum_{Y_1^{10} \geq n > Y_1} a_f(n) n^{s_1 + w - k} \right\} Y^w \Gamma(w) dw \right|^2 dt$$

$$\begin{aligned}
& + \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} T^{5+20\epsilon_1} Y^{-2} Y_1^{-10+40\epsilon_1} \\
& \ll (\log T)^2 \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} \frac{T^{4+20\epsilon_1}}{Y^2} \int_{T-(\log T)^2}^{2T+(\log T)^2} \sum_U \left| \sum_{(U,2U)} a_f(n) n^{s_1+\Re w-k} \right|^2 dt \\
& + \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} T^{5+20\epsilon_1} Y^{-2} Y_1^{-10+40\epsilon_1} \\
& \ll (\log T)^2 \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} \frac{T^{4+20\epsilon_1}}{Y^2} \sum_U \sum_{(U \leq n \leq 2U)} \frac{|a_f(n)|^2}{n^{2(2k-1-(k-\frac{3}{2}+2\epsilon_1))}} n \\
& + \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} T^{5+20\epsilon_1} Y^{-2} Y_1^{-10+40\epsilon_1} \\
& \ll (\log T)^2 \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} \frac{T^{4+20\epsilon_1}}{Y^2} \sum_U \sum_{(U \leq n \leq 2U)} \frac{(d(n))^2 n^{k-1}}{n^{2(k-(\frac{k}{2}-1+2\epsilon_1))}} \\
& + \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} T^{5+20\epsilon_1} Y^{-2} Y_1^{-10+40\epsilon_1} \\
& \ll (\log T)^2 \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} \frac{T^{4+20\epsilon_1}}{Y^2} \sum_{j=0}^{C \log Y_1} (\log Y_1)^3 (2^j Y_1)^{-1+4\epsilon_1} \\
& + \left(\frac{2\pi}{N}\right)^{-4+4\epsilon_1} T^{5+20\epsilon_1} Y^{-2} Y_1^{-10+40\epsilon_1} \\
& \ll (\log T)^2 N^{4-4\epsilon_1} T^{4+20\epsilon_1} Y^{-2} Y_1^{-1+4\epsilon_1} (\log Y_1)^3 \\
(4.4) \quad & + N^{4-4\epsilon_1} T^{5+20\epsilon_1} Y^{-2} Y_1^{-10+40\epsilon_1}.
\end{aligned}$$

Also we have

$$\begin{aligned}
I_1 &= \frac{1}{2\pi i} \int_{\substack{\Re w = -1+2\epsilon_1, \\ |v| \leq (\log T)^2}} \chi(s_1+w) \left\{ \sum_{n \leq Y_1} a_f(n) n^{s_1+w-k} \right\} Y^w \Gamma(w) dw \\
&= \frac{1}{2\pi i} \int_{\substack{\Re w = -\frac{1}{2}, \\ |v| \leq (\log T)^2}} \chi(s_1+w) \left\{ \sum_{n \leq Y_1} a_f(n) n^{s_1+w-k} \right\} Y^w \Gamma(w) dw \\
(4.5) \quad &+ O\left((NT)^2 (\log Y_1)^2 Y^{-\frac{1}{2}} e^{-C \log T}\right)
\end{aligned}$$

by moving the line of integration to $\Re w = -\frac{1}{2}$. Notice that, if $-1+2\epsilon_1 \leq u \leq -\frac{1}{2}$, then $\frac{k}{2}-1+2\epsilon_1 \leq \Re(s_1+w) \leq \frac{k}{2}-\frac{1}{2}$ and $-1-\frac{k}{2}+2\epsilon_1 \leq \Re(s_1+w-k) \leq -\frac{k}{2}-\frac{1}{2}$. Therefore, for $T \leq t \leq 2T$, we have

$$|\chi(s_1+w)| \ll \left(\frac{NT}{2\pi}\right)^{k-2(2\epsilon_1+\frac{k}{2}-1)} \ll (NT)^2.$$

Also, we have

$$\sum_{n \leq Y_1} a_f(n) n^{s_1+w-k} \ll \sum_{n \leq Y_1} d(n) n^{\frac{k-1}{2}+\frac{k-1}{2}-k} \ll_k (\log Y_1)^2,$$

$$Y^w \ll Y^{-\frac{1}{2}} \quad \text{and} \quad \Gamma(w) \ll e^{-C(\log T)^2}$$

on horizontal lines. Again using Montgomery- Vaughan Theorem and the inequality (2.10), we obtain

$$\begin{aligned}
\int_T^{2T} |I_1|^2 dt &\ll (\log T)^2 \left(\frac{2\pi}{N}\right)^{-2} \frac{T^2}{Y} \int_{T-(\log T)^2}^{2T+(\log T)^2} \left| \sum_{n \leq Y_1} a_f(n) n^{s_1 + \Re w - k} \right|^2 dt \\
&+ (NT)^4 (\log Y_1)^4 Y^{-1} e^{-C(\log T)^2} \\
&\ll (\log T)^2 N^2 \frac{T^2}{Y} \sum_{n \leq Y_1} \frac{|a_f(n)|^2 n}{n^{2(\frac{k}{2} + \frac{1}{2})}} \\
&+ (NT)^4 (\log Y_1)^4 Y^{-1} e^{-C(\log T)^2} \\
&\ll (\log T)^2 \frac{T^3}{Y} \sum_{n \leq Y_1} \frac{(d(n))^2 n^k}{n^{k+1}} \\
&+ (NT)^4 (\log Y_1)^4 Y^{-1} e^{-C(\log T)^2} \\
&\ll (\log T)^2 N^2 \frac{T^2}{Y} (\log Y_1)^3 \\
(4.6) \quad &+ (NT)^4 (\log Y_1)^4 Y^{-1} e^{-C(\log T)^2}.
\end{aligned}$$

Now, we have

$$S = \sum_{n=1}^{\infty} \frac{a_f(n)}{n^{s_1}} e^{-n/Y}.$$

$$\begin{aligned}
\int_T^{2T} |S|^2 dt &= \sum_{n=1}^{\infty} \frac{|a_f(n)|^2}{n^k} e^{-\frac{2n}{Y}} (T + O(n)) \\
&= T \sum_{n=1}^{\infty} \frac{|a_f(n)|^2}{n^k} e^{-\frac{2n}{Y}} \\
&+ O\left(\sum_{n \leq Y/2} \frac{|a_f(n)|^2}{n^k} n + \sum_{n > Y/2} \frac{|a_f(n)|^2}{n^k} n \left(\frac{Y}{n}\right)^{2\alpha+2} \right) \\
(4.7) \quad &= T \sum_{n=1}^{\infty} \frac{|a_f(n)|^2}{n^k} e^{-\frac{2n}{Y}} + O(Y(\log Y)^3)
\end{aligned}$$

with a suitable α .

From (4.1) to (4.7), we obtain,

$$\int_T^{2T} |L_f(s_1)|^2 dt = T \sum_{n=1}^{\infty} \frac{|a_f(n)|^2}{n^k} e^{-\frac{2n}{Y}}$$

$$\begin{aligned}
& + O\left((\log T)^2 N^{4-4\epsilon_1} Y^{-2} T^4 Y_1^{-1+4\epsilon_1} (\log Y_1)^3\right) \\
& + O\left(N^{4-4\epsilon_1} T^5 Y^{-2} Y_1^{-10+40\epsilon_1}\right) + O\left(N^2 Y^{-1} T^2 (\log Y_1)^3 (\log T)^2\right) \\
& + O\left(Y (\log Y)^3\right) + O\left(Y^{1+2\epsilon} N^{4+2\epsilon} e^{-C(\log T)^2}\right) \\
& + O\left(N^4 (\log Y_1)^4 Y^{-1} e^{-C(\log T)^2}\right) \\
(4.8) \quad & = T \sum_{n=1}^{\infty} \frac{|a_f(n)|^2}{n^k} e^{-\frac{2n}{Y}} + O(E_1 + E_2 + E_3 + E_4 + E_5 + E_6) \quad \text{say,}
\end{aligned}$$

where E_i are with obvious notion. It is clear from lemma 3.1 that

$$(4.9) \quad \sum_f' \frac{1}{\langle f, f \rangle} E_i \ll E_i \quad (i = 1, 2, 3, 4, 5, 6).$$

Therefore, we get

$$\begin{aligned}
(4.10) \quad \sum_f' \frac{1}{\langle f, f \rangle} \int_T^{2T} |L_f(s_1)|^2 dt & = T \sum_f' \frac{1}{\langle f, f \rangle} \sum_{n=1}^{\infty} \frac{|a_f(n)|^2}{n^k} e^{-\frac{2n}{Y}} \\
& + O(E_1 + E_2 + E_3 + E_4 + E_5 + E_6).
\end{aligned}$$

We observe that

$$\begin{aligned}
S_1 & =: \sum_f' \frac{1}{\langle f, f \rangle} \sum_{n=1}^{\infty} \frac{|a_f(n)|^2}{n^k} e^{-\frac{2n}{Y}} \\
& = \sum_{n=1}^{\infty} \frac{e^{-\frac{2n}{Y}}}{n} \sum_f' \frac{|a_f(n)|^2}{\langle f, f \rangle n^{k-1}} \\
& \ll \sum_{n=1}^{\infty} \frac{e^{-\frac{2n}{Y}} (d(n))^2}{n} \sum_f' \frac{1}{\langle f, f \rangle} \quad (\text{by (2.10)}) \\
& \ll_k \sum_{n=1}^{\infty} \frac{e^{-\frac{2n}{Y}} (d(n))^2}{n} \quad (\text{from lemma 3.1}) \\
& \ll_k \sum_{n \leq \frac{Y}{2}} \frac{(d(n))^2}{n} \left(1 + O\left(\frac{n}{Y}\right)\right) + \sum_{n \leq \frac{Y}{2}} \frac{(d(n))^2 Y}{n} \\
(4.11) \quad & \ll_k (\log Y)^3.
\end{aligned}$$

Hence, we obtain

$$\begin{aligned}
(4.12) \quad \sum_f' \frac{1}{\langle f, f \rangle} \int_T^{2T} |L_f(s_1)|^2 dt & \ll_k T (\log Y)^3 \\
& + E_1 + E_2 + E_3 + E_4 + E_5 + E_6.
\end{aligned}$$

First, we choose Y such that E_3 and E_4 to be of the same order. This suggests us to choose $Y = NT$. This means that

$$\max(E_3, E_4) \ll NT \left\{ (\log Y_1)^3 (\log T)^2 + (\log Y)^3 \right\}.$$

Now, we are forced to choose $Y_1 = NT$ and we notice that

$$E_1 \ll NT (\log Y_1)^3 (\log T)^2.$$

Hence, for this choice $Y = Y_1 = NT$, we get

$$(4.13) \quad \sum_f \frac{1}{\langle f, f \rangle} \int_T^{2T} |L_f(s_1)|^2 dt \ll_k T(\log(NT))^3 + NT(\log(NT))^3(\log T)^2 \\ + N^{5+10\epsilon} e^{-C(\log T)^2} + N^3(\log(NT))^4 e^{-C(\log T)^2},$$

which proves the theorem, since, on the r.h.s of (4.13), the first and the last terms are dominated by the other two terms.

- [1] R. Balasubramanian, *An improvement of a theorem of Titchmarsh on the mean-square of $|\zeta(\frac{1}{2} + it)|$* , Proc. London Math. Soc., (3) **36** (1978), 540-576.
- [2] R. Balasubramanian and K. Ramachandra, *An alternative approach to a theorem of Tom Meurman*, Acta Arith., **55** (1990), 351-364.
- [3] K. Chandrasekharan and R. Narasimhan, *Functional equations with multiple Gamma factors and the average order of arithmetical functions*, Ann. Math. **76** (1962), 93-136.
- [4] K. Chandrasekharan and R. Narasimhan, *The approximate functional equation for a class of zeta-functions*, Math Ann., **152** (1963), 30-64.
- [5] P. Deligne, *Formes modulaires et représentations ℓ - adiques*, Sém. Bourbaki, (1968/69), exposés 355.
- [6] P. Deligne, *La conjecture de Weil-I* Inst.Hautes Études Sci.Pub. Math **43** (1974), 273-307.
- [7] P.X. Gallagher, *Local mean value and density estimates for Dirichlet L- functions*, Indag Math., **37** (1975), 259-264.
- [8] D. Goldfeld and P. Sarnak, *Sums of Kloosterman sums*, Invent. Math., **71** (1983), 251-293.
- [9] A. Good, *Approximative funktionalgleichungen und Mittelwertsätze für Dirichletreihen, die Spitzenformen assoziiert sind*, Comment. Math. Helvetici., **50** (1975), 327-361.
- [10] A. Good, *The square mean of Dirichlet series associated with cusp forms*, Mathematika., **29** (1982), 278-295.

- [11] D.R. Heath-Brown, *The twelfth power moment of the Riemann zeta-function*, Quart. J. Math (Oxford Ser) (2) **29** (1978), 443-462.
- [12] E. Hecke, *Über die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichung*, Math. Ann. **112** (1936), 664-699.
- [13] E. Hecke, *Über Dirichlet-Reihen mit Funktionalgleichung und ihre Nullstellen auf der Mittleregeraden*, München Akad.Sitzungsber. II, **8** (1937), 73-95.
- [14] A. Ivić, *The Riemann zeta-function*, Wiley (1985).
- [15] A. Ivić, *Large values of certain number theoretic error terms*, Acta Arith., **56** (1990), 135-159.
- [16] A. Ivić and Y. Motohashi, *The mean square of the error term for the fourth power moment of the zeta-function*, Proc. London Math. Soc., (3) **69** (1994), 309-329.
- [17] H. Iwaniec, *Fourier coefficients of cusp forms and the Riemann zeta-function*, Bordeaux Sémin. Théorie des nombres, exp no: **18** (1979/80), 1-36.
- [18] H. Iwaniec, *Topics in classical automorphic forms*, Vol **17**, American Math. Soc., Providence, Rhode Island (1997).
- [19] M. Jutila and Y. Motohashi, *Mean value estimates for exponential sums and L- functions : a spectral - theoretic approach*, J. Reine Angew Math., **459** (1995), 61-87.
- [20] M. Jutila, *The fourth moment of Riemann's zeta-function and the additive divisor problem*, Analytic number theory, Vol 2 (Allerton Park, IL, 1995) 517-536, Progr. Math., **139**, Birkhauser Boston, Boston, MA, 1996.
- [21] M. Jutila, *Mean values of Dirichlet series via Laplace transforms*, Analytic number theory (Kyoto, 1996), 169-207, London Math. Soc., Lecture Notes Ser., **247** (1997), Cambridge Univ. Press, Cambridge 1997.
- [22] S. Kanemitsu, A. Sankaranarayanan and Y. Tanigawa, *A mean value theorem for Dirichlet series and a general divisor problem*, Monatsh. Math., **136** (2002), 17-34.
- [23] K. Matsumoto and T. Meurman, *The mean-square of the Riemann zeta-function in the critical strip-II*, Acta Arith., **68** (1994), 369-382.
- [24] T. Meurman, *A generalisation of Atkinson's formula to L- functions*, Acta Arith., **47** (1986), 351-370.

- [25] H.L. Montgomery, *Topics in multiplicative number theory*, (Springer, Berlin) (1971).
- [26] H.L. Montgomery and R.C. Vaughan, *Hilbert's inequality*, J. London Math. Soc., (2) **8** (1974) , 73-82.
- [27] A. Perelli, *General L- functions*, Ann. Mat. Pura Appl., (4) **130** (1982), 287-306.
- [28] K. Ramachandra, *A simple proof of the mean fourth power estimate for $\zeta(\frac{1}{2} + it)$ and $L(\frac{1}{2} + it, \chi)$* , Annali. della. Scuola Normale Superiore di Pisa, Classe di Sci., Ser IV, **1** (1974), 81-97.
- [29] K. Ramachandra, *Application of a theorem of Montgomery and Vaughan to the zeta-function*, J London Math. Soc.,**10** (1975), 482-486.
- [30] K. Ramachandra, *Some problems of analytic number Theory-I*, Acta Arith., **31** (1976), 313-324.
- [31] K. Ramachandra, *Some remarks on a theorem of Montgomery and Vaughan*, J.Number Theory **11** (1979), 465-471.
- [32] K. Ramachandra, *Progress towards a conjecture on the mean-value of Titchmarsh series*, in: Recent Progress in Analytic Number Theory, H. Halberstam and C. Hooley (eds.), **1** Academic Press, (1981), 303-318.
- [33] M. Ram Murty, *The analytic rank of $J_0(N)(\mathbf{Q})$* Canadian Math. Soc., Conf. Proc., **15** (1995), 263-277.
- [34] M. Ram Murty and V. Kumar Murty, *Mean values of derivatives of modular L- series*, Ann Math., **133** (1991), 447-475.
- [35] R.A. Rankin, *Contributions to the theory of Ramanujan's function $\tau(n)$ and similar arithmetical functions - I* , Proc. Cambridge. Phil. Soc., **35** (1939), 351-356.
- [36] R.A. Rankin, *Contributions to the theory of Ramanujan's function $\tau(n)$ and similar arithmetical functions - II* , Proc. Cambridge. Phil. Soc., **35** (1939), 357-372.
- [37] A. Sankaranarayanan, *Fundamental properties of symmetric square L- functions-I*, Illinois J. Math., **46** (2002), 23-43.
- [38] A. Selberg, *Contributions to the theory of the Riemann zeta-function*, in: Collected Papers, Vol I, Springer, (1989), 214-280.

- [39] E.C. Titchmarsh, *The theory of the Riemann zeta-function*, 2nd Edition (edited by D.R.Heath-Brown), Clarendon Press (Oxford) (1986).

PRESENT ADDRESS : INSTITUTO DE MATEMÁTICAS, UNAM, MORELIA CAMPUS, AP. POSTAL 61-3 (XANGARI), CP 58 089, MORELIA, MICHOACAN, MEXICO .

E-mail address : `sank@matmor.unam.mx`

PERMANENT ADDRESS : SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH , HOMI BHABHA ROAD, MUMBAI - 400 005, INDIA.

E-mail address : `sank@math.tifr.res.in`

(Accepted on 03-09-2003)