



HAL
open science

Heat semigroup and singular PDEs

Ismaël Bailleul, Frederic Bernicot

► **To cite this version:**

Ismaël Bailleul, Frederic Bernicot. Heat semigroup and singular PDEs. Journal of Functional Analysis, 2016, 270 (9), pp.3344-3452. hal-01110466v2

HAL Id: hal-01110466

<https://hal.science/hal-01110466v2>

Submitted on 28 May 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Heat semigroup and singular PDEs

I. BAILLEUL¹ and F. BERNICOT²

(with an Appendix by F. Bernicot & D. Frey)

ABSTRACT. We provide in this work a semigroup approach to the study of singular PDEs, in the line of the paracontrolled approach developed recently by Gubinelli, Imkeller and Perkowski. Starting from a heat semigroup, we develop a functional calculus and introduce a paraproduct based on the semigroup, for which commutator estimates and Schauder estimates are proved, together with their paracontrolled extensions. This machinery allows us to investigate singular PDEs in potentially unbounded Riemannian manifolds under mild geometric conditions. As an illustration, we study the generalized parabolic Anderson model equation and prove, under mild geometric conditions, its well-posed character in Hölder spaces, in small time on a potentially unbounded 2-dimensional Riemannian manifold, for an equation driven by a coloured noise, and for all times for the linear parabolic Anderson model equation in 2-dimensional unbounded manifolds. This machinery can be extended to an even more singular setting and deal with Sobolev spaces rather than Hölder spaces.

CONTENTS

Introduction	2
1.1 State of the art	2
1.2 A generalized parabolic Anderson model	3
Functional calculus adapted to the heat semigroup	5
2.1 Heat semigroup on a doubling space	5
2.2 Time derivatives and Carré du champ of the semigroup	10
2.3 Hölder and Besov spaces through the heat semigroup	12
Paraproduct and commutator estimates in Hölder spaces	16
3.1 Paraproducts based on the semigroup	16
3.2 Paraproduct estimates	18
3.3 Commutator estimates	20
3.4 Paralinearization and composition estimates	25
3.5 Schauder estimates	29
Paracontrolled calculus	30
4.1 Paracontrolled distributions	32
4.2 Schauder estimate for paracontrolled distributions	34
The (generalized) parabolic Anderson Model in dimension 2	36
5.1 Local well-posedness result for generalized PAM	36
5.2 Global well-posedness result for linear PAM	38
5.3 Renormalization for a weighted noise	41
Heat kernel and technical estimates	48
Extension of the theory	54
B.1 Regularity assumptions	54
B.2 Functional calculus and gradient estimates in Hölder and Sobolev spaces	57
B.3 Paraproduct and commutator estimates in Hölder-Sobolev spaces	62
B.4 Resolution of PAM in such a 2-dimensional setting	72

¹I.B. was partly supported by the ANR project "Retour Post-doctorant", no. 11-PDOC-0025; I.B. also thanks the U.B.O. for their hospitality, part of this work was written there.

²F. Bernicot's research is partly supported by ANR projects AFoMEN no. 2011-JS01-001-01 and HAB no. ANR-12-BS01-0013.

Introduction

1.1. **State of the art** Following the recent breakthrough of Hairer [32] and Gubinelli, Imkeller, Perkowski [29], there has been recently a tremendous activity in the study of parabolic singular partial differential equations (PDEs), such as the KPZ equation

$$(\partial_t - \partial_x^2)u = (\partial_x u)^2 + \xi,$$

the stochastic quantization equation

$$(\partial_t - \Delta)u = -u^3 + \xi,$$

or the Parabolic Anderson Model equation

$$(\partial_t - \Delta)u = F(u)\xi$$

in all of which ξ stands for a space or space-time white noise. Each of these equations involves, under the form of a product, a term which does not make sense a priori, given the expected regularity of the solution in terms of the regularity of the noise ξ . Hairer's theory of regularity structures is built on the insights of earlier works [33, 37, 34] on $(1+1)$ -dimensional space-time problems where he used the framework of rough paths theory, under the form of Gubinelli's controlled paths, to make sense of previously ill-posed singular PDEs and give a meaningful solution theory. Rough paths theory was used in this approach as a framework for studying the properties in the 1-dimensional *space variable* of potential solutions. However, the very notion of a rough path is intimately linked with the 1-dimensional time axis that parametrizes paths.

To by-pass this barrier, both the theory of regularity structures and the paracontrolled approach developed in [29] take as a departure point the fact that, like in rough paths theory, to make sense of the equation, one needs to enrich the noise ξ into a finite collection of objects/distributions, and that one should try and describe the potential solution of a singular PDE in terms of that enriched noise. The latter depends on the equation under study and plays in the theory of regularity structures the role played by polynomials in the usual C^k world to give local descriptions of functions under the form of Taylor expansions at every space-time point. The description of a solution in the paracontrolled approach is of a different nature and rests on a global comparison with the solution to a linear equation $((\partial_t - \Delta)u = \xi$, in the above examples) via the use of Bony's paraproduct. In both approaches, the use of an ansatz for the solution space allows for fixed point arguments to give a robust solution theory where the solution becomes a continuous function of all the parameters of the problem.

So far, both theories have only been formulated and tested on some singular PDEs on the torus, to the exception of the works [59, 60] on singular perturbations of the Navier-Stokes equation on \mathbb{R}^3 , and the forthcoming work [36] on the parabolic Anderson model equation in \mathbb{R}^3 . We introduce in the present work a functional analytic setting in which we are able to extend the paracontrolled approach of [29] to investigate singular PDEs of the form

$$(\partial_t + L)u = F(u, \xi)$$

for a nonlinear term $F(u, \xi)$, on potentially unbounded (Riemannian or even sub-Riemannian) manifolds or graphs. (The change of sign $-$ to $+$ in the operator is irrelevant.) This is a priori far from obvious as the main analytic tool used in the paracontrolled approach in the torus involves some tools from Fourier analysis that do not make sense on manifolds or graphs. We develop to that end a functional calculus adapted to the heat semigroup associated with the operator $(\partial_t + L)$, which we use to define a paraproduct enjoying the same regularity properties as its Euclidean analogue. Such paraproducts adapted to a semigroup, as well as

a parilinearization theory, have already been studied in recent works [8, 12]. However, the irregular character of the noises ξ involved in the above motivating equations requires us to improve the definition of such paraproducts so as to build a framework where to consider regularity with a negative exponent; such an extension will be provided here. Building on these tools, one can set up, as in [29], a framework where to investigate the well-posed character of a whole class of parabolic singular PDEs. It is especially nice that all the objects in our framework are defined uniquely in terms of semigroups, unlike the notions of Hölder spaces used in the theory of regularity structures that involve a metric structure unrelated to the equation under study. As a by-product, we are able to handle some general classes of operators L whose treatment seem to be beyond the present-day scope of the theory of regularity structures, as illustrated in some examples given in section 2.1.

It is unclear presently how one can adapt the different notions and tools of the theory of regularity structures to extend them to a (Lipschitz) manifold or graph setting, or to other second order operators (other than the Laplace operator), as well as to work with Sobolev spaces (instead of Hölder spaces). Apart from the very definition of a regularity structure on a manifold, the existence of the reconstruction operator in this setting seems in particular challenging, as its proof in \mathbb{R}^d involves some deep results on wavelets that were not proved so far to hold true on generic manifolds, not even on all open sets of \mathbb{R}^d . Their extension to a non-smooth setting also seems highly non-trivial. So it comes as a good news that one can use some reasonably elaborate theory of semigroups to implement the alternative machinery of paracontrolled calculus in that setting; as described below, it also allows us to have much flexibility on the operator L and also on the geometry of the ambient space. Roughly speaking, we could say that the point of view of the theory of regularity structures relies on the metric and differential properties of the underlying space, while the present extension of the paracontrolled calculus corresponds to a functional point of view adapted to the operator L involved in the parabolic singular PDEs. We link here these two sides of the medal by requiring from the heat semigroup $(e^{-tL})_{t>0}$ to have a kernel together with its gradient, that satisfies pointwise Gaussian bounds; this describes in some sense the link between the functional calculus and the ambient space, with its metric and its differential geometry.

Moreover, we will detail in Appendix B, how this approach can be used in the context of Sobolev spaces (instead of Hölder spaces). From a technical point of view, it is a bit more difficult since Sobolev spaces involve simultaneously all the frequencies, whereas for Hölder spaces we can work at a fixed frequency scale. We do not know how such extension could also be implemented through the regularity structure's theory.

Motivated by this observation, the first part of this work is devoted to a precise study of the so-called paracontrolled calculus in a very abstract setting, given by a doubling ambient space, equipped with a self-adjoint operator $-L$ generating a semigroup with Gaussian bounds for its kernel and its gradient. A suitable definition of paraproducts is given, and the main rules of calculus for paracontrolled distributions are described. This general theory is all we need to study a number of parabolic singular PDEs.

1.2. A generalized parabolic Anderson model As an illustration, we shall study the generalized parabolic Anderson model equation (gPAM)

$$\partial_t u + Lu = F(u) \xi, \quad u(0) = u_0,$$

on some possibly unbounded 2-dimensional Riemannian manifold M satisfying some mild geometric conditions. One can take as operator the Laplace-Beltrami operator or some sub-elliptic diffusion operator; see section 2.1 for examples. The nonlinearity F is \mathcal{C}_b^3 , and ξ stands above for a coloured Gaussian noise with weight in $L^2 \cap L^\infty$ – see definition 5.5. The following results involve some Hölder spaces \mathcal{C}^γ , with negative exponents γ , that are defined in section

2.3 in terms only of the semigroup $(P_t)_{t \geq 0}$ generated by $-L$. We denote by X the solution to the equation

$$\partial_t X + LX = \xi,$$

given by the formula $X(t) := \int_0^t P_{t-s}(\xi) ds$. The following theorem is given in more precise form in theorems 5.1 and 5.4. There is no need right now to understand precisely what the 'resonant' term $\Pi(X, \zeta)$ below is; together with ζ it forms the above mentioned enriched noise that makes the theory so efficient.

Theorem 1.1. Let $\alpha \in (\frac{2}{3}, 1)$, an initial data $u_0 \in \mathcal{C}^{2\alpha}$, a nonlinearity $F \in C_b^3$, and a time horizon $T > 0$ be fixed. Assume that $\zeta \in \mathcal{C}^{\alpha-2}$.

- (a) **Local well-posedness for (gPAM).** If the resonant term $\Pi(X, \zeta)$ is well-defined as a continuous function from $[0, T]$ to $\mathcal{C}^{\alpha-2}$, then for a small enough time horizon T , the generalized PAM equation

$$(1.1) \quad \partial_t u + Lu = F(u) \zeta, \quad u(0) = u_0$$

has a unique solution in some function space.

- (b) **Global well-posedness for (PAM).** If the resonant term $\Pi(X, \zeta)$ is well-defined as a continuous function from \mathbb{R}_+ to a weighted version of $\mathcal{C}^{\alpha-2}$, then the PAM equation

$$\partial_t u + Lu = u \zeta, \quad u(0) = u_0$$

has a unique global in time solution in some function space.

The implementation of this result in the case where ζ is a random Gaussian noise takes the following form, for a precise version of which we refer to theorem 5.7.

Theorem 1.2. Let ξ stand for a time-independent weighted noise in space, and set $\xi^\varepsilon := P_\varepsilon \xi$, and $X^\varepsilon(t) = \int_0^t P_{t-s}(\xi^\varepsilon) ds$.

- (a) The pair $(\xi^\varepsilon, X^\varepsilon)$ converges in probability in some space to some extended noise (ζ, X) , with $\zeta = \xi$, and $\Pi(X, \zeta)$ well-defined in the above sense.
- (b) Furthermore, if u^ε stands for the solution of the renormalized equation

$$(1.2) \quad \partial_t u^\varepsilon + Lu^\varepsilon = F(u^\varepsilon) \xi^\varepsilon - c^\varepsilon F'(u^\varepsilon) F(u^\varepsilon), \quad u^\varepsilon(0) = u_0$$

where $c^\varepsilon(\cdot) := \mathbb{E}[\Pi(L^{-1}\xi^\varepsilon, \xi^\varepsilon)(\cdot)]$ is a deterministic real-valued function on M , then u^ε converges in probability to the solution u of equation (1.1) associated with (ζ, X) , in some space whose definition depends on whether or not F is linear.

We have organized our work as follows. Section 2 presents the functional setting in which our theory is set. The main geometrical assumptions on the geometric background are given in section 2.1, where examples are given; these assumptions involve the properties of the heat kernel of the semigroup $(e^{-tL})_{t \geq 0}$ generated by L . A family of operators is introduced in section 2.2, which will play in the sequel the role played by Fourier projectors in the classical Littlewood-Paley theory. We introduce in section 2.3 a scale of Hölder spaces, defined uniquely in terms of the semigroup $(e^{-tL})_{t \geq 0}$. A paraproduct is introduced in section 3.1 and is shown in section 3.2 to enjoy the same continuity properties as its Euclidean analogue. A crucial commutator estimate between paraproduct and resonant terms is proved in section 3.3, together with some parilinearization and composition estimates in section 3.4. Following [29], we then introduce in section 4.1 what plays the role in our setting of paracontrolled distributions, and prove some fundamental Schauder estimates in section 3.5. Sections 2 to 4 give us all the material needed to investigate singular PDEs from the point of view of paracontrolled distributions. Section 5 is dedicated to the proof of theorems 1.1 and 1.2.

We end this work by an Appendix (Appendix B), jointly written with Dorothee Frey, in which we aim to explain how we can weaken an assumption of Lipschitz regularity of the heat kernel (Lip) which we make in the main body of this work, in terms of more geometrical properties, and show that one can prove results in Sobolev spaces similar to those proved in the main body of that work in Hölder spaces.

We collect here a number of **notations** that will be used throughout that work.

- For a ball B of radius r and a real $\lambda > 0$, denote by λB the ball concentric with B and with radius λr . Finally, we will use $u \lesssim v$ to say that there exists a constant C (independent of the important parameters) such that $u \leq Cv$ and $u \simeq v$ to say that $u \lesssim v$ and $v \lesssim u$. We also adopt the non-conventional notation γ_a for the classical gamma function, defined for $a > 0$ by the formula

$$\gamma_a := \int_0^\infty x^a e^{-x} \frac{dx}{x};$$

the capital letter Γ will be used to denote the carré du champ operator of some other operator.

- For $p \in [1, \infty]$ and every $f \in L^p$, the L^p -norm (implicitly with respect to the measure μ) is denoted by $\|f\|_p$. For $p, q \in [1, \infty]$ and an operator T acting from L^p to L^q , we write $\|T\|_{p \rightarrow q}$ for its norm.
- For an integer $k \geq 0$, we write C_b^k for the set of functions continuously differentiable k -times $f : \mathbb{R} \rightarrow \mathbb{R}$, equipped with the norm

$$\|f\|_{C_b^k} := \|f\|_\infty + \sup_{1 \leq i \leq k} \|f^{(i)}\|_\infty.$$

2

Functional calculus adapted to the heat semigroup

As announced in the introduction, this section is dedicated to setting the functional framework where we shall set our study. Section 2.1 sets the geometrical framework needed for what we want to do, in terms of a semigroup. We introduce in section 2.2 some operators that will play the role of 'localizers' in frequency space. These operators are used in section 2.3 to define a scale of Hölder spaces which will be instrumental in the sequel.

2.1. Heat semigroup on a doubling space Let (M, d) be a locally compact separable metrisable space, equipped with a Borel measure μ , finite on compact sets and strictly positive on any non-empty open set. Given a ball $B(x, r)$ of center x and radius r , the notation $V(x, r)$ will stand in the sequel for $\mu(B(x, r))$. To make things concrete, the space (M, d) will mainly be for us smooth Riemannian manifold or a (possibly infinite) metric graph. We shall assume that the metric measure space (M, d, μ) satisfies the following **volume doubling property**

$$(VD) \quad V(x, 2r) \lesssim V(x, r),$$

for all $x \in M$ and positive r , which can be stated equivalently under the form

$$(2.1) \quad V(x, r) \lesssim \left(\frac{r}{s}\right)^\nu V(x, s),$$

for some positive scaling factor ν , for all $x \in M$, and all $0 < s \leq r$; it implies the inequality

$$V(x, r) \lesssim \left(\frac{d(x, y) + r}{s}\right)^\nu V(y, s),$$

for any two points x, y in M and $0 < s \leq r$. (Another easy consequence of the volume doubling property is that balls with a non-empty intersection and comparable radii have comparable measures.)

Let also be given a non-negative self-adjoint operator L on $L^2(M, \mu)$ with dense domain $\mathcal{D}_2(L) \subset L^2(M, \mu)$. Denote by \mathcal{E} its associated quadratic form, defined by the formula

$$\mathcal{E}(f, g) := \int_M f L g \, d\mu,$$

on a domain \mathcal{F} which contains $\mathcal{D}_2(L)$. We shall assume that the Dirichlet form \mathcal{E} is strongly local and regular; we refer the reader to the books [23, 31] of Fukushima & co. and Gyrya–Saloff-Coste for precise definitions and background on Dirichlet forms. These two properties will be obviously satisfied in the examples we shall work with. It follows from these conditions that the operator L generates a strongly continuous semigroup $(e^{-tL})_{t>0}$ of contractions on $L^2(M, \mu)$ which is conservative, in the sense that $e^{-tL}\mathbf{1} = \mathbf{1}$, for all $t \geq 0$; see e.g. Subsection 2.2.7 in the book [31]. We shall also assume that the semigroup $(e^{-tL})_{t>0}$ has a kernel, given for all positive times t by a non-negative measurable real-valued function p_t on $M \times M$, such that

$$(e^{-tL}f)(x) = \int_M p_t(x, y) f(y) \, d\mu(y),$$

for μ -almost all x in M , and every $f \in \mathcal{D}_2(L)$. The kernel p_t is called the **heat kernel** associated with L . We assume that it satisfies for all $0 < t \leq 1$ and μ -almost all x, y , the following typical upper estimates

$$p_t(x, y) \lesssim \frac{1}{\sqrt{V(x, \sqrt{t})V(y, \sqrt{t})}}.$$

Under the volume doubling condition (VD), the previous estimate self-improves into a Gaussian upper estimate for the heat kernel and its time derivatives

$$(UE) \quad \left| \partial_t^a p_t(x, y) \right| \lesssim \frac{t^{-a}}{\sqrt{V(x, \sqrt{t})V(y, \sqrt{t})}} \exp\left(-c \frac{d(x, y)^2}{t}\right).$$

that holds for all integers a , all $0 < t \leq 1$, and μ -almost every $x, y \in M$; see for instance the article [25, Theorem 1.1] for the Riemannian case, and the work [17, Section 4.2] for a metric measure space setting. We also assume that the heat kernel satisfies the following Lipschitz condition³

$$(Lip) \quad \left| p_t(x, y) - p_t(z, y) \right| \lesssim \left(\frac{d(x, z)}{\sqrt{t}} \right) \frac{1}{\sqrt{V(x, \sqrt{t})V(y, \sqrt{t})}} \exp\left(-c \frac{d(x, y)^2}{t}\right).$$

It follows classically from the Gaussian estimates (UE) and the doubling property that the heat semigroup $(e^{-tL})_{t>0}$ is uniformly bounded on $L^p(M, \mu)$ for every $p \in [1, \infty]$ and strongly continuous for $p \in [1, \infty)$. Last, note that $(e^{-tL})_{0 < t \leq 1}$ is, under these conditions, bounded

³ In the regularity structures theory or Euclidean paracontrolled theory, regularity at any order may be considered because of the implicit use of the very nice differential geometry of the Euclidean space, or the torus. In our current and far more general framework, since we only have assumptions on the heat kernel and its gradient, it is natural to expect to be able to quantify regularity of some objects, up until order 1, and not more. That is why in the different statements proved in the next sections some extra mild conditions on the regularity exponents will appear, as compared with their Euclidean analogue. Since we aim to work with the optimal / minimal setting, these new limitations cannot be removed and we are restricted to study regularity at order at most 1, including negative orders; this is not restrictive as far as applications are concerned.

analytic on $L^p(M, \mu)$, for every $1 < p < +\infty$, which means in particular that the time-derivatives $((tL)^n e^{-tL})_{0 < t \leq 1}$ are bounded on $L^p(M, \mu)$ uniformly in $0 < t \leq 1$, for every integer $n \geq 0$; see [52].

Here are four representative classes of **examples** of doubling metric measure spaces and Dirichlet forms satisfying the above conditions. We emphasize (as it can be seen by the list of examples) that we have much flexibility in terms of the operator L as well as in terms of the underlying space (M, d, μ) .

- (a) **Markov chains.** Let X be a countable set equipped with a Markov chain, specified by a symmetric Markov kernel $k : X \times X \rightarrow \mathbb{R}_+$, and let m be a non-negative function on X , used to define a measure \mathfrak{m} on X , with density m with respect to the counting measure μ . Denote by $\langle \cdot, \cdot \rangle_{\mathfrak{m}}$ the scalar product on $\ell^2(\mathfrak{m})$. Consider also for integers $n \geq 1$ the iterated kernel k^n , defined recursively by $k^n(x, y) := \int k^{n-1}(x, z)k(z, y) \mu(dz)$. Denoting by K the symmetric Markov operator with kernel k , the formula

$$\begin{aligned} \mathcal{E}(f, g) &= \frac{1}{2} \sum_{x, y \in X} k_{xy} (f_x - f_y)(g_x - g_y) \\ &= \sum_{x \in X} f_x \frac{1}{m_x} \left(g_x - \sum_{y \in X} k_{xy} g_y \right) m_x \\ &= \langle f, Lg \rangle_{\mathfrak{m}} \end{aligned}$$

associated with the non-negative self-adjoint operator

$$(Lg)(x) = \frac{1}{m_x} \left(g_x - \sum_{y \in X} k_{xy} g_y \right) = g(x) - (Kg)(x),$$

defines a (strongly local) regular Dirichlet form and allows us to generate the continuous heat semigroup $(e^{-tL})_{t \geq 0}$. (The above sum in x is implicitly restricted to those x for which $m_x > 0$, so there is no loss of generality in assuming that $m > 0$.) The map k induces a distance d on X by setting

$$d(x, y) := \min \{ n \geq 1 ; \exists z_0, \dots, z_n, \text{ with } z_0 = x, z_n = y \text{ and } b(z_i, z_{i+1}) > 0, \text{ for } i = 0..n-1 \},$$

for $y \neq x$. Following Grigor'yan's result [26], one can give growth conditions on the \mathfrak{m} -volume of d -balls that ensure the conservative character of the semigroup generated by L in $\ell^2(\mathfrak{m})$. Then it is classical that getting Gaussian upper estimates for the semigroup $(e^{-tL})_{t \geq 0}$ is very closely related to getting discrete-time versions of Gaussian estimates for the iterated Markov chains $(K^n)_{n \geq 1}$, and similarly for the Lipschitz regularity of their kernels. Usually, given such a discrete framework, one prefers to work with the discrete-time Markov chains rather than the continuous heat semigroup. To obtain upper Gaussian estimates and a Lipschitz regularity for the iterated Markov chains on a graphs is the topic of a huge literature to which we refer the reader; see for instance by Hebisch and Saloff-Coste [38] for discrete groups and by Ischiwata [41] for an extension to nilpotent covering graphs and more recently [42] for a perturbation of these previous results. For example, the regular graphs \mathbb{Z}^d and $(\mathbb{Z}/N\mathbb{Z})^d$ have heat semigroups satisfying the Gaussian estimates (UE) and the Lipschitz property (Lip). Needless to say, for a (large) finite graph (X, E) , with edge set E , and $b_{xy} = 1$ if $(x, y) \in E$, and $m_x = \sum_{y \in X} b_{xy}$, the previous results hold with the graph distance in the role of d .

- (b) **Second order differential operators on Riemannian manifolds.** Let (M, d, μ) be a doubling possibly non-compact complete Riemannian manifold with Ricci curvature

bounded from below. Then the heat semigroup $(e^{-t\Delta})_{t \geq 0}$ generated by the Riemannian Laplace operator satisfies both the upper Gaussian estimates (UE) and the Lipschitz regularity (Lip) for small time $0 < t \leq 1$, and for every time $t > 0$ if the Ricci curvature is nonnegative; see [56] and [45] for references. Particular examples are every smooth compact Riemannian manifolds, or unbounded Riemannian manifolds with pinched negative Ricci curvature, such as hyperbolic spaces.

Even on the Euclidean space \mathbb{R}^d , we may consider a second order divergence form operator $L = -\operatorname{div}(A\nabla)$ given by a map A taking values in real symmetric matrices and satisfying usual ellipticity (or accretivity) condition. Then if A is Hölder continuous, it is known that $-L$ generates a self-adjoint semigroup with (UE) and (Lip); see [4]. Similarly, consider an open (and bounded) subset $\Omega \subset \mathbb{R}^d$ and consider for L , the self-adjoint Laplace operator associated with (Dirichlet or Neumann) boundary conditions. There is an extensive literature to describe assumptions on Ω such that (UE) and (Lip) are satisfied. The present scope may well be beyond the present scope of regularity structures, for which the Green function of the operator need to satisfy some regularity assumptions that were not proven to hold true under a sole Hölder continuity assumption for A , and whose formulation on a manifold is a real problem outside the real of Lie groups or homogeneous spaces. The theory developed below works in that relatively minimal setting.

The estimates (UE) and (Lip) also hold when working on a convex or C^2 -regular bounded subset of the Euclidean space, with L given by Laplace operator with Neumann boundary conditions; see [58].

- (c) **Sub-elliptic left invariant diffusions on groups.** Let G be a unimodular connected Lie group, endowed with its left-right Haar measure μ . Consider a family $\mathbf{X} := \{X_1, \dots, X_\ell\}$ of left-invariant vector fields on G satisfying Hörmander condition. They define a class of admissible paths γ_\bullet , characterized by the existence, for each of them, of measurable functions a_1, \dots, a_k such that one has

$$\gamma'(t) = \sum_{i=1}^k a_i(t) X_i(\ell(t)).$$

The length of such a curve is defined as

$$|\gamma| := \frac{1}{2} \int_0^1 \left(\sum_{i=1}^{\ell} |a_i(t)|^2 \right)^{\frac{1}{2}} dt,$$

and the (Carnot-Caratheodory) distance $d(x, y)$ between any two points x, y of G is defined as the infimum of the lengths of all admissible curves joining x to y . We then consider the sublaplacian Δ defined by

$$\Delta := - \sum_{i=1}^k X_i^2.$$

Then the operator Δ generates a heat semigroup satisfying both the upper Gaussian estimates (UE) and the Lipschitz regularity (Lip) for small time $t \in (0, 1)$; see for instance Chapter 8 in the book [57]. If the group is nilpotent then it is also globally doubling [30] and so the heat semigroup satisfies the Gaussian upper bound (UE) and enjoys the Lipschitz property (Lip) for every $t > 0$; see [55, 50]. Particular examples of such groups, are stratified Lie groups and so Heisenberg groups. For such Heisenberg-type Lie groups, a kind of Fourier transform may be defined involving their irreducible

unitary representations, which can be used to define an analog of the Euclidean para-products / paradifferential calculus, such as done is [24]. We shall see, as a by-product of the present work, that the structure of heat semigroup is sufficient to construct similar tools with greater scope.

- (d) The general case given by a subelliptic operator is more difficult. Let (M, d, μ) be a complete and smooth connected manifold endowed with a self-adjoint smooth locally subelliptic diffusion operator L satisfying $L1 = 0$. Then Baudoin and Garofalo introduced in [7] a property, called “a generalized curvature-dimension inequality”, which has to be thought as a lower bound on a sub-Riemannian generalization of the Ricci tensor. Under such a condition, the heat kernel generated by L satisfies (UE) as well as (Lip) (see [49]). We refer the reader to [7] for some examples of such sub-elliptic settings and the fact that the heat kernel also satisfies in that case some Gaussian lower bound.

Throughout that work, a point $o \in M$ will be fixed, that we shall use to define a class of test functions, together with its ‘dual’ class of distributions.

Definition 2.1. We define a Fréchet space of test functions setting

$$\mathcal{S}_o := \left\{ f \in \bigcap_{n \geq 0} \mathcal{D}_2(L^n); \forall a_1, a_2 \in \mathbb{N}, \left\| (1 + d(o, \cdot))^{a_1} L^{a_2} f \right\|_2 < \infty \right\},$$

with

$$\|f\| := \sup_{a_1, a_2 \in \mathbb{N}} 1 \wedge \left\| (1 + d(o, \cdot))^{a_1} L^{a_2} f \right\|_2.$$

A **distribution** is a continuous linear functional on \mathcal{S}_o ; we write \mathcal{S}'_o for the set of all distributions.

(We point out that the arbitrary choice of point $o \in M$ is only relevant in the case of a unbounded ambient space M ; even in that case, the space \mathcal{S}_o does not depend on o , for o ranging inside a bounded subset of M .) Every bounded function defines for instance an element of \mathcal{S}'_o . Examples of test functions are provided by the $p_t(x, \cdot)$, for every fixed $x \in M$ and $0 < t \leq 1$. Indeed for integers a_1, a_2 , the upper bound (UE) gives

$$\begin{aligned} \left| (1 + d(o, y))^{a_1} (L^{a_2} p_t(x, \cdot))(y) \right| &\lesssim \frac{t^{-a_2}}{V(x, \sqrt{t})} (1 + d(o, y))^{a_1} e^{-c \frac{d(x, y)^2}{t}} \\ &\lesssim \frac{t^{-a_2}}{V(x, \sqrt{t})} (1 + d(o, x))^{a_1} e^{-c' \frac{d(x, y)^2}{t}} \end{aligned}$$

for some positive constants c and c' . Note that the heat semigroup acts not only on functions, but also on distributions, by setting

$$\langle (e^{tL} \phi), f \rangle := \langle \phi, (e^{tL} f) \rangle$$

for $\phi \in \mathcal{S}'_o$ and $f \in \mathcal{S}_o$. (We refer the reader to [15] and [46] for more details on the extension of the semigroup to distributions.)

For a linear operator T acting from \mathcal{S}_o to \mathcal{S}'_o , it will be useful below, to denote by K_T its Schwarz kernel, characterized by the identity

$$\langle T(f), g \rangle = \int K_T(x, y) f(y) g(x) \mu(dy) \mu(dx).$$

giving an integral representation for every $f, g \in \mathcal{S}_o$.

2.2. Time derivatives and Carré du champ of the semigroup

Let us introduce here a family of operators that will play the role in our setting of the Fourier multipliers used in the classical Littlewood-

Paley theory, that localize a function in frequency space. These will be the building blocks used to define a convenient paraproduct for our needs, as done below in section 3.1.

Definition 2.2. Given a fixed positive integer a , set

$$(2.2) \quad Q_t^{(a)} := (tL)^a e^{-tL}$$

and

$$(2.3) \quad P_t^{(a)} := \phi_a(tL), \quad \text{where} \quad \phi_a(x) := \frac{1}{\gamma_a} \int_x^\infty s^a e^{-s} \frac{ds}{s}, \quad x \geq 0,$$

for every $t > 0$.

So we have for instance $P_t^{(1)} = e^{-tL}$, and $Q_t^{(1)} = tL e^{-tL}$. The two families of operators $(P_t^{(a)})_{t>0}$ and $(Q_t^{(a)})_{t>0}$ are defined so as to have

$$(2.4) \quad t \partial_t P_t^{(a)} = tL \phi_a'(tL) = -\gamma_a^{-1} Q_t^{(a)},$$

so $Q_t^{(a)} = (-1)^a t^a \partial_t^a e^{-tL}$, and $P_t^{(a)} = p_a(tL) e^{-tL}$, for some polynomial p_a of degree $a - 1$, with $p_a(0) = 1$. The analyticity of the semigroup provides a direct control of the operators $P_t^{(a)}$ and $Q_t^{(a)}$.

Proposition 2.3. For any integer $a \geq 0$, operators $P_t^{(a)}$ and $Q_t^{(a)}$ have a kernel satisfying Gaussian estimates (UE), and the Lipschitz regularity (Lip); as a consequence, they are bounded in every L^p spaces for $p \in [1, \infty]$, uniformly with respect to $t \in (0, 1]$.

Following the above interpretation of the operators $Q^{(a)}$ and $P^{(a)}$, the following Calderón reproducing formula provides a decomposition of a function f in $L^p(M, \mu)$ into a low frequency part and a high frequency part very similar to the Littlewood-Paley decomposition of a distribution in terms of frequencies; see e.g. [6].

Proposition 2.4 (Calderón reproducing formula). Given $p \in (1, +\infty)$ and $f \in L^p(M, \mu)$, we have

$$\lim_{t \rightarrow 0^+} P_t^{(a)} f = f \quad \text{in } L^p(M, \mu)$$

for every positive integer a , and so

$$(2.5) \quad f = \gamma_a^{-1} \int_0^1 Q_t^{(a)} f \frac{dt}{t} + P_1^{(a)}(f).$$

PROOF – One knows from theorem 3.1 in [20], that the operator L has a bounded H^∞ functional calculus in $L^p(M, \mu)$ under the volume doubling condition on (M, d, μ) , and the assumption that the heat kernel satisfies the upper estimate (UE). Since this implies in particular sectoriality of L in $L^p(M, \mu)$, Theorem 3.8 in [18] yields the decomposition of $L^p(M, \mu)$ into nullspace and range of L . Using this decomposition, the Convergence Lemma implies for every $f \in L^p(M, \mu)$

$$\begin{aligned} f &= \lim_{t \rightarrow 0} P_t^{(a)} f = - \int_0^1 \partial_t P_t^{(a)} f \frac{dt}{t} + P_1^{(a)}(f) \\ &= \gamma_a^{-1} \int_0^1 Q_t^{(a)} f \frac{dt}{t} + P_1^{(a)}(f), \end{aligned}$$

where the limit is taken in $L^p(M, \mu)$ and where we have used identity (2.4); see e.g. [3, Theorem D] or [43, Lemma 9.13].

▷

We shall also make an extensive use in the sequel of the square-root of L , given by its **carré du champ** operator Γ , defined for all $(f, g) \in \mathcal{D}_2(L) \times \mathcal{D}_2(L)$ as a bilinear operator satisfying the identity

$$\mathcal{E}(f, g) := \int_M f L(g) d\mu = \int_M g L(f) d\mu = \int_M \Gamma(f, g) d\mu.$$

It is also given by the explicit formula

$$\Gamma(f, g) = -\frac{1}{2} \left(L(fg) - fL(g) - gL(f) \right);$$

we shall write $\mathcal{D}_2(\Gamma) \subset L^2$ for its domain, which contains $\mathcal{D}_2(L)$. As a shorthand, we write $\Gamma(f)$ for $\Gamma(f, f)^{\frac{1}{2}}$ in the sequel, which can be thought as the length of the intrinsic gradient of f . It follows from the conservative property of L and its non-negative character, that the bilinear map Γ is positive and satisfies the identity

$$\|\Gamma(f)\|_{L^2}^2 = \int_M \Gamma(f, f) d\mu = \int_M f L(f) d\mu = \mathcal{E}(f, f).$$

According to the Beurling-Deny-Le Jan formula, the carré du champ satisfies a *Leibniz rule*

$$(2.6) \quad \Gamma(fg, h) = f \Gamma(g, h) + g \Gamma(f, h),$$

for all $f, g, h \in \mathcal{D}_2(\Gamma)$, and a *chain rule*

$$(2.7) \quad L(F(f)) = F'(f) L(f) + F''(f) \Gamma(f, f).$$

for every function $F \in C_b^2(\mathbb{R})$ and every $f \in \mathcal{D}_2(L)$; the function $F(f)$ is automatically in $\mathcal{D}_2(L)$ – see e.g. [23, Section 3.2] and [53, Appendix] for these points.

The following pointwise and L^p -estimate for the intrinsic gradient of the semigroup will be used several times in a crucial way; its proof is given in Appendix A. It says that the carré du champ of the semigroup satisfies also some Gaussian pointwise estimates, as given by the following claim.

Proposition 2.5. The following inequality holds

$$(2.8) \quad \left| (\sqrt{t}\Gamma)(e^{-tL}f)(x_0) \right| \lesssim \int_M \frac{1}{\sqrt{V(x_0, \sqrt{t})V(y, \sqrt{t})}} \exp\left(-c \frac{d(x_0, y)^2}{t}\right) |f(y)| d\mu(y),$$

for every $t > 0$, every function $f \in L^2$, and almost every $x_0 \in M$. Consequently, we have

$$\sup_{t>0} \left\| (\sqrt{t}\Gamma)(e^{-tL} \cdot) \right\|_{p \rightarrow p} < \infty,$$

for every $p \in [1, \infty]$. We may replace the semigroup e^{-tL} in the above equations by any of the operators $P_t^{(a)}$, for any $a \geq 0$.

2.3. Hölder and Besov spaces through the heat semigroup

Let us recall as a start that given a parameter $\sigma \in (0, 1]$, a bounded function $f \in L^\infty$ is said to belong to the Hölder space Λ^σ if

$$\|f\|_{\Lambda^\sigma} := \|f\|_\infty + \sup_{0 < d(x,y) \leq 1} \frac{|f(x) - f(y)|}{d(x,y)^\sigma} < \infty.$$

Recall on the other hand the definition of the inhomogeneous Besov spaces associated to a semigroup; they were precisely studied in several works, such as [15] or [27], to name but a few. We shall make an extensive use of these spaces.

Definition 2.6. Fix a positive integer a , an exponent $p, q \in (1, \infty)$, and $\sigma \in \mathbb{R}$. A distribution $f \in \mathcal{S}'_o$, is said to belong to the **Besov space** $B_{p,q}^\sigma$ if

$$\|f\|_{B_{p,q}^\sigma} := \|e^{-L}f\|_p + \left(\int_0^1 t^{-q\frac{\sigma}{2}} \left\| Q_t^{(a)} f \right\|_p^q \frac{dt}{t} \right)^{1/q} < \infty.$$

This definition of the space does not depend on the integer $a \geq 1$, provided a is big enough.

We refer the reader to [15] for details about such spaces and a proof of the fact that they do not depend on the parameter a used to define them, provided a is sufficiently large with respect to s . The limiting case $p = q = \infty$ leads to the following definition.

Definition 2.7. Let a positive integer a be given. For $\sigma \in (-\infty, 2)$, a distribution $f \in \mathcal{S}'_o$ is said to belong to the space \mathcal{C}^σ if

$$\|f\|_{\mathcal{C}^\sigma} := \left\| e^{-L}f \right\|_\infty + \sup_{0 < t \leq 1} \left\| Q_t^{(a)} f \right\|_\infty t^{-\frac{\sigma}{2}} < \infty.$$

This definition of the space does not depend on the integer $a \geq 1$.

We give in Appendix A a simple and self-contained proof that the space \mathcal{C}^σ does not depend on a , and that any two norms $\|\cdot\|_{\mathcal{C}^\sigma}$, defined with two different values of a , are equivalent. The following proposition justifies that we call the spaces \mathcal{C}^σ Hölder space, for all $\sigma < 2$, possibly non-positive.

Proposition 2.8. For $\sigma \in (0, 1)$, the spaces Λ^σ and \mathcal{C}^σ are the same and the two corresponding norms are equivalent.

We give here a complete proof of this proposition as it provides an elementary illustration of how the properties of the operators $Q_t^{(a)}$ are used to make actual computations. This kind of reasoning and computations will be used repeatedly in the sequel, when working with our paraproduct. Recall that the operators $Q_t^{(a)}$ have kernels $K_{Q_t^{(a)}}$ satisfying Gaussian pointwise estimates, by proposition 2.3.

PROOF – We divide the proof in two steps, by showing successively that Λ^σ is continuously injected in \mathcal{C}^σ , and that, conversely, \mathcal{C}^σ is continuously injected in Λ^σ .

Step 1 – $\Lambda^\sigma \hookrightarrow \mathcal{C}^\sigma$. Note first that since the Hölder space Λ^σ is made up of bounded functions, it is included in \mathcal{S}'_o . Fix an integer $a \geq 1$; then for every $t \in (0, 1)$, we have

$$\left(Q_t^{(a)} f \right)(x) = \left(Q_t^{(a)} (f - f(x)) \right)(x) = \int K_{Q_t^{(a)}}(x, z) (f(z) - f(x)) \mu(dz).$$

For the points $z \in M$, with $d(x, z) \leq \sqrt{t} < 1$, we have

$$|f(z) - f(x)| \leq d(x, z)^\sigma \|f\|_{\Lambda^\sigma} \leq t^{\frac{\sigma}{2}} \|f\|_{\Lambda^\sigma}$$

so that

$$\begin{aligned} \left| \int_{d(x,z) \leq \sqrt{t}} K_{Q_t^{(a)}}(x,z)(f(z) - f(x)) \mu(dz) \right| &\leq t^{\frac{\sigma}{2}} \|f\|_{\Lambda^\sigma} \int |K_{Q_t^{(a)}}(x,z)| \mu(dz) \\ &\lesssim t^{\frac{\sigma}{2}} \|f\|_{\Lambda^\sigma}, \end{aligned}$$

since $Q_t^{(a)}$ has a kernel satisfying Gaussian pointwise bounds. The same bounds show that

$$\begin{aligned} \left| \int_{\sqrt{t} \leq d(x,z) \leq 1} K_{Q_t^{(a)}}(x,z)(f(z) - f(x)) \mu(dz) \right| &\leq \|f\|_{\Lambda^\sigma} \left(\int_{\sqrt{t} \leq d(x,z) \leq 1} |K_{Q_t^{(a)}}(x,z)| d(x,z)^\sigma \mu(dz) \right) \\ &\lesssim t^{\frac{\sigma}{2}} \|f\|_{\Lambda^\sigma} \int_{\sqrt{t} \leq d(x,z) \leq 1} \frac{1}{V(x, \sqrt{t})} e^{-c \frac{d(x,z)^2}{t}} \left(\frac{d(x,z)}{\sqrt{t}} \right)^\sigma \mu(dz) \\ &\lesssim t^{\frac{\sigma}{2}} \|f\|_{\Lambda^\sigma}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \left| \int_{1 \leq d(x,z)} K_{Q_t^{(a)}}(x,z)(f(z) - f(x)) \mu(dz) \right| &\leq \|f\|_\infty \left(\int_{1 \leq d(x,z)} |K_{Q_t^{(a)}}(x,z)| \mu(dz) \right) \\ &\lesssim e^{-c/t} \|f\|_{\Lambda^\sigma} \\ &\lesssim t^{\frac{\sigma}{2}} \|f\|_{\Lambda^\sigma}, \end{aligned}$$

so it comes that the inequality

$$\left| (Q_t^{(a)} f)(x) \right| \lesssim t^{\frac{\sigma}{2}} \|f\|_{\Lambda^\sigma}$$

holds uniformly in $t \in (0, 1)$, and for every $x \in M$, which proves that $\|f\|_{\mathcal{C}^\sigma} \lesssim \|f\|_{\Lambda^\sigma}$.

Step 2 – $\mathcal{C}^\sigma \hookrightarrow \Lambda^\sigma$. Let $f \in \mathcal{C}^\sigma$ be given. Using the decomposition of the identity provided by Calderón reproducing formula

$$f = e^{-L} f - \int_0^1 Q_t^{(1)} f \frac{dt}{t},$$

we first deduce that f is bounded, with

$$\|f\|_\infty \lesssim \|f\|_{\mathcal{C}^\sigma} \left(1 + \int_0^1 t^{\frac{\sigma}{2}} \frac{dt}{t} \right) \lesssim \|f\|_{\mathcal{C}^\sigma}.$$

Moreover, for any two points x, y , with $0 < d(x, y) \leq 1$, we have

$$\begin{aligned} f(x) - f(y) &= \left\{ (e^{-L} f)(x) - (e^{-L} f)(y) \right\} - \left\{ \int_0^1 \left((Q_t^{(1)} f)(x) - (Q_t^{(1)} f)(y) \right) \frac{dt}{t} \right\} \\ &= \left\{ e^{-L} f(x) - e^{-L} f(y) \right\} - \left\{ (Q_1^{(1)} f)(x) - (Q_1^{(1)} f)(y) \right\} \\ &\quad - \int_0^1 \left\{ (Q_t^{(2)} f)(x) - (Q_t^{(2)} f)(y) \right\} \frac{dt}{t}. \end{aligned}$$

One can use the Lipschitz regularity (Lip) of the heat kernel to bound the first term in the above sum, giving

$$\begin{aligned} \left| e^{-L} f(x) - e^{-L} f(y) \right| &\lesssim \int |p_1(x, z) - p_1(y, z)| |f(z)| \mu(dz) \\ &\lesssim d(x, y) \|f\|_\infty. \end{aligned}$$

As similar bounds hold for Le^{-L} , by analyticity of the heat kernel, the second term admits a similar upper bound. Let now focus on the third term, using a similar reasoning and noting that $Q_t^{(2)} = 16 Q_{t/2}^{(2)} Q_{t/2}^{(2)}$. So, for $d(x, y) \leq \sqrt{t}$, we can write

$$\begin{aligned} \left| (Q_t^{(2)} f)(x) - (Q_t^{(2)} f)(y) \right| &\lesssim \int \left| K_{Q_{t/2}^{(2)}}(x, z) - K_{Q_{t/2}^{(2)}}(y, z) \right| \left| K_{Q_{t/2}^{(2)}} f(z) \right| \mu(dz) \\ &\lesssim \frac{d(x, y)}{\sqrt{t}} \left\| K_{Q_{t/2}^{(2)}} f \right\|_{\infty} \\ &\lesssim \frac{d(x, y)}{\sqrt{t}} t^{\frac{\sigma}{2}} \|f\|_{\mathcal{C}^{\sigma}}. \end{aligned}$$

If $\sqrt{t} \leq d(x, y)$, then we directly have

$$\left| (Q_t^{(2)} f)(x) - (Q_t^{(2)} f)(y) \right| \lesssim \left\| Q_t^{(2)} f \right\|_{\infty} \lesssim t^{\frac{\sigma}{2}} \|f\|_{\mathcal{C}^{\sigma}}.$$

Hence,

$$\begin{aligned} \left| \int_0^1 \left\{ (Q_t^{(2)} f)(x) - (Q_t^{(2)} f)(y) \right\} \frac{dt}{t} \right| &\lesssim \left(\int_0^{d(x, y)^2} t^{\frac{\sigma}{2}} \frac{dt}{t} + \int_{d(x, y)^2}^1 \left(\frac{d(x, y)}{\sqrt{t}} \right) t^{\frac{\sigma}{2}} \frac{dt}{t} \right) \|f\|_{\mathcal{C}^{\sigma}} \\ &\lesssim d(x, y)^{\sigma} \|f\|_{\mathcal{C}^{\sigma}}, \end{aligned}$$

since $\sigma \in (0, 1)$. Consequently, we have obtained

$$|f(x) - f(y)| \lesssim d(x, y)^{\sigma} \|f\|_{\mathcal{C}^{\sigma}}$$

uniformly for every $x \neq y$ with $d(x, y) \leq 1$, so indeed $\|f\|_{\Lambda^{\sigma}} \lesssim \|f\|_{\mathcal{C}^{\sigma}}$. ▷

Our main example of \mathcal{C}^{σ} distribution, with negative Hölder exponent σ , will be given by typical realizations of a (possibly weighted) noise over (M, μ) – see Proposition 5.6. To prove that regularity property, it will be convenient to assume that the metric measure space (M, d, μ) has the following property, called **Ahlfors regularity**. There exists a positive constant c_1 such that

$$V(x, 1) \geq c_1,$$

for all $x \in M$, which, by the doubling property, implies that we have

$$(2.9) \quad V(x, r) \geq c_1 r^{\nu},$$

for some positive exponent ν , all $x \in M$ and all $0 < r \leq 1$. (The constant ν is d on a d -dimensional manifold.) This is a relatively weak assumption that is essentially satisfied in a Riemannian setting for closed manifolds without boundary and injectivity radius bounded below by a positive constant. Under that additional assumption, we have the following Besov embedding, proved in Appendix A.

Lemma 2.9 (Besov embedding). *Given $-\infty < \sigma < 2$, and $1 < p < \infty$, we have the following continuous embeddings.*

$$B_{p,p}^{\sigma} \hookrightarrow B_{p,\infty}^{\sigma} \hookrightarrow B_{\infty,\infty}^{\sigma - \frac{\nu}{p}} = \mathcal{C}^{\sigma - \frac{\nu}{p}}$$

Besov embedding can be used in a very efficient way to investigate the regularity properties of random Gaussian fields, as will be illustrated in section 5.3.

Remark 2.10. *Let us point out here that our Hölder spaces C^σ , with $\sigma < 0$, coincide in the Euclidean setting with those used by Hairer [32]. Indeed, on the Euclidean space it is known that to define Besov spaces or Hölder spaces through Littlewood-Paley functionals, we may chose any good Fourier multipliers satisfying suitable conditions; the latter are satisfied by the derivatives $(Q_t^{(a)})_t$ of the heat semigroup. So our spaces correspond to the standard inhomogeneous spaces defined by any Littlewood-Paley functionals. From wavelet or frame characterization (see for instance [47]), we then conclude that our Hölder space coincides with those used in [32] or [36].*

Before turning to the definition of our paraproduct, we close this section with two continuity properties involving the Hölder spaces C^σ , which we shall use in the sequel.

Proposition 2.11. For any $\sigma \in (-\infty, 2)$, and every integer $a \geq 0$, we have

$$\left\| P_1^{(a)} f \right\|_\infty \lesssim \|f\|_{C^\sigma}.$$

PROOF – We have by construction $P_1^{(a)} = (1 + \alpha_1 L + \dots + \alpha_{a-1} L^{a-1}) e^{-L}$, for some coefficients $\alpha_1, \dots, \alpha_{a-1}$. As we have by definition $\|e^{-L} f\|_\infty \lesssim \|f\|_{C^\sigma}$, and $L^\ell e^{-L} = Q_1^{(\ell)}$, for $\ell = 1 \dots (a-1)$, we see that $\|L^\ell e^{-L} f\|_\infty \lesssim \|f\|_{C^\sigma}$, since we have seen above that we can choose the parameter a in the definition of the Hölder space. \triangleright

Proposition 2.12. For $\sigma \in (-\infty, 1)$, we have

$$\sup_{t \in (0,1]} t^{-\frac{\sigma}{2}} \left\| (\sqrt{t} \Gamma) (e^{-tL} f) \right\|_\infty \lesssim \|f\|_{C^\sigma}.$$

The same conclusion holds with any of the operators $P_t^{(a)}$ in the role of e^{-tL} .

PROOF – Given $t \in (0, 1]$, use Calderón reproducing formula to write

$$\left| (\sqrt{t} \Gamma) (e^{-tL} f) \right| \lesssim \int_0^1 \left| (\sqrt{t} \Gamma) (e^{-tL} Q_s^{(1)} f) \right| \frac{ds}{s} + \left| (\sqrt{t} \Gamma) (e^{-(1+t)L} f) \right|.$$

We divide the integration interval in the above-right hand side into $(0, t)$ and $[t, 1]$ to bound that term. For $s < t$, we have $e^{-tL} Q_s^{(1)} = \frac{s}{t/2+s} e^{-t/2L} Q_{s+t/2}^{(1)}$, so we can use Proposition 2.5 to get

$$\begin{aligned} \left\| (\sqrt{t} \Gamma) (e^{-tL} Q_s^{(1)} f) \right\|_\infty &\lesssim \frac{s}{t} \left\| (\sqrt{t} \Gamma) (e^{-t/2L}) \right\|_{\infty \rightarrow \infty} \left\| Q_{s+t/2}^{(1)} f \right\|_\infty \\ &\lesssim \frac{s}{t} t^{\frac{\sigma}{2}} \|f\|_{C^\sigma}. \end{aligned}$$

Similarly for $t \leq s$, then $e^{-tL} Q_s^{(1)} = e^{-sL/2} \frac{s}{t+s/2} Q_{s/2+t}^{(1)}$, and we have

$$\begin{aligned} \left\| (\sqrt{t} \Gamma) (e^{-tL} Q_s^{(1)} f) \right\|_\infty &\lesssim \left(\frac{t}{s} \right)^{\frac{1}{2}} \left\| (\sqrt{s} \Gamma) (e^{-s/2L}) \right\|_{\infty \rightarrow \infty} \left\| Q_{s/2+t}^{(1)} f \right\|_\infty \\ &\lesssim \left(\frac{t}{s} \right)^{\frac{1}{2}} s^{\frac{\sigma}{2}} \|f\|_{C^\sigma}. \end{aligned}$$

Similar computations give the estimate

$$\left\| (\sqrt{t} \Gamma) (e^{-(1+t)L} f) \right\|_\infty \lesssim \sqrt{t} \|f\|_{C^\sigma}.$$

We conclude by integrating with respect to $s \in (0, 1)$, using here the fact that $\sigma < 1$. \triangleright

Paraproduct and commutator estimates in Hölder spaces

3.1. Paraproducts based on the semigroup Bony's paraproduct machinery has its roots in the Littlewood-Paley decomposition of any distribution f as a sum of smooth functions $\Delta_i f$ localized in the frequency space, so a product fg of any two distributions can formally be decomposed as

$$(3.1) \quad fg = \sum_{i,j} \Delta_i f \Delta_j g = \sum_{|i-j| \geq 2} \Delta_i f \Delta_j g + \sum_{|i-j| \leq 1} \Delta_i f \Delta_j g =: (1) + \Pi(f, g)$$

into a sum of products of two functions oscillating on different scales, and an a priori resonant term $\Pi(f, g)$. This decomposition draws its usefulness from some relatively elementary a priori estimates that show that the term (1) above makes sense and is well-controlled under extremely general conditions, while the resonant term $\Pi(f, g)$ can be shown to define a continuous map from $\mathcal{C}^\alpha \times \mathcal{C}^\beta$ to $\mathcal{C}^{\alpha+\beta}$, provided $\alpha + \beta > 0$. These estimates rely crucially on some properties inherited from the very definition of the Littlewood-Paley blocks as Fourier projectors. These properties cannot be grasped so easily in our semigroup setting; however, we shall use the operators $P_t^{(a)}, Q_t^{(a)}$ and $\sqrt{t}\Gamma$ or $(tL)P_t^{(a)}$ as frequency projectors, with $P_t^{(a)}$ projecting on frequencies lower than or equal to $t^{-\frac{1}{2}}$, and $Q_t^{(a)}, \sqrt{t}\Gamma$ or $(tL)P_t^{(a)}$ as localizing at frequencies of order $t^{-\frac{1}{2}}$. This will be our main guide in the definition of our paraproduct given below. This paraproduct will depend on a choice of a positive integer-valued parameter b that can be tuned on demand in any given problem. To clarify notations, we shall repeatedly use below the notation $f \cdot g$ for the (usual) product of two functions.

Rather than starting with Bony's decomposition (3.1), we take as a starting point Calderon's reproducing formula

$$(3.2) \quad \begin{aligned} fg &= \lim_{t \rightarrow 0} P_t^{(b)} \left(P_t^{(b)} f \cdot P_t^{(b)} g \right) = - \int_0^1 t \partial_t \left\{ P_t^{(b)} \left(P_t^{(b)} f \cdot P_t^{(b)} g \right) \right\} \frac{dt}{t} + \Delta_{-1}(f, g) \\ &= \frac{1}{\gamma_b} \int_0^\infty \left\{ P_t^{(b)} \left(Q_t^{(b)} f \cdot P_t^{(b)} g \right) + P_t^{(b)} \left(P_t^{(b)} f \cdot Q_t^{(b)} g \right) + Q_t^{(b)} \left(P_t^{(b)} f \cdot P_t^{(b)} g \right) \right\} \frac{dt}{t} + \Delta_{-1}(f, g), \end{aligned}$$

where

$$\Delta_{-1}(f, g) := P_1^{(b)} \left(P_1^{(b)} f \cdot P_1^{(b)} g \right)$$

stands for the 'low-frequency part' of the product of f and g , and where we implicitly make the necessary assumptions on f and g for the above formula to make sense.

Guided by the above heuristic argument about the role of the operators $P_t^{(a)}, Q_t^{(a)}$, etc. as frequency projectors, we decompose the terme involving the product of $P_t^{(a)} f$ and $P_t^{(a)} g$, by using the definition of the carré du champ operator Γ

$$L(\phi_1 \cdot \phi_2) = L(\phi_1) \phi_2 + L(\phi_2) \phi_1 - 2\Gamma(\phi_1, \phi_2)$$

and write

$$\begin{aligned} &Q_t^{(b)} \left(P_t^{(b)} f \cdot P_t^{(b)} g \right) \\ &= Q_t^{(b-1)} \left((tL)P_t^{(b)} f \cdot P_t^{(b)} g \right) + Q_t^{(b-1)} \left(P_t^{(b)} f \cdot (tL)P_t^{(b)} g \right) - 2Q_t^{(b-1)} \Gamma \left(\sqrt{t}P_t^{(b)} f, \sqrt{t}P_t^{(b)} g \right) \\ &=: B_g(f) + B_f(g) + R(f, g). \end{aligned}$$

If one rewrites identity (3.2) under the form

$$fg =: \int_0^1 \left\{ (1) + (2) + (3) \right\} \frac{dt}{t} + \Delta_{-1}(f, g)$$

with obvious notations, this suggest to decompose it as

$$fg = \int_0^1 \left(\{(1) + B_g(f)\} + \{(2) + B_f(g)\} + R(f, g) \right) \frac{dt}{t} + \Delta_{-1}(f, g)$$

and to identify the integral of the terms into brackets in the above formula as paraproducts, and by defining the resonant term as the integral of $R(f, g)$. This is what was done in [12] where this notion of paraproduct, introduced in [8], was shown to have nice continuity properties in Hölder spaces \mathcal{C}^α , provided only deals with positive exponents α . Given our needs to deal with negative exponents, a refinement of this decomposition seems to be needed to get some continuity properties for negative exponent as well. We thus use the carré du champ formula in each term (1) and (2), and write

$$\begin{aligned} (1) &= (tL)P_t^{(b)} \left(Q_t^{(b-1)} f \cdot P_t^{(b)} g \right) + \left\{ 2P_t^{(b)}(t\Gamma) \left(Q_t^{(b-1)} f, P_t^{(b)} g \right) - P_t^{(b)} \left(Q_t^{(b-1)} f \cdot (tL)P_t^{(b)} g \right) \right\} \\ &=: A_g(f) + S(f, g), \end{aligned}$$

with $S(f, g)$ the sum of the two terms into bracket, and

$$(2) = A_f(g) + S(g, f).$$

Note that the functions $A_f(g), S(f, g), \dots$ all depend implicitly on time. This decomposition leads to the following definition.

Definition 3.1. Given an integer $b \geq 2$ and $f \in \bigcup_{s \in (0,1)} \mathcal{C}^s$ and $g \in J^\infty$, we define their paraproduct by the formula

$$\begin{aligned} \Pi_g^{(b)}(f) &= \frac{1}{\gamma_b} \int_0^1 \left\{ A_g(f) + B_g(f) \right\} \frac{dt}{t} \\ &= \frac{1}{\gamma_b} \int_0^1 \left\{ (tL)P_t^{(b)} \left(Q_t^{(b-1)} f \cdot P_t^{(b)} g \right) + Q_t^{(b-1)} \left((tL)P_t^{(b)} f \cdot P_t^{(b)} g \right) \right\} \frac{dt}{t}. \end{aligned}$$

The well-posed character of this integral is proved in proposition 3.3 below. With this notation, Calderon's formula becomes

$$fg = \Pi_g^{(b)}(f) + \Pi_f^{(b)}(g) + \Pi^{(b)}(f, g) + \Delta_{-1}(f, g)$$

with the 'low-frequency part'

$$\Delta_{-1}(f, g) := P_1^{(b)} \left(P_1^{(b)} f \cdot P_1^{(b)} g \right)$$

and the 'resonant term'

$$\begin{aligned} \Pi^{(b)}(f, g) &= \frac{1}{\gamma_b} \int_0^1 \left\{ S(f, g) + S(g, f) + R(f, g) \right\} \frac{dt}{t} \\ &= \frac{1}{\gamma_b} \int_0^1 \left\{ -P_t^{(b)} \left(Q_t^{(b-1)} f \cdot (tL)P_t^{(b)} g \right) + 2P_t^{(b)} \Gamma \left(\sqrt{t} Q_t^{(b-1)} f, \sqrt{t} P_t^{(b)} g \right) \right\} \frac{dt}{t} \\ &\quad + \frac{1}{\gamma_b} \int_0^1 \left\{ -P_t^{(b)} \left((tL)P_t^{(b)} f \cdot Q_t^{(b-1)} g \right) + 2P_t^{(b)} \Gamma \left(\sqrt{t} P_t^{(b)} f, \sqrt{t} Q_t^{(b-1)} g \right) \right\} \frac{dt}{t} \\ &\quad - \frac{1}{\gamma_b} \int_0^1 2Q_t^{(b-1)} \Gamma \left(\sqrt{t} P_t^{(b)} f, \sqrt{t} P_t^{(b)} g \right) \frac{dt}{t}. \end{aligned}$$

Note that we have $\Pi_1^{(b)}(\cdot) = \text{Id}$, as a consequence of our choice of renormalizing constant.

3.2. Paraproduct estimates We prove in this paragraph the basic continuity estimates satisfied by the maps defined by the low frequency part, the paraproduct and the resonant term. The low-frequency part is easily bounded.

Proposition 3.2. Fix an integer $b \geq 2$. For any $\alpha, \beta \in \mathbb{R}$ and every $\gamma > 0$ we have for every $f \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$

$$(3.3) \quad \|\Delta_{-1}(f, g)\|_{\mathcal{C}^\gamma} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}.$$

PROOF – Consider the collection $(Q_s^{(a)})_{0 < s \leq 1}$ for a large enough integer $a \geq \gamma$. Then

$$Q_s^{(a)} \Delta_{-1}(f, g) = Q_s^{(a)} P_1^{(b)} \left(P_1^{(b)} f \cdot P_1^{(b)} g \right).$$

Since $s \leq 1$, we have $Q_s^{(a)} P_t^{(b)} = \gamma_a^{-1} s^a e^{-sL} L^a P_1^{(b)}$, with the operator $L^a P_1^{(b)}$ bounded on L^∞ . We obtain the conclusion from Proposition 2.11 as we have

$$\begin{aligned} \|Q_s^{(a)} \Delta_{-1}(f, g)\|_\infty &\lesssim s^a \|P_1^{(b)} f\|_\infty \|P_1^{(b)} g\|_\infty \\ &\lesssim s^\gamma \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}. \end{aligned}$$

▷

The continuity properties of the paraproduct are given by the following statement; they are the exact analogue of their classical counterpart, based on Littlewood-Paley decomposition, as can be found for instance in the textbook [6] of Bahouri, Chemin and Danchin.

Proposition 3.3. Fix an integer $b \geq 2$. For any $\alpha \in (-2, 1)$ and $f \in \mathcal{C}^\alpha$, we have

- for every $g \in L^\infty$

$$(3.4) \quad \left\| \Pi_g^{(b)}(f) \right\|_{\mathcal{C}^\alpha} \lesssim \|g\|_\infty \|f\|_{\mathcal{C}^\alpha}$$

- for every $g \in \mathcal{C}^\beta$ with $\beta < 0$ and $\alpha + \beta \in (-2, 1)$

$$(3.5) \quad \left\| \Pi_g^{(b)}(f) \right\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|g\|_{\mathcal{C}^\beta} \|f\|_{\mathcal{C}^\alpha}.$$

Remark 3.4. The range $(-2, 1)$ for the regularity exponent could seem unusual, since in the standard Euclidean theory such continuities hold for every $\alpha \in \mathbb{R}$. However, as explained in footnote 3, the restriction $\alpha < 1$ comes from our optimal / minimal setting where we only assume gradient estimates on the heat kernel. In another hand, the restriction $\alpha > -2$ can be explained as follows. In the Euclidean theory, nice Fourier multipliers can be used to have a 'perfect' frequency decomposition and the study of paraproducts mainly relies on the following rule: the spectrum of the product of two functions is included into the sum of the two spectrums, which comes from the group structure through the Fourier representation of the convolution.

In our setting, the frequency decomposition involving the heat semigroup (as in the Calderón reproducing formula) is not so perfect and above all, the previous rule on the spectrum does not hold, at least not in such a 'perfect' sense. That is why it appears this new limitation $\alpha > -2$, which is inherent to the semigroup approach.

PROOF – Recall that

$$\Pi_g^{(b)}(f) = \frac{1}{\gamma_b} \int_0^1 (tL) P_t^{(b)} \left(Q_t^{(b-1)} f \cdot P_t^{(b)} g \right) + Q_t^{(b-1)} \left((tL) P_t^{(b)} f \cdot P_t^{(b)} g \right) \frac{dt}{t}.$$

Set $c := b - 1 \geq 1$, and given $s \in (0, 1]$, consider $Q_s^{(c)} \Pi_g(f)$. For $s \leq t$, we use that

$$Q_s^{(c)} (tL) P_t^{(b)} = \left(\frac{s}{t}\right)^c (tL)^{c+1} P_t^{(b)} e^{-sL} \quad \text{and} \quad Q_s^{(c)} Q_t^{(b-1)} = \left(\frac{s}{t}\right)^c Q_t^{(b-1+c)} e^{-sL},$$

and for $t \leq s$ that

$$Q_s^{(c)}(tL)P_t^{(b)} = \frac{t}{s}Q_s^{(c+1)}P_t^{(b)} \quad \text{and} \quad Q_s^{(c)}Q_t^{(b-1)} = \frac{t}{s}Q_s^{(c+1)}Q_t^{(b-2)}.$$

Hence, with the uniform L^∞ -boundedness of Q_t, P_t operators, we have

$$\begin{aligned} \left\| Q_s^{(c)}\Pi_g^{(b)}(f) \right\|_\infty &\lesssim \int_0^s \frac{t}{s} \left\| Q_t^{(b-1)}f \right\|_\infty \left\| P_t^{(b)}g \right\|_\infty + \frac{t}{s} \left\| (tL)P_t^{(b)}f \right\|_\infty \left\| P_t^{(b)}g \right\|_\infty \frac{dt}{t} \\ &\quad + \int_s^1 \left(\frac{s}{t} \right)^c \left\| Q_t^{(b-1)}f \right\|_\infty \left\| P_t^{(b)}g \right\|_\infty + \left(\frac{s}{t} \right)^c \left\| (tL)P_t^{(b)}f \right\|_\infty \left\| P_t^{(b)}g \right\|_\infty \frac{dt}{t}. \end{aligned}$$

Since $f \in \mathcal{C}^\alpha$ we have

$$\left\| Q_t^{(b-1)}f \right\|_\infty + \left\| (tL)P_t^{(b)}f \right\|_\infty \lesssim t^{\frac{\alpha}{2}} \|f\|_{\mathcal{C}^\alpha}.$$

Moreover, if $g \in L^\infty$ then

$$\left\| P_t^{(b)}g \right\|_\infty \lesssim \|g\|_\infty$$

and if $g \in \mathcal{C}^\beta$ with $\beta < 0$ then

$$\begin{aligned} \left\| P_t^{(b)}g \right\|_\infty &\leq \int_t^1 \left\| Q_u^{(b)}g \right\|_\infty \frac{du}{u} + \left\| P_1^{(b)}(g) \right\|_\infty \\ &\lesssim \left(\int_t^1 \frac{du}{u^{1-\beta/2}} + 1 \right) \|g\|_{\mathcal{C}^\beta} \lesssim t^{\frac{\beta}{2}} \|g\|_{\mathcal{C}^\beta}. \end{aligned}$$

As a consequence, we deduce the following bounds.

- If $g \in L^\infty$ then

$$\begin{aligned} \left\| Q_s^{(c)}\Pi_g^{(b)}(f) \right\|_\infty &\lesssim \left(\int_0^s \left(\frac{t}{s} \right) t^{\frac{\alpha}{2}} \frac{dt}{t} + \int_s^1 \left(\frac{s}{t} \right)^c t^{\frac{\alpha}{2}} \frac{dt}{t} \right) \|f\|_{\mathcal{C}^\alpha} \|g\|_\infty \\ &\lesssim s^{\frac{\alpha}{2}} \|f\|_{\mathcal{C}^\alpha} \|g\|_\infty, \end{aligned}$$

since $\alpha \in (-2, 1)$ and $c \geq 1$. This holds for every $s > 0$ which yields (3.4).

- If $g \in \mathcal{C}^\beta$ with $\alpha + \beta \in (-2, 1)$ then

$$\begin{aligned} \left\| Q_s^{(c)}\Pi_g^{(b)}(f) \right\|_\infty &\lesssim \left(\int_0^s \left(\frac{t}{s} \right) t^{\frac{\alpha+\beta}{2}} \frac{dt}{t} + \int_s^1 \left(\frac{s}{t} \right)^c t^{\frac{\alpha+\beta}{2}} \frac{dt}{t} \right) \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \\ &\lesssim s^{\frac{\alpha+\beta}{2}} \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}, \end{aligned}$$

since $2c \geq 1 > \alpha + \beta > -2$. This holds for every $s > 0$ which yields (3.5). ▷

Proposition 3.5. Fix an integer $b > 2$. For any $\alpha, \beta \in (-\infty, 1)$ with $\alpha + \beta > 0$, for every $f \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$, we have the continuity estimate

$$\left\| \Pi^{(b)}(f, g) \right\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}.$$

PROOF – We recall that

$$\begin{aligned} \Pi^{(b)}(f, g) &= \frac{1}{\gamma_b} \int_0^1 -P_t^{(b)} \left(Q_t^{(b-1)}f \cdot (tL)P_t^{(b)}g \right) + 2P_t^{(b)}\Gamma \left(\sqrt{t}Q_t^{(b-1)}f, \sqrt{t}P_t^{(b)}g \right) \frac{dt}{t} \\ &\quad + \frac{1}{\gamma_b} \int_0^1 -P_t^{(b)} \left((tL)P_t^{(b)}f \cdot Q_t^{(b-1)}g \right) + 2P_t^{(b)}\Gamma \left(\sqrt{t}P_t^{(b-1)}f, \sqrt{t}Q_t^{(b)}g \right) \frac{dt}{t} \\ &\quad + \frac{1}{\gamma_b} \int_0^1 2Q_t^{(b-1)}\Gamma \left(\sqrt{t}P_t^{(b)}f, \sqrt{t}P_t^{(b)}g \right) \frac{dt}{t}. \end{aligned}$$

Fix an integer $c = b - 1 \geq 1$, and consider the function $Q_s^{(c)} \Pi^{(b)}(f, g)$, for every $s \in (0, 1]$. It is given by an integral over $(0, 1)$, which we split into (I) an integral over $(0, s)$, and (II) an integral over $(s, 1)$. Since $f \in \mathcal{C}^\alpha$, the use of Proposition 2.12, with $\alpha < 1$, yields the estimate

$$\begin{aligned} & \left\| Q_t^{(b-1)} f \right\|_\infty + \left\| \sqrt{t} \Gamma(Q_t^{(b-1)} f) \right\|_\infty + \left\| (tL) P_t^{(b)} f \right\|_\infty + \left\| \sqrt{t} \Gamma(Q_t^{(b-1)} f) \right\|_\infty \\ & + \left\| \sqrt{t} \Gamma(P_t^{(b)} f) \right\|_\infty \lesssim t^{\frac{\alpha}{2}} \|f\|_{\mathcal{C}^\alpha}; \end{aligned}$$

a similar estimate holds with g in place of f , and β in place of α . Using the uniform L^∞ -boundedness of the different approximation operators, we get for the first part

$$\begin{aligned} \left\| Q_s^{(c)}(I) \right\|_\infty & \lesssim \left(\int_0^s t^{\frac{\alpha+\beta}{2}} \frac{dt}{t} \right) \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \\ & \lesssim s^{\frac{\alpha+\beta}{2}} \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}, \end{aligned}$$

where we used the strict inequality $\alpha + \beta > 0$. For the second part, we observe that for $t > s$ then

$$Q_s^{(c)} P_t = \left(\frac{s}{t} \right)^c e^{-sL} (tL)^c P_t \quad \text{and} \quad Q_s^{(c)} Q_t^{(b-1)} = \left(\frac{s}{t} \right)^c Q_t^{(c+b-1)} e^{-sL}.$$

So we get for the second part

$$\begin{aligned} \left\| Q_s^{(c)}(II) \right\|_\infty & \lesssim \left(\int_s^1 t^{\frac{\alpha+\beta}{2}} \left(\frac{s}{t} \right)^c \frac{dt}{t} \right) \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \\ & \lesssim s^{\frac{\alpha+\beta}{2}} \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}, \end{aligned}$$

where we used the fact that $2c \geq 2 > \alpha + \beta$. ▷

3.3. Commutator estimates The following commutator estimate gives sense to the difference of two terms in a framework where none of them makes sense separately, as it does not fit the conditions put forward in proposition 3.5. We fix an integer $b \geq 2$ in this section and write $\Pi(f, g)$ for $\Pi^{(b)}(f, g)$, and $\Pi_g(f)$ for $\Pi_g^{(b)}(f)$.

Proposition 3.6. Consider the a priori unbounded trilinear operator

$$C(f, g, h) := \Pi^{(b)}\left(\Pi_g^{(b)}(f), h\right) - g \Pi^{(b)}(f, h),$$

on \mathcal{S}'_o . Let α, β, γ be Hölder regularity exponents with $\alpha \in (-1, 1)$, $\beta \in (0, 1)$ and $\gamma \in (-\infty, 1]$. If

$$0 < \alpha + \beta + \gamma \quad \text{and} \quad \alpha + \gamma < 0$$

then, setting $\delta := (\alpha + \beta) \wedge 1 + \gamma$, we have

$$(3.6) \quad \left\| C(f, g, h) \right\|_{\mathcal{C}^\delta} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma},$$

for every $f \in \mathcal{C}^\alpha$, $g \in \mathcal{C}^\beta$ and $h \in \mathcal{C}^\gamma$; so the commutator defines a trilinear map from $\mathcal{C}^\alpha \times \mathcal{C}^\beta \times \mathcal{C}^\gamma$ to \mathcal{C}^δ .

PROOF – Note first that the paraproduct $\Pi_g(f)$ is given, up to a multiplicative constant, by the sum of two terms of the form

$$\mathcal{A}(f, g) = \int_0^1 Q_t^1 \left(Q_t^2 f \cdot P_t g \right) \frac{dt}{t},$$

and the resonant part $\Pi(f, g)$ by the sum of five terms of tone of the following forms

$$(3.7) \quad \mathcal{R}(f, g) = \int_0^1 \mathcal{P}_t^1 \Gamma \left(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 g \right) \frac{dt}{t},$$

or

$$\mathcal{R}(f, g) = \int_0^1 \mathcal{P}_t \left((tL) \mathcal{P}_t f \cdot \mathcal{Q}_t g \right) \frac{dt}{t},$$

or

$$\mathcal{R}(f, g) = \int_0^1 \mathcal{P}_t^1 \left(\mathcal{Q}_t f \cdot (tL) \mathcal{P}_t^2 g \right) \frac{dt}{t},$$

where the operators

- $\mathcal{Q}_t, \mathcal{Q}_t^j$ are of the form $(tL)^{b-1} p(tL) e^{-tL}$ with a polynomial function p ,
- $\mathcal{P}_t, \mathcal{P}_t^j$ are of the form $p(tL) e^{-tL}$ with a polynomial function p .

(Note also that terms of the for $\psi(tL)$ are a posteriori of the form $\phi(tL)$.) So it suffices to focus on a generic term of the form

$$D(f, g, h) := \mathcal{R}(\mathcal{A}(f, g), h) - g \mathcal{R}(f, h)$$

and prove the continuity estimate (3.6) for it. We focus on the case where \mathcal{R} has form (3.7), the treatment of the other cases being similar and somewhat easier. We split the proof of the commutator estimate (3.6) for D in two steps, and introduce an intermediate quantity

$$\mathcal{S}(f, g, h) := \int_0^1 \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot P_t g \right) \frac{dt}{t}$$

for which we shall prove that we have both

$$(3.8) \quad \|g \mathcal{R}(f, h) - \mathcal{S}(f, g, h)\|_{C^\delta} \lesssim \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma}$$

and

$$(3.9) \quad \|D(f, g, h) - \mathcal{S}(f, g, h)\|_{C^\delta} \lesssim \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma}.$$

Step 1 – proof of (3.8). We first prove a weaker version of the continuity estimate (3.8), under the form of the inequality

$$(3.10) \quad \|g \mathcal{R}(f, h) - \mathcal{S}(f, g, h)\|_\infty \lesssim \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma}.$$

As a start, remark that we have

$$(3.11) \quad (g \mathcal{R}(f, h) - \mathcal{S}(f, g, h))(x) = \int_0^1 \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) \frac{dt}{t},$$

for μ -almost every $x \in M$. Since $g \in C^\beta$, with $\beta \in (0, 1)$, we have

$$\begin{aligned} \|P_t g - g\|_\infty &\lesssim \int_0^t \|Q_s g\|_\infty \frac{ds}{s} \\ &\lesssim \left(\int_0^t s^{\beta/2} \frac{ds}{s} \right) \|g\|_{C^\beta} \lesssim t^{\beta/2} \|g\|_{C^\beta}, \end{aligned}$$

so we have

$$\begin{aligned} |P_t g(y) - g(x)| &\leq |P_t g(y) - g(y)| + |g(y) - g(x)| \\ &\lesssim t^{\beta/2} \|g\|_{C^\beta} + d(x, y)^\beta \|g\|_{C^\beta} \\ &\lesssim \left(t^{\beta/2} + d(x, y)^\beta \right) \|g\|_{C^\beta}, \end{aligned}$$

for every $x, y \in M$. Coming back to equation (3.11) and using Gaussian pointwise estimates for the kernel of \mathcal{P}_t^2 , together with Proposition 2.12, we have

$$\begin{aligned} & \left| \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) \right| \\ & \lesssim \left\{ \int_M \frac{1}{V(x, \sqrt{t})} \exp \left(-c \frac{d(x, y)^2}{t} \right) |g(x) - P_t g(y)| d\mu(y) \right\} \|\sqrt{t}\Gamma(\mathcal{P}_t^2 f)\|_\infty \|\sqrt{t}\Gamma(\mathcal{P}_t^3 h)\|_\infty \\ & \lesssim \left\{ \int_M \frac{1}{V(x, \sqrt{t})} \exp \left(-c \frac{d(x, y)^2}{t} \right) \left(t^{\beta/2} + d(x, y)^\beta \right) d\mu(y) \right\} t^{\frac{\alpha}{2}} t^{\frac{\gamma}{2}} \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma} \end{aligned}$$

The continuity estimate (3.10) comes from integrating with respect to time, taking into account the fact that $\alpha + \beta + \gamma > 0$.

Let then estimate the regularity of $g\mathcal{R}(f, h) - \mathcal{S}(f, g, h)$. For $x, y \in M$, with $d(x, y) \leq$, write

$$(g(x)\mathcal{R}(f, h) - \mathcal{S}(f, g, h))(x) - (g(y)\mathcal{R}(f, h) - \mathcal{S}(f, g, h))(y) =: U + V$$

with U defined by the formula

$$\int_0^{d(x, y)^2} \left\{ \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) - \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \cdot (g(y) - P_t g) \right) (y) \right\} \frac{dt}{t},$$

and V is defined by the formula

$$\int_{d(x, y)^2}^1 \left\{ \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) - \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \cdot (g(y) - P_t g) \right) (y) \right\} \frac{dt}{t}.$$

By repeating, the argument used in the proof of (3.10), we easily bound U by the quantity

$$\begin{aligned} U & \lesssim \left(\int_0^{d(x, y)^2} t^{(\alpha+\beta+\gamma)/2} \frac{dt}{t} \right) \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma} \\ & \lesssim d(x, y)^\delta \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma}. \end{aligned}$$

For the second part, we use

$$|V| \leq A + B$$

with A equal to

$$\left| \int_{d(x, y)^2}^1 \left\{ \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) - \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (y) \right\} \frac{dt}{t} \right|$$

and

$$B := \int_{d(x, y)^2}^1 |g(x) - g(y)| \cdot \left| \mathcal{P}_t^1 \left(\Gamma(\sqrt{t}\mathcal{P}_t^2 f, \sqrt{t}\mathcal{P}_t^3 h) \right) (y) \right| \frac{dt}{t}.$$

The last quantity is bounded by

$$\begin{aligned} B & \lesssim d(x, y)^\beta \|g\|_{C^\beta} \int_{d(x, y)^2}^1 \left\| \sqrt{t}\Gamma(\mathcal{P}_t^2 f) \right\|_\infty \left\| \sqrt{t}\Gamma(\mathcal{P}_t^3 h) \right\|_\infty \frac{dt}{t} \\ & \lesssim d(x, y)^\beta \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma} \int_{d(x, y)^2}^1 t^{\frac{\alpha+\gamma}{2}} \frac{dt}{t} \\ & \lesssim d(x, y)^\delta \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma}. \end{aligned}$$

For the quantity A , we use the Lipschitz regularity (Lip) of the heat kernel to get the upper bound,

$$\begin{aligned}
& \int_{d(x,y)^2}^1 \left\{ \int_M \frac{d(x,y)}{\sqrt{t}V(x,\sqrt{t})} \exp\left(-c\frac{d(x,z)^2}{t}\right) |g(x) - P_t g(z)| \mu(dz) \right\} \left\| \sqrt{t}\Gamma(\mathcal{P}_t^2 f) \right\|_\infty \left\| \sqrt{t}\Gamma(\mathcal{P}_t^3 h) \right\|_\infty \frac{dt}{t} \\
& \lesssim \left\{ \int_{d(x,y)^2}^1 \int_M \frac{d(x,y)}{\sqrt{t}V(x,\sqrt{t})} e^{-c\frac{d(x,z)^2}{t}} \left(d(x,z)^\beta + t^{\beta/2} \right) \mu(dz) t^{(\alpha+\gamma)/2} \frac{dt}{t} \right\} \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma} \\
& \lesssim \left(\int_{d(x,y)^2}^1 \frac{d(x,y)}{\sqrt{t}} t^{(\alpha+\beta+\gamma)/2} \frac{dt}{t} \right) \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma} \\
& \lesssim d(x,y)^{\alpha+\beta+\gamma} \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma},
\end{aligned}$$

where we have used the fact that $\alpha + \beta + \gamma \in (0, 1)$. The combination of all the previous estimates yields

$$\begin{aligned}
& \left| \left\{ g\mathcal{R}(f, h) - \mathcal{S}(f, g, h) \right\}(x) - \left\{ g\mathcal{R}(f, h) - \mathcal{S}(f, g, h) \right\}(y) \right| \\
& \leq |U| + A + B \lesssim d(x, y)^\delta \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma},
\end{aligned}$$

which concludes the proof of the continuity estimate (3.8).

Step 2 – proof of (3.9). Given the collection $(Q_r := Q_r^{(1)})_{r \in (0,1]}$ of operators, we need to prove that we have

$$(3.12) \quad \left\| Q_r \left(\mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right) \right\|_\infty \lesssim r^{\delta/2}.$$

for every $r \in (0, 1]$, and where

$$(3.13) \quad \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) = \int_0^1 \mathcal{P}_t^1 \Gamma \left(\sqrt{t} \left\{ \int_0^1 \mathcal{P}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^3 g \right) \frac{ds}{s} - P_t g \cdot \mathcal{P}_t^2 f \right\}, \sqrt{t} \mathcal{P}_t^3 h \right) \frac{dt}{t}.$$

The notation is confusing and we have to be careful: when Γ acts $P_t g \cdot \mathcal{P}_t^2 f$, it is thought to only acts on the variable of $\mathcal{P}_t^2 f$ (the variable of $P_t g$ is frozen). We shall bound above the absolute value of the Γ term in the integral, which is of the form $\Gamma(p, q)$, by $\Gamma(p)\Gamma(q)$ – recall we write $\Gamma(p)$ for $\sqrt{\Gamma(p, p)}$. Set for that purpose

$$A_t(f, g) := \sqrt{t}\Gamma \left(\int_0^1 \mathcal{P}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^3 g \right) \frac{ds}{s} - P_t g \mathcal{P}_t^2 f \right).$$

We have for almost every $x \in M$

$$\begin{aligned}
A_t(f, g)(x) & \leq \sqrt{t}\Gamma \mathcal{P}_t^2 \left(\int_0^1 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^3 g \right) (x) \frac{ds}{s} - P_t g(x) \cdot f \right) (x) \\
& \leq \int_0^1 \sqrt{t}\Gamma \mathcal{P}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \left(\mathcal{P}_s^3 g - P_t g(x) \right) \right) (x) \frac{ds}{s} + |P_t g(x)| \sqrt{t}\Gamma[\mathcal{P}_t^2 \mathcal{P}_1 f](x),
\end{aligned}$$

where we used the property

$$\int_0^1 \mathcal{Q}_s^1 \mathcal{Q}_s^2 \frac{ds}{s} = \text{Id} - \mathcal{P}_1,$$

for some \mathcal{P}_1 operator. As in Step 1, the fact that $\beta > 0$ implies

$$\begin{aligned}
\left| \mathcal{P}_s^3 g(y) - P_t g(x) \right| & \leq \left| \mathcal{P}_s^3 g(y) - g(y) \right| + |g(y) - g(x)| + |g(x) - P_t g(x)| \\
& \lesssim \left(s^{\beta/2} + t^{\beta/2} + d(x, y)^\beta \right) \|g\|_{C^\beta} \lesssim \left(\max(s, t)^{\beta/2} + d(x, y)^\beta \right) \|g\|_{C^\beta}.
\end{aligned}$$

Moreover, it follows from Lemma A.5 below, on the composition of Gaussian pointwise estimates, that the operator $\sqrt{t}\Gamma\left(\mathcal{P}_t^2\mathcal{Q}_s^1\right)$ has pointwise Gaussian estimates at the scale $\max(s, t)$ with an extra factor $\left(\frac{\min(s, t)}{\max(s, t)}\right)^{\frac{1}{2}}$; so if one sets $\tau := \max(s, t)$, we have

$$\begin{aligned} & \sqrt{t}\Gamma\mathcal{P}_t^2\mathcal{Q}_s^1\left(\mathcal{Q}_s^2f(\mathcal{P}_s^3g - P_tg(x))\right)(x) \\ & \lesssim \left(\frac{\min(s, t)}{\max(s, t)}\right)^{\frac{1}{2}} \left\{ \int_M \frac{1}{V(x, \sqrt{\tau})} e^{-c\frac{d(x, y)^2}{\tau}} \left(\tau^{\frac{\beta}{2}} + d(x, y)^\beta\right) d\mu(y) \right\} \|\mathcal{Q}_s^2f\|_\infty \|g\|_{C^\beta} \\ & \lesssim \left(\frac{\min(s, t)}{\max(s, t)}\right)^{\frac{1}{2}} \tau^{\frac{\beta}{2}} s^{\frac{\alpha}{2}} \|f\|_{C^\alpha} \|g\|_{C^\beta} \\ & \lesssim \left(\frac{\min(s, t)}{\max(s, t)}\right)^{\frac{1}{2}} \max(s, t)^{\frac{\beta}{2}} s^{\frac{\alpha}{2}} \|f\|_{C^\alpha} \|g\|_{C^\beta}. \end{aligned}$$

Integrating in s , and taking into account the fact that $\alpha > -1$ and $\alpha + \beta < 1$, we obtain for $A_t(f, g)$ the estimate

$$\begin{aligned} \|A_t(f, g)\|_\infty & \lesssim \left\{ \int_0^t \left(\frac{s}{t}\right)^{\frac{1}{2}} t^{\beta/2} s^{\alpha/2} \frac{ds}{s} + \int_t^1 \left(\frac{t}{s}\right)^{\frac{1}{2}} s^{\beta/2} s^{\alpha/2} \frac{ds}{s} + \sqrt{t} \right\} \|f\|_{C^\alpha} \|g\|_{C^\beta} \\ & \lesssim t^{\frac{\alpha+\beta}{2}} \|f\|_{C^\alpha} \|g\|_{C^\beta}. \end{aligned}$$

Observe that in the case where $\alpha + \beta \geq 1$, we get

$$\|A_t(f, g)\|_\infty \lesssim t^{\frac{1}{2}} \|f\|_{C^\alpha} \|g\|_{C^\beta},$$

Coming back to identity (3.13), we have

$$\left| \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right| \leq \int_0^1 \mathcal{P}_t^1 \left(A_t(f, g) \cdot \sqrt{t}\Gamma(\mathcal{P}_t^4 h) \right) \frac{dt}{t},$$

and since $\alpha + \beta + \gamma > 0$, it follows that

$$\begin{aligned} \left\| \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right\|_\infty & \lesssim \left(\int_0^1 t^{(\alpha+\beta+\gamma)/2} \frac{dt}{t} \right) \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma} \\ & \lesssim \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma}. \end{aligned}$$

Moreover, taking into account that we have $Q_r^{(1)}\mathcal{P}_t^1 = \frac{r}{t}\overline{Q}_t$ for $t \geq r$, and $\alpha + \beta + \gamma < 1$, we see that the estimate (3.12) holds true

$$\begin{aligned} & \left\| Q_r^{(1)} \left(\mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right) \right\|_\infty \\ & \lesssim \int_0^r \|A_t(f, g)\|_\infty \left\| \sqrt{t}\Gamma(\mathcal{P}_t^4 h) \right\|_\infty \frac{dt}{t} + \int_r^1 \frac{r}{t} \|A_t(f, g)\|_\infty \left\| \sqrt{t}\Gamma(\mathcal{P}_t^4 h) \right\|_\infty \frac{dt}{t} \\ & \lesssim \left(\int_0^r t^{(\alpha+\beta+\gamma)/2} \frac{dt}{t} + \int_r^1 r t^{(\alpha+\beta+\gamma-2)/2} \frac{dt}{t} \right) \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma} \\ & \lesssim r^{\frac{\delta}{2}} \|f\|_{C^\alpha} \|g\|_{C^\beta} \|h\|_{C^\gamma}. \end{aligned}$$

▷

3.4. Paralinearization and composition estimates Two ingredients are needed to turn the machinery of paraproducts into an efficient tool.

To understand how nonlinear functions act on Hölder functions \mathcal{C}^α , with $0 < \alpha < 1$, and to understand how one can compose two paraproducts. The first point is the object of the following analogue of Bony's classical result on paralinearization [13], while the second point is dealt with by theorem 3.8 below.

Theorem 3.7. Let fix an integer $b \geq 2$, $\alpha \in (0, 1)$, and consider a nonlinearity $F \in C_b^3$. Then for every $f \in \mathcal{C}^\alpha$, we have $F(f) \in \mathcal{C}^\alpha$ and

$$R_F(f) := F(f) - \Pi_{F'(f)}^{(b)}(f) \in \mathcal{C}^{2\alpha}.$$

More precisely

$$\left\| F(f) - \Pi_{F'(f)}^{(b)}(f) \right\|_{\mathcal{C}^{2\alpha}} \lesssim \|F\|_{C_b^3} \left(1 + \|f\|_{\mathcal{C}^\alpha}^2\right).$$

If $F \in C_b^4$ then the remainder term $R_F(f)$ is Lipschitz with respect to f , in so far far as we have

$$\left\| R_F(f) - R_F(g) \right\|_{\mathcal{C}^{2\alpha}} \lesssim \|F\|_{C_b^4} \left(1 + \|f\|_{\mathcal{C}^\alpha} + \|g\|_{\mathcal{C}^\alpha}\right)^2 \|f - g\|_{\mathcal{C}^\alpha}.$$

PROOF – First using the Leibniz rule for the operator L , we know that for $h \in \mathcal{C}^\alpha$ then

$$L(F(h)) = F'(h)L(h) + F''(h)\Gamma(h)^2.$$

Now, since the semigroup is continuous at $t = 0$, we have

$$F(f) = \lim_{t \rightarrow 0} P_t^{(b)} F(P_t^{(b)} f),$$

so we can write

$$\begin{aligned} F(f) &= - \int_0^1 \frac{d}{dt} P_t^{(b)} F(P_t^{(b)} f) dt + P_1^{(b)} F(P_1^{(b)} f) \\ &= \frac{1}{\gamma_b} \int_0^1 \left\{ Q_t^{(b)} (F(P_t^{(b)} f)) + P_t^{(b)} \left(Q_t^{(b)} f \cdot F'(P_t^{(b)} f) \right) \right\} \frac{dt}{t} + P_1^{(b)} F(P_1^{(b)} f). \end{aligned}$$

Using the relation $Q_t^{(b)} = Q_t^{(b-1)}(tL)$, together with the chain rule

$$L\left(F(P_t^{(b)} f)\right) = F'(P_t^{(b)} f)LP_t^{(b)} f + F''(P_t^{(b)} f)\Gamma(P_t^{(b)} f)^2,$$

we get

$$Q_t^{(b)} \left(F(P_t^{(b)} f) \right) = Q_t^{(b-1)} \left((tL)P_t^{(b)} f \cdot F'(P_t^{(b)} f) \right) + Q_t^{(b-1)} \left(F''(P_t^{(b)} f) \cdot t\Gamma(P_t^{(b)} f)^2 \right).$$

Note here the identity

$$\begin{aligned} P_t^{(b)} \left(Q_t^{(b)} f \cdot F'(P_t^{(b)} f) \right) &= (tL)P_t^{(b)} \left(Q_t^{(b-1)} f \cdot F'(P_t^{(b)} f) \right) - P_t^{(b)} \left(Q_t^{(b-1)} f \cdot tLF'(P_t^{(b)} f) \right) \\ &\quad - 2P_t^{(b)} t\Gamma \left(Q_t^{(b-1)} f, F'(P_t^{(b)} f) \right). \end{aligned}$$

So we have

$$F(f) - \Pi_{F'(f)}^{(b)}(f) =: (a) + (b) + (c) + (d) + (e) + (f)$$

with

$$\begin{aligned}
(a) &:= P_1^{(b)} F(P_1^{(b)} f), \\
(b) &:= \frac{1}{\gamma_b} \int_0^1 Q_t^{(b-1)} \left\{ (tL) P_t^{(b)} f \cdot \left(F'(P_t^{(b)} f) - P_t^{(b)} (F'(f)) \right) \right\} \frac{dt}{t} \\
(c) &:= \frac{1}{\gamma_b} \int_0^1 (tL) P_t^{(b)} \left\{ Q_t^{(b-1)} f \cdot \left(F'(P_t^{(b)} f) - P_t^{(b)} (F'(f)) \right) \right\} \frac{dt}{t} \\
(d) &:= \frac{1}{\gamma_b} \int_0^1 Q_t^{(b-1)} \left\{ F''(P_t^{(b)} f) \cdot t\Gamma(P_t^{(b)} f)^2 \right\} \frac{dt}{t} \\
(e) &:= -\frac{1}{\gamma_b} \int_0^1 P_t^{(b)} \left\{ Q_t^{(b-1)} f \cdot tL F'(P_t^{(b)} f) \right\} \frac{dt}{t} \\
(f) &:= -\frac{2}{\gamma_b} \int_0^1 P_t^{(b)} t\Gamma(Q_t^{(b-1)} f, F'(P_t^{(b)} f)) \frac{dt}{t}.
\end{aligned}$$

We are now going to control each of these terms in the Hölder space $\mathcal{C}^{2\alpha}$.

Step 1 – term (a). Since $f \in \mathcal{C}^\alpha$, we know that $P_1^{(b)} f \in L^\infty$, so $F(P_1^{(b)} f)$ is also bounded. From Proposition 2.11, we get

$$\|(a)\|_{\mathcal{C}^{2\alpha}} \lesssim \left\| F(P_1^{(b)} f) \right\|_\infty \lesssim \|f\|_{\mathcal{C}^\alpha}.$$

Step 2 – terms (b), (c). The following quantity appears in these two terms

$$\begin{aligned}
\left| F'(P_t^{(b)} f) - P_t^{(b)} (F'(f)) \right| &\lesssim \left\| F'(f) - F'(P_t^{(b)} f) \right\|_\infty + \left\| F'(f) - P_t^{(b)} (F'(f)) \right\|_\infty \\
&\lesssim \|F''\|_\infty \left\| f - P_t^{(b)} f \right\|_\infty + \left\| F'(f) - P_t^{(b)} (F'(f)) \right\|_\infty \\
&\lesssim \|F''\|_\infty \int_0^t \|Q_s^{(b)} f\|_\infty \frac{ds}{s} + \int_0^t \|Q_s^{(b)} (F'(f))\|_\infty \frac{ds}{s} \\
&\lesssim \|F''\|_\infty \left(\int_0^t s^{\alpha/2} \frac{ds}{s} \right) \|f\|_{\mathcal{C}^\alpha} + \left(\int_0^t s^{\alpha/2} \frac{ds}{s} \right) \|F'(f)\|_{\mathcal{C}^\alpha} \\
(3.14) \quad &\lesssim t^{\frac{\alpha}{2}} \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha};
\end{aligned}$$

we used along the way the characterization of Hölder space, for $0\alpha < 1$, given by Proposition 2.8, to see that

$$\|F'(f)\|_{\mathcal{C}^\alpha} \lesssim \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}.$$

Using this estimate (3.14), we deduce the following bound. Uniformly for every $s \in (0, 1)$, we have

$$\begin{aligned}
\left\| Q_s^{(1)}(b) \right\|_\infty &\lesssim \left(\int_0^1 \left\| Q_s^{(1)} Q_t^{(b-1)} \right\|_{\infty \rightarrow \infty} t^{\alpha/2} \left\| F'(P_t^{(b)} f) - P_t^{(b)} (F'(f)) \right\|_\infty \frac{dt}{t} \right) \|f\|_{\mathcal{C}^\alpha} \\
&\lesssim \left(\int_0^s t^\alpha \frac{dt}{t} + \int_s^1 \frac{s}{t} t^\alpha \frac{dt}{t} \right) \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}^2 \\
&\lesssim t^\alpha \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}^2,
\end{aligned}$$

where we used that $\left\| Q_s^{(1)} Q_t^{(b-1)} \right\|_{L^\infty \rightarrow L^\infty} \lesssim \frac{\min(s,t)}{\max(s,t)}$. That yields

$$\|(b)\|_{\mathcal{C}^{2\alpha}} \lesssim \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}^2,$$

and a similar inequality holds also for the third term (c).

Step 3 – terms (d), (e) and (f). We quickly sketch the boundedness of each of these three terms. Using Proposition 2.12, we get a bound uniform in $s \in (0, 1)$, of the form

$$\begin{aligned} \|Q_s^{(1)}(d)\|_\infty &\lesssim \left(\int_0^1 \|Q_s^{(1)} Q_t^{(b-1)}\|_{\infty \rightarrow \infty} \left\| \sqrt{t} \Gamma(P_t^{(b)} f) \right\|_\infty^2 \frac{dt}{t} \right) \|F''\|_\infty \\ &\lesssim \left(\int_0^s t^\alpha \frac{dt}{t} + \int_s^1 \frac{s}{t} t^\alpha \frac{dt}{t} \right) \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}^2 \\ &\lesssim t^\alpha \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}^2. \end{aligned}$$

Similarly

$$\begin{aligned} \|Q_s^{(1)}(f)\|_\infty &\lesssim \left(\int_0^1 \|Q_s^{(1)} P_t^{(b)}\|_{\infty \rightarrow \infty} \left\| \sqrt{t} \Gamma(Q_t^{(b-1)} f) \right\|_\infty \left\| \sqrt{t} \Gamma(F'(P_t^{(b)} f)) \right\|_\infty \frac{dt}{t} \right) \\ &\lesssim \left(\int_0^s t^\alpha \frac{dt}{t} + \int_s^1 \frac{s}{t} t^\alpha \frac{dt}{t} \right) \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}^2 \\ &\lesssim t^\alpha \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}^2, \end{aligned}$$

where we used the Leibniz rule

$$\left\| \sqrt{t} \Gamma(F'(P_t^{(b)} f)) \right\|_\infty \leq \|F''\|_\infty \left\| \sqrt{t} \Gamma(P_t^{(b)} f) \right\|_\infty \lesssim t^{\alpha/2} \|F''\|_\infty \|f\|_{\mathcal{C}^\alpha}.$$

For the remaining last term (e), we can still using the Leibniz rule and get

$$\left\| t L F'(P_t^{(b)} f) \right\|_\infty \leq \left(\|F''\|_\infty + \|F'''\|_\infty \right) \left(\left\| (tL) P_t^{(b)} f \right\|_\infty + \left\| \sqrt{t} \Gamma(P_t^{(b)} f) \right\|_\infty^2 \right)$$

which then yields

$$\begin{aligned} \|Q_s^{(1)}(e)\|_\infty &\lesssim \int_0^1 \|Q_s^{(1)} P_t^{(b)}\|_{\infty \rightarrow \infty} \left\| Q_t^{(b-1)} f \right\|_\infty \left\| t L F'(P_t^{(b)} f) \right\|_\infty \frac{dt}{t} \\ &\lesssim \left(\int_0^s t^\alpha \frac{dt}{t} + \int_s^1 \frac{s}{t} t^\alpha \frac{dt}{t} \right) \|F\|_{\mathcal{C}_b^3} \|f\|_{\mathcal{C}^\alpha}^2 \\ &\lesssim t^\alpha \|F\|_{\mathcal{C}_b^3} \|f\|_{\mathcal{C}^\alpha}^2. \end{aligned}$$

By combining the previous estimates, we conclude that we have

$$\|(d)\|_{\mathcal{C}^{2\alpha}} + \|(e)\|_{\mathcal{C}^{2\alpha}} + \|(f)\|_{\mathcal{C}^{2\alpha}} \lesssim \|F\|_{\mathcal{C}_b^3} \|f\|_{\mathcal{C}^\alpha} (1 + \|f\|_{\mathcal{C}^\alpha}),$$

which ends the proof of the estimate of the remainder. The Lipschitz regularity of the remainder term is proved by very similar arguments which we leave to the reader. \triangleright

Let us now examine the composition of two paraproducts. Note that for $u \in \mathcal{C}^\alpha$ and $v \in \mathcal{C}^\beta$, with $\alpha \in (0, 1)$, $\beta \in (0, \alpha]$, we have $uv \in \mathcal{C}^\beta$.

Theorem 3.8. Fix an integer $b \geq 2$, $\alpha \in (0, 1)$, $\beta \in (0, \alpha]$ and consider $u \in \mathcal{C}^\alpha$ and $v \in \mathcal{C}^\beta$. Then for every $f \in \mathcal{C}^\alpha$, we have

$$\Pi_u^{(b)} \left(\Pi_v^{(b)}(f) \right) - \Pi_{uv}^{(b)}(f) \in \mathcal{C}^{\alpha+\beta}$$

with

$$\left\| \Pi_u^{(b)} \left(\Pi_v^{(b)}(f) \right) - \Pi_{uv}^{(b)}(f) \right\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|f\|_{\mathcal{C}^\alpha} \|u\|_{\mathcal{C}^\alpha} \|v\|_{\mathcal{C}^\beta}.$$

PROOF – We leave a detailed proof to the reader and we just sketch it, since it is similar and easier than the proof of Theorem 3.7. Following Proposition 3.3, we know that the two terms $\Pi_u^{(b)}\left(\Pi_v^{(b)}(f)\right)$ and $\Pi_{uv}^{(b)}(f)$ belong to \mathcal{C}^α . The idea is to use the \mathcal{C}^β -regularity of v to gain the same regularity in the difference.

Indeed, adopting the notations used above, the paraproduct $\Pi_g^{(b)}(f)$ is given, up to a multiplicative constant, by two terms with the form

$$I(f, g) = \int_0^1 \mathcal{Q}_t^1 \left(\mathcal{Q}_t^2 f \cdot \mathcal{P}_t^1 g \right) \frac{dt}{t},$$

where in \mathcal{Q}_t^1 and \mathcal{Q}_t^2 we have at least a term (tL) to the power 1. Let us focus on this form. Then we have

$$I(I(f, v), u) = \int_0^1 \int_0^1 \mathcal{Q}_t^1 \left(\mathcal{Q}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^1 v \right) \cdot \mathcal{P}_t^1 u \right) \frac{ds dt}{st}$$

and

$$I(f, vu) = \int_0^1 \int_0^1 \mathcal{Q}_t^1 \left(\mathcal{Q}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \right) \cdot \mathcal{P}_t^1 (vu) \right) \frac{ds dt}{st},$$

where we have used the normalization $\Pi_1^{(b)} = \text{Id}$, which means here that $I(f, vu) = I(I(f, 1), vu)$. Then using the \mathcal{C}^β -regularity of v and the fact that \mathcal{Q}_t^i involves at least a power 1 of (tL) , one can check that uniformly in $s, t \in (0, 1)^2$, we have

$$\begin{aligned} & \left\| \mathcal{Q}_t^1 \left(\mathcal{Q}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^1 v \right) \cdot \mathcal{P}_t^1 u \right) - \mathcal{Q}_t^1 \left(\mathcal{Q}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \right) \cdot \mathcal{P}_t^1 (vu) \right) \right\|_\infty \\ & \lesssim \frac{\min(s, t)}{\max(s, t)} s^{\frac{\alpha}{2}} (s+t)^{\frac{\beta}{2}} \|f\|_{\mathcal{C}^\alpha} \|v\|_{\mathcal{C}^\beta} \|u\|_{\mathcal{C}^\alpha}. \end{aligned}$$

So integrating in $s \in (0, 1)$ yields for $\alpha + \beta < 2$

$$\begin{aligned} & \int_0^1 \left\| \mathcal{Q}_t^1 \left(\mathcal{Q}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^1 g \right) \cdot \mathcal{P}_t^1 u \right) - \mathcal{Q}_t^1 \left(\mathcal{Q}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f \right) \cdot \mathcal{P}_t^1 (vu) \right) \right\|_\infty \frac{ds}{s} \\ & \lesssim t^{\frac{\alpha+\beta}{2}} \|f\|_{\mathcal{C}^\alpha} \|v\|_{\mathcal{C}^\beta} \|u\|_{\mathcal{C}^\alpha}. \end{aligned}$$

Then as previously we check that for every $\tau \in (0, 1)$ we have

$$\begin{aligned} & \left\| \mathcal{Q}_\tau^1 \left(I(I(f, v), u) - I(f, vu) \right) \right\|_\infty \\ & \lesssim \left(\int_0^1 \frac{\min(\tau, t)}{\max(\tau, t)} t^{(\alpha+\beta)/2} \frac{dt}{t} \right) \|f\|_{\mathcal{C}^\alpha} \|v\|_{\mathcal{C}^\beta} \|u\|_{\mathcal{C}^\alpha} \\ & \lesssim \tau^{\frac{\alpha+\beta}{2}} \|f\|_{\mathcal{C}^\alpha} \|v\|_{\mathcal{C}^\beta} \|u\|_{\mathcal{C}^\alpha}, \end{aligned}$$

since $\alpha + \beta < 2$. That allows us to conclude that

$$\left\| I(I(f, v), u) - I(f, vu) \right\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|f\|_{\mathcal{C}^\alpha} \|v\|_{\mathcal{C}^\beta} \|u\|_{\mathcal{C}^\alpha}.$$

▷

3.5. Schauder estimates Proposition 3.10 gives an elementary proof in our setting of a Schauder-type estimate about the regularizing character of the convolution operation with the operators $P_s^{(b)}$. Its paracontrolled analogue, given in section 4.2 provides a crucial ingredient in the study of parabolic singular PDEs, from the point of view of paracontrolled distributions.

Definition 3.9. For $\alpha \in (0, 2)$ and $T > 0$, we set

$$C_T \mathcal{C}^\alpha := \left\{ f \in L^\infty(\mathcal{S}'_o), \|f\|_{C_T \mathcal{C}^\alpha} := \sup_{t \in [0, T]} \|f(t)\|_{\mathcal{C}^\alpha} < \infty \right\}$$

and

$$C_T^{\alpha/2} L^\infty := \left\{ f \in L^\infty([0, T] \times M), \|f\|_{C_T^{\alpha/2} L^\infty} := \sup_{\substack{s \neq t \\ 0 \leq s, t \leq T}} \frac{\|f(t) - f(s)\|_\infty}{|t - s|^{\frac{\alpha}{2}}} < \infty \right\}.$$

We then define the space

$$\mathcal{L}_T^\alpha := C_T \mathcal{C}^\alpha \cap C_T^{\alpha/2} L^\infty.$$

Proposition 3.10. Consider an integer $b \geq 0$ and $\beta \in \mathbb{R}$. For every $T > 0$ and $v \in C_T \mathcal{C}^\beta$ then $V(t) := \int_0^t P_{t-s}^{(b)} v(s) ds$ belongs to $C_T \mathcal{C}^{\beta+2}$ with for every $t \in [0, T]$

$$\|V(t)\|_{\mathcal{C}^{\beta+2}} \lesssim (1+T) \sup_{s \in [0, t]} \|v(s)\|_{\mathcal{C}^\beta}.$$

Moreover if $-2 < \beta < 0$ then we also have

$$\|V\|_{C_T^{\frac{\beta+2}{2}} L^\infty} \lesssim \|v\|_{C_T \mathcal{C}^\beta}.$$

PROOF – We consider another integer $c \geq |\beta|/2 + 1$ and a parameter $\tau \in (0, 1]$. Then

$$Q_\tau^{(c)}(V(t)) = \int_0^t Q_\tau^{(c)} P_{t-s}^{(b)} v(s) ds.$$

We have

$$\|Q_\tau^{(c)} P_{t-s}^{(b)} v(s)\|_\infty \lesssim \left(\frac{\tau}{\tau + t - s} \right)^c \|Q_{\tau+t-s}^{(c)} v(s)\|_\infty \lesssim \left(\frac{\tau}{\tau + t - s} \right)^c (\tau + t - s)^{\frac{\beta}{2}} \|v(s)\|_{\mathcal{C}^\beta}.$$

So by integrating, it comes

$$\begin{aligned} \|Q_\tau^{(c)}(V(t))\|_\infty &\lesssim \left\{ \int_0^t \left(\frac{\tau}{\tau + t - s} \right)^c (\tau + t - s)^{\frac{\beta}{2}} ds \right\} \sup_{s \in [0, t]} \|v(s)\|_{\mathcal{C}^\beta} \\ &\lesssim \tau^{\frac{\beta}{2}+1} \sup_{s \in [0, t]} \|v(s)\|_{\mathcal{C}^\beta}. \end{aligned}$$

This holds uniformly in $\tau \in (0, 1]$ and so one concludes the proof of the first statement with the global inequality

$$\|V(t)\|_\infty \lesssim \left\{ \int_0^t (\dots) ds \right\} \|v\|_{C_t \mathcal{C}^\beta} \lesssim T \|v\|_{C_T \mathcal{C}^\beta}.$$

For the second statement, we note that for $s < t \leq T$ we have

$$\begin{aligned} V(t) - V(s) &= \left(P_{t-s}^{(b)} - \text{Id} \right) (V(s)) + \int_s^t P_{t-r}^{(b)} (v(r)) dr \\ &= \int_0^{t-s} Q_r^{(a)} V(s) \frac{dr}{r} + \int_s^t P_{t-r}^{(b)} (v(r)) dr. \end{aligned}$$

We have

$$\begin{aligned} \left\| \int_0^{t-s} Q_r^{(a)} V(s) \frac{dr}{r} \right\|_\infty &\lesssim \left(\int_0^{t-s} r^{\frac{\beta}{2}+1} \frac{dr}{r} \right) \|V(s)\|_{C^{\frac{\beta}{2}}} \\ &\lesssim (t-s)^{\frac{\beta}{2}+1} \|V(s)\|_{C^{\beta+2}} \end{aligned}$$

and since $\beta < 0$, we also have

$$\begin{aligned} \left\| \int_s^t P_{t-r}^{(b)}(v(r)) dr \right\|_{L^\infty} &\lesssim \int_s^t \left(\int_{t-r}^1 \|Q_\tau^{(b)} v(r)\|_\infty \frac{d\tau}{\tau} + \|P_1^{(b)}(v(r))\|_\infty \right) dr \\ &\lesssim \int_s^t \left(\|v(r)\|_{C^\beta} \int_{t-r}^1 \tau^{\frac{\beta}{2}} \frac{d\tau}{\tau} + \|P_1^{(N)}(v(r))\|_\infty \right) dr \\ &\lesssim (t-s)^{\frac{\beta}{2}+1} \sup_{r \in [0,t]} \|v(r)\|_{C^\beta}, \end{aligned}$$

where we used $\frac{\beta}{2} + 1 \in (0, 1)$.

▷

Corollary 3.11. For a fixed integer $b \geq 0$ and $\alpha \in (0, 2)$, the map

$$(Jf)(t) := \int_0^t P_{t-s}^{(b)}(f(s)),$$

defined on $CC^{\alpha-2}$, satisfies

$$\|Jf\|_{\mathcal{L}_T^\alpha} \lesssim (1+T) \|f\|_{C_T C^{\alpha-2}},$$

uniformly in $T > 0$.

Remark 3.12. Observe that in Proposition 3.10 the weight $(1+T)$ can be weakened, up to a little loss on the regularity exponent. Indeed, the exact same proof allows us for some $\varepsilon \in (0, 1)$ to prove

$$\|V(t)\|_{C^{\beta+2-2\varepsilon}} \lesssim T^\varepsilon \sup_{s \in [0,t]} \|v(s)\|_{C^\beta}$$

and

$$\|V\|_{C_T^{(\beta+2-2\varepsilon)/2} L^\infty} \lesssim T^\varepsilon \|v\|_{C_T C^\beta}.$$

So

$$\|Jf\|_{\mathcal{L}_T^\alpha} \lesssim T^\varepsilon \|f\|_{C_T C^{\alpha-2+2\varepsilon}}.$$

We refer the reader to Proposition 5.3 for a detailed proof of a more difficult statement, where we show how we can improve the bound $(1+T)$ up to a small loss on the regularity.

Paracontrolled calculus

The ideas of paracontrolled calculus, as introduced in [29], have their roots in Gubinelli's notion of controlled path [28]. The latter provides an alternative formulation of Lyons' rough paths theory that offers a simple approach to the core of the theory, while rephrasing it in a very useful Banach setting. Let us have here a glimpse at this field, as a guide for what we shall be doing in this section and the next one. We refer the reader to [21] for a very nice and pedagogical introduction to the subject, assuming only here that she/he knows only the very definition of a (weak geometric) α -Hölder rough path, for some $\frac{1}{3} < \alpha \leq \frac{1}{2}$.

Assume we are given an \mathbb{R}^ℓ -valued (weak geometric) α -Hölder rough path

$$\mathbf{X} = ((X_{ts}, \mathbb{X}_{ts}))_{0 \leq s \leq t \leq T},$$

with $X_{ts} \in \mathbb{R}^\ell$ and $\mathbb{X}_{ts} \in \mathbb{R}^\ell \otimes \mathbb{R}^\ell$, and a map $\sigma \in C(\mathbb{R}^d, L(\mathbb{R}^\ell, \mathbb{R}^d))$. Following Lyons, an \mathbb{R}^d -valued path x_\bullet is said to solve the rough differential equation

$$(4.1) \quad dx_t = \sigma(x_t) \mathbf{X}(dt)$$

if one has

$$(4.2) \quad x_t - x_s = \sigma(x_s) X_{ts} + \sigma'(x_s) \sigma(x_s) \mathbb{X}_{ts} + O(|t - s|^a)$$

for all $0 \leq s \leq t \leq T$, for some constant $a > 1$. (If $X_{ts} = h_t - h_s$, and $\mathbb{X}_{ts} = \int_s^t (h_r - h_s) \otimes dh_r$, for some \mathbb{R}^ℓ -valued C^1 control h , equation (4.2) is nothing but a second order Taylor expansion for the solution to the controlled differential equation $\dot{x}_t = \sigma(x_t) \dot{h}_t$.) Gubinelli's crucial remark was to notice that for a path x_\bullet to satisfy equation (4.2), it needs to be controlled by X in the sense that one has

$$(4.3) \quad x_t - x_s = x'_s X_{ts} + O(|t - s|^{2\alpha}),$$

for some $L(\mathbb{R}^\ell, \mathbb{R}^d)$ -valued α -Hölder path x'_\bullet , here $x'_s = \sigma(x_s)$. The point of this remark is that, somewhat conversely, if we are given an $L(\mathbb{R}^\ell, \mathbb{R}^d)$ -valued α -Hölder path z_\bullet controlled by \mathbf{X} , then there exists a unique \mathbb{R}^d -valued path y_\bullet whose increments satisfy

$$y_t - y_s = z_s X_{ts} + z'_s \mathbb{X}_{ts} + O(|t - s|^a),$$

for some exponent $a > 1$. With a little bit of abuse, we write $\int_0^\bullet z_s \mathbf{X}(ds)$ for that path y_\bullet – this path depends not only on z but rather on (z, z') . This path depends continuously on (z, z') and \mathbf{X} in the right topologies. Given an \mathbb{R}^d -valued path x_\bullet controlled by \mathbf{X} , and σ sufficiently regular, *the $L(\mathbb{R}^\ell, \mathbb{R}^d)$ -valued path $z_s := \sigma(x_s)$ is controlled by \mathbf{X} , with a control of the size of (z, z') given in terms of the size of (x, x')* . So, for a path x_\bullet to solve the rough differential equation (4.1), it is necessary and sufficient that it satisfies

$$x_t - x_s = \int_s^t \sigma(x_r) \mathbf{X}(dr),$$

for all $0 \leq s \leq t \leq T$, that is, x_\bullet is a fixed point of the continuous map

$$x_\bullet \mapsto \int_0^\bullet \sigma(x_r) \mathbf{X}(dr),$$

from the space of paths controlled by \mathbf{X} to itself. (Note that we indeed need the full rough path \mathbf{X} to define that map, and not just X .) The well-posed character of equation (4.1) is then shown by proving that this map is a contraction if one works on a sufficiently small time interval.

Our present setting will not differ much from the above description. We aim in the sequel at solving equations of the form

$$(\partial_t + \Delta)u = F(u) \zeta,$$

for some distribution ζ . Comparing this equation with (4.1), the role of the rough path will be played in that setting by a pair $\mathbf{X} = (\zeta, Z)$ of distributions, with ζ in the role of dX_t , with $\Pi(Z, \zeta)$, well-defined, somehow in the role of $d\mathbb{X}_t$, and $(\partial_t + \Delta)$ in the role of $\frac{d}{dt}$. The elementary insight that the/a solution u should behave at small space scales as ζ is turned into the definition of a distribution controlled by \mathbf{X} , as given in definition 4.1 below, using the paraproduct as a means of comparison, for writing a first order Taylor expansion of u similar to identity (4.3). The crucial point of this definition is that one can make sense of the product $F(u) \zeta$, in that controlled setting, see theorem 4.2, which provides an analogue of the right

hand side of identity (4.2) defining there $\sigma(x_s) \mathbf{X}(ds)$. To run formally the above argument, we shall need to see how controlled distributions are transformed by a nonlinear map; this is the content of theorem 4.3. Some problems intrinsically linked with the multidimensional setting of the problem are dealt with in section 4.2, where a version of Schauder theorem is proved for paracontrolled distributions.

4.1. Paracontrolled distributions We fix throughout that section and the next one an integer $b \geq 2$.

Definition 4.1. Let $\alpha \in (-2, 1)$ and $\beta > 0$ be given, together with $X \in \mathcal{C}^\alpha$. A pair of **distributions** $(f, g) \in \mathcal{C}^\alpha \times \mathcal{C}^\beta$ is said to be **paracontrolled by** X if $(f, g)^\sharp := f - \Pi_g^b(X)$ satisfies

$$(f, g)^\sharp \in \mathcal{C}^{\alpha+\beta}.$$

In such a case, we write $(f, g) \in b_\alpha^\beta(X)$ and define the norm

$$\|(f, g)\|_{b_\alpha^\beta} := \|(f, g)^\sharp\|_{\mathcal{C}^{\alpha+\beta}} + \|g\|_{\mathcal{C}^\beta}.$$

If $Y \in \mathcal{C}^\alpha$ and $(h, k) \in b_\alpha^\beta(Y)$ then we also write

$$d_{b_\alpha^\beta}((f, g), (h, k)) := \|g - k\|_{\mathcal{C}^\beta} + \|(f, g)^\sharp - (h, k)^\sharp\|_{\mathcal{C}^{\alpha+\beta}}.$$

Note that this choice of norm allows to compare paracontrolled distributions associated with *different* model distributions X and Y . Following the terminology of [29], the function g is called the **derivative of** f , and the term $(f, g)^\sharp$, the **remainder**; one should think of the decomposition

$$f = \Pi_g^b(X) + (f, g)^\sharp$$

as a kind of first order Taylor formula for f , in terms of regularity properties. The notion of derivative depends of course on which model distribution is used. As a first step towards completing the above program, the following statement gives an analogue in our setting of the right hand side of identity (4.3) defining $\sigma(x_s) \mathbf{X}(ds)$ in the rough paths context. It is motivated by the following simple regularity analysis based on propositions 3.3 and 3.5, giving regularity conditions for the well-posed character of terms if the form $\Pi_u^{(b)}(v)$ or $\Pi^{(b)}(u, v)$. Given $f \in \mathcal{C}^\alpha$ and $v \in \mathcal{C}^\gamma$, with $0 < \alpha < 1$ and $\gamma < 0$, we have from Calderon's identity the formal identity

$$fv = \Pi_f^{(b)}(v) + \Pi_v^{(b)}(f) + \Pi^{(b)}(f, v),$$

where the only term that is potentially undefined is the diagonal term $\Pi^{(b)}(f, v)$. If however, f is controlled by X , with derivative $g \in \mathcal{C}^\beta$, we can write

$$\Pi^{(b)}(f, v) = \Pi^{(b)}\left(\Pi_g^{(b)}(X), v\right) + \Pi^{(b)}((f, g)^\sharp, v),$$

with $\Pi^{(b)}\left(\Pi_g^{(b)}(X), v\right)$ well-defined if $\alpha + \beta + \gamma > 0$. So, writing

$$\Pi^{(b)}\left(\Pi_g^{(b)}(X), v\right) = C(X, g, v) + g\Pi^{(b)}(X, v),$$

we finally see that the only undefined term in the above a priori decomposition of fv is the term $\Pi^{(b)}(X, v)$, in that controlled setting. The following theorem turns that elementary regularity analysis into a constructive recipe for defining fv .

Theorem 4.2. Fix an integer $b \geq 1$. Let $\alpha \in (0, 1)$, $\beta \in (0, \alpha]$ and $\gamma < 0$ be such that

$$\alpha + \beta + \gamma \in (0, 1), \quad \alpha + \gamma < 0 < \beta.$$

Let $X \in \mathcal{C}^\alpha$, $v \in \mathcal{C}^\gamma$, $\mathfrak{p} \in \mathcal{C}^{\alpha+\gamma}$ be such that there exist sequences of smooth functions $(X_n)_{n \geq 0}, (v_n)_{n \geq 0}$ converging to X and v , in \mathcal{C}^α and \mathcal{C}^γ respectively, with the property that $\Pi^{(b)}(X_n, v_n)$ converges to \mathfrak{p} in $\mathcal{C}^{\alpha+\gamma}$. Then the application

$$(f, g) \cdot v := \Pi_f^{(b)}(v) + \Pi_v^{(b)}(f) + \Pi^{(b)}((f, g)^\sharp, v) + C(X, g, v) + g \mathfrak{p}$$

defines a trilinear operator which satisfies

$$(4.4) \quad \|(f, g) \cdot v - \Pi_f^{(b)}(v)\|_{\mathcal{C}^{\alpha+\gamma}} \lesssim \|(f, g)\|_{b_\alpha^\beta(X)} \left\{ \|v\|_{\mathcal{C}^\gamma} + \|X\|_{\mathcal{C}^\alpha} \|v\|_{\mathcal{C}^\gamma} + \|\mathfrak{p}\|_{\mathcal{C}^{\alpha+\gamma}} \right\}.$$

So $((f, g) \cdot v, f) \in b_\gamma^\alpha(v)$. Furthermore, this operation is locally Lipschitz in the sense that we have, with obvious notations where Y, w, \mathfrak{q} have the same role as X, v, \mathfrak{p} respectively,

$$\begin{aligned} & \left\| ((f, g) \cdot v, f)^\sharp - ((h, k) \cdot w, h)^\sharp \right\|_{\mathcal{C}^{\alpha+\gamma}} \\ & \lesssim C_M \left\{ d_{b_\alpha^\beta}((f, g), (h, k)) + \|X - Y\|_{\mathcal{C}^\alpha} + \|v - w\|_{\mathcal{C}^\gamma} + \|\mathfrak{p} - \mathfrak{q}\|_{\mathcal{C}^{\alpha+\gamma}} \right\}, \end{aligned}$$

where C_M is a positive constant with polynomial growth in

$$M := \max \left\{ \|X\|_{\mathcal{C}^\alpha}, \|v\|_{\mathcal{C}^\gamma}, \|\mathfrak{p}\|_{\mathcal{C}^{\alpha+\gamma}}, \|Y\|_{\mathcal{C}^\alpha}, \|w\|_{\mathcal{C}^\gamma}, \|\mathfrak{q}\|_{\mathcal{C}^{\alpha+\gamma}}, \|(f, g)\|_{b_\alpha^\beta(X)}, \|(h, k)\|_{b_\alpha^\beta(Y)} \right\}.$$

By definition of the commutator, we note that if v is smooth and $v = \Pi^{(b)}(f, g)$ then $g\Pi^{(b)}(u, v)$ is well-defined, by Proposition 3.5, and a simple computation yields

$$(f, g) \cdot v = fv.$$

So this new operation “ \cdot ” allows us to extend, in some specific situations, the pointwise multiplication between a function and a distribution.

PROOF – Let $(f, g) \in b_\alpha^\beta(X)$, with $f, X \in \mathcal{C}^\alpha$, $g \in \mathcal{C}^\beta$ and $(f, g)^\sharp \in \mathcal{C}^{\alpha+\beta}$. Let us examine each terms of

$$(f, g) \cdot v = \Pi_f^{(b)}(v) + \Pi_v^{(b)}(f) + \Pi^{(b)}((f, g)^\sharp, v) + C(X, g, v) + g \mathfrak{p}.$$

By Proposition 3.3, we have $\Pi_f^{(b)}(v) \in \mathcal{C}^\gamma$ and $\Pi_v^{(b)}(f) \in \mathcal{C}^{\alpha+\gamma}$. Proposition 3.5 yields that $\Pi^{(b)}((f, g)^\sharp, v) \in \mathcal{C}^{\alpha+\beta+\gamma}$. Applying Proposition 3.6 with $\alpha + \gamma < 0$, implies that $C(X, g, v) \in \mathcal{C}^\delta$, with $\delta = \min(\alpha + \beta, 1)$. Since $\alpha + \gamma < \beta$, then $g \mathfrak{p} \in \mathcal{C}^{\alpha+\gamma}$. Each of these terms belong also in \mathcal{C}^γ and (4.4) holds since $\alpha < 1$.

We let the reader check the Lipschitz inequality for this operation, in terms of (f, g) , X , v and \mathfrak{p} .

▷

Combining the above fact with the parilinearization formula, we are able to study the action of a nonlinearity on paracontrolled distributions, giving us the equivalent of the elementary fact that, in the above classical controlled setting for rough differential equations, the image by some map σ of a path (x, x') controlled by some reference rough path \mathbf{X} is again controlled by \mathbf{X} , and has a size given in terms of the size of (x, x') , under reasonable regularity conditions on σ .

Theorem 4.3. Let $\alpha \in (0, 1)$ and $\beta \in (0, \alpha]$. Let $X \in \mathcal{C}^\alpha$, $(f, g) \in b_\alpha^\beta(X)$ and $F \in C_b^4$. Then $(F(f), F'(f)g)$ belongs to $b_\alpha^\beta(X)$, and

$$\left\| (F(f), F'(f)g) \right\|_{b_\alpha^\beta(X)} \lesssim \|F\|_{C_b^3} (1 + \|(f, g)\|_{b_\alpha^\beta(X)}^2) (1 + \|X\|_{\mathcal{C}^\alpha}^2).$$

Moreover, this operation is locally Lipschitz in the sense that we have, with the same notations as above,

$$d_{b_\alpha^\beta} \left((F(f), F'(f)g), (F(h), F'(h)k) \right) \lesssim C_M \|F\|_{C_b^4}^3 \left(d_{b_\alpha^\beta}((f, g), (h, k)) + \|X - Y\|_{C^\alpha} \right),$$

where C_M is a constant with a polynomial growth in

$$M := \max \left\{ \|X\|_{C^\alpha}, \|Y\|_{C^\alpha}, \|(f, g)\|_{b_\alpha^\beta(X)}, \|(h, k)\|_{b_\alpha^\beta(Y)} \right\}.$$

PROOF – Consider $f \in \mathcal{C}^\alpha$ and so $F(f) \in \mathcal{C}^\alpha$ (since F is Lipschitz). We know that $F'(f)g \in \mathcal{C}^\beta$, since $F'(f) \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$. Using the notations of Theorem 3.7, we have

$$F(f) - \Pi_{F'(f)g}^{(b)}(X) = \Pi_{F'(f)}^{(b)}(f) - \Pi_{F'(f)g}^{(b)}(X) + R_F(f)$$

with $R_F(f) \in \mathcal{C}^{2\alpha} \subset \mathcal{C}^{\alpha+\beta}$. Since $f = (f, g)^\sharp + \Pi_g^{(b)}(X)$, we have

$$\Pi_{F'(f)}^{(b)}(f) = \Pi_{F'(f)}^{(b)} \Pi_g^{(b)}(X) + \Pi_{F'(f)}^{(b)}((f, g)^\sharp)$$

with $\Pi_{F'(f)}^{(b)}((f, g)^\sharp) \in \mathcal{C}^{\alpha+\beta}$ (due to Proposition 3.3). So

$$F(f) - \Pi_{F'(f)g}^{(b)}(X) \in \Pi_{F'(f)}^{(b)} \Pi_g^{(b)}(X) - \Pi_{F'(f)g}^{(b)}(X) + \mathcal{C}^{\alpha+\beta}.$$

Using Theorem 3.8, we deduce that

$$F(f) - \Pi_{F'(f)g}^{(b)}(X) \in \mathcal{C}^{\alpha+\beta},$$

which concludes the proof of $(F(f), F'(f)g) \in b_\alpha^\beta(X)$. We let the reader to check the Lipschitz inequality for this operation, in terms of (f, g) and X . \triangleright

4.2. Schauder estimate for paracontrolled distributions The above definition of a paracontrolled distribution is adapted to a time-independent setting. To deal with the time-dependent setting needed to handle the parabolic equations considered in practical examples, we use an adapted notion. Recall the definition of the space \mathcal{L}_T^α given in definition 3.9.

Definition 4.4. Let $\alpha \in \mathbb{R}$ and $\beta > 0$ be such that $\alpha + \beta \in (0, 2)$, and fix $X \in \mathcal{L}_T^\alpha$, for some $T > 0$. A pair of distributions $(f, g) \in \mathcal{L}_T^\alpha \times \mathcal{L}_T^\beta$ is said to be **paracontrolled by X** if

$$(f, g)^\sharp := f - \Pi_g^{(b)}(X) \in C_T \mathcal{C}^{\alpha+\beta} \cap C_T^{\beta/2} L^\infty.$$

In such a case, we write $(f, g) \in \mathcal{L}_{\alpha, T}^\beta(X)$ and define the norm

$$\|(f, g)\|_{b_{\alpha, T}^\beta} := \|(f, g)^\sharp\|_{C_T \mathcal{C}^{\alpha+\beta}} + \|(f, g)^\sharp\|_{C_T^{\beta/2} L^\infty} + \|g\|_{\mathcal{L}_T^\beta}.$$

If $Y \in \mathcal{L}_T^\alpha$ and $(h, k) \in \mathcal{L}_{\alpha, T}^\beta(Y)$, then we also write

$$d_{b_{\alpha, T}^\beta}((f, g), (h, k)) := \|g - k\|_{\mathcal{L}_T^\beta} + \|(f, g)^\sharp - (h, k)^\sharp\|_{C_T \mathcal{C}^{\alpha+\beta} \cap C_T^{\beta/2} L^\infty}.$$

Remark 4.5. We just point out that the previous definition is weaker than the property $(f, g)^\sharp \in \mathcal{L}_T^{\alpha+\beta}$. Indeed in $\mathcal{L}_T^{\alpha+\beta}$ the assumed time-regularity is stronger. Unfortunately, as in [29], we will not be able to solve the fix point (associated to PAM equation) with this stronger norm, but only with the one defined previously.

Theorem 4.6. Let $\beta \in (0, 1)$, $\alpha \in (0, 2 - \beta)$ and a fixed positive time horizon T be given. For some $v \in C_T \mathcal{C}^{\alpha-2}$, let X be the solution on $[0, T)$ of

$$\mathcal{L}X := (\partial_t + L)X = v$$

with $X|_{t=0} = 0$. Consider $g \in \mathcal{L}_T^\beta$, $h \in C_T \mathcal{C}^{\alpha+\beta-2}$ and $f_0 \in \mathcal{C}^{\alpha+\beta}$, and denote by f the solution to the initial value problem

$$\mathcal{L}f = h + \Pi_g^{(b)}(v)$$

with $f|_{t=0} = f_0$. Then we have $(f, g) \in b_{\alpha, T}^\beta(X)$ and

$$\|(f, g)\|_{b_{\alpha, T}^\beta} \lesssim \|f_0\|_{\mathcal{C}^{\alpha+\beta}} + (1 + T) \left\{ \|g\|_{\mathcal{L}_T^\beta} (1 + \|v\|_{C_T \mathcal{C}^{\alpha-2}}) + \|h\|_{C_T \mathcal{C}^{\alpha+\beta-2}} \right\}.$$

Moreover, the map $(v, X, g, h, f_0) \mapsto (f, g)$ is locally Lipschitz.

PROOF – Since $v \in C_T \mathcal{C}^{\alpha-2}$ then Corollary 3.11 yields that $X \in \mathcal{L}_T^\alpha$. Moreover $g \in \mathcal{L}_T^\beta$ and f is the solution of

$$\mathcal{L}f = h + \Pi_g^{(b)}(v)$$

with $h \in C_T \mathcal{C}^{\alpha+\beta-2} \subset C_T \mathcal{C}^{\alpha-2}$ and $\Pi_g^{(b)}(v) \in C_T \mathcal{C}^{\alpha-2}$, by Proposition 3.3, so Corollary 3.11 yields that $f \in \mathcal{L}_T^\alpha$. So it remains us to check that

$$(4.5) \quad (f, g)^\sharp := f - \Pi_g^{(b)}(X) \in C_T \mathcal{C}^{\alpha+\beta} \cap C_T^{\beta/2} L^\infty.$$

Let us derive an equation for this quantity:

$$\begin{aligned} \mathcal{L}(f, g)^\sharp &= \mathcal{L}f - \mathcal{L}\Pi_g^{(b)}(X) = h + \Pi_g^{(b)}(v) - \mathcal{L}\Pi_g^{(b)}(X) \\ &= h + \Pi_g^{(b)}(\mathcal{L}X) - \mathcal{L}\Pi_g^{(b)}(X) = h + [\mathcal{L}, \Pi_g^{(b)}](X). \end{aligned}$$

By definition $h \in C_T \mathcal{C}^{\alpha+\beta-2}$. Moreover we have seen that $\Pi_g^{(b)}(v) \in C_T \mathcal{C}^{\alpha-2}$ and similarly $\Pi_g^{(b)}(X) \in C_T \mathcal{C}^\alpha$ so that $\mathcal{L}\Pi_g^{(b)}(X) \in C_T \mathcal{C}^{\alpha-2}$. By studying the difference (which consist to commute the paraproduct $\Pi_g^{(b)}$ with \mathcal{L}) with introducing an intermediate time-space paraproduct, as done in [29, Lemma 5.1] – whose proof can easily be extended to our setting, we obtain that

$$J[\mathcal{L}, \Pi_g^{(b)}](X) \in C_T \mathcal{C}^{\alpha+\beta} \cap C_T^{\beta/2} L^\infty,$$

where J is the resolution of heat equation (see Corollary 3.11). We invite the reader to check the Lipschitz inequality for this operation, in terms of v, X, g, h, f_0 . \triangleright

With this result in hands, we now have all the theoretical apparatus needed to study some examples of singular parabolic PDEs. We have chosen to illustrate our machinery on what may be one of the simplest examples of such an equation, the generalized parabolic Anderson equation, (gPAM), that was already handled in the 2-dimensional torus both by Hairer in [32] using his theory of regularity structures, and by Gubinelli, Imkeller and Perkowski in [29], using their Fourier-based paracontrolled approach. This choice is motivated by the fact that only one (probabilistic) renormalization is needed to implement the paracontrolled machinery, while further renormalizations are needed in the stochastic quantization or KPZ equations. So the reader can see in the next section the machinery at work without being overwhelmed by side probabilistic matters.

The (generalized) parabolic Anderson Model in dimension 2

This section is devoted to the study in our abstract setting of the (generalized) parabolic Anderson Model, in dimension 2. The **setting** is described in Section 2. The space (M, d, μ) is a space of homogeneous type, equipped with a semigroup $(e^{-tL})_{t>0}$ satisfying the regularity assumptions (UE) and (Lip). In the next two subsections, we restrict our attention to the dimension $\nu = 2$. Let us insist here on the fact that even in this modest setting, the above semigroup approach offers some results that seem to be beyond the present scope of the theory of regularity structures, in so far as we are for instance allowed to work in various underlying spaces and even in the Euclidean space with operators L of the form $\operatorname{div}(A\nabla)$, with A Hölder continuous – see example 2 in section 2.1. (We are also able to deal with unbounded manifolds by working with weighted noises; which can also be done with regularity structures in \mathbb{R}^d , as testified by the work [35] of Hairer and Labbé on the linear PAM equation in \mathbb{R}^3 , and their elementary approach [36] to that equation in \mathbb{R}^2 .) The first two subsections are dedicated to proving some local and global in time well-posedness results, for the deterministic (gPAM) and (PAM) equations respectively. To turn that machinery into an efficient tool for investigating stochastic PDEs in which the singular term involves a Gaussian noise, we need to lift this noise into an enriched distribution; this step requires a probabilistic limit procedure generically called a *renormalization step*. It is performed in section 5.3, in the geometric framework of a potentially unbounded manifold and a coloured noise.

5.1. Local well-posedness result for generalized PAM Fix the integer $b \geq 2$, which allows us to consider the corresponding paraproducts.

Theorem 5.1. Let $\alpha \in (\frac{2}{3}, 1)$ be given, and $\alpha' < \alpha$ be close enough to α to have $2\alpha + \alpha' - 2 > 0$; let also a large enough integer $b \geq 1$, and a finite positive time horizon T be given. Let also fix an initial data $u_0 \in \mathcal{C}^{2\alpha}$, and a nonlinearity $F \in \mathcal{C}_b^3$. Given $\zeta \in \mathcal{C}^{\alpha-2}$, denote by $Z := J(\zeta)$ the solution of the heat equation defined in Corollary 3.11, and assume that the resonant term $\Pi^{(b)}(Z, \zeta)$ is well-defined in $C_T \mathcal{C}^{\alpha-2}$. Then the generalized PAM

$$\partial_t u + Lu = F(u)\zeta, \quad u(0) = u_0$$

has a unique solution $(u, u') \in b_{\alpha, T}^{\alpha'}(Z)$ with $u' = F(u)$, provided T is small enough.

Since, we have established in the previous sections the main estimates of the paracontrolled calculus, we can this result by following the same proof as in [29], as extended here to our more abstract setting.

PROOF – Fix $\alpha \in (2/3, 1)$ with $\alpha' < \alpha$ (close enough to α such that $2\alpha + \alpha' - 2 > 0$) and $K, T > 0$. The singular perturbation $\zeta \in \mathcal{C}^{\alpha-2}$ is fixed and Proposition 5.3 shows that the resolution of the heat equation $Z := J(\zeta)$ is well defined and $Z \in \mathcal{L}_T^\alpha$. Consider $A(T, K)$, the set of couple of distributions $(u, u') \in C_T \mathcal{C}^\alpha \times C_T \mathcal{C}^{\alpha'}$ such that

$$(u, u') \in b_{\alpha, T}^{\alpha'}(Z), \quad \|(u, u')\|_{b_{\alpha, T}^{\alpha'}} \leq K, \quad u'(0) = F(u_0) \text{ and } (u, u')^\sharp(0) = u_0.$$

We define on $A(T, K)$ the map γ_T as follows. For $(u, u') \in A(T, K)$, we set $\gamma_T(u, u') = (v, F(u))$ with v the solution of

$$\partial_t v + Lv = (F(u), F'(u)u') \cdot \zeta, \quad v(0) = u_0.$$

Then since $(u, u') \in b_{\alpha, T}^{\alpha'}(Z)$ and $Z \in \mathcal{L}_T^\alpha$, Theorem 4.3 implies that $(F(u), F'(u)u') \in b_{\alpha, T}^{\alpha'}(Z)$ with

$$\left\| (F(u), F'(u)u') \right\|_{b_{\alpha, T}^{\alpha'}(Z)} \lesssim \|F\|_{C_b^3} \left(1 + \|(u, u')\|_{b_{\alpha, T}^{\alpha'}}^2 \right) (1 + \|X\|_{C_T C^\alpha}^2).$$

So since we assume that the resonant term $\Pi^{(b)}(Z, \zeta) \in C_T C^{\alpha-2}$ is well defined, then Theorem 4.2 (with $\gamma = \alpha - 2$) allows us to define $(F(u), F'(u)u') \cdot \zeta$ such that

$$\left((F(u), F'(u)u') \cdot \zeta, F(u) \right) \in b_{\alpha-2, T}^{\alpha'}(\zeta)$$

(since we have $\alpha' + 2\alpha - 2 > 0$) with

$$\begin{aligned} & \left\| (F(u), F'(u)u') \cdot \zeta - \Pi_{F(u)}^b(\zeta) \right\|_{C_T C^{2\alpha-2}} \\ & \lesssim \left\| (F(u), F'(u)u') \right\|_{b_{\alpha, T}^{\alpha'}(Z)} \left(\|\zeta\|_{C^{\alpha-2}} + \|Z\|_{C_T C^\alpha} \|\zeta\|_{C^{\alpha-2}} + \|\Pi^{(b)}(Z, \zeta)\|_{C_T C^{2\alpha-2}} \right). \end{aligned}$$

We have the decomposition

$$(F(u), F'(u)u') \cdot \zeta = \left((F(u), g - F'(u)u') \cdot \zeta - \Pi_{F(u)}^b(\zeta) \right) + \Pi_{F(u)}^b(\zeta).$$

Using Theorem 4.6 with the comment following theorem 4.2 and the fact that by definition $F(u)\zeta := (F(u), F'(u)u') \cdot \zeta$, we deduce that the solution v of the equation

$$\partial_t v + Lv = (F(u), F'(u)u') \cdot \zeta, \quad v(0) = u_0$$

with initial condition $u_0 \in \mathcal{C}^{2\alpha}$, satisfies

$$(v, F(u)) \in b_{\alpha, T}^{\alpha'}(Z),$$

with $\left\| (v, F(u)) \right\|_{b_{\alpha, T}^{\alpha'}(Z)}$ bounded above by

$$\|u_0\|_{\mathcal{C}^{2\alpha}} + T^\varepsilon \left(\|F(u)\|_{\mathcal{L}_T^\alpha} (1 + \|\zeta\|_{C^{\alpha-2}}) + \left\| (F(u), g - F'(u)u') \cdot \zeta - \Pi_{F(u)}^b(\zeta) \right\|_{C_T C^{2\alpha-2}} \right),$$

where $\varepsilon := (\alpha - \alpha')/2 > 0$. At the end, by combining all the previous estimates we conclude that $(v, F(u))$ belongs to $b_{\alpha, T}^{\alpha'}(Z)$ and has a $b_{\alpha, T}^{\alpha'}(Z)$ -norm bounded above by

$$\|u_0\|_{\mathcal{C}^{2\alpha}} + T^\varepsilon \left(\|F\|_{C_b^3} \|u\|_{\mathcal{L}_T^\alpha} (1 + \|\zeta\|_{C^{\alpha-2}}) + \|F\|_{C_b^3} (1 + \|(u, u')\|_{b_{\alpha, T}^{\alpha'}}^2) (1 + \|Z\|_{C_T C^\alpha}^2) (\star) \right),$$

with

$$(\star) := \|\zeta\|_{C^{\alpha-2}} + \|F\|_{C_b^3} \|u\|_{C_T C^\alpha} \|\zeta\|_{C^{\alpha-2}} + \|\Pi^{(b)}(Z, \zeta)\|_{C_T C^{2\alpha-2}}.$$

So then we conclude that for a large enough K and a small enough time T then γ_T maps $A(T, K)$ into $A(T, K)$. Moreover, we also have for (u, u') and (v, v') contained in $A(T, K)$ that

$$d_{b_{\alpha, T}^{\alpha'}} \left(\gamma_T(u, u'), \gamma_T(v, v') \right) \lesssim T^\varepsilon \left(\|u - v\|_{\mathcal{L}_T^\alpha} + d_{b_{\alpha, T}^{\alpha'}}((u, u'), (v, v')) \right)$$

with implicit constants depending only on K , Z and ζ . So, as ε is positive, for a small enough T the map γ_T defines a contraction of $A(T, K)$. We may then apply Picard iteration theorem to find a unique fixed point of γ_T . Since it is easy to check (as detailed in [29]) that a solution of (gPAM) has to be in $A(T, K)$ (at least for a small enough T), this shows the local in time well-posed character of the equation. \triangleright

5.2. Global well-posedness result for linear PAM We focus in this subsection on the linear (PAM) equation and prove a global in time well-posedness result in that setting. With that aim in mind, we define a weighted (in time) version of the previous functional spaces of paracontrolled distributions.

Definition 5.2. Given $\lambda \geq 1$ and $\alpha \in (-2, 2)$, set

$$\mathcal{E}_\lambda \mathcal{C}^\alpha := \left\{ f \in L_{loc}^\infty(\mathcal{S}'_0), \|f\|_{\mathcal{E}_\lambda \mathcal{C}^\alpha} := \sup_{t \geq 0} e^{-\lambda t} \|f(t)\|_{\mathcal{C}^\alpha} < \infty \right\}$$

and

$$\mathcal{E}_\lambda^{\alpha/2} L^\infty := \left\{ f \in L_{loc}^\infty(\mathcal{S}'_0), \|f\|_{\mathcal{E}_\lambda^{\alpha/2} L^\infty} := \sup_{\substack{s \neq t \\ 0 \leq s, t \leq 1}} e^{-\lambda s} \frac{\|f(t) - f(s)\|_\infty}{|t - s|^{\frac{\alpha}{2}}} < \infty \right\}.$$

We then define the space $\mathcal{L}_\lambda^\alpha := \mathcal{E}_\lambda \mathcal{C}^\alpha \cap \mathcal{E}_\lambda^{\alpha/2} L^\infty$ and similarly the space $b_{\alpha, \lambda}^\beta$.

Following the reasoning of Theorem 5.1, we aim to obtain in the linear situation some global in time results. One of the main ingredient used is given by the Schauder estimates, through Proposition 3.10 or Corollary 3.11. We now give an extension of these estimates with the above exponentially weighted spaces.

Proposition 5.3. Consider an integer $a \geq 0$, $\beta \in (-2, 0)$ and $\lambda \geq 1$. For every $\varepsilon \in (0, 1)$ and $v \in \mathcal{E}_\lambda \mathcal{C}^\beta$ the function $V(t) := \int_0^t P_{t-s}^{(a)} v(s) ds$ belongs to $\mathcal{E}_\lambda \mathcal{C}^{\beta+2-2\varepsilon}$ and satisfies the λ -uniform bounds

$$\|V\|_{\mathcal{E}_\lambda \mathcal{C}^{\beta+2-2\varepsilon}} \lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}.$$

and

$$\|V\|_{\mathcal{E}_\lambda^{(\beta+2-2\varepsilon)/2} L^\infty} \lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}.$$

Consequently,

$$\|V\|_{\mathcal{L}_\lambda^{(\beta+2-2\varepsilon)/2}} \lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}.$$

PROOF – We adapt the proof of Proposition 3.10 and add an extra new argument to consider the exponential weight in time. So consider another integer $b \geq |\beta|/2 + 1$ and a parameter $\tau \in (0, 1]$. Then

$$Q_\tau^{(b)}(V(t)) = \int_0^t Q_\tau^{(b)} P_{t-s}^{(a)} v(s) ds.$$

Hence,

$$\begin{aligned} \left\| Q_\tau^{(b)} P_{t-s}^{(a)} v(s) \right\|_\infty &\lesssim \left(\frac{\tau}{\tau + t - s} \right)^b \left\| Q_{\tau+t-s}^{(b)} v(s) \right\|_\infty \\ &\lesssim \left(\frac{\tau}{\tau + t - s} \right)^b (\tau + t - s)^{\beta/2} \|v(s)\|_{\mathcal{C}^\beta}. \end{aligned}$$

So by integrating, it comes

$$\begin{aligned} e^{-\lambda t} \left\| Q_\tau^{(b)}(V(t)) \right\|_\infty &\lesssim \left(\int_0^t e^{-\lambda(t-s)} \left(\frac{\tau}{\tau + t - s} \right)^b (\tau + t - s)^{\beta/2} ds \right) \sup_{s \in [0, t]} e^{-\lambda s} \|v(s)\|_{\mathcal{C}^\beta} \\ &\lesssim \left(\int_0^t e^{-\lambda(t-s)} \left(\frac{\tau}{\tau + t - s} \right)^b (\tau + t - s)^{\beta/2} ds \right) \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}. \end{aligned}$$

Let us just consider the integral term (temporarily denoted by I). If $t \leq \tau$ then

$$\begin{aligned} I &\leq \tau^{\beta/2} \int_0^t e^{-\lambda(t-s)} ds \lesssim \tau^{\beta/2} \frac{1 - e^{-\lambda t}}{\lambda} \lesssim \tau^{\beta/2} (\lambda t)^{(1-\varepsilon)} \lambda^{-1} \\ &\lesssim \tau^{\beta/2+1-\varepsilon} \lambda^{-\varepsilon}. \end{aligned}$$

If $t \geq \tau$ then

$$\begin{aligned} I &\leq \tau^{\beta/2} \int_{t-\tau}^t e^{-\lambda(t-s)} ds + \int_0^{t-\tau} e^{-\lambda(t-s)} \left(\frac{\tau}{t-s} \right)^b (t-s)^{\beta/2} ds \\ &\lesssim \tau^{\beta/2} \frac{1 - e^{-\lambda\tau}}{\lambda} + \tau^{\beta/2+1-\varepsilon} \int_0^{t-\tau} e^{-\lambda(t-s)} (t-s)^{\varepsilon-1} ds \\ &\lesssim \tau^{\beta/2} \frac{1 - e^{-\lambda\tau}}{\lambda} + \tau^{\beta/2+1-\varepsilon} \lambda^{-\varepsilon} \left(\int_0^\infty e^{-x} x^{\varepsilon-1} dx \right) \\ &\lesssim \tau^{\beta/2+1-\varepsilon} \lambda^{-\varepsilon}. \end{aligned}$$

So in both situations, we deduce that uniformly in $\lambda \geq 1$ and $t > 0$, it comes

$$e^{-\lambda t} \left\| Q_\tau^{(b)}(V(t)) \right\|_\infty \lesssim \tau^{\beta/2+1-\varepsilon} \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}$$

and similarly

$$e^{-\lambda t} \left\| P_1[V(t)] \right\|_\infty \lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}.$$

Consequently, we deduce that for every $t \geq 0$

$$e^{-\lambda t} \|V(t)\|_{\mathcal{C}^{\beta+2-2\varepsilon}} \lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta},$$

which yields

$$\|V\|_{\mathcal{E}_\lambda \mathcal{C}^{\beta+2-2\varepsilon}} \lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}.$$

For the second statement, for $s < t$ we have

$$\begin{aligned} V(t) - V(s) &= (P_{t-s}^{(a)} - \text{Id})[V(s)] + \int_s^t P_{t-r}^{(a)}[v(r)] dr \\ &= \int_0^{t-s} Q_r^{(a)} V(s) \frac{dr}{r} + \int_s^t P_{t-r}^{(a)}[v(r)] dr. \end{aligned}$$

So

$$\begin{aligned} e^{-\lambda t} \left\| \int_0^{t-s} Q_r^{(a)} V(s) \frac{dr}{r} \right\|_\infty &\lesssim \left(\int_0^{t-s} r^{\beta/2+1} \frac{dr}{r} \right) e^{-\lambda t} \|V(s)\|_{\mathcal{C}^{\beta+2}} \\ &\lesssim (t-s)^{\beta/2+1} e^{-\lambda(t-s)} e^{-\lambda(t-s)} \|V\|_{\mathcal{E}_\lambda \mathcal{C}^{\beta+2}} \\ &\lesssim (t-s)^{\beta/2+1} (\lambda(t-s))^{-\varepsilon} \|V\|_{\mathcal{E}_\lambda \mathcal{C}^{\beta+2}} \\ &\lesssim (t-s)^{\beta/2+1-\varepsilon} \lambda^{-\varepsilon} \|V\|_{\mathcal{E}_\lambda \mathcal{C}^{\beta+2}} \end{aligned}$$

and also (since $\beta < 0$)

$$\begin{aligned}
e^{-\lambda t} \left\| \int_s^t P_{t-r}^{(a)} [v(r)] dr \right\|_{L^\infty} &\lesssim \int_s^t e^{-\lambda t} \left(\int_{t-r}^1 \|Q_\tau^{(a)} v(r)\|_\infty \frac{d\tau}{\tau} + \|P_1^{(a)} [v(r)]\|_\infty \right) dr \\
&\lesssim \int_s^t e^{-\lambda t} \left(\|v(r)\|_{\mathcal{C}^\beta} \int_{t-r}^1 \tau^{\beta/2} \frac{d\tau}{\tau} + \|P_1^{(a)} [v(r)]\|_{L^\infty} \right) dr \\
&\lesssim \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta} \int_s^t e^{-\lambda(t-r)} \left(\int_{t-r}^1 \tau^{\beta/2} \frac{d\tau}{\tau} + 1 \right) dr \\
&\lesssim \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta} \int_s^t e^{-\lambda(t-r)} \left((t-r)^{\beta/2} + 1 \right) dr \\
&\lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta} (t-s)^{\beta/2+1-\varepsilon},
\end{aligned}$$

where we used $\beta/2 + 1 \in (0, 1)$. So we conclude to

$$\|V\|_{\mathcal{E}_\lambda^{(\beta+2)/2-\varepsilon} L^\infty} \lesssim \lambda^{-\varepsilon} \|v\|_{\mathcal{E}_\lambda \mathcal{C}^\beta}.$$

▷

Theorem 5.4. Let $\alpha \in (\frac{2}{3}, 1)$ be given, and $\alpha' < \alpha$ be close enough to α to have $2\alpha + \alpha' - 2 > 0$; let also choose a large enough integer $b \geq 1$, and fix an initial data $u_0 \in \mathcal{C}^{2\alpha}$ with some $\lambda \geq 1$. Assume that $\zeta \in \mathcal{C}^{\alpha-2}$ and that the resonant term $\Pi^{(b)}(Z, \zeta)$ is well-defined in $\mathcal{E}_\lambda \mathcal{C}^{\alpha-2}$ where $Z = J(\zeta)$ is the solution of the heat equation $(\partial_t + L)Z = \zeta$, with null initial condition. Then if λ is large enough, the linear PAM equation

$$\partial_t u + Lu = u \zeta, \quad u(0) = u_0$$

has a unique global in time solution $(u, u') \in b_{\alpha, \lambda}^{\alpha'}(Z)$ with $u' = u$.

PROOF – Consider $\alpha \in (2/3, 1)$, K and $\lambda \geq 1$ parameters (which will be fixed later). Since $\zeta \in \mathcal{C}^{\alpha-2}$, Proposition 5.3 implies that $Z = J(\zeta) \in \mathcal{L}_\lambda^\alpha$. For some $\alpha' \in (0, \alpha)$ (close enough to α such that $2\alpha + \alpha' - 2 > 0$), consider the set $A(\lambda, K)$ defined as

$$\left\{ (u, u') \in \mathcal{E}_\lambda \mathcal{C}^\alpha \times \mathcal{E}_\lambda \mathcal{C}^{\alpha'} ; (u, u') \in b_{\alpha, \lambda}^{\alpha'}(Z), \|(u, u')\|_{b_{\alpha, \lambda}^{\alpha'}} \leq K, u'(0) = u_0, (u, u')^\#(0) = u_0 \right\}.$$

We define on $A(\lambda, K)$ the map γ_λ as follows: for $(u, u') \in A(\lambda, K)$, we set $\gamma_\lambda(u, u') = (v, u)$ with v the solution of

$$\partial_t v + Lv = (u, u') \cdot \zeta, \quad v(0) = u_0.$$

Then by the same considerations, as detailed for Theorem 5.1 (with some simplifications getting around the parilinearization step since here we only consider the linear situation) and using Proposition 5.3 instead of Proposition 3.10, we get the following: for $(u, u') \in A(\lambda, K)$ then $\gamma_\lambda(u, u')$ belongs to $b_{\alpha, \lambda}^{\alpha'}(Z)$ and satisfies (uniformly in $\lambda \geq 1$) with $\varepsilon := \alpha - \alpha' > 0$

$$\|\gamma_\lambda(u, u')\|_{b_{\alpha, T}^{\alpha'}(Z)} \lesssim \|u_0\|_{\mathcal{C}^{2\alpha}} + \lambda^{-\varepsilon} \left(\|u\|_{\mathcal{L}_\lambda^\alpha} (1 + \|\zeta\|_{\mathcal{C}^{\alpha-2}}) + \|(u, u')\|_{b_{\alpha, T}^{\alpha'}} (\star) \right),$$

with

$$(\star) := \left(\|\zeta\|_{\mathcal{C}^{\alpha-2}} + \|Z\|_{\mathcal{E}_\lambda \mathcal{C}^\alpha} \|\zeta\|_{\mathcal{C}^{\alpha-2}} + \|\Pi^{(b)}(Z, \zeta)\|_{\mathcal{C}_T \mathcal{C}^{2\alpha-2}} \right).$$

Since $\varepsilon > 0$, we deduce that for K and λ large enough then γ_λ maps $A(\lambda, K)$ into $A(\lambda, K)$. Moreover, we also have for (u, u') and (v, v') contained in $A(\lambda, K)$

$$d_{b_{\alpha, \lambda}^{\alpha'}} \left(\gamma_\lambda(u, u'), \gamma_\lambda(v, v') \right) \lesssim \lambda^{-\varepsilon} \left(\|u - v\|_{\mathcal{L}_\lambda^\alpha} + d_{b_{\alpha, \lambda}^{\alpha'}}((u, u'), (v, v')) \right)$$

with implicit constants depending only on K , Z and ζ . So for a large enough $\lambda \geq 1$ then γ_λ defines a contraction on $A(\lambda, K)$. We may then apply Picard iteration theorem to find a fixed point of γ_λ which yields a global (in time) existence and uniqueness of solution. \triangleright

5.3. Renormalization for a weighted noise We cannot expect to work in the Besov spaces used above when working in unbounded ambient spaces and with a spatial white noise; so weights need to be introduced, with a choice to be made. We can either put the weight in the Hölder spaces and still consider a uniform white noise, or we can put the weight on the noise and consider a coloured noise with values in unweighted Hölder spaces. The first approach has been recently implemented by Hairer and Labbé in a forthcoming work (see [36]) on the linear (PAM) equation in \mathbb{R}^3 . We chose to work with the second option here, partly motivated by exploring this unexplored question, partly because it seems to us that spatial white noise in an unbounded space has more something of a mathematical abstraction than of a model for real-life phenomena.

Definition 5.5. Let ω be an $L^2(\mu)$ weight on M ; the **noise with weight** ω is the centered Gaussian process ξ indexed by $L^2(\omega\mu)$, such that for every continuous function $f \in L^2(\omega\mu)$ we have

$$(5.1) \quad \mathbb{E}[\xi(f)^2] = \int f^2(x) \omega(x) \mu(dx).$$

Let us define the following notation. For $t > 0$, we denote by \mathcal{G}_t the Gaussian kernel

$$\mathcal{G}_t(x, y) := \frac{1}{V(x, \sqrt{t})} e^{-c \frac{d(x, y)^2}{t}};$$

it also depends on the positive constant c , although we do not mention it in the notation for convenience. Since, we will have to "commute" in some sense the Gaussian kernels with the weight ω defining the colour of the noise, it seems natural to make the following assumption. We **assume** the existence of some implicit constant such that for every $t \in (0, 1]$ and every $x, y \in M$, we have

$$(5.2) \quad \mathcal{G}_t(x, y) \omega(y) \lesssim \omega(x) \mathcal{G}_t(x, y).$$

Recall the definition of **Ahlfors regularity** of a measure μ on a metric space (M, d) , given in section 2.3 before the Sobolev embedding theorem 2.9, and quantified in equation (2.9). In that setting, it is relatively elementary to use the latter and prove by classical means that a coloured noise, as defined above, has a realization that takes almost-surely its values in some Hölder space.

Proposition 5.6. Assume that (M, d, μ) is Ahlfors regular and let ξ be a noise on M , with weight $\omega \in L^1 \cap L^\infty$ satisfying the assumption (5.2). Then, for every $\sigma < -\frac{\nu}{2}$, there exists a version of ξ , still denoted by the same symbol, which takes almost surely its values in \mathcal{C}^σ .

PROOF – It suffices from general principles and lemma 2.9 to check that the two expectations

$$\mathbb{E} \left(\int_{\mathbb{R}^d} |e^{-L} f|^p(x) \mu(dx) \right)$$

and

$$(\star) := \mathbb{E} \left(\int_0^1 t^{-p \frac{\sigma}{2}} \left\| Q_t^{(a)} f \right\|_p^p \frac{dt}{t} \right)$$

are finite for every $p > 2$. We show how to deal with the second expectation, the first one being easier to treat with similar arguments. Starting from the fact that $(Q_t^{(a)} f)(x)$ is, for

every $x \in M$, a Gaussian random variable with covariance the $L^2(\omega\mu)$ -norm of $K_{Q_t^{(a)}}(x, \cdot)$, the equivalence of Gaussian moments with (5.1) give the upper bound

$$\begin{aligned} \mathbb{E} \left[\left| Q_t^{(a)} f \right|^p(x) \right] &\lesssim \mathbb{E} \left[\left| Q_t^{(a)} f \right|^2(x) \right]^{\frac{p}{2}} \\ &\lesssim \left| \int K_{Q_t^{(a)}}(x, z)^2 \omega(z) \mu(dz) \right|^{\frac{p}{2}}. \end{aligned}$$

Using the Gaussian bounds for the kernel of $Q_t^{(a)}$ with property (5.2) and Ahlfors regularity, this implies for $t \in (0, 1)$

$$\mathbb{E} \left[\left| Q_t^{(a)} f \right|^p(x) \right] \lesssim \omega(x)^{\frac{p}{2}} t^{-\nu \frac{p}{4}}.$$

Hence, it follows

$$(\star) \lesssim \int_0^1 \int_M \omega(x)^{\frac{p}{2}} t^{-p \frac{\nu}{4}} t^{-p \frac{\sigma}{2}} \mu(dx) \frac{dt}{t} \lesssim \|\omega\|_{\frac{p}{2}}$$

if $\sigma < -\frac{\nu}{2}$. We conclude since $\omega \in L^1 \cap L^\infty \subset L^{\frac{p}{2}}$. ▷

Let ξ be a coloured noise, with weight ω , and define for every $s > 0$, a function $g_s : M \rightarrow \mathbb{R}$, by the formula

$$g_s(x) := \mathbb{E} \left[\Pi(e^{-sL}\xi, \xi)(x) \right];$$

so that we formally have

$$\int_0^\infty g_s(x) ds = \mathbb{E} \left[\Pi(L^{-1}\xi, \xi)(x) \right].$$

An explicit computation can be used in the case of the torus and the white noise to show that this integral diverges; see [29]. A similar computation can be done in our setting with the help of a highly non-trivial estimate on the kernel of the operators $Q_s^{(1)}$, showing that the above integral also diverges at almost all points x of M . These facts justifies that we consider the modified integral (5.3) below. Even though we shall only use here theorem 5.7 in a 2-dimensional setting, we prove it in the optimal range of homogeneous dimensions $d \in [2, 4)$, for use in forthcoming works. Denote by ϑ the function $J(\xi)$ solution to the linear equation $(\partial_t + \Delta)\vartheta = \xi$.

Theorem 5.7 (Renormalization). Assume that (M, d, μ) is locally Ahlfors regular, with homogeneous dimension $d = \nu \in [2, 4)$. Consider ξ a weighted noise with weight $\omega \in L^1 \cap L^\infty$ satisfying assumptions (5.2). For some integer $a \geq 0$, set

$$(5.3) \quad (\vartheta \diamond \xi)(t) := \int_0^t \left\{ \Pi(e^{-sL}\xi, \xi)(x) - g_s(x) \right\} ds,$$

where we recall that $g_s(x) := \mathbb{E} \left[\Pi(e^{-sL}\xi, \xi)(x) \right]$, and we write Π for $\Pi^{(a)}$. Consider one of the following time functional space $\mathcal{F} = C_T$ (for some arbitrary $T < \infty$) or $\mathcal{F} = \mathcal{E}_\lambda$, for some arbitrary $\lambda \geq 1$. Then for every $\alpha \in \left(1 - \frac{d}{4}, 2 - \frac{d}{2}\right)$ and $p \in (1, \infty)$, we have

$$\mathbb{E} \left[\left\| \vartheta \diamond \xi \right\|_{\mathcal{F}C^{2\alpha-2}}^p \right] < \infty.$$

Moreover, by considering for $\varepsilon \in (0, 1)$, the regularized versions $\xi^\varepsilon := e^{-\varepsilon L}\xi$, and $\vartheta^\varepsilon := e^{-\varepsilon L}\vartheta$, and $c^\varepsilon := \int_0^\infty \mathbb{E} \left[\Pi(e^{-sL}\xi^\varepsilon, \xi^\varepsilon) \right] ds$, then for every $p \in [1, \infty)$, we have

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[\left\| \vartheta \diamond \xi - (\Pi^{(b)}(\vartheta^\varepsilon, \xi^\varepsilon) - c^\varepsilon) \right\|_{\mathcal{F}C^{2\alpha-2}}^p \right] = 0.$$

Remarks 5.8. • In particular, if the ambient space M is bounded (and so of finite measure) then the constant weight $\omega \equiv 1$ satisfies (5.2) and belongs to $L^1 \cap L^\infty$. So the previous results can be applied to white noise.

- In Proposition 5.6 as well as in Theorem 5.7, we do not need $\omega \in L^1 \cap L^\infty$. Indeed, what is really needed is $\omega \in L^p$ for sufficiently large and finite exponent p .
- Consider a fix point $o \in M$, then any weight of the form $\omega(x) = (1 + d(x, o))^{-M}$ for $M > d/2$ satisfies the assumption (5.2) and belongs to $L^\infty \cap L^2$.

PROOF – By definition of white noise with colour ω , we know that if T, T' are two self-adjoint operators then for every $y, z \in M$

$$(5.4) \quad \mathbb{E}[(T\xi)(y)(T'\xi)(z)] = \int K_T(u, y)K_{T'}(u, z)\omega(u)\mu(du).$$

Moreover if T and T' are self-adjoint operators, with a kernel pointwisely bounded by Gaussian kernels at scale $t, t' \in (0, 1]$: for almost every $x, y \in M$

$$|K_T(x, y)| \lesssim \mathcal{G}_t(x, y) \quad \text{and} \quad |K_{T'}(x, y)| \lesssim \mathcal{G}_{t'}(x, y)$$

then we deduce by (5.4) and Assumption (5.2) that

$$(5.5) \quad \begin{aligned} \mathbb{E}[T\xi(y)T'\xi(z)] &= \int K_T(u, y)K_{T'}(u, z)\omega(u)\mu(du) \lesssim \int \mathcal{G}_t(u, y)\mathcal{G}_{t'}(u, z)\omega(u)\mu(du) \\ &\lesssim \omega(y) \int \mathcal{G}_t(u, y)\mathcal{G}_{t'}(u, z)\mu(du) \lesssim \omega(y)\mathcal{G}_{t+t'}(y, z), \end{aligned}$$

where we used Lemma A.5. Fix now an integer $b \geq 2 + \frac{d}{2}$ and for $r \in (0, 1]$ and $s > 0$, we define the quantity

$$\begin{aligned} A(r, s) &:= \int_0^1 \int_0^1 \left(\frac{r}{r+t_1}\right)^b \left(\frac{r}{r+t_2}\right)^b \cdots \\ &\quad \cdots \left(\frac{t_1 t_2}{(t_1+s)(t_2+s)}\right)^{\frac{1}{2}} (s+t_1+t_2)^{-\frac{d}{2}} (r+t_1+t_2)^{-\frac{d}{2}} \frac{dt_1}{t_1} \frac{dt_2}{t_2}, \end{aligned}$$

We set the function

$$\Xi_s := \Pi^{(a)}(e^{-sL}\xi, \xi) - g_s,$$

and we claim that for every $r \in (0, 1]$, $s > 0$ and every $x \in M$ then

$$(5.6) \quad \mathbb{E} \left[|Q_r^{(b)}\Xi_s(x)|^2 \right] \lesssim A(r, s)\omega(x)^2.$$

Step 1 – Proof of (5.6). The resonant (or diagonal) part of the paraproduct $\Pi^{(a)}$ is given by five terms, of the form

$$\mathcal{R}^1(f, g) = \int_0^1 \mathcal{P}_t((tL)\mathcal{P}_t^1 f \cdot \mathcal{Q}_t g) \frac{dt}{t} \quad \text{or} \quad \mathcal{R}^2(f, g) = \int_0^1 \mathcal{P}_t(\mathcal{Q}_t f \cdot (tL)\mathcal{P}_t^1 g) \frac{dt}{t},$$

or

$$\mathcal{R}^3(f, g) = \int_0^1 \mathcal{P}_t \Gamma(\sqrt{t}\mathcal{P}_t^1 f, \sqrt{t}\mathcal{P}_t^2 g) \frac{dt}{t}$$

where

- $\mathcal{P}_t, \mathcal{P}_t^1$ and \mathcal{P}_t^2 are operators of the form $p(tL)e^{-tL}$ with p a polynomial function;
- \mathcal{Q}_t is of the form $(tL)^{a-1}p(tL)e^{-tL}$ with a polynomial function p .

So both of these operators have a kernel with Gaussian pointwise estimates and we only have to deal with these three generic quantities.

Let us focus on a term of the first form and study

$$\Xi_s^1 := \mathcal{R}^1(e^{-sL}\xi, \xi)(x) - g_s^1(x) \quad \text{with} \quad g_s^1(x) := \mathbb{E}\left[II^1(e^{-sL}\xi, \xi)(x)\right].$$

Due to the covariance rule of Gaussian variables, we have for T, U, T', U' self-adjoint operators (using (5.4)) and every $y, z \in M$

$$\begin{aligned} & \mathbb{E}\left(T\xi(y)U\xi(y)T'\xi(z)U'\xi(z) - \mathbb{E}[T\xi(y)U\xi(y)]\mathbb{E}[T'\xi(z)U'\xi(z)]\right) \\ &= \mathbb{E}[T\xi(y)T'\xi(z)]\mathbb{E}[U\xi(y)U'\xi(z)] + \mathbb{E}[T\xi(y)U'\xi(z)]\mathbb{E}[U\xi(y)T'\xi(z)]. \end{aligned}$$

Hence $\mathbb{E}\left[|Q_r^{(b)}\Xi_s^1(x)|^2\right]$ is equal to

$$\begin{aligned} & \int_0^1 \int_0^1 \left\{ (Q_r^{(b)}\mathcal{P}_{t_1} \otimes Q_r^{(b)}\mathcal{P}_{t_2}) \left(\mathbb{E}\left[(t_1L)\mathcal{P}_{t_1}^1 e^{-sL}\xi(\bullet)(t_2L)\mathcal{P}_{t_2}^1 e^{-sL}\xi(\bullet)\right] \mathbb{E}\left[\mathcal{Q}_{t_1}\xi(\bullet)\mathcal{Q}_{t_2}\xi(\bullet)\right] \right) (x, x) \right. \\ & \left. + (Q_r^{(b)}\mathcal{P}_{t_1} \otimes Q_r^{(b)}\mathcal{P}_{t_2}) \left(\mathbb{E}\left[(t_1L)\mathcal{P}_{t_1}^1 e^{-sL}\xi(\bullet)\mathcal{Q}_{t_2}\xi(\bullet)\right] \mathbb{E}\left[\mathcal{Q}_{t_1}\xi(\bullet)(t_2L)\mathcal{P}_{t_2}^1 e^{-sL}\xi(\bullet)\right] \right) (x, x) \right\} \frac{dt_1}{t_1} \frac{dt_2}{t_2}, \end{aligned}$$

where we use the notation $f(\bullet, \bullet)$ for a function of two variables, with $(fg)(\bullet, \bullet)$ standing for the map $(y, z) \mapsto f(y, z)g(y, z)$. Moreover, to shorten notations, we shall use below the notation dm for the measure $\mu(dy)\mu(dz)\frac{dt_1}{t_1}\frac{dt_2}{t_2}$. By applying (5.5), it follows

$$\mathbb{E}\left[|Q_r^{(b)}\Xi_s^1(x)|^2\right] \lesssim J_1 + J_2$$

with

$$J_1 := \int \left|K_{Q_r^{(b)}\mathcal{P}_{t_1}}(x, y)\right| \left|K_{Q_r^{(b)}\mathcal{P}_{t_2}}(x, z)\right| \omega(y)\omega(z) \frac{t_1}{t_1+s} \frac{t_2}{t_2+s} \mathcal{G}_{t_1+t_2+s}(y, z) \mathcal{G}_{t_1+t_2}(y, z) dm$$

and

$$J_2 := \int \left|K_{Q_r^{(b)}\mathcal{P}_{t_1}}(x, y)\right| \left|K_{Q_r^{(b)}\mathcal{P}_{t_2}}(x, z)\right| \omega(y)\omega(z) \frac{t_1}{t_1+s} \frac{t_2}{t_2+s} \mathcal{G}_{t_1+t_2+s}(y, z) \mathcal{G}_{t_1+t_2+s}(y, z) dm.$$

Let us first explain how we can estimate the kernel of $Q_r^{(b)}\mathcal{P}_{t_1}$. Using the notation $\mathcal{P}_{t_1} = p(t_1L)e^{-t_1L}$ for some polynomial function p , it comes

$$\begin{aligned} Q_r^{(b)}\mathcal{P}_{t_1} &= \left(\frac{r}{r+\frac{1}{2}t_1}\right)^b \left((r+\frac{1}{2}t_1)L\right)^b e^{-rL} p(t_1L) e^{-t_1L} \\ &= \left(\frac{r}{r+\frac{1}{2}t_1}\right)^b Q_{r+\frac{1}{2}t_1}^{(b)} p(t_1L) e^{-\frac{1}{2}t_1L}, \end{aligned}$$

so since $r+\frac{1}{2}t_1 \simeq r+t_1$, $Q_{r+\frac{1}{2}t_1}^{(b)}$ has a kernel with Gaussian bounds at the scale $r+\frac{1}{2}t_1$ and $p(t_1L)e^{-\frac{1}{2}t_1L}$ at the scale t_1 , it follows by Lemma A.5 that $Q_r^{(b)}\mathcal{P}_{t_1}$ has a kernel pointwise bounded by \mathcal{G}_{r+t_1} with an extra factor $\left(\frac{r}{r+t_1}\right)^b$. Coming back to estimate the first term J_1 . We have the upper bound

$$\begin{aligned} J_1 &\lesssim \int \left| \frac{r^2}{(r+t_1)(r+t_2)} \right|^b \mathcal{G}_{r+t_1}(x, y) \mathcal{G}_{r+t_2}(x, z) \omega(y)\omega(z) \frac{t_1}{t_1+s} \frac{t_2}{t_2+s} \mathcal{G}_{t_1+t_2+s}(y, z) \mathcal{G}_{t_1+t_2}(y, z) dm \\ &\lesssim \omega(x)^2 \int \left| \frac{r^2}{(r+t_1)(r+t_2)} \right|^b \frac{t_1}{t_1+s} \frac{t_2}{t_2+s} \mathcal{G}_{r+t_1}(x, y) \mathcal{G}_{r+t_2}(x, z) \mathcal{G}_{t_1+t_2+s}(y, z) \mathcal{G}_{t_1+t_2}(y, z) dm, \end{aligned}$$

where we used Assumption 5.2. Due to Lemma A.5 with Ahlfors regularity (2.9), we have

$$\begin{aligned} & \iint \mathcal{G}_{r+t_1}(x, y) \mathcal{G}_{r+t_2}(x, z) \mathcal{G}_{t_1+t_2+s}(y, z) \mathcal{G}_{t_1+t_2}(y, z) \mu(dy) \mu(dz) \\ & \lesssim (t_1 + t_2 + s)^{-\frac{d}{2}} \iint \mathcal{G}_{r+t_1}(x, y) \mathcal{G}_{r+t_2}(x, z) \mathcal{G}_{t_1+t_2}(y, z) \mu(dy) \mu(dz) \\ & \lesssim (t_1 + t_2 + s)^{-\frac{d}{2}} (r + t_1 + t_2)^{-\frac{d}{2}}. \end{aligned}$$

Hence,

$$\begin{aligned} J_1 & \lesssim \omega(x)^2 \int_0^1 \int_0^1 \left| \frac{r^2}{(r+t_1)(r+t_2)} \right|^b \frac{t_1}{t_1+s} \frac{t_2}{t_2+s} (s+t_1+t_2)^{-\frac{d}{2}} (r+t_1+t_2)^{-\frac{d}{2}} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \\ & \lesssim \omega(x)^2 A(r, s). \end{aligned}$$

The second term J_2 can be similarly bounded, which concludes the proof of (5.6) for Ξ^1 . The corresponding term Ξ^2 with \mathcal{R}^2 can be estimated in the same way. So it remains us now to focus on the last and third term with

$$\mathcal{R}^3(f, g) = \int_0^1 \mathcal{P}_t \Gamma(\mathcal{P}_t^1 f, \mathcal{P}_t^2 g) \frac{dt}{t}$$

and $\Xi_s^3 := \mathcal{R}^3(e^{-sL}\xi, \xi)(x) - g_s^3(x)$. Following the exact same reasoning we have

$$\mathbb{E} \left(|Q_r^{(b)} \Xi_s^3(x)|^2 \right) \lesssim K_1 + K_2$$

with K_1 equal to

$$\int \left| K_{Q_r^{(b)} P_{t_1}}(x, y) \right| \left| K_{Q_r^{(b)} P_{t_2}}(x, z) \right| \omega(y) \omega(z) \left| \frac{t_1 t_2}{(t_1+s)(t_2+s)} \right|^{\frac{1}{2}} \mathcal{G}_{t_1+t_2+s}(y, z) \mathcal{G}_{t_1+t_2}(y, z) dv$$

and K_2 equal to

$$\int \left| K_{Q_r^{(b)} P_{t_1}}(x, y) \right| \left| K_{Q_r^{(b)} P_{t_2}}(x, z) \right| \omega(y) \omega(z) \left| \frac{t_1 t_2}{(t_1+s)(t_2+s)} \right|^{\frac{1}{2}} \mathcal{G}_{t_1+t_2+s}(y, z) \mathcal{G}_{t_1+t_2+s}(y, z) dv.$$

Following the same computations, gives us that both K_1 and K_2 are bounded as follows

$$K_1 + K_2 \lesssim \omega(x)^2 A(r, s),$$

which concludes the proof of (5.6).

Step 2 – Conclusion. We refer the reader to Lemma A.6 for a precise control of quantity A. Combining (5.6) with Lemma A.6 gives

$$(5.7) \quad \mathbb{E} \left[|Q_r^{(b)} \Xi_s(x)| \right] \lesssim \mathbb{E} \left[|Q_r^{(b)} \Xi_s(x)|^2 \right]^{\frac{1}{2}} \lesssim \omega(x) \left(\frac{r}{s+r} \right)^{\frac{1}{2}} (rs)^{-\frac{d}{4}} \left(1 + \log \left(\frac{s+r}{s} \right) \right)^{\frac{1}{2}}.$$

We then consider

$$(\vartheta \diamond \xi)(t) := \int_0^t \left(\Pi(e^{-sL}\xi, \xi)(x) - g_s(x) \right) ds.$$

We refer the reader to Definition 2.6 for the definition of Besov spaces (involving an integer b). So for all $0 \leq s < t$, it comes

$$\begin{aligned} \mathbb{E} \left[\left\| \vartheta \diamond \xi(t) - \vartheta \diamond \xi(s) \right\|_{B_{2p,2p}^{2\alpha-2}}^{2p} \right] &= \int_0^1 r^{-p(2\alpha-2)} \int_M \mathbb{E} \left(\left| Q_r^{(b)} [\vartheta \diamond \xi(t) - \vartheta \diamond \xi(s)] \right|^{2p} \right) \mu(dx) \frac{dr}{r} \\ &\lesssim \int_0^1 r^{-p(2\alpha-2)} \int_M \left(\int_s^t \mathbb{E} \left(\left| Q_r^{(b)} \Xi(\tau)(x) \right| \right) d\tau \right)^{2p} \mu(dx) \frac{dr}{r} \\ &\lesssim \|\omega\|_{2p}^{2p} \int_0^1 r^{-p(2\alpha-2)} \left(\int_s^t \left(\frac{r}{r+\tau} \right)^{\frac{1}{2}} (r\tau)^{-\frac{d}{4}} \left(1 + \log \left(\frac{\tau+r}{\tau} \right) \right)^{\frac{1}{2}} d\tau \right)^{2p} \frac{dr}{r}, \end{aligned}$$

where we have used Gaussian hypercontractivity; see [22]. So it comes, by Minkowski inequality,

$$\begin{aligned} \mathbb{E} \left[\left\| \vartheta \diamond \xi(t) - \vartheta \diamond \xi(s) \right\|_{B_{2p,2p}^{2\alpha-2}}^{2p} \right] &\lesssim \|\omega\|_{2p}^{2p} \left(\int_s^t \left(\int_0^1 r^{-2p(\alpha-1)} \left(\frac{r}{r+\tau} \right)^p (r\tau)^{-\frac{dp}{2}} \left(1 + \log \left(\frac{\tau+r}{\tau} \right) \right)^p \frac{dr}{r} \right)^{\frac{1}{2p}} d\tau \right)^{2p}. \end{aligned}$$

We have

$$\begin{aligned} &\int_0^1 r^{-2p(\alpha-1)} \left(\frac{r}{r+\tau} \right)^p (r\tau)^{-\frac{dp}{2}} \left(1 + \log \left(\frac{\tau+r}{\tau} \right) \right)^p \frac{dr}{r} \\ &\lesssim \int_0^\tau r^{-2p(\alpha-1)} \left(\frac{r}{\tau} \right)^p (r\tau)^{-\frac{dp}{2}} \frac{dr}{r} + \int_\tau^1 r^{-2p(\alpha-1)} (r\tau)^{-\frac{dp}{2}} \left(1 + \log \left(\frac{r}{\tau} \right) \right)^p \frac{dr}{r} \\ &\lesssim \tau^{-2p(\alpha-1)-dp}, \end{aligned}$$

since

$$-2(\alpha-1) - \frac{d}{2} < 0 < -2(\alpha-1) + 1 - \frac{d}{2}$$

which is equivalent to

$$1 - \frac{d}{4} < \alpha < \frac{3}{2} - \frac{d}{4}.$$

Observe that this last condition is satisfied since $d \in [2, 4)$ and so $1 - \frac{d}{4} < \alpha < 2 - \frac{d}{2} \leq \frac{3}{2} - \frac{d}{4}$. Then because of $\alpha < 2 - \frac{d}{2}$, it yields $-(\alpha-1) - \frac{d}{2} > -1$ and so

$$\begin{aligned} \mathbb{E} \left[\left\| \vartheta \diamond \xi(t) - \vartheta \diamond \xi(s) \right\|_{B_{2p,2p}^{2\alpha-2}}^{2p} \right] &\lesssim \|\omega\|_{2p}^{2p} \left(\int_s^t \tau^{-(\alpha-1)-\frac{d}{2}} d\tau \right)^{2p} \\ &\lesssim \|\omega\|_{2p}^{2p} |s-t|^{-2p(\alpha-2)-dp}. \end{aligned}$$

We can then use Kolmogorov's continuity criterion to deduce that for every $T < \infty$ and $\lambda \geq 1$, we have

$$\mathbb{E} \left[\left\| \vartheta \diamond \xi \right\|_{C_T B_{2p,2p}^{2\alpha-2}}^{2p} \right] + \mathbb{E} \left[\left\| \vartheta \diamond \xi \right\|_{\mathcal{E}_\lambda B_{2p,2p}^{2\alpha-2}}^{2p} \right] < \infty.$$

And using Besov embedding (due to Ahlfors regularity (2.9), see Lemma A.4), we know that

$$B_{2p,2p}^{2\alpha-2} \hookrightarrow B_{2p,\infty}^{2\alpha-2} \hookrightarrow B_{\infty,\infty}^{2\alpha-2-d/2p} = \mathcal{C}^{2\alpha-2-d/2p}.$$

So for every $p \geq 1$ and every $\alpha \in (1 - \frac{d}{2}, 2 - \frac{d}{2})$

$$\mathbb{E} \left[\left\| \vartheta \diamond \xi \right\|_{C_T \mathcal{C}^{2\alpha-2-d/2p}}^{2p} \right] + \mathbb{E} \left[\left\| \vartheta \diamond \xi \right\|_{\mathcal{E}_\lambda \mathcal{C}^{2\alpha-2-d/2p}}^{2p} \right] < \infty,$$

which allows us to conclude.

Step 3 – Second part of the statement. The second part of the statement about the approximation results can be similarly obtained, we only have to include some additional factors coming from

$$\xi - \xi^\varepsilon = (1 - e^{-\varepsilon L})\xi = \int_0^\varepsilon Q_\sigma^{(1)}\xi \frac{d\sigma}{\sigma}.$$

A careful examination shows that in the previous reasoning, ξ may be replaced by $Q_\sigma^{(1)}\xi$ and the difference involves some extra factors of the type

$$\frac{s\sigma}{(s+\sigma)^2}, \quad \frac{\sigma t_1}{(\sigma+t_1)^2} \quad \text{or} \quad \frac{\sigma t_2}{(\sigma+t_2)^2}.$$

In these three situations (by replacing ξ with $Q_\sigma^{(1)}\xi$), the same estimates hold with a quantity $A_\sigma(r, s)$ satisfying (instead of (A.9)), for $\eta > 0$ as small as we want

$$(5.8) \quad A_\sigma(r, s) \lesssim \left(\frac{r}{s+r}\right) (rs)^{-d/2} \left(1 + \log\left(\frac{s+r}{s}\right)\right) \left(\frac{\sigma}{\min(r, s)}\right)^\eta.$$

Then we let the reader to check that since all the conditions on the exponents are “open conditions”, then the previous reasoning can be reproduced, up to a small loss of regularity. So with

$$F_\varepsilon := \vartheta \diamond \xi - (\Pi^{(b)}(\vartheta^\varepsilon, \xi^\varepsilon) - c^\varepsilon)$$

we get for a sufficiently small $\eta > 0$

$$\begin{aligned} \mathbb{E} \left[\|F_\varepsilon(t) - F_\varepsilon(s)\|_{B_{2p, 2p}^{2\alpha-2}}^{2p} \right] &\lesssim \left(\int_0^\varepsilon \int_s^t \left(\frac{\sigma}{\tau}\right)^\eta \tau^{-(\alpha-1)-d/2} d\tau \frac{d\sigma}{\sigma} \right)^{2p} \\ &\lesssim |s-t|^{-2p(\alpha-2)-dp} \left(\frac{\varepsilon}{|s-t|}\right)^{2p\eta}, \end{aligned}$$

which allows us to conclude as previously using Besov embedding. \triangleright

PROOF OF THEOREM 1.2 – Let us fix the (coloured) white noise ξ and its regularized version $\xi^\varepsilon := e^{-\varepsilon L}\xi$. As in Theorem 5.7 or Theorem 1.2, let us consider the function

$$c^\varepsilon := \int_0^\infty \mathbb{E} [\Pi(e^{-sL}\xi^\varepsilon, \xi^\varepsilon)] ds = \mathbb{E} [\Pi(L^{-1}\xi^\varepsilon, \xi^\varepsilon)].$$

In order to make appear this term in the equation, we can introduce a suitable correction term in the regularized problems and we are conducted (as detailed in [29]) to study the following renormalized PDE

$$\partial_t u^\varepsilon + Lu^\varepsilon = F(u^\varepsilon)\xi^\varepsilon - c^\varepsilon F'(u^\varepsilon)F(u^\varepsilon).$$

We then follow the exact same approach as for Theorem 5.1 (or Theorem 5.4 for the global estimates with the spaces \mathcal{E}_λ), adapted to this modified PDE. So we only detail the required modification. We cannot use $\Pi^{(b)}(X^\varepsilon, \xi^\varepsilon)$ but we have to replace it by $\Pi^{(b)}(X^\varepsilon, \xi^\varepsilon) - c^\varepsilon$ which belongs to $C_T \mathcal{C}^{2\alpha-2}$.

Using the arguments of Theorem 4.2 (with $\gamma = \alpha - 2$), it allows us to define the following “product” for $(u, u') \in d_\alpha^\alpha(X^\varepsilon)$

$$\left((F(u), F'(u)u') \cdot \xi^\varepsilon \right) - c^\varepsilon u' F'(u) \in b_{\alpha-2, T}^{\alpha'}(\zeta)$$

uniformly with respect to ε (since an upper bound involves only $\|\Pi^{(b)}(X^\varepsilon, \xi^\varepsilon) - c^\varepsilon\|_{C_T C^{2\alpha-2}}$) with

$$\begin{aligned} & \left\| (F(u), F'(u)u') \cdot \xi^\varepsilon - \Pi_{F(u)}^b(\xi^\varepsilon) - c^\varepsilon u' F'(u) \right\|_{C_T C^{2\alpha-2}} \\ & \lesssim \left\| (F(u), F'(u)u') \right\|_{b_{\alpha,T}^{\alpha'}} \left(\|\xi^\varepsilon\|_{C^{\alpha-2}} + \|X^\varepsilon\|_{C_T C^\alpha} \|\xi^\varepsilon\|_{C^{\alpha-2}} + \|\Pi^{(b)}(X^\varepsilon, \xi^\varepsilon) - c^\varepsilon\|_{C_T C^{2\alpha-2}} \right). \end{aligned}$$

Then we conclude as in Theorem 5.1 (or Theorem 5.4), by using a fixed point theorem. \triangleright

A

Heat kernel and technical estimates

We gather in this Appendix a number of propositions whose proofs were not given in the course of the paper, so as to keep focused on the most essential aspects of our work. These proofs are given here.

We start by proving the following pointwise and L^p -estimate for the gradient of the heat semigroup.

Proposition A.1. Assume that (M, d, μ) is a doubling space equipped with a semigroup satisfying (UE) and (Lip). Then for every $t > 0$, $x_0 \in M$ and every function $f \in L^2$ we have

$$(A.1) \quad \left| (\sqrt{t}\Gamma)(e^{-tL}f)(x_0) \right| \lesssim \int_M \frac{1}{\sqrt{V(x_0, \sqrt{t})V(y, \sqrt{t})}} \exp\left(-c \frac{d(x_0, y)^2}{t}\right) |f(y)| \mu(dy).$$

Let us first introduce the following notation: for a function $f \in L_{loc}^2$ and a ball $B \subset M$, we write $\text{Osc}_B(f)$ for the L^2 oscillation of f on B defined by

$$\text{Osc}_B(f) := \left(\int_B |f - \text{Avg}_B(f)|^2 d\mu \right)^{1/2},$$

where $\text{Avg}_B(f) = \int_B f d\mu$ stands for the average of f on the ball B .

PROOF – Fix the function $f \in L^2$ and consider $g = e^{-tL}f$. By L^2 -Caccioppoli inequality (see Lemma below), we have for every x_0 and $r > 0$ that

$$\left(\int_{B(x_0, r)} \Gamma(g)^2 d\mu \right)^{\frac{1}{2}} \lesssim \frac{1}{r} \text{Osc}_{B(x_0, 2r)}(g) + \left(\int_{B(x_0, 2r)} |Lg|^2 d\mu \right)^{1/2}.$$

So if x_0 is a Lebesgue point of $\Gamma(g)^2$ and $|Lg|^2$ (which is the case for almost every point $x_0 \in M$) then taking the limit for $r \rightarrow 0$ yields

$$(A.2) \quad \Gamma(g)(x_0) \lesssim \liminf_{r \rightarrow 0} \frac{1}{r} \text{Osc}_{B(x_0, 2r)}(g) + |Lg(x_0)|.$$

Since $(tL)e^{-tL}$ has a kernel satisfying the Gaussian upper estimates (UE) (by analyticity), we deduce that

$$(A.3) \quad |tLg(x_0)| \lesssim \int_M \frac{1}{\sqrt{V(x_0, \sqrt{t})V(y, \sqrt{t})}} \exp\left(-c \frac{d(x_0, y)^2}{t}\right) |f(y)| \mu(dy).$$

Using Lipschitz regularity (Lip) for the heat kernel and doubling property, it comes for $x, z \in B(x_0, 2r)$ with $r \leq \sqrt{t}$

$$\begin{aligned} |g(x) - g(z)| &= |e^{-tL}f(x) - e^{-tL}f(z)| \\ &\lesssim \left(\frac{d(x, z)}{\sqrt{t}} \right) \int_M \frac{1}{\sqrt{V(x_0, \sqrt{t})V(y, \sqrt{t})}} \exp\left(-c \frac{d(x, y)^2}{t}\right) |f(y)| \mu(dy). \end{aligned}$$

Hence, uniformly with respect to $r \in (0, \sqrt{t})$ we obtain

$$(A.4) \quad \frac{1}{r} \text{Osc}_{B(x_0, 2r)}(g) \lesssim \frac{1}{\sqrt{t}} \int_M \frac{1}{\sqrt{V(x_0, \sqrt{t})V(y, \sqrt{t})}} \exp\left(-c \frac{d(x, y)^2}{t}\right) |f(y)| \mu(dy).$$

By combining these last inequalities (A.3) and (A.4) into (A.2), one concludes to (A.1). \triangleright

Lemma A.2 (Cacciopoli inequality). For every ball B of radius $r > 0$ and every function $f \in \mathcal{D}_2(L)$ we have

$$\left(\int_B \Gamma(f)^2 d\mu \right)^{\frac{1}{2}} \lesssim \frac{1}{r} \text{Osc}_{2B}(f) + r \left(\int_{2B} |Lf|^2 d\mu \right)^{\frac{1}{2}}.$$

Before to check this inequality, let us first recall some consequences of the Gaussian upper estimates (UE). Under (UE), we know that a scale-invariant local Sobolev inequality holds, more precisely

$$\|f\|_q^2 \lesssim |B|^{\frac{2}{q}-1} (\|f\|_2^2 + r^2 \mathcal{E}(f, f)),$$

for every ball B of radius $r > 0$, every $f \in \mathcal{D}_2(\Gamma)$ supported in B and for some $q > 2$. This inequality was introduced in [51] and was shown, under (VD), to be equivalent to (UE) in the Riemannian setting. The equivalence was stated in our more general setting in [54]. See also [14] for many reformulations of local Sobolev inequalities, an alternative proof of the equivalence with (UE), and more references.

Such a local Sobolev inequality also implies a following relative Faber-Krahn inequality (see for instance [39, Theorem 2.5], as well as [14, Section 3.3]): for every ball B with a small enough radius $r > 0$, every function $f \in \mathcal{D}_2(\Gamma)$ supported in B then

$$(A.5) \quad \|f\|_2 \lesssim r \|\Gamma(f)\|_2.$$

PROOF OF LEMMA A.2 – We refer to [9, Lemma A.1] for such a result for harmonic function: if $u \in \mathcal{D}_2(L)$ is harmonic on $2B$ (which means $L(u) = 0$ on $2B$) then

$$(A.6) \quad \left(\int_B \Gamma(u)^2 d\mu \right)^{\frac{1}{2}} \lesssim \frac{1}{r} \text{Osc}_{2B}(u).$$

Now consider $f \in \mathcal{D}_2(L)$. By [9, Lemma 4.6], it is known that there exists $u \in \mathcal{D}_2(L)$ harmonic on $2B$ such that $f - u \in \mathcal{D}(\Gamma)$ is supported on the ball $2B$. By the support property, it follows

$$\|\Gamma(f - u)\|_2^2 = \int (f - u)L(f - u) d\mu = \int (f - u)L(f) d\mu.$$

So using Faber-Krahn inequality (A.5) we obtain

$$(A.7) \quad \|f - u\|_2 \lesssim r \|\Gamma(f - u)\|_{L^2(2B)}$$

and so

$$\|\Gamma(f - u)\|_2^2 \lesssim \|f - u\|_2 \|L(f)\|_{L^2(2B)} \lesssim r \|\Gamma(f - u)\|_2 \|L(f)\|_{L^2(2B)},$$

which yields

$$(A.8) \quad \|\Gamma(f - u)\|_2 \lesssim r \|L(f)\|_{L^2(2B)}.$$

Then we split

$$\|\Gamma(f)\|_2 \leq \|\Gamma(f - u)\|_2 + \|\Gamma(u)\|_2 \lesssim r \|L(f)\|_{L^2(2B)} + \|\Gamma(u)\|_2$$

and then use (A.6) to get

$$\begin{aligned} \left(\int_B \Gamma(f)^2 d\mu \right)^{\frac{1}{2}} &\lesssim r \left(\int_{2B} |L(f)|^2 d\mu \right)^{\frac{1}{2}} + \frac{1}{r} \text{Osc}_{2B}(u) \\ &\lesssim r \left(\int_{2B} |L(f)|^2 d\mu \right)^{\frac{1}{2}} + \frac{1}{r} \text{Osc}_{2B}(f) + \frac{1}{r} \left(\int_{2B} |f - u|^2 d\mu \right)^{\frac{1}{2}} \\ &\lesssim r \left(\int_{2B} |L(f)|^2 d\mu \right)^{\frac{1}{2}} + \frac{1}{r} \text{Osc}_{2B}(f), \end{aligned}$$

where we used again (A.7) and (A.8) at the last step. \triangleright

We also give a proof of the following basic important fact about the Hölder spaces \mathcal{C}^σ defined in definition 2.7.

Proposition A.3. For $\sigma < 2$, the Hölder spaces \mathcal{C}^σ do not depend on the parameter a used to define them, and the two norms on \mathcal{C}^σ corresponding to two different parameters a, a' , are equivalent.

PROOF – Given two positive integers a and a' , consider the two spaces \mathcal{C}_a^σ and $\mathcal{C}_{a'}^\sigma$, and their corresponding norms. Fix $t \in (0, 1]$. If $a' \geq a$, then writing

$$Q_t^{(a')} = 2^{a'} Q_{t/2}^{(a')} e^{-tL/2} = 2^{a'} Q_{t/2}^{(a)} Q_{t/2}^{(a'-a)}$$

and using the fact that the operators $Q_{t/2}^{(a'-a)}$ are uniformly bounded on L^∞ , we get

$$\|\cdot\|_{\mathcal{C}_a^\sigma} \lesssim \|\cdot\|_{\mathcal{C}_{a'}^\sigma}.$$

If now $a' < a$, write

$$Q_t^{(a')} = \frac{1}{\gamma_{a-a'}} \int_0^1 Q_t^{(a')} Q_s^{(a-a')} \frac{ds}{s} + Q_t^{(a')} P_1^{(a-a')}.$$

For $s \leq t$, we have

$$Q_t^{(a')} Q_s^{(a-a')} = \left(\frac{s}{t}\right)^{a-a'} Q_{t+s}^{(a)} \left(\frac{t}{t+s}\right)^a$$

so that for $f \in \mathcal{C}_a^\sigma$

$$\begin{aligned} \left\| Q_t^{(a')} Q_s^{(a-a')} f \right\|_\infty &\lesssim \left(\frac{s}{t}\right)^{a-a'} \left\| Q_{t+s}^{(a)} f \right\|_\infty \\ &\lesssim \left(\frac{s}{t}\right)^{a-a'} t^{\frac{\sigma}{2}} \|f\|_{\mathcal{C}_a^\sigma}. \end{aligned}$$

For $t \leq s$, we have

$$Q_t^{(a')} Q_s^{(a-a')} = \left(\frac{s}{t}\right)^{a-a'} Q_{t+s}^{(a)} \left(\frac{t}{t+s}\right)^a$$

so that

$$\begin{aligned} \left\| Q_t^{(a')} Q_s^{(a-a')} f \right\|_\infty &\lesssim \left(\frac{t}{s} \right)^{a'} \left\| Q_{t+s}^{(a)} f \right\|_\infty \\ &\lesssim \left(\frac{t}{s} \right)^{a'} s^{\frac{\sigma}{2}} \|f\|_{C_a^\sigma}, \end{aligned}$$

and similarly

$$\left\| Q_t^{(a')} P_1^{(a-a')} f \right\|_\infty \lesssim t^{a'} \|f\|_{C_a^\sigma}.$$

Then by integrating (and since $a', a - a' \geq 1 > \frac{\sigma}{2}$) we have

$$\begin{aligned} \left\| Q_t^{(a')} f \right\|_\infty &\lesssim \left(\int_0^t \left(\frac{s}{t} \right)^{a-a'} \frac{ds}{s} \right) t^{\frac{\sigma}{2}} \|f\|_{C_a^\sigma} + \left(\int_t^1 \left(\frac{t}{s} \right)^{a'} s^{\frac{\sigma}{2}} \frac{ds}{s} \right) \|f\|_{C_a^\sigma} + t^{a'} \|f\|_{C_a^\sigma} \\ &\lesssim t^{\frac{\sigma}{2}} \|f\|_{C_a^\sigma}, \end{aligned}$$

which concludes the proof that

$$\| \cdot \|_{C_{a'}^\sigma} \lesssim \| \cdot \|_{C_a^\sigma}.$$

▷

The following lemma provides a useful way of proving that a distribution is Hölder; it was used in sections 2.3 and 5.3 to investigate the almost sure regularity properties of white noise and the renormalized paraproduct dealt with in theorem 5.7. We recall that Besov spaces were defined in Definition 2.6.

Lemma A.4. Assume that the metric measure space (M, d, μ) is Ahlfors regular (see (2.9)), with exponent ν . Then, given $-\infty < \sigma < 2$, and $1 < p < \infty$, we have the continuous embeddings

$$B_{p,p}^\sigma \hookrightarrow B_{p,\infty}^\sigma \hookrightarrow B_{\infty,\infty}^{\sigma - \frac{\nu}{p}} = C^{\sigma - \frac{\nu}{p}}.$$

PROOF – The first embedding is a direct application of the following fact. For $s \in (0, 1)$ and an integer $a \geq 2$ then

$$Q_s^{(a)} f = \frac{2}{s} \int_{s/2}^s Q_t^{(a)} \left(\frac{s}{t} \right)^a e^{-(s-t)L} f dt.$$

Since the semigroup is uniformly bounded on L^p , we get

$$\left\| Q_s^{(a)} f \right\|_p \lesssim \int_{s/2}^s \left\| Q_t^{(a)} f \right\|_p \frac{dt}{t}$$

and by Hölder inequality

$$\left\| Q_s^{(a)} f \right\|_p \lesssim \left(\int_{s/2}^s \left\| Q_t^{(a)} f \right\|_p^p \frac{dt}{t} \right)^{1/p} \lesssim s^{\frac{\sigma}{2}} \|f\|_{B_{p,p}^\sigma}.$$

The second embedding comes from the following elementary fact. For $t \in (0, 1)$, let T a linear operator with a kernel, pointwisely bounded by a Gaussian kernel \mathcal{G}_t at scale t , then with Ahlfors regularity (2.9), we have

$$\|T\|_{L^p \rightarrow L^\infty} \lesssim t^{-\frac{\nu}{2p}}.$$

So for $s \in (0, \frac{1}{2})$, applying to $T = Q_s^{(a)}$ we obtain since $Q_{2s}^{(2a)} = 2^{2a} Q_s^{(a)} Q_s^{(a)}$

$$\left\| Q_{2s}^{(2a)} f \right\|_\infty \lesssim s^{-\frac{\nu}{2p}} \left\| Q_s^{(a)} f \right\|_{L^p} \lesssim s^{-\frac{\nu}{2p} + \frac{\sigma}{2}} \|f\|_{B_{p,\infty}^\sigma},$$

which proves the embedding $B_{p,\infty}^\sigma \hookrightarrow B_{\infty,\infty}^{\sigma - \frac{\nu}{p}}$.

▷

The next three lemmas were used in the proof of the renormalization theorem 5.7.

Lemma A.5. For every $t > 0$, set \mathcal{G}_t the Gaussian kernel at scale t

$$\mathcal{G}_t(x, y) := \frac{1}{V(x, \sqrt{t})} e^{-c \frac{d(x, y)^2}{t}},$$

where we forget the dependence with respect to the constant c in the notation. Then for $s > t > 0$ and every $x, z \in M$, we have

$$\int \mathcal{G}_s(x, y) \mathcal{G}_t(y, z) \mu(dy) \lesssim \mathcal{G}_s(x, z) \lesssim \frac{1}{V(x, \sqrt{s}) + V(z, \sqrt{s})}.$$

PROOF – By considering that $\mathcal{G}_t(\cdot, z)$ belongs to L^1 , uniformly in s , we directly obtain that

$$\int \mathcal{G}_s(x, y) \mathcal{G}_t(y, z) \mu(dy) \lesssim \sup_{y \in M} \mathcal{G}_s(x, y) \lesssim \frac{1}{V(x, \sqrt{s})}.$$

Moreover,

$$\begin{aligned} \exp(-d(x, y)^2/s) \cdot \exp(-d(y, z)^2/t) &\leq \exp(-d(x, y)^2/s) \cdot \exp(-d(y, z)^2/s) \\ &\leq \exp(-d(x, z)^2/(2s)). \end{aligned}$$

So in the product $\mathcal{G}_s(x, y) \mathcal{G}_t(y, z)$, we may factorize an exponential decay and so for some implicit constants, we have

$$\int \mathcal{G}_s(x, y) \mathcal{G}_t(y, z) d\mu(y) \lesssim \mathcal{G}_s(x, z).$$

▷

Lemma A.6. For $r \in (0, 1]$, $s > 0$ and $d \geq 2$, let us consider the quantity

$$\begin{aligned} A(r, s) := &\int_0^1 \int_0^1 \left(\frac{r}{r+t_1} \right)^b \left(\frac{r}{r+t_2} \right)^b \dots \\ &\dots \left(\frac{t_1 t_2}{(t_1+s)(t_2+s)} \right)^{\frac{1}{2}} (s+t_1+t_2)^{-\frac{d}{2}} (r+t_1+t_2)^{-\frac{d}{2}} \frac{dt_1}{t_1} \frac{dt_2}{t_2}, \end{aligned}$$

where $b \geq 2 + \frac{d}{2}$ is an integer. Then we have

$$(A.9) \quad A(r, s) \lesssim \left(\frac{r}{s+r} \right) (rs)^{-\frac{d}{2}} \left(1 + \log \left(\frac{s+r}{s} \right) \right).$$

PROOF – The two variables t_1, t_2 play a symmetric role so we may restrict our attention to the double integral under the condition $t_2 \leq t_1$. The part A_1 of the double integral where

$t_2 \leq t_1 \leq r$ gives

$$\begin{aligned}
A_1(r, s) &= \int_0^r \int_0^{t_1} \left(\frac{t_1 t_2}{(t_1 + s)(t_2 + s)} \right)^{\frac{1}{2}} r^{-\frac{d}{2}} (t_1 + s)^{-\frac{d}{2}} \frac{dt_2}{t_2} \frac{dt_1}{t_1} \\
&\lesssim \int_0^r \left(\frac{t_1}{t_1 + s} \right)^{\frac{1}{2}} (t_1 + s)^{-\frac{d}{2}} r^{-\frac{d}{2}} \left(\frac{t_1}{t_1 + s} \right)^{\frac{1}{2}} \left(1 + \log \left(\frac{s + t_1}{s} \right) \right) \frac{dt_1}{t_1} \\
&\lesssim r^{-\frac{d}{2}} \int_0^r \left(\frac{t_1}{(t_1 + s)^{d/2+1}} \right) \left(1 + \log \left(\frac{s + t_1}{s} \right) \right) \frac{dt_1}{t_1} \\
&\lesssim r^{-\frac{d}{2}} \left\{ s^{-d/2} - \left(1 + \log \left(\frac{s + r}{s} \right) \right) (s + r)^{-\frac{d}{2}} \right\} \\
&\lesssim (sr)^{-\frac{d}{2}} \frac{r}{s + r} \left(1 + \log \left(\frac{s + r}{s} \right) \right),
\end{aligned}$$

where we used the basic inequality

$$\int_0^{t_1} \frac{dt_2}{\sqrt{t_2(t_2 + s)}} \lesssim \left(\frac{t_1}{s + t_1} \right)^{\frac{1}{2}} \left(1 + \log \left(\frac{s + t_1}{s} \right) \right)$$

which can be easily checked by splitting into the two cases $t_1 \leq s$ and $s \leq t_1$. The second part A_2 of the double integral where $t_2 \leq r \leq t_1$ is controlled as follows

$$\begin{aligned}
A_2(r, s) &= \int_r^1 \int_0^r \left(\frac{r}{r + t_1} \right)^b \left(\frac{t_1 t_2}{(t_1 + s)(t_2 + s)} \right)^{\frac{1}{2}} t_1^{-\frac{d}{2}} (t_1 + s)^{-\frac{d}{2}} \frac{dt_2}{t_2} \frac{dt_1}{t_1} \\
&\lesssim \int_r^1 \left(\frac{r}{t_1} \right)^b \left(\frac{t_1}{t_1 + s} \right)^{\frac{1}{2}} \left(\frac{r}{r + s} \right)^{\frac{1}{2}} \left(1 + \log \left(\frac{s + r}{s} \right) \right) (t_1 + s)^{-\frac{d}{2}} t_1^{-\frac{d}{2}} \frac{dt_1}{t_1} \\
&\lesssim \frac{r}{r + s} (r(r + s))^{-\frac{d}{2}} \left(1 + \log \left(\frac{s + r}{s} \right) \right),
\end{aligned}$$

where we used Lemma A.7. The third and last part A_3 of the double integral where $r \leq t_2 \leq t_1$ satisfies

$$\begin{aligned}
A_3(r, s) &= \int_r^1 \int_r^{t_1} \left(\frac{r}{t_1} \right)^b \left(\frac{r}{t_2} \right)^b \left(\frac{t_1 t_2}{(t_1 + s)(t_2 + s)} \right)^{\frac{1}{2}} t_1^{-\frac{d}{2}} (t_1 + s)^{-\frac{d}{2}} \frac{dt_2}{t_2} \frac{dt_1}{t_1} \\
&\lesssim \int_r^1 \left(\frac{r}{t_1} \right)^b \left(\frac{t_1}{t_1 + s} \right)^{\frac{1}{2}} \left(\frac{r}{r + s} \right)^{\frac{1}{2}} (t_1 + s)^{-\frac{d}{2}} t_1^{-\frac{d}{2}} \frac{dt_1}{t_1}.
\end{aligned}$$

We then use again Lemma A.7, to obtain

$$A_3(r, s) \lesssim \frac{r}{r + s} (rs)^{-\frac{d}{2}}.$$

Inequality (A.9) comes by combining the above three estimates. ▷

Lemma A.7. For every $r, t \in (0, 1)$ and any $0 < \rho < \varepsilon$

$$\int_r^1 \left(\frac{t}{t + s} \right)^\rho t^{-\varepsilon} \frac{dt}{t} \lesssim \left(\frac{r}{s + r} \right)^\rho r^{-\varepsilon}.$$

PROOF – Indeed if $r \geq s$ then for every $t \in (r, 1)$ we have $t \simeq t + s$ and so

$$\int_r^1 \left(\frac{t}{t + s} \right)^\rho t^{-\varepsilon} \frac{dt}{t} \lesssim \int_r^1 t^{-\varepsilon} \frac{dt}{t} \simeq r^{-\varepsilon}.$$

Now if $s \geq r$, we split the integral in two terms and we have

$$\begin{aligned} \int_r^1 \left(\frac{t}{t+s} \right)^\rho t^{-\varepsilon} \frac{dt}{t} &\lesssim \int_r^s \left(\frac{t}{s} \right)^\rho t^{-\varepsilon} \frac{dt}{t} + \int_s^1 t^{-\varepsilon} \frac{dt}{t} \\ &\lesssim \left(\frac{r}{s} \right)^\rho r^{-\varepsilon} + s^{-\varepsilon} \lesssim \left(\frac{r}{s} \right)^\rho r^{-\varepsilon}, \end{aligned}$$

where we used $\varepsilon > A\rho$.

▷

B

Extension of the theory

Consider as above a doubling metric measure space (M, d, μ) equipped with a heat semigroup satisfying the upper gaussian estimates (UE). We aim in this appendix at explaining how one can get the same conclusions as in the above main body of work

- (a) by weakening the Lipschitz regularity assumption on the heat kernel (Lip), assuming only some integrated estimates of the gradient of the heat kernel;
- (b) by developing the theory of paracontrolled calculus in Sobolev spaces rather than in Hölder spaces. By Sobolev embedding, Sobolev spaces are included in some Hölder spaces, so it will be interesting to understand if starting from an initial data belonging to some Sobolev space, the solution of renormalized singular PDEs will lives in this same scale of Sobolev spaces. From a technical point of view, it is a bit more difficult since Sobolev spaces involve simultaneously all the frequencies, whereas for Hölder spaces we can work at a fixed frequency scale.

We give in section B.1 the regularity assumptions on the heat kernel under which we shall work here, and reformulate and extend in section B.2 the main continuity estimates on the operators $P_t^{(a)}$, $Q_t^{(a)}$ and Γ needed to extend the paraproduct machinery to the present setting. The latter, together with some crucial commutator estimates in Hölder and Sobolev spaces, is investigated in section B.3. The last and short section B.4 describes how these results can be used to extend the results of section 5 to our optimal regularity setting.

This appendix was written jointly with Dorothee Frey.

B.1. Regularity assumptions Rather than assuming the Lipschitz property (Lip) used above we shall assume here that the gradient / carré du champ operator Γ satisfies some L^q estimates and the L^q -de Giorgi property recalled below in sections B.1.1 and B.1.2. We shall also assume that it satisfies a scale-invariant Poincaré inequality recalled in section B.1.3.

B.1.1. L^q -estimates of the gradient of the semigroup Given $q_0 > 2$, the following uniform L^{q_0} -boundedness of the gradient (or “carré du champ”) of the semi-group, was introduced in [2]

$$(G_{p_0}) \quad \sup_{t>0} \|\sqrt{t}\Gamma e^{-tL}\|_{q_0 \rightarrow q_0} < +\infty.$$

By definition of the carré du champ operator, (G_2) holds trivially. It is known in that case that this global L^2 -inequality can be improved into localized estimates, via L^2 -Davies-Gaffney estimates. For every subset $E, F \subset M$ and every $t > 0$, we have

$$\|e^{-tL}\|_{L^2(E) \rightarrow L^2(F)} + \sqrt{t} \|\Gamma e^{-tL}\|_{L^2(E) \rightarrow L^2(F)} \lesssim e^{-c \frac{d^2(E,F)}{t}}.$$

Assuming the volume doubling condition (VD) and the Gaussian upper bound (UE) for the heat kernel, one can interpolate the estimate (G_{q_0}) with the above L^2 -Davies-Gaffney estimates and deduce that (G_q) holds for every $q \in [2, q_0]$. More precisely, for every subset $E, F \subset M$ and every $t > 0$, the inequality

$$\|e^{-tL}\|_{L^q(E) \rightarrow L^q(F)} + \sqrt{t} \|\Gamma e^{-tL}\|_{L^q(E) \rightarrow L^q(F)} \lesssim e^{-c_q \frac{d^2(E,F)}{t}}$$

holds for some positive constant c_q , only depending on $q \in [2, q_0]$. Following [2, Proposition 1.10], the latter estimate can be reformulated in terms of integral estimates of the gradient of the heat kernel. Denoting by p_t the kernel of e^{-tL} , we have

$$\sqrt{t} \|\Gamma_x p_t(\cdot, y)\|_q \lesssim [V(y, \sqrt{t})]^{-(1-\frac{1}{q})},$$

for μ -almost all $y \in M$ and all positive times. By interpolation with the L^2 -Gaffney estimates, there exists a positive constant c such that

$$(B.1) \quad \sqrt{t} \left\| e^{c \frac{d(\cdot, y)^2}{t}} \Gamma_x p_t(\cdot, y) \right\|_q \lesssim [V(y, \sqrt{t})]^{-(1-\frac{1}{q})}$$

holds for μ -almost all $y \in M$ and all positive times.

We refer the reader to [2] for more details about Property (G_p) and the link with the boundedness of the Riesz transform; see also [11] and references therein for more details.

B.1.2. L^q -de Giorgi Property The so-called "de Giorgi property", or "Dirichlet property", on the growth of the Dirichlet integral for harmonic functions was introduced by De Giorgi in [19], for second order divergence form differential operators on \mathbb{R}^n , with real coefficients. In de Giorgi's work, this property prescribes a(n at most) linear growth rate for the L^2 -average of gradients of harmonic functions. This property was subsequently used in many works and in various situations in order to prove Hölder regularity for solutions of inhomogeneous elliptic equations and systems. An L^q -version was recently introduced in [9], and we refer the reader to that work for more details about it.

Definition B.1 (L^q -de Giorgi property). Given $q \in [1, +\infty)$ and $\theta \in (0, 1)$, we say that the operator Γ satisfies the inequality $(DG_{q,\theta})$ if it satisfies the following estimate. For every positive $r \leq R$, every pair of concentric balls B_r, B_R with radii r and R , respectively, and for every function $f \in \mathcal{D}$, one has

$$(DG_{q,\theta}) \quad \left(\int_{B_r} |\Gamma f|^q d\mu \right)^{\frac{1}{q}} \lesssim \left(\frac{R}{r} \right)^\theta \left\{ \left(\int_{B_R} |\Gamma f|^q d\mu \right)^{\frac{1}{q}} + R \|Lf\|_{L^\infty(B_R)} \right\}.$$

We sometimes omit the parameter θ , and write (DG_q) if $(DG_{q,\theta})$ is satisfied for some $\theta \in (0, 1)$.

As we always have

$$\left(\int_{B_r} |\Gamma f|^q d\mu \right)^{\frac{1}{q}} \lesssim \left(\frac{|B_R|}{|B_r|} \right)^{\frac{1}{q}} \left(\int_{B_R} |\Gamma f|^q d\mu \right)^{\frac{1}{q}}$$

for every $f \in \mathcal{D}$ and $0 < r < R$, if the space is doubling, with dimension ν , the inequality $(DG_{q,\theta})$ holds for every $q > \nu$, with $\theta = \frac{\nu}{q} < 1$.

B.1.3. Poincaré inequality Last, we shall assume that the carré du champ operator Γ satisfies the following scale-invariant Poincaré inequality

$$(P_2) \quad \left(\int_B \left| f - \int_B f d\mu \right|^2 d\mu \right)^{\frac{1}{2}} \lesssim r \left(\int_B \Gamma(f)^2 d\mu \right)^{\frac{1}{2}},$$

for every $f \in \mathcal{D}_2(L)$ and every ball B of radius r . We refer the reader to [9] for a precise study of the connection between Poincaré inequality, L^p -gradient estimates and de Giorgi property. Let us just point out that if Γ satisfies the above Poincaré inequality and the gradient estimate (G_{p_0}) , then there exists a parameter $\theta \in (0, 1)$ such that the inequality $(DG_{p,\theta})$ holds for every $p \in [2, p_0]$. Note also that in the first and main part of this work, we assumed an upper Gaussian pointwise estimates for the gradient of the heat kernel equivalent to (G_∞) . This assumption yields the Poincaré inequality (P_2) , the integrated gradient estimate (G_{q_0}) for every $q_0 \in [2, \infty]$, and also de Giorgi property $(DG_{q,\theta})$ for every $\theta \in (0, 1)$ and every $q \in [2, \infty)$.

Our aim in this appendix will thus be to weaken the (G_∞) assumption made above into a combination of (G_{q_0}) and $(DG_{q,\theta})$, for some exponent q_0, q and θ .

In the first and main part of this work, the paracontrolled calculus and its application to the 2-dimensional parabolic Anderson model equation was studied under the assumptions (UE) and (G_∞) that the heat kernel and its gradient satisfy pointwise Gaussian upper bounds. The aim of this section is to weaken the latter condition. Here are **examples** where the operator Γ satisfies only the properties (P_2) , (G_q) and (DG_q) for some $q > 2$, and where (G_∞) does not hold.

- (a) **Conical manifolds.** Consider a compact Riemannian manifold N of dimension $n-1 \geq 1$, and define $M := (0, \infty) \times N$ as the conical manifold whose basis is N . It is known that M is a doubling manifold of dimension n which satisfies (UE). Moreover, as shown by Li in [44], the operator Γ satisfies (G_q) if and only if

$$q < q(N) := \left(\frac{1}{2} - \sqrt{\left(\frac{1}{2} - \frac{2}{n} \right)^2 + \frac{\lambda_1}{n}} \right)^{-1}$$

where λ_1 is the first non-vanishing eigenvalue of the Laplace operator on N . As an example, if we consider $N = r\mathbb{S}^1$ the circle of radius $r > 1$, then

$$q(N) = \frac{2r}{r-1}.$$

So theorem B.20 below allows us to solve the PAM equation on M for r sufficiently close to 1.

- (b) **Elliptic perturbation of the Laplacien.** On the Euclidean space \mathbb{R}^d , or any non-compact doubling Riemannian manifold satisfying Poincaré inequality (P_2) and the Gaussian bound (UE), we may consider a second order divergence form operator $L = -\operatorname{div}(A\nabla)$ given by a map A taking values in real symmetric matrices and satisfying the usual ellipticity condition. Then if A is Hölder continuous, it is known that $-L$ generates a self-adjoint semigroup with (UE) and Gaussian pointwise bounds for the gradient of the semigroup (G_∞) ; see [4]. In such a case we may apply the results proved in the first part. Following Auscher's work [1], we know that the combination of property (UE) with Hölder regularity of the heat kernel is stable under L^∞ perturbation. So fix A_0 a Hölder continuous map with values in real symmetric matrices and satisfying usual ellipticity condition. Then for every $Q > 2$, and any positive Θ , there exists a positive constant ε such that for any map A on the state space, with values in the space of real symmetric matrices, and such that $\|A - A_0\|_\infty \leq \varepsilon$, the operator $L = -\operatorname{div}(A\nabla)$

satisfies (G_{q_0}) for some $q_0 > Q$, and has de Giorgi property $(DG_{q,\theta})$, for $\theta = d/q < \Theta$ and some $Q \leq q < q_0$. In such a situation, we may apply Theorem B.20 and deduce that we can solve the PAM equation in such a 2-dimensional context.

- (c) **Lipschitz domain with Neumann boundary conditions.** Similarly, consider an open and bounded subset $\Omega \subset \mathbb{R}^2$ and consider for L the self-adjoint Laplace operator associated with Neumann boundary conditions. Then by a change of variable, this situation is very similar to the previous one: if the boundary is sufficiently close (in a Lipschitz sense) to a smooth set (at least of regularity C^2), then we can solve the PAM equation.

B.2. Functional calculus and gradient estimates in Hölder and Sobolev spaces We start this section by quantifying the localization properties of the operators $P_t^{(a)}$ and $Q_t^{(a)}$, and their gradients, in L^p spaces, before turning to the gradient estimates of the heat semigroup in the intrinsic Hölder and Sobolev spaces in section B.2.2.

B.2.1. Localization properties of the approximation operators $P_t^{(a)}$ and $Q_t^{(a)}$ As we know, for every integer $a \geq 0$, the operators $P_t^{(a)}$ and $Q_t^{(a)}$ have a kernel satisfying Gaussian estimates (UE). The above regularity assumptions $(G_q), (DG_q), (P_2)$ on the gradient operator actually imply much more.

Lemma B.2. Let p_t stands for the kernel of e^{-tL} or $P_t^{(a)}, Q_t^{(a)}$ for any integer $a \geq 1$. Under $(DG_{q,\theta})$ and (G_{q_0}) with Poincaré inequality (P_2) for some $2 \leq q < q_0$, we have the following Hölder regularity estimate for the heat kernel. For every $\eta \in (0, 1 - \theta]$, $t > 0$ and almost every $x, y, z \in M$

$$|p_t(x, z) - p_t(y, z)| \lesssim \left(\frac{d(x, y)}{\sqrt{t}} \right)^\eta V(z, \sqrt{t})^{-1} e^{-c \frac{d(x, z)^2}{t}}.$$

We only sketch the proof and refer the reader to [9] for details.

PROOF – We follow the argument of Morrey's inequality, which relies oscillation estimates to some gradient bounds. Let $x, y \in M$ be Lebesgue points for $f = p_t(\cdot, z)$ with $d(x, y) \leq \sqrt{t}$, otherwise there is nothing to be done. Let $B_i(x) = B(x, 2^{-i}d(x, y))$, for $i \in \mathbb{N}$. Note that for all $i \in \mathbb{N}$, $B_i(x) \subset B_0(x)$. By Poincaré's inequality, this yields

$$\left| f(x) - \int_{B_0(x)} f d\mu \right| \lesssim \sum_{i \geq 0} 2^{-i} d(x, y) \left(\int_{B_i(x)} |\Gamma f|^q d\mu \right)^{\frac{1}{q}}.$$

By considering $B_{\sqrt{t}}$ a ball of radius \sqrt{t} containing both x, y , $(DG_{q,\theta})$ yields

$$\left(\int_{B_i(x)} |\Gamma f|^q d\mu \right)^{\frac{1}{q}} \lesssim \left(\frac{\sqrt{t}}{2^{-i}d(x, y)} \right)^\theta \left[\left(\int_{B_{\sqrt{t}}} |\Gamma f|^q d\mu \right)^{\frac{1}{q}} + \sqrt{t} \|Lf\|_{L^\infty(B_{\sqrt{t}})} \right].$$

Since $f = p_t(\cdot, z)$, by (G_{q_0}) and (UE) we know that

$$\left(\int_{B_{\sqrt{t}}} |\sqrt{t}\Gamma f|^q d\mu \right)^{\frac{1}{q}} + \|tLf\|_{L^\infty(B_{\sqrt{t}})} \lesssim V(z, \sqrt{t})^{-1} e^{-c \frac{d(x, z)^2}{t}}$$

so we can conclude the proof by summing over i , since $\theta \in (0, 1)$. ▷

Under the sole assumption (UE) that the kernels of the operators $P_t^{(a)}$ and $Q_t^{(a)}$ have Gaussian upper bounds, these operators are bounded in every L^p space for $p \in [1, \infty]$, uniformly with respect to $t \in (0, 1]$. Moreover, for every $p_1, p_2 \in [1, \infty]$ and $t > 0$, they satisfy the following L^{p_1} - L^{p_2} off-diagonal estimates at scale \sqrt{t} , which quantify the localization properties of these operators. For every ball B_1, B_2 of radius \sqrt{t} , and for every function $f \in L^{p_1}(B_1)$, we have

$$\left(\int_{B_2} |P_t^{(a)} f|^{p_2} d\mu \right)^{1/p_2} + \left(\int_{B_2} |Q_t^{(a)} f|^{p_2} d\mu \right)^{1/p_2} \lesssim e^{-c \frac{d(B_1, B_2)^2}{t}} \left(\int_{B_1} |f|^{p_1} d\mu \right)^{1/p_1}.$$

One can refine this estimate by using off-diagonal estimates, such as done in [10, Lemma 2.5, Lemma 2.6].

Proposition B.3. Assume (G_{q_0}) for some $q_0 > 2$.

- (i) For every non-negative integer a and every $p \in [2, q_0)$, the operators $\Gamma P_t^{(a)}$ and $\Gamma Q_t^{(a)}$ satisfy L^2 - L^p the following off-diagonal estimates at the scale \sqrt{t} . For every ball B_1, B_2 of radius \sqrt{t} and every function $f \in L^2(B_1)$, we have

$$\left(\int_{B_2} |\sqrt{t} \Gamma Q_t^{(a)} f|^p d\mu \right)^{\frac{1}{p}} + \left(\int_{B_2} |\sqrt{t} \Gamma P_t^{(a)} f|^p d\mu \right)^{\frac{1}{p}} \lesssim e^{-c \frac{d(B_1, B_2)^2}{t}} \left(\int_{B_1} |f|^2 d\mu \right)^{\frac{1}{2}}.$$

It follows in that we have

$$\sup_{t>0} \left\{ \left\| (\sqrt{t} \Gamma)(P_t^{(a)} \cdot) \right\|_{p \rightarrow p} + \left\| (\sqrt{t} \Gamma)(Q_t^{(a)} \cdot) \right\|_{p \rightarrow p} \right\} < \infty$$

for every $p \in [2, q_0)$.

- (ii) For every positive real number a and every positive t , the operator $Q_t^{(a)}$ is an integral operator with kernel $k_t^{(a)}$ that satisfies the inequality

$$(B.2) \quad \left| k_t^{(a)}(x, y) \right| \lesssim \frac{1}{V(x, \sqrt{t})^\lambda V(y, \sqrt{t})^{1-\lambda}} \left(1 + \frac{d^2(x, y)}{t} \right)^{-a}$$

for all $\lambda \in [0, 1]$ and μ -almost all $x, y \in M$. As a consequence the operator $Q_t^{(a)}$ satisfies the following L^{p_1} - L^{p_2} off-diagonal bounds of order a at scale \sqrt{t} , for every $p_1, p_2 \in [1, +\infty]$. Given any balls B_1, B_2 of radius \sqrt{t} , and any function $f \in L^{p_1}(B_1)$, we have

$$\left(\int_{B_2} |Q_t^{(a)} f|^{p_2} d\mu \right)^{\frac{1}{p_2}} \lesssim \left(1 + \frac{d(B_1, B_2)^2}{t} \right)^{-a} \left(\int_{B_1} |f|^{p_1} d\mu \right)^{\frac{1}{p_1}}.$$

Besides these localization property in the physical space, the approximation operators $Q_t^{(a)}$ satisfy some orthogonality properties, which will be of crucial use in proving the continuity properties of the paraproduct and resonant operators below, and which can be viewed as an analog of the Littlewood-Paley theory, as made clear in [10, Proposition 2.13 and Lemma 2.15].

Lemma B.4. Let a be a positive real number. Set

$$\tilde{Q}_t := (t\mathcal{L})^{\frac{a}{2}} e^{-\frac{t}{2}\mathcal{L}} = 2^{\frac{a}{2}} Q_{\frac{t}{2}}^{\frac{a}{2}},$$

so that $Q_t^{(a)} = \tilde{Q}_t^2$. Assume the Gaussian upper bound (UE) holds. Let also $F : (0, +\infty) \times M \rightarrow \mathbb{R}$ be a measurable function and write $F_t(x)$ for $F(t, x)$. Then for every $p \in (1, +\infty)$, one has

$$\left\| \int_0^{+\infty} Q_t^{(a)} F_t \frac{dt}{t} \right\|_p \lesssim \left\| \left(\int_0^{+\infty} |\tilde{Q}_t F_t|^2 \frac{dt}{t} \right)^{1/2} \right\|_p,$$

whenever the right hand side has a meaning and is finite. If $F = f$ does not depend on t , we have the following L^p -boundedness of the vertical square function

$$\left\| \int_0^{+\infty} |Q_t^{(a)} f| \frac{dt}{t} \right\|_p \simeq \|f\|_p.$$

B.2.2. Gradient estimates in Hölder and Sobolev spaces As said above, we shall now work in the following setting, strictly weaker than the geometrical setting used in the first five sections of this work.

Regularity assumptions

- (i) The metric measure space (M, d, μ) is doubling and the semigroup satisfies the Gaussian bound (UE).
- (ii) The gradient operator Γ satisfies (G_{q_0}) and $(DG_{q,\theta})$ for some $2 \leq q < q_0 \leq \infty$, and the scale-invariant Poincaré inequality.

If $q_0 = q = 2$, we require that the L^2 Davies-Gaffney estimates hold instead of $(DG_{q,\theta})$. As we shall see below, one can extend the machinery of paracontrolled calculus to that setting in Hölder and Sobolev spaces. Recall the definition of the spaces Λ^σ and \mathcal{C}^σ given in section 2.3. The parameter θ is involved in the property $(DG_{q,\theta})$. The following embedding is proved as Proposition 2.8 by using the fact proved in Lemma B.2 that the heat kernel is Hölder continuous, with exponent $1 - \theta$, instead of its Lipschitz character.

Proposition B.5. For $\sigma \in (0, 1)$, the space Λ^σ is continuously embedded into \mathcal{C}^σ . If $\sigma \in (0, 1 - \theta)$, the two spaces are the same with equivalent norms.

Sobolev spaces are naturally defined in terms of L as follows.

Definition B.6. Fix an exponent $p \in (1, \infty)$, and $s \in \mathbb{R}$. A distribution $f \in \mathcal{S}'_o$, is said to belong to the inhomogeneous **Sobolev space** $W^{s,p}$ if

$$\|f\|_{W^{s,p}} := \left\| (1 + L)^{\frac{s}{2}} f \right\|_p \simeq \|e^{-L} f\|_p + \left\| (1 + L)^{\frac{s}{2}} f \right\|_p < \infty.$$

Proposition B.7. For $\sigma \in (-\infty, 1 - \theta)$ and $f \in \mathcal{C}^\sigma$, we have

$$\sup_{x \in M} \left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} \lesssim t^{\frac{\sigma}{2}} \|f\|_{\mathcal{C}^\sigma}.$$

The same conclusion holds with any of the operators $P_t^{(a)}$, with an integer $a \geq 1$, in the role of e^{-tL} .

PROOF – Consider $b \geq 1$, and write

$$\sqrt{t} \Gamma e^{-tL} f = \int_0^1 \sqrt{t} \Gamma Q_s^{(b)} e^{-tL} f \frac{ds}{s} + \sqrt{t} \Gamma P_1^{(b)} e^{-tL} f.$$

For $s \leq t$, we have

$$\begin{aligned} \left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma Q_s^{(b)} e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} &= \left(\frac{s}{s+t} \right)^b \left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma Q_{s+t}^{(b)} f|^q d\mu \right)^{\frac{1}{q}} \\ &\lesssim \left(\frac{s}{t} \right)^b \sum_{\ell \geq 0} \gamma_\ell \left(\int_{2^\ell B(x, \sqrt{t})} |Q_{(s+t)/2}^{(b)} f|^q d\mu \right)^{\frac{1}{q}}, \end{aligned}$$

where γ_ℓ are exponentially decreasing coefficients and where we used L^q - L^q off-diagonal estimates of $\Gamma e^{-(s+t)L/2}$ (at the scale $\sqrt{s+t} \simeq \sqrt{t}$) with the relation

$$Q_{s+t}^{(b)} = 2^b e^{-(s+t)L/2} Q_{(s+t)/2}^{(b)}.$$

So we have

$$\begin{aligned} \left(\int_{B(x, \sqrt{t})} |\sqrt{t} L^{1/2} Q_s^{(b)} e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} &\lesssim \left(\frac{s}{t} \right)^b \sum_{\ell \geq 0} \gamma_\ell \|Q_{(s+t)/2}^{(b)} f\|_\infty \\ &\lesssim \left(\frac{s}{t} \right)^b t^{\frac{\sigma}{2}} \|f\|_{\mathcal{C}^\sigma}, \end{aligned}$$

and we can integrate this inequality on the interval $s \in (0, t)$. For $s \geq t$, we use Property $(DG_{q, \theta})$ to have

$$\begin{aligned} \left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma Q_s^{(b)} e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} &\lesssim \left(\frac{t}{s} \right)^{(1-\theta)/2} \left(\int_{B(x, \sqrt{s})} |\sqrt{s} \Gamma Q_s^{(b)} e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} \\ &\quad + \left(\frac{t}{s} \right)^{(1-\theta)/2} \left\| Q_s^{(b+1)} e^{-tL} f \right\|_{L^\infty(B(x, \sqrt{s}))} \\ &\lesssim \left(\frac{t}{s} \right)^{(1-\theta)/2} s^{\frac{\sigma}{2}} \|f\|_{\mathcal{C}^\sigma}, \end{aligned}$$

where we have used $Q_s^{(b)} = 2^b Q_{s/2}^{(b/2)} Q_{s/2}^{(b/2)}$ with L^q - L^q (resp. L^q - L^∞) off-diagonal estimates for $\Gamma Q_{s/2}^{(b/2)}$ (resp. $(sL)Q_{s/2}^{(b/2)}$), provided b is large enough. This inequality can be then integrated along $s \in (t, 1)$ as soon as $\theta + \sigma < 1$.

We perform the same analysis for the term $\sqrt{t} \Gamma P_1^{(b)} e^{-tL} f$, which gives

$$\begin{aligned} \left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma P_1^{(b)} e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} &\lesssim t^{(1-\theta)/2} \left(\int_{B(x, 1)} |\Gamma P_1^{(b)} e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} \\ &\quad + t^{(1-\theta)/2} \left\| L P_1^{(b)} e^{-tL} f \right\|_{L^\infty(B(x, 1))} \\ &\lesssim t^{(1-\theta)/2} \|f\|_{\mathcal{C}^\sigma}. \end{aligned}$$

The conclusion follows from this inequality since $t \in (0, 1)$ and $\sigma < 1 - \theta$. ▷

Proposition B.8. For $\alpha \in (0, 1 - \theta)$ and $0 < 2\delta < 1 - \theta - \alpha$, we have uniformly in $x \in M$ and $t > 0$

$$\left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} \lesssim \mathcal{M} \left[(tL)^{\alpha/2} Q_{t/2}^{(\delta)} f \right],$$

where \mathcal{M} is the Hardy-Littlewood maximal function. The same conclusion holds with any of the operators $P_t^{(a)}$, with an integer $a \geq 1$, in the role of e^{-tL} and also by replacing $B(x, \sqrt{t})$ by any bigger ball $B(x, K\sqrt{t})$ and the estimates are uniform with respect to $K \geq 1$.

PROOF – We write (for a chosen large enough)

$$\sqrt{t} \Gamma e^{-tL} = \sqrt{t} \Gamma L^{-\alpha/2} e^{-tL} L^{\alpha/2} f = \sqrt{t} \int_0^\infty \Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f \frac{ds}{s^{1-\alpha/2}}.$$

For $s < t$, we then write

$$Q_s^{(a)} e^{-tL} = \left(\frac{s}{t}\right)^a Q_t^{(a)} e^{-sL} = \left(\frac{2s}{t}\right)^a Q_{t/2}^{(a-\delta)} e^{-sL} Q_{t/2}^{(\delta)}$$

and using L^1 - L^q off-diagonal estimates of the carré du champ of the semigroup, this yields

$$\left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f|^q d\mu \right)^{\frac{1}{q}} \lesssim \left(\frac{s}{t}\right)^a \mathcal{M} \left[L^{\alpha/2} Q_{t/2}^{(\delta)} f \right](x).$$

For $t \leq s$ we have by $(DG_{q, \theta})$

$$\begin{aligned} \left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f|^q d\mu \right)^{\frac{1}{q}} &\lesssim \left(\frac{s}{t}\right)^{(\theta-1)/2} \left(\int_{B(x, \sqrt{s})} |\sqrt{s} \Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f|^q d\mu \right)^{\frac{1}{q}} \\ &\quad + \left(\frac{s}{t}\right)^{(\theta-1)/2} \left\| Q_s^{(a+1)} e^{-tL} L^{\alpha/2} f \right\|_{L^\infty(B(x, \sqrt{s}))} \\ &\lesssim \left(\frac{s}{t}\right)^{(\theta-1)/2+\delta} \mathcal{M} \left[L^{\alpha/2} Q_t^{(\delta)} f \right](x), \end{aligned}$$

where we used that $Q_s^{(a)} e^{-tL} = (s/t)^\delta Q_s^{(a-\delta)} Q_t^{(\delta)}$ with L^1 - L^q (resp. L^1 - L^∞) off-diagonal estimates for $\Gamma Q_s^{(a-\delta)}$ (resp. $Q_s^{(a+1-\delta)}$), provided a is large enough.

Hence,

$$\begin{aligned} \left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} &\lesssim \left[\int_0^t \left(\frac{s}{t}\right)^a \frac{ds}{s^{1-\alpha/2}} + \int_t^1 \left(\frac{s}{t}\right)^{(\theta-1)/2+\delta} \frac{ds}{s^{1-\alpha/2}} \right] \mathcal{M} \left[L^{\alpha/2} Q_t^{(\delta)} f \right](x) \\ &\lesssim \mathcal{M} \left[(tL)^{\alpha/2} Q_t^{(\delta)} f \right](x), \end{aligned}$$

due to a large enough and $2\delta < 1 - \theta - \alpha$. We let the reader check the straightforward modifications that are required to deal with a bigger ball $B(x, K\sqrt{t})$, and that the estimates are uniform with respect to $K \geq 1$. ▷

Replacing the L^1 - L^q off-diagonal estimates by L^p - L^q estimates, the same proof as above leads to the following result.

Proposition B.9. Assume the local Ahlfors regularity. Suppose $p \in [1, \infty)$ and $\alpha \in (-\infty, 1 - \theta + \frac{\nu}{p})$, and $f \in W^{\alpha, p}$. Then, uniformly in $x \in M$ and $t \in (0, 1]$,

$$\left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} \lesssim t^{-\frac{\nu}{2p} + \frac{\alpha}{2}} \|f\|_{W^{\alpha, p}}.$$

The same conclusion holds with any of the operators $P_t^{(a)}$, with an integer $a \geq 1$, in the role of e^{-tL} .

PROOF – As previously, we write

$$\sqrt{t} \Gamma e^{-tL} = \sqrt{t} \int_0^1 \Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f \frac{ds}{s^{1-\alpha/2}} + \sqrt{t} \Gamma P_1^{(a)} e^{-tL} f,$$

for a a large enough integer. For $s < t$, we then have

$$\left(\int_{B(x, \sqrt{t})} |\sqrt{t} \Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f|^q d\mu \right)^{\frac{1}{q}} \lesssim \left(\frac{s}{t}\right)^a t^{-\nu/(2p)} \|f\|_{W^{\alpha, p}}$$

and for $t \leq s$ we have by $(DG_{q,\theta})$

$$\begin{aligned} \left(\int_{B(x,\sqrt{t})} |\sqrt{t}\Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f|^q d\mu \right)^{\frac{1}{q}} &\lesssim \left(\frac{s}{t} \right)^{(\theta-1)/2} \left(\int_{B(x,\sqrt{s})} |\sqrt{s}\Gamma Q_s^{(a)} e^{-tL} L^{\alpha/2} f|^q d\mu \right)^{\frac{1}{q}} \\ &\quad + \left(\frac{s}{t} \right)^{(\theta-1)/2} \left\| Q_s^{(a+1)} e^{-tL} L^{\alpha/2} f \right\|_{L^\infty(B(x,\sqrt{s}))} \\ &\lesssim \left(\frac{s}{t} \right)^{(\theta-1)/2} s^{-\nu/(2p)} \|f\|_{W^{\alpha,p}}. \end{aligned}$$

For the low frequency part, we have

$$\left(\int_{B(x,\sqrt{t})} |\sqrt{t}\Gamma P_1^{(a)} e^{-tL} f|^q d\mu \right)^{\frac{1}{q}} \lesssim t^{(1-\theta)/2} \|f\|_{W^{\alpha,p}}.$$

Hence,

$$\begin{aligned} &\left(\int_{B(x,\sqrt{t})} |\sqrt{t}\Gamma e^{-tL} f|^p d\mu \right)^{1/p} \\ &\lesssim \left[\int_0^t \left(\frac{s}{t} \right)^a \frac{ds}{s^{1-\alpha/2}} + \int_t^1 \left(\frac{s}{t} \right)^{(\theta-1)/2-\nu/(2p)} \frac{ds}{s^{1-\alpha/2}} + t^{(1-\theta)/2} \right] t^{-\nu/(2p)} \|f\|_{W^{\alpha,p}} \\ &\lesssim t^{-\frac{\nu}{2p} + \frac{\alpha}{2}} \|f\|_{W^{\alpha,p}}, \end{aligned}$$

due to a large enough and $\alpha < 1 - \theta + \frac{\nu}{p}$. ▷

B.3. Paraproduct and commutator estimates in Hölder-Sobolev spaces

This subsection is devoted to the statement / proofs of the main estimates about Paraproducts and commutators, in the current more general framework.

B.3.1. Paraproduct estimates

We state in this paragraph the basic continuity estimates satisfied by the maps defined by the low frequency part, the paraproduct and the resonant terms – see Subsection 3.1 for the precise definition of these quantities and for detailed proofs. The low-frequency part is easily bounded.

Proposition B.10. Fix an integer $b \geq 2$. For any $\alpha, \beta \in \mathbb{R}$ and every $\gamma > 0$ we have for every $f \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$

$$(B.3) \quad \|\Delta_{-1}(f, g)\|_{\mathcal{C}^\gamma} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}.$$

If the space (M, d, μ) is locally Alhfors regular, then for every $\alpha, \beta, \gamma \in \mathbb{R}$ and $p \in [1, \infty)$, we have for every $f \in W^{\alpha,p}$ and $g \in W^{\beta,p}$

$$(B.4) \quad \|\Delta_{-1}(f, g)\|_{W^{\gamma,p}} \lesssim \|f\|_{W^{\alpha,p}} \|g\|_{W^{\beta,p}}.$$

The continuity properties of the paraproduct are given by the following statement.

Proposition B.11. Fix an integer $b \geq 2$. For any $\alpha \in (-2, 1)$ and $f \in \mathcal{C}^\alpha$, we have

- for every $g \in L^\infty$

$$(B.5) \quad \left\| \Pi_g^{(b)}(f) \right\|_{\mathcal{C}^\alpha} \lesssim \|g\|_\infty \|f\|_{\mathcal{C}^\alpha}$$

- for every $g \in \mathcal{C}^\beta$ with $\beta < 0$ and $\alpha + \beta \in (-2, 1)$

$$(B.6) \quad \left\| \Pi_g^{(b)}(f) \right\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|g\|_{\mathcal{C}^\beta} \|f\|_{\mathcal{C}^\alpha}.$$

The proof is already given for Proposition 3.3 – and only relies on (UE) (which is also assumed here). We then state the analog in Sobolev spaces.

Proposition B.12. Assume local Ahlfors regularity. Fix an integer $b \geq 2$ and $p \in [1, \infty)$. For any $\alpha \in (-2, 1)$ and $f \in W^{\alpha, p}$, we have

- for every $g \in W^{\beta, p}$ with $\frac{\nu}{p} < \beta < 1$

$$(B.7) \quad \left\| \Pi_g^{(b)}(f) \right\|_{W^{\alpha, p}} \lesssim \|g\|_{W^{\beta, p}} \|f\|_{W^{\alpha, p}}$$

- for every $g \in W^{\beta, p}$ with $\beta < \frac{\nu}{p}$ and $\alpha + \beta - \frac{\nu}{p} \in (-2, 1)$

$$(B.8) \quad \left\| \Pi_g^{(b)}(f) \right\|_{W^{\alpha + \beta - \frac{\nu}{p}, p}} \lesssim \|g\|_{W^{\beta, p}} \|f\|_{W^{\alpha, p}}.$$

Even if the proof is not very difficult, we give the details here in order to explain how to use the L^p -orthogonality property put forward in Lemma B.4.

PROOF – Recall that

$$\Pi_g^{(b)}(f) = \frac{1}{\gamma_b} \int_0^1 (tL)P_t^{(b)} \left(Q_t^{(b-1)} f \cdot P_t^{(b)} g \right) + Q_t^{(b-1)} \left((tL)P_t^{(b)} f \cdot P_t^{(b)} g \right) \frac{dt}{t}.$$

With $s = \alpha + (\beta - \frac{\nu}{p})_- > -2$, Lemma B.4 yields

$$\|\Pi_g^{(b)}(f)\|_{W^{s, p}} \lesssim \left\| \left(\int_0^1 t^{-s} \left| Q_t^{(b-1)} f \cdot P_t^{(b)} g \right|^2 + t^{-s} \left| (tL)P_t^{(b)} f \cdot P_t^{(b)} g \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_p.$$

If $\beta > \nu/p$ (and so $s = \alpha$) then uniformly with respect to $t > 0$ we have due to the local Ahlfors regularity (which allows us to use a Sobolev embedding, see [10, Lemma 10.5])

$$\|P_t^{(b)} g\|_{\infty} \lesssim \|g\|_{\infty} \lesssim \|g\|_{W^{\beta, p}}$$

and so

$$\begin{aligned} \|\Pi_g^{(b)}(f)\|_{W^{s, p}} &\lesssim \left\| \left(\int_0^1 t^{-s} \left| Q_t^{(b-1)} f \right|^2 + t^{-s} \left| (tL)P_t^{(b)} f \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_p \|g\|_{W^{\beta, p}} \\ &\lesssim \|f\|_{W^{s, p}} \|g\|_{W^{\beta, p}}, \end{aligned}$$

where we used again Lemma B.4. If $\beta < \frac{\nu}{p}$ (and so $s = \alpha + \beta - \frac{\nu}{p}$), then

$$\begin{aligned} \|P_t^{(b)} g\|_{\infty} &\lesssim \int_t^1 \|Q_s^{(b)} g\|_{\infty} \frac{ds}{s} + \|P_1^{(b)} g\|_{\infty} \\ &\lesssim \int_t^1 s^{\beta/2} \|Q_s^{(b-\beta/2)} L^{\beta/2} g\|_{\infty} \frac{ds}{s} + \|g\|_{W^{\beta, p}} \\ &\lesssim \left(1 + \int_t^1 s^{(\beta-\frac{\nu}{p})/2} \frac{ds}{s} \right) \|g\|_{W^{\beta, p}} \\ &\lesssim t^{(\beta-\frac{\nu}{p})/2} \|g\|_{W^{\beta, p}}. \end{aligned}$$

Hence, we conclude with Lemma B.4 since

$$\begin{aligned} \|\Pi_g^{(b)}(f)\|_{W^{s, p}} &\lesssim \left\| \left(\int_0^1 t^{-\alpha} \left| Q_t^{(b-1)} f \right|^2 + t^{-\alpha} \left| (tL)P_t^{(b)} f \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_p \|g\|_{W^{\beta, p}} \\ &\lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}}. \end{aligned}$$

▷

Proposition B.13. Fix an integer $b > 2$. For any $\alpha, \beta \in (-\infty, 1 - \theta)$ with $\alpha + \beta > 0$, for every $f \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$, we have the continuity estimate

$$\left\| \Pi^{(b)}(f, g) \right\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}.$$

PROOF – We only study the most difficult term in the resonant term $\Pi^{(b)}(f, g)$, which takes the form

$$(B.9) \quad A(f, g) := \int_0^1 P_t^{(b-1)} \Gamma \left(\sqrt{t} P_t^{(b-1)} f, \sqrt{t} P_t^{(b-1)} g \right) \frac{dt}{t}.$$

$P_t^{(b-1)}$ satisfies L^1 - L^∞ off-diagonal estimates at order N (N can be chosen arbitrarily large, since b is an integer) and so

$$\left| P_t^{(b-1)}(h)(x) \right| \lesssim \sum_{\ell \geq 0} 2^{-\ell N} \left(\int_{2^\ell B(x, \sqrt{t})} |h| d\mu \right).$$

With $h = \sqrt{t} \Gamma P_t^{(b-1)} f \cdot \sqrt{t} \Gamma P_t^{(b-1)} g$ and Hölder's inequality, we deduce that

$$A(f, g)(x) \lesssim \sum_{\ell \geq 0} 2^{-\ell N} \int_0^1 \left(\int_{B(x, 2^\ell \sqrt{t})} |\sqrt{t} \Gamma P_t^{(b-1)} f|^2 d\mu \right)^{1/2} \left(\int_{B(x, 2^\ell \sqrt{t})} |\sqrt{t} \Gamma P_t^{(b-1)} g|^2 d\mu \right)^{1/2} \frac{dt}{t}.$$

We then conclude as previously, with Proposition B.7. \triangleright

We then give the analog estimate in Sobolev spaces.

Proposition B.14. Assume the local Ahlfors regularity. Fix an integer $b > 2$ and $p \in (1, \infty)$. For any $\alpha, \beta \in (-\infty, 1 - \theta)$ with $\alpha + \beta > \frac{\nu}{p}$, for every $f \in W^{\alpha, p}$ and $g \in W^{\beta, p}$, we have the continuity estimate

$$\left\| \Pi^{(b)}(f, g) \right\|_{W^{\alpha+\beta-\frac{\nu}{p}, p}} \lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}}.$$

PROOF – Again, we only study the most difficult term $A(f, g)$ defined in (B.9). With $s := \alpha + \beta - \frac{\nu}{p} > 0$, we have by Lemma B.4

$$\|L^{s/2} A(f, g)\|_p \lesssim \left\| \left(\int_0^1 t^{-s} \left| (tL)^{s/2} P_t^{(b-1)} \Gamma \left(\sqrt{t} P_t^{(b-1)} f, \sqrt{t} P_t^{(b-1)} g \right) \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_p.$$

Since $s > 0$, $(tL)^{s/2} P_t^{(b-1)}$ satisfies L^1 - L^∞ off-diagonal estimates at order $\frac{s}{2}$ (see Proposition B.3) and so

$$\left| (tL)^{s/2} P_t^{(b-1)}(h)(x) \right| \lesssim \sum_{\ell \geq 0} 2^{-\ell \frac{s}{2}} \left(\int_{2^\ell B(x, \sqrt{t})} |h| d\mu \right).$$

With $h = \sqrt{t} \Gamma P_t^{(b-1)} f \cdot \sqrt{t} \Gamma P_t^{(b-1)} g$ and Hölder's inequality, we deduce that $\|L^{\frac{s}{2}} A(f, g)\|_p$ is bounded by

$$(B.10) \quad \sum_{\ell \geq 0} 2^{-\ell s} \left\| \left(\int_0^1 t^{-s} \left(\int_{B(x, 2^\ell \sqrt{t})} |\sqrt{t} \Gamma P_t^{(b-1)} f|^2 d\mu \right) \left(\int_{B(x, 2^\ell \sqrt{t})} |\sqrt{t} \Gamma P_t^{(b-1)} g|^2 d\mu \right) \frac{dt}{t} \right)^{1/2} \right\|_p.$$

Then using Proposition B.9 with the Ahlfors regularity, we have

$$\left(\int_{B(x, 2^\ell \sqrt{t})} |\sqrt{t} \Gamma P_t^{(b-1)} g|^2 d\mu \right)^{1/2} \lesssim t^{-\nu/(2p)} t^{\beta/2} \|g\|_{W^{\beta, p}}.$$

By Proposition B.8, we get

$$\left(\int_{B(x, 2^\ell \sqrt{t})} |\sqrt{t} \Gamma P_t^{(b-1)} f|^2 \right)^{1/2} \lesssim \mathcal{M}[(tL)^{\alpha/2} Q_t^{(\delta)} f](x),$$

for some $\delta > 0$. Hence

$$\begin{aligned} \|L^{s/2} A(f, g)\|_p &\lesssim \|g\|_{W^{\beta, p}} \sum_{\ell \geq 0} 2^{-\ell s} \left\| \left(\int_0^1 t^{-s} |\mathcal{M}[(tL)^{\alpha/2} Q_t^{(\delta)} f]|^2 t^{\beta-\nu/p} \frac{dt}{t} \right)^{1/2} \right\|_p \\ &\lesssim \|g\|_{W^{\beta, p}} \left\| \left(\int_0^1 |\mathcal{M}[Q_t^{(\delta)} L^{\alpha/2} f]|^2 \frac{dt}{t} \right)^{1/2} \right\|_p. \end{aligned}$$

Using the Fefferman-Stein inequality (on the maximal function) with the L^p -boundedness of the square function (see Lemma B.4), we deduce that

$$\|L^{s/2} A(f, g)\|_p \lesssim \|L^{\alpha/2} f\|_p \|g\|_{W^{\beta, p}} \lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}}.$$

By a similar reasoning, we have

$$\begin{aligned} \|A(f, g)\|_p &\lesssim \int_0^1 \left\| P_t^{(b-1)} \Gamma \left(\sqrt{t} P_t^{(b-1)} f, \sqrt{t} P_t^{(b-1)} g \right) \right\|_p \frac{dt}{t} \\ &\lesssim \|g\|_{W^{\beta, p}} \int_0^1 \left\| \mathcal{M}[Q_t^{(\delta)} L^{\alpha/2} f] \right\|_p t^{-\nu/(2p)} t^{\beta/2} \frac{dt}{t} \\ &\lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}} \left(\int_0^1 t^{s/2} \frac{dt}{t} \right) \\ &\lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}}, \end{aligned}$$

where we used that $s > 0$ and the L^p -boundedness of the approximation operators. That concludes the proof of the estimate

$$\|A(f, g)\|_{W^{s, p}} \lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}}.$$

Since the resonant part $\Pi^{(b)}$ can be split into a finite number of terms similar to $A(f, g)$, we then deduce the Sobolev boundedness of the resonant part. \square

B.3.2. Commutator estimates We now focus on the adaptation of the commutator estimates given above in Proposition 3.6.

Proposition B.15. Consider the a priori unbounded trilinear operator

$$C(f, g, h) := \Pi^{(b)} \left(\Pi_g^{(b)}(f), h \right) - g \Pi^{(b)}(f, h),$$

on \mathcal{S}'_o . Let α, β, γ be Hölder regularity exponents with $\alpha \in (-1, 1 - \theta)$, $\beta \in (0, 1 - \theta)$ and $\gamma \in (-\infty, 1]$. If

$$0 < \alpha + \beta + \gamma \quad \text{and} \quad \alpha + \gamma < 0$$

then, setting $\delta := (\alpha + \beta) \wedge (1 - \theta) + \gamma$, we have

$$(B.11) \quad \|C(f, g, h)\|_{\mathcal{C}^\delta} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma},$$

for every $f \in \mathcal{C}^\alpha$, $g \in \mathcal{C}^\beta$ and $h \in \mathcal{C}^\gamma$; so the commutator defines a trilinear map from $\mathcal{C}^\alpha \times \mathcal{C}^\beta \times \mathcal{C}^\gamma$ to \mathcal{C}^δ .

PROOF – We refer to the proof of Proposition 3.6 for details and we keep the same notations. So it suffices to focus on a generic term of the form

$$D(f, g, h) := \mathcal{R}(\mathcal{A}(f, g), h) - g \mathcal{R}(f, h)$$

and prove the continuity estimate (B.11) for it. As previously done, we split the proof of the commutator estimate (B.11) for D in two steps, and introduce an intermediate quantity

$$\mathcal{S}(f, g, h) := \int_0^1 \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot P_t g \right) \frac{dt}{t},$$

for which we shall prove that we have both

$$(B.12) \quad \|g \mathcal{R}(f, h) - \mathcal{S}(f, g, h)\|_{\mathcal{C}^\delta} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma}$$

and

$$(B.13) \quad \|D(f, g, h) - \mathcal{S}(f, g, h)\|_{\mathcal{C}^\delta} \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma}.$$

Step 1 – proof of (B.12). This part is very similar to Step 1 of Proposition 3.6, so we only point out the modifications. Using Gaussian pointwise estimates for the kernel of \mathcal{P}_t^1 , we have for almost every $x \in M$

$$\begin{aligned} \left| \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) \right| &\lesssim \sum_{\ell \geq 0} e^{-c4^\ell} \left(\sup_{d(x, y) \leq 2^\ell \sqrt{t}} |g(x) - P_t g(y)| \right) \dots \\ &\dots \left(\int_{2^\ell B(x, \sqrt{t})} |\sqrt{t} \Gamma(\mathcal{P}_t^2 f)|^2 d\mu \right)^{1/2} \left(\int_{2^\ell B(x, \sqrt{t})} |\sqrt{t} \Gamma(\mathcal{P}_t^3 h)|^2 d\mu \right)^{1/2}. \end{aligned}$$

By using the \mathcal{C}^β -regularity of g as well as Proposition B.7 to estimate the L^2 averages, we get

$$\begin{aligned} \left| \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) \right| &\lesssim \left(\sum_{\ell \geq 0} e^{-c4^\ell} (4^\ell t)^{\beta/2} t^{\alpha/2} t^{\gamma/2} \right) \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma} \\ &\lesssim t^{(\alpha+\beta+\gamma)/2} \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma}. \end{aligned}$$

Consequently, the continuity estimate (B.12) in L^∞ comes from integrating with respect to time, taking into account the fact that $\alpha + \beta + \gamma > 0$.

Then to estimate the regularity of $g \mathcal{R}(f, h) - \mathcal{S}(f, g, h)$, one can exactly reproduce the same reasoning as for Proposition 3.6 by using the Hölder regularity of the heat kernel (Lemma B.2), which involves the condition $\alpha + \beta + \gamma < 1 - \theta$ (since $\beta < 1 - \theta$ and $\alpha + \gamma < 0$).

Step 2 – proof of (B.13). Given the collection $(Q_r := Q_r^{(1)})_{r \in (0, 1]}$ of operators, we need to prove that we have

$$(B.14) \quad \left\| Q_r \left(\mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right) \right\|_\infty \lesssim r^{\delta/2}.$$

for every $r \in (0, 1]$, and where

$$(B.15) \quad \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) = \int_0^1 \mathcal{P}_t^1 \Gamma \left(\sqrt{t} \left\{ \int_0^1 \mathcal{P}_t^2 \mathcal{Q}_s^1 (\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^3 g) \frac{ds}{s} - P_t g \cdot \mathcal{P}_t^2 f \right\}, \sqrt{t} \mathcal{P}_t^3 h \right) \frac{dt}{t}.$$

We are going to follow the same argument as for Proposition 3.6 and we only detail the modifications. So we set

$$A_t(f, g) := \sqrt{t} \Gamma \left(\int_0^1 \mathcal{P}_t^2 \mathcal{Q}_s^1 (\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^3 g) \frac{ds}{s} - P_t g \mathcal{P}_t^2 f \right).$$

and using the L^1 - L^∞ off-diagonal estimates of \mathcal{P}_t^1 , we then deduce that for almost every $x_0 \in M$

$$(B.16) \quad \begin{aligned} & \left| \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h)(x_0) \right| \\ & \lesssim \sum_{\ell \geq 0} \int_0^1 e^{-c4^\ell} \left(\int_{B(x_0, 2^\ell \sqrt{t})} |A_t(f, g)|^2 d\mu \right)^{1/2} \left(\int_{B(x_0, 2^\ell \sqrt{t})} |\sqrt{t}\Gamma \mathcal{P}_t^4 h|^2 d\mu \right)^{1/2}. \end{aligned}$$

Using a suitable normalization of the operators $\int_0^1 \mathcal{Q}_s^1 \mathcal{Q}_s^2 f \frac{ds}{s} = f - \mathcal{P}_1 f$ for some operator \mathcal{P}_1 , it yields for every $x \in M$

$$A_t(f, g)(x) \leq \int_0^1 \sqrt{t}\Gamma \mathcal{P}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f (\mathcal{P}_s^3 g - P_t g(x)) \right)(x) \frac{ds}{s} + |P_t g(x)| \sqrt{t}\Gamma [\mathcal{P}_t^2 \mathcal{P}_1 f](x).$$

This quantity will then be integrated on $B_\ell := B(x_0, 2^\ell \sqrt{t})$, so we first aim to replace $P_t g(x)$ by $\int_{B_\ell} P_t g d\mu$. Observe that

$$(B.17) \quad \begin{aligned} A_t(f, g)(x) & \leq \int_0^1 \sqrt{t}\Gamma \mathcal{P}_t^2 \mathcal{Q}_s^1 \left(\mathcal{Q}_s^2 f (\mathcal{P}_s^3 g - \int_{B_\ell} P_t g d\mu) \right)(x) \frac{ds}{s} \\ & \quad + \left| P_t g(x) - \int_{B_\ell} P_t g d\mu \right| \sqrt{t}\Gamma \mathcal{P}_t^2 f(x) + |P_t g(x)| \sqrt{t}\Gamma [\mathcal{P}_t^2 \mathcal{P}_1 f](x). \end{aligned}$$

As before, we use $\beta > 0$ and the \mathcal{C}^β regularity of g to have

$$\left| P_t g(x) - \int_{B_\ell} P_t g d\mu \right| \lesssim (2^\ell \sqrt{t})^\beta \|g\|_{\mathcal{C}^\beta},$$

and uniformly in $y \in M$

$$\left| \mathcal{P}_s^3 g(y) - \int_{B(x_0, \sqrt{t})} P_t g d\mu \right| \lesssim (\max(s, t)^{\beta/2} + d(x_0, y)^\beta) \|g\|_{\mathcal{C}^\beta}.$$

Moreover, it follows from the composition of L^2 off-diagonal estimates (corresponding to a L^2 analog of Lemma A.5 – Part1, see also [10, Lemma 2.5]), that the operator $\sqrt{t}\Gamma \left(\mathcal{P}_t^2 \mathcal{Q}_s^1 \right)$ satisfies L^2 off-diagonal estimates at the scale $\max(s, t)$ with an extra factor $\left(\frac{\min(s, t)}{\max(s, t)} \right)$; so if one sets $\tau := \max(s, t)$, we have with (B.17)

$$\begin{aligned} & \left(\int_{B(x_0, 2^\ell \sqrt{t})} |A_t(f, g)|^2 d\mu \right)^{1/2} \\ & \lesssim \int_0^1 \sum_{k \geq 0} \left(\frac{\min(s, t)}{\max(s, t)} \right) e^{-c4^k} (4^{k+\ell} \tau)^{\beta/2} \left(\int_{B(x_0, 2^{k+\ell} \sqrt{\tau})} |\mathcal{Q}_s^2 f|^2 d\mu \right)^{1/2} \|g\|_{\mathcal{C}^\beta} \frac{ds}{s} \\ & \quad + \left(\int_{B(x_0, 2^\ell \sqrt{t})} |\sqrt{t}\Gamma \mathcal{P}_t^2 f|^2 d\mu \right)^{1/2} (2^\ell \sqrt{t})^\beta \|g\|_{\mathcal{C}^\beta} + t^{(1-\theta)/2} \|g\|_\infty \|f\|_{\mathcal{C}^\alpha} \\ & \lesssim 2^{\ell\beta} \left[\int_0^1 \left(\frac{\min(s, t)}{\max(s, t)} \right) s^{\frac{\alpha}{2}} \tau^{\frac{\theta}{2}} \frac{ds}{s} + t^{\frac{\alpha}{2}} \right] \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \\ & \lesssim 2^{\ell\beta} t^{\frac{\alpha+\theta}{2}} \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}, \end{aligned}$$

where we used Proposition B.7 and the fact that $\alpha > -1$ and $\alpha + \beta \leq 1 - \theta$ to estimate the integral over s . Observe that in the case where $\alpha + \beta \geq 1 - \theta$, we get

$$\left(\int_{B(x_0, 2^\ell \sqrt{t})} |A_t(f, g)|^2 d\mu \right)^{1/2} \lesssim 2^{\ell\beta} t^{\frac{1-\theta}{2}} \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}.$$

Coming back to the identity (B.16), with Proposition B.7 we have

$$\begin{aligned} \left| \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right|(x_0) &\lesssim \left(\int_0^1 t^{(\alpha+\beta+\gamma)/2} \frac{dt}{t} \right) \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma} \\ &\lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma}, \end{aligned}$$

since $\alpha + \beta + \gamma > 0$, uniformly for every $x_0 \in M$. We then conclude to

$$\left\| \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right\|_\infty \lesssim \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta} \|h\|_{\mathcal{C}^\gamma}.$$

Moreover, taking into account that we have $Q_r^{(1)} \mathcal{P}_t^1 = \frac{r}{t} \overline{Q}_t$ for $t \geq r$, we see that the estimate (B.14) holds true (see the proof of Proposition 3.6). \triangleright

We then aim to have a similar commutator estimate in Sobolev spaces.

Proposition B.16. Assume the local Ahlfors regularity. Let α, β, γ be regularity exponents and $p \in (1, \infty)$ with $\alpha \in (-1, 1 - \theta)$, $\beta \in (\nu/p, 1 - \theta)$ and $\gamma \in (-\infty, 1]$. If

$$\frac{2\nu}{p} < \alpha + \beta + \gamma \quad \text{and} \quad \alpha + \gamma < \frac{\nu}{p}$$

then, setting $\delta := (\alpha + \beta - \frac{\nu}{p}) \wedge 1 + \gamma - \nu/p > 0$ and assume that $2\delta > \beta - \nu/p$. We have

$$(B.18) \quad \left\| C(f, g, h) \right\|_{W^{\delta, p}} \lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}} \|h\|_{W^{\gamma, p}},$$

for every $f \in W^{\alpha, p}$, $g \in W^{\beta, p}$ and $h \in W^{\gamma, p}$; so the commutator defines a trilinear map from $W^{\alpha, p} \times W^{\beta, p} \times W^{\gamma, p}$ to $W^{\delta, p}$.

We follow the exact same proof as previously, so we keep the same notations and only focus on the modifications.

PROOF – Consider a generic term of the form

$$D(f, g, h) := \mathcal{R}(\mathcal{A}(f, g), h) - g \mathcal{R}(f, h)$$

and prove the continuity estimate (B.18) for it. Aiming that, we split into two terms by introducing the quantity

$$\mathcal{S}(f, g, h) := \int_0^1 \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot P_t g \right) \frac{dt}{t}.$$

for which we shall prove that we have both

$$(B.19) \quad \left\| g \mathcal{R}(f, h) - \mathcal{S}(f, g, h) \right\|_{W^{\delta, p}} \lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}} \|h\|_{W^{\gamma, p}}$$

and

$$(B.20) \quad \left\| D(f, g, h) - \mathcal{S}(f, g, h) \right\|_{W^{\delta, p}} \lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}} \|h\|_{W^{\gamma, p}}.$$

Step 1 – proof of (B.19). We first prove a weaker version of the continuity estimate (B.19), under the form of the inequality

$$(B.21) \quad \left\| g \mathcal{R}(f, h) - \mathcal{S}(f, g, h) \right\|_p \lesssim \|f\|_{W^{\alpha, p}} \|g\|_{W^{\beta, p}} \|h\|_{W^{\gamma, p}}.$$

As previously, we have

$$(B.22) \quad (g \mathcal{R}(f, h) - \mathcal{S}(f, g, h))(x) = \int_0^1 \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right)(x) \frac{dt}{t},$$

for μ -almost every $x \in M$. Since $g \in \mathcal{C}^\beta$, with $\beta > \nu/p$ then $g \in W^{\beta, p} \subset \mathcal{C}^{\beta - \nu/p}$ and so

$$\|P_t g - g\|_\infty \lesssim t^{(\beta - \nu/p)/2} \|g\|_{W^{\beta, p}}.$$

Hence

$$(B.23) \quad |P_t g(y) - g(x)| \lesssim \left(\sqrt{t} + d(x, y)\right)^{\beta - \nu/p} \|g\|_{W^{\beta, p}},$$

for every $x, y \in M$. Coming back to equation (B.22) and using Gaussian pointwise estimates for the kernel of \mathcal{P}_t^1 , we have for almost every $x \in M$

$$\begin{aligned} & \left| \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) \right| \\ & \lesssim t^{(\beta - \nu/p)/2} \|g\|_{W^{\beta, p}} \sum_{\ell \geq 0} e^{-c4^\ell} \left(\int_{2^\ell B(x, \sqrt{t})} |\sqrt{t} \Gamma(\mathcal{P}_t^2 f)|^2 d\mu \right)^{1/2} \left(\int_{2^\ell B(x, \sqrt{t})} |\sqrt{t} \Gamma(\mathcal{P}_t^3 h)|^2 d\mu \right)^{1/2}. \end{aligned}$$

So using Propositions B.8 and B.7, we deduce that

$$\left| \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) \right| \lesssim \mathcal{M}[L^{\alpha/2} f](x) \|g\|_{W^{\beta, p}} \|h\|_{W^{\beta, p}} t^{(\alpha + \beta + \gamma)/2 - \nu/p}.$$

Then the continuity estimate (B.21) comes from integrating with respect to time, taking into account the fact that $\alpha + \beta + \gamma > \frac{2\nu}{p}$.

Let us then estimate the regularity of $F := g\mathcal{R}(f, h) - \mathcal{S}(f, g, h)$. It is known (see [16, Section 2.1.1], [5, Section 5.2] or [10, Proposition 9.7]) that

$$\|F\|_{W^{\delta, p}} \lesssim \|F\|_p + \|S_\delta(F)\|_p,$$

where S_δ is the Strichartz functional of index $\delta \in (0, 1)$:

$$S_\delta(F)(x) := \left(\int_0^1 r^{-2\delta} \left(\int_{B(x, r)} |F(x) - F(y)| d\mu(y) \right)^2 \frac{dr}{r} \right)^{1/2}.$$

Fix $r > 0$ and two points $x, y \in M$ with $d(x, y) \leq r$. Then as previously, we write

$$\begin{aligned} F(x) - F(y) &= (g(x)\mathcal{R}(f, h) - \mathcal{S}(f, g, h))(x) - (g(y)\mathcal{R}(f, h) - \mathcal{S}(f, g, h))(y) \\ &=: U + V \end{aligned}$$

with U defined by the formula

$$\int_0^{r^2} \left\{ \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) - \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(y) - P_t g) \right) (y) \right\} \frac{dt}{t},$$

and V by

$$\int_{r^2}^1 \left\{ \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) - \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(y) - P_t g) \right) (y) \right\} \frac{dt}{t}.$$

By repeating previous arguments, we easily bound U as follows

$$U \lesssim \left(\int_0^{r^2} t^{(\alpha + \beta + \gamma)/2 - \nu/p} \left[\mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](x) + \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](y) \right] \frac{dt}{t} \right) \|g\|_{W^{\beta, p}} \|h\|_{W^{\gamma, p}},$$

for some $\varepsilon > 0$ satisfying $\alpha + \varepsilon < 1 - \theta$.

For the second part, we use

$$|V| \leq A + B$$

with A equal to

$$\left| \int_{r^2}^1 \left\{ \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (x) - \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \cdot (g(x) - P_t g) \right) (y) \right\} \frac{dt}{t} \right|$$

and

$$B := \int_{r^2}^1 |g(x) - g(y)| \cdot \left| \mathcal{P}_t^1 \left(\Gamma(\sqrt{t} \mathcal{P}_t^2 f, \sqrt{t} \mathcal{P}_t^3 h) \right) (y) \right| \frac{dt}{t}.$$

The last quantity is bounded (following the same estimates as previously since $g \in \mathcal{C}^{\beta-\nu/p}$) by

$$B \lesssim r^{\beta-\nu/p} \left(\int_{r^2}^1 t^{(\alpha+\gamma-\nu/p)/2} \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](y) \frac{dt}{t} \right) \|g\|_{W^{\beta,p}} \|h\|_{W^{\gamma,p}}.$$

For the quantity A , we combine the previous argument with the Hölder regularity of the heat kernel, Lemma B.2, to get the upper bound

$$A \lesssim \left(\int_{r^2}^1 \left(\frac{r}{\sqrt{t}} \right)^{1-\theta} t^{(\alpha+\beta+\gamma)/2-\nu/p} \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](y) \frac{dt}{t} \right) \|g\|_{W^{\beta,p}} \|h\|_{W^{\gamma,p}}.$$

The combination of all the previous estimates yields

$$\begin{aligned} |F(x) - F(y)| &\leq |U| + A + B \\ &\lesssim \left[\left(\int_0^{r^2} t^{\delta/2} \left[\mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](x) + \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](y) \right] \frac{dt}{t} \right) \right. \\ &\quad \left. + \left(\int_{r^2}^1 \left(\frac{r}{\sqrt{t}} \right)^{1-\theta} t^{\delta/2} \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](y) \frac{dt}{t} \right) \right. \\ &\quad \left. + \left(\int_{r^2}^1 r^{\beta-\nu/p} t^{(\alpha+\gamma-\nu/p)/2} \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f](y) \frac{dt}{t} \right) \right] \|g\|_{W^{\beta,p}} \|h\|_{W^{\gamma,p}}. \end{aligned}$$

This estimate holds uniformly for every $y \in B(x, r)$ and so can be averaged on this ball. We then conclude by Hardy's inequality (with $\delta > 0$, $\delta < 1 - \theta$ and $\beta > \nu/p$) that

$$S_\delta(F) \lesssim \left(\int_0^1 \left| \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f] \right|^2 \frac{dt}{t} \right)^{1/2} \|g\|_{W^{\beta,p}} \|h\|_{W^{\gamma,p}}.$$

Using Fefferman-Stein's inequality and the L^p -boundedness of the vertical square function (see Lemma B.4), we then deduce that

$$\|F\|_{W^{\delta,p}} \lesssim \|F\|_p + \|S_\delta(F)\|_p \lesssim \|f\|_{W^{\alpha,p}} \|g\|_{W^{\beta,p}} \|h\|_{W^{\gamma,p}},$$

which concludes the proof of the continuity estimate (B.19).

Step 2 – proof of (B.20). We need to prove that we have

$$(B.24) \quad (*) := \left\| L^{\delta/2} \left(\mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) \right) \right\|_p \lesssim \|f\|_{W^{\alpha,p}} \|g\|_{W^{\beta,p}} \|h\|_{W^{\gamma,p}},$$

where

$$(B.25) \quad \mathcal{R}(\mathcal{A}(f, g), h) - \mathcal{S}(f, g, h) = \int_0^1 \mathcal{P}_t^1 \Gamma \left(\sqrt{t} \left\{ \int_0^1 \mathcal{P}_t^2 \mathcal{Q}_s^1 (\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^3 g) \frac{ds}{s} - \mathcal{P}_t g \cdot \mathcal{P}_t^2 f \right\}, \sqrt{t} \mathcal{P}_t^4 h \right) \frac{dt}{t}.$$

Using Lemma B.4 with the L^1 - L^∞ off-diagonal estimates of \mathcal{P}_t^1 , we deduce that quantity (*) is estimated by

$$(*) \lesssim \sum_{\ell \geq 0} 2^{-\ell \delta} \left\| \left(\int_0^1 t^{-\delta} \left(\int_{B(x_0, 2^\ell \sqrt{t})} |A_t(f, g)|^2 d\mu \right) \left(\int_{B(x_0, 2^\ell \sqrt{t})} |\sqrt{t} \Gamma \mathcal{P}_t^3 h|^2 d\mu \right) \frac{dt}{t} \right)^{1/2} \right\|_p,$$

with

$$A_t(f, g) := \sqrt{t} \Gamma \left(\int_0^1 \mathcal{P}_t^2 \mathcal{Q}_s^1 (\mathcal{Q}_s^2 f \cdot \mathcal{P}_s^3 g) \frac{ds}{s} - \mathcal{P}_t g \mathcal{P}_t^2 f \right).$$

Following the reasoning in the previous proof of Proposition B.15, by combining with Proposition B.8 we can obtain that for almost every x_0

$$\begin{aligned} \left(\int_{B(x_0, 2^\ell \sqrt{t})} |A_t(f, g)|^2 d\mu \right)^{1/2} &\lesssim 2^{\ell(\beta-\nu/p)} t^{(\beta-\nu/p)/2} \mathcal{M}[Q_t^{(\varepsilon)}(tL)^{\alpha/2} f](x_0) \|g\|_{W^{\beta,p}} \\ &\quad + t^{(1-\theta)/2} \mathcal{M}[Q_1^{(\varepsilon)} L^{\alpha/2} f](x_0) \|g\|_{W^{\beta,p}}. \end{aligned}$$

Hence, since $2\delta > \beta - \nu/p$ we obtain

$$(*) \lesssim \|g\|_{W^{\beta,p}} \|h\|_{W^{\gamma,p}} \left\{ \left\| \left(\int_0^1 \mathcal{M}[Q_t^{(\varepsilon)} L^{\alpha/2} f]^2 \frac{dt}{t} \right)^{1/2} \right\|_p + \|f\|_p \right\}.$$

We then conclude to (B.24) by the Fefferman-Stein inequality and Lemma B.4. \triangleright

B.3.3. Composition estimates The above continuity estimates are the main estimates used in the main part of this work to prove the parilinearisation and composition estimates for paraproduct. We state these results here in Hölder and Sobolev spaces under our relaxed assumptions and leave the reader the task of checking that the proofs of section 3.4 are easily adapted.

Theorem B.17. Fix an integer $b \geq 2$ and a nonlinearity $F \in C_b^3$.

(a) Let $\alpha \in (0, 1 - \theta)$ be given. For every $f \in C^\alpha$, we have $F(f) \in C^\alpha$ and

$$R_F(f) := F(f) - \Pi_{F'(f)}^{(b)}(f) \in C^{2\alpha}.$$

More precisely,

$$\left\| F(f) - \Pi_{F'(f)}^{(b)}(f) \right\|_{C^{2\alpha}} \lesssim \|F\|_{C_b^3} \left(1 + \|f\|_{C^\alpha}^2 \right).$$

If $F \in C_b^4$, then the remainder term $R_F(f)$ is Lipschitz with respect to f , in so far as we have

$$\left\| R_F(f) - R_F(g) \right\|_{C^{2\alpha}} \lesssim \|F\|_{C_b^4} \left(1 + \|f\|_{C^\alpha} + \|g\|_{C^\alpha} \right)^2 \|f - g\|_{C^\alpha}.$$

(b) Fix $p \in (1, \infty)$. For every $\alpha \in (\nu/p, 1 - \theta)$ and every $f \in W^{\alpha,p}$, we have $F(f) \in W^{\alpha,p}$ and

$$R_F(f) := F(f) - \Pi_{F'(f)}^{(b)}(f) \in W^{2\alpha-\nu/p,p}.$$

More precisely

$$\left\| F(f) - \Pi_{F'(f)}^{(b)}(f) \right\|_{W^{2\alpha-\nu/p,p}} \lesssim \|F\|_{C_b^3} \left(1 + \|f\|_{W^{\alpha,p}}^2 \right).$$

If $F \in C_b^4$, then the remainder term $R_F(f)$ is Lipschitz with respect to f .

Let us now examine the composition of two paraproducts. Note that for $u \in C^\alpha$ and $v \in C^\beta$, with $\alpha \in (0, 1)$, $\beta \in (0, \alpha]$, we have $uv \in C^\beta$.

About the composition of paraproducts, Theorem 3.8 in Hölder spaces still holds since it only relies on the Gaussian estimate (UE); its Sobolev counterpart also holds.

Theorem B.18. (a) Fix an integer $b \geq 2$, $\alpha \in (0, 1)$, $\beta \in (0, \alpha]$ and consider $u \in C^\alpha$ and $v \in C^\beta$. Then for every $f \in C^\alpha$, we have

$$\Pi_u^{(b)} \left(\Pi_v^{(b)}(f) \right) - \Pi_{uv}^{(b)}(f) \in C^{\alpha+\beta}$$

with

$$\left\| \Pi_u^{(b)} \left(\Pi_v^{(b)}(f) \right) - \Pi_{uv}^{(b)}(f) \right\|_{C^{\alpha+\beta}} \lesssim \|f\|_{C^\alpha} \|u\|_{C^\alpha} \|v\|_{C^\beta}.$$

- (b) Fix an integer $b \geq 2$ and $p \in (1, \infty)$. For $\alpha \in (0, 1)$ and $\beta \in (\nu/p, \alpha]$, consider $u \in W^{\alpha,p}$ and $v \in W^{\beta,p}$. Then for every $f \in W^{\alpha,p}$, we have

$$\Pi_u^{(b)} \left(\Pi_v^{(b)}(f) \right) - \Pi_{uv}^{(b)}(f) \in W^{\alpha+\beta-2\nu/p,p}$$

with

$$\left\| \Pi_u^{(b)} \left(\Pi_v^{(b)}(f) \right) - \Pi_{uv}^{(b)}(f) \right\|_{W^{\alpha+\beta-2\nu/p,p}} \lesssim \|f\|_{W^{\alpha,p}} \|u\|_{W^{\alpha,p}} \|v\|_{W^{\beta,p}}.$$

B.3.4. Schauder estimates Proposition 3.10 gives an elementary proof in our setting of a Schauder-type estimate about the regularizing character of the convolution operation with the operators $P_s^{(b)}$. The same properties hold in our minimal setting since they only rely on Gaussian property (UE) and the semigroup structure, together with a Sobolev version which we state here without proof as it can be proved along the lines of proof of proposition 3.10. (Another approach can be also obtained by interpolating between the trivial case $\varepsilon = 1$ and the limit case $\varepsilon = 0$. The latter case $\varepsilon = 0$, corresponds exactly to the so-called L^p maximal regularity which has been the topic of a huge literature, see for example [40] where the Gaussian upper estimates (UE) are used.)

Proposition B.19. (a) Consider $\beta \in \mathbb{R}$ and $\varepsilon \in (0, 1)$. For every $T > 0$ and $v \in C_T C^\beta$, the function $V(t) := \int_0^t P_{t-s}^{(b)} v(s) ds$ belongs to $C_T C^{\beta+2-2\varepsilon}$ with

$$\|V(t)\|_{C^{\beta+2-2\varepsilon}} \lesssim T^\varepsilon \sup_{s \in [0,t]} \|v(s)\|_{C^\beta}$$

and

$$\|V\|_{C_T^{(\beta+2-2\varepsilon)/2} L^\infty} \lesssim T^\varepsilon \|v\|_{C_T C^\beta}.$$

So

$$\|Jf\|_{\mathcal{L}_T^\alpha} \lesssim T^\varepsilon \|f\|_{C_T C^{\alpha-2+2\varepsilon}}.$$

- (b) Consider $\beta \in \mathbb{R}$, $p \in (1, \infty)$ and $\varepsilon \in (0, 1)$. For every $T > 0$ and $v \in C_T W^{\beta,p}$, the function $V(t) := \int_0^t P_{t-s}^{(b)} v(s) ds$ belongs to $C_T W^{\beta+2-2\varepsilon,p}$ with

$$\|V(t)\|_{W^{\beta+2-2\varepsilon,p}} \lesssim T^\varepsilon \sup_{s \in [0,t]} \|v(s)\|_{W^{\beta,p}}$$

and

$$\|V\|_{C_T^{(\beta+2-2\varepsilon)/2} L^p} \lesssim T^\varepsilon \|v\|_{C_T W^{\beta,p}}.$$

B.4. Resolution of PAM in such a 2-dimensional setting Building on the estimates proved in this Appendix, it is elementary to introduce and study paracontrolled distributions in Hölder and Sobolev spaces along the lines of Subsections 4.1 and 4.2, in the present extended setting. Its application to the parabolic Anderson model equation (PAM) is also almost straightforward as we only need to check that the renormalization procedure explained in details in subsection 5.3 under the (Lip) assumption can be run here as well. This is indeed the case if the exponent q_0 in the gradient assumption (G_{q_0}) is large enough, as this assumption yields some " L^{q_1} -Gaussian" estimates for every $q_1 < q_0$.

Let us compute, as an example, an integral of type

$$I_{s,t} := \int \Gamma_x p_t(x, y) \Gamma_x p_s(x, z) d\mu(x),$$

where p_t is the heat kernel of e^{-tL} , and $s < t$. By (B.1) with the local Ahlfors regularity, there exists a positive constant c such that we have

$$\begin{aligned} I_{s,t} &\lesssim t^{-\frac{\nu}{2q_1}} s^{-\frac{\nu}{2q_1}} \left(\int e^{c\frac{d(x,y)^2}{t}} e^{c\frac{d(x,z)^2}{s}} d\mu(x) \right)^{1-\frac{2}{q_1}} \\ &\lesssim t^{-\frac{\nu}{2q_1}} s^{-\frac{\nu}{2q_1}} \left(\int \mathcal{G}_t(x, y) \mathcal{G}_s(x, z) d\mu(x) \right)^{1-\frac{2}{q_1}} \\ &\lesssim t^{-\frac{\nu}{2q_1}} s^{-\frac{\nu}{2q_1}} \mathcal{G}_{t+s}(y, z)^{\frac{\nu}{2} \left(1-\frac{2}{q_1}\right)}, \end{aligned}$$

where we used Lemma A.5. So with respect to Subsection 5.3, where (Lip) was assumed and where $I_{s,t}$ would be estimated by $G_{s+t}(y, z)$, we now have the estimate

$$I_{s,t} \lesssim \left(\frac{(t+s)^2}{ts} \right)^{\frac{\nu}{2q_1}} G_{t+s}(y, z),$$

involving an extra factor $\left(\frac{(t+s)^2}{ts} \right)^{\frac{\nu}{2q_1}}$. Since all the conditions on the exponents were open conditions in Subsection 5.3, we may allow a small loss if it is small enough. As a consequence, we deduce that if q_1 can be chosen large enough then we may adapt and repeat the renormalization procedure of the white noise in Hölder and Sobolev spaces. The latter condition on q_1 is equivalent to taking q_0 big enough.

We summarize this result under the following form, which gives an analogue of theorem 1.2.

Theorem B.20. Assume the local Ahlfors regularity of dimension 2, as well as (P_2) , (G_{q_0}) and $(DG_{q,\theta})$ for q_0 large enough and θ small enough. Fix $p > 2$ a large enough exponent.

Let ξ stand for a time-independent weighted noise in space, and set $\xi^\varepsilon := P_\varepsilon \xi$, and $X^\varepsilon(t) = \int_0^t P_{t-s}(\xi^\varepsilon) ds$.

- (a) The pair $(\xi^\varepsilon, X^\varepsilon)$ converges in probability in some space (in the Hölder scaling $(C^s)_s$ or Sobolev scaling $(W^{s,p})$) to some extended noise (ζ, X) , with $\zeta = \xi$, and $\Pi(X, \zeta)$ well-defined in the above sense.
- (b) Furthermore, if u^ε stands for the solution of the renormalized equation

$$(B.26) \quad \partial_t u^\varepsilon + Lu^\varepsilon = F(u^\varepsilon) \xi^\varepsilon - c^\varepsilon F'(u^\varepsilon) F(u^\varepsilon), \quad u^\varepsilon(0) = u_0$$

where $c^\varepsilon(\cdot) := \mathbb{E} \left[\Pi(L^{-1} \xi^\varepsilon, \xi^\varepsilon)(\cdot) \right]$ is a deterministic real-valued function on M , then u^ε converges in probability to the solution u of $(gPAM)$ associated with (ζ, X) , in some space whose definition depends on whether or not F is linear.

References

- [1] P. Auscher, Regularity theorems and heat kernel for elliptic operators, *J. London Math. Soc.*, **54** (1996), no. 2, 284–296.
- [2] P. Auscher, T. Coulhon, X. T. Duong, and S. Hofmann, Riesz transform on manifolds and heat kernel regularity, *Ann. Sci. Ecole Norm. Sup.*, **37** (2004), no. 4, 911–957.
- [3] D. Albrecht, X. T. Duong and A. McIntosh, Operator theory and harmonic analysis, in *Instructional Workshop on Analysis and Geometry, Part III (Canberra, 1995)*, Proc. Centre Math. Appl. Austral. Nat. Univ. 34 (1996), 77–136.

- [4] P. Auscher, A. McIntosh and P. Tchamitchian, Noyau de la chaleur d'opérateurs elliptiques complexes, *Math. Research Lett.* **1** (1994), 37–45.
- [5] N. Badr, F. Bernicot, and E. Russ, Algebra properties for Sobolev spaces-applications to semilinear PDEs on manifolds, *J. Anal. Math.*, **118**, no.2 (2012), 509–544.
- [6] H. Bahouri, R. Danchin and J.Y. Chemin, Fourier analysis and nonlinear partial differential equations, *Grundlehren der Mathematischen Wissenschaften* 343, (2011).
- [7] F. Baudoin and N. Garofalo, Curvature-dimension inequalities and Ricci lower bounds for sub-Riemannian manifolds with transverse symmetries, *J. Eur. Math. Soc.* (2015).
- [8] F. Bernicot, A T(1)-Theorem in relation to a semigroup of operators and applications to new paraproducts, *Trans. Amer. Math. Soc.* **364** (2012), 257–294.
- [9] F. Bernicot, T. Coulhon and D. Frey, Gaussian heat kernel bounds through elliptic Moser iteration, *submitted*, arXiv:1407.3906.
- [10] F. Bernicot, T. Coulhon and D. Frey, Sobolev algebra through Heat semigroup, *submitted*, arXiv:1505.01442.
- [11] F. Bernicot and D. Frey, Riesz transforms through reverse Hölder and Poincaré inequalities, *submitted*, arXiv:1503.02508.
- [12] F. Bernicot and Y. Sire, Propagation of low regularity for solutions of nonlinear PDEs on a Riemannian manifold with a sub-Laplacian structure, *Ann. I. H. Poincaré - AN* **30** (2013), 935–958.
- [13] J.M. Bony, Calcul symbolique et propagation des singuliers pour les équations aux dérivées partielles non linéaires, *Ann. Sci. Eco. Norm. Sup.* **114** (1981), 209–246.
- [14] S. Boutayeb, T. Coulhon and A. Sikora, A new approach to pointwise heat kernel upper bounds on doubling metric measure spaces, preprint 2013, arXiv:1311.0367.
- [15] H-Q. Bui, X. T. Duong and L. Yan, Calderón reproducing formulas and new Besov spaces associated with operators, *Adv. in Math.* **229** (2012), 2449–2502.
- [16] T. Coulhon, E. Russ, and V. Tardivel-Nachef, Sobolev algebras on Lie groups and Riemannian manifolds, *Amer. J. of Math.*, **123** (2001), 283–342.
- [17] T. Coulhon and A. Sikora, Gaussian heat kernel bounds via Phragmén-Lindelöf theorem, *Proc. London Math. Soc.* **96** (2008), 507–544.
- [18] M. Cowling, I. Doust, A. McIntosh and A. Yagi, Banach space operators with a bounded H^∞ functional calculus, *J. Austral. Math. Soc. Ser. A* **60** (1996), 51–89.
- [19] E. De Giorgi, Sulla differenziabilità e analiticità delle estremali degli integrali multipli regolari, *Mem. Accad. Sci. Torino Cl. Sci. Fis. Mat. Nat.*, **3** (1957), no. 3, 25–43.
- [20] X.-T. Duong and D. W. Robinson, Semigroup kernels, Poisson bounds, and holomorphic functional calculus, *J. Funct. Anal.* **142**(1) (1996), 89–128.
- [21] P. Friz and M. Hairer, A course on rough paths (with an introduction to regularity structures), Springer, 2014.
- [22] P. Friz and N. Victoir, *Multidimensional stochastic processes as rough paths; Theory and Applications*. Cambridge University Press, 2010.
- [23] M. Fukushima, Y. Oshima and M. Takeda, *Dirichlet forms and symmetric Markov processes*. De Gruyter Studies in Mathematics 19, Walter de Gruyter, Berlin, 1994.
- [24] I. Gallagher and Y. Sire, Besov algebras on Lie groups of polynomial growth, *Studia Math.* **212** (2012), no. 2, 119–139.
- [25] A. Grigor'yan, Gaussian upper bounds for the heat kernel on arbitrary manifolds, *J. Diff. Geom.* **45** (1997), 33–52.
- [26] A. Grigor'yan, Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds, *Bull. Amer. Math. Soc.* **36** (1999), 135–249.
- [27] A. Grigor'yan and L. Liu, Heat kernel and Lipschitz-Besov spaces, *Forum Math.* (2015).
- [28] M. Gubinelli, Controlling rough paths, *J. Funct. Anal.* **2016**, no. 1, (2004), 86–140.
- [29] M. Gubinelli, P. Imkeller and N. Perkowski, Paracontrolled distributions and singular PDEs, Preprint 2012, arXiv:1210.2684.
- [30] Y. Guivar'ch, Croissance polynomiale et périodes des fonctions harmoniques, *Bull. Soc. Math. France* **101** (1973), 333–379.
- [31] P. Gyrya and L. Saloff-Coste, *Neumann and Dirichlet heat kernels in inner uniform domains*, Astérisque **33** (2011), Soc. Math. France.
- [32] M. Hairer, A theory of regularity structures, *Invent. Math.*, 198 (2014), no. 2, 269–504.
- [33] M. Hairer, Rough Stochastic PDEs, *Commun. Pure Appl. Math.*, 64 (2011), no. 11, 1547–1585.
- [34] M. Hairer, Solving the KPZ equation, *Annals of Maths*, 178 (2013), no. 2, 559–664.
- [35] M. Hairer and C. Labbé, Multiplicative stochastic heat equations on the whole space, Preprint, arXiv:arXiv:1504.07162.

- [36] M. Hairer and C. Labbé, A simple construction of the continuum parabolic Anderson model on \mathbb{R}^2 , Preprint, arXiv:1501.00692.
- [37] M. Hairer and H. Weber, Rough Burgers-like equations with multiplicative noise, *Probab. Theory Rel. Fields*, 155 (2013), no. 1, 71–126.
- [38] W. Hebisch and L. Saloff-Coste, Gaussian estimates for Markov chains and random walks on groups, *Ann. Probab.* **21** (1993), no. 2, 673–709.
- [39] W. Hebisch and L. Saloff-Coste, On the relation between elliptic and parabolic Harnack inequalities, *Ann. Inst. Fourier* **51** (2001), no. 5, 1437–1481.
- [40] M. Hieber and J. Prüss, Heat kernels and maximal L^p - L^q estimates for parabolic evolution equations, *Comm. in Partial Differential Equation* **22** (1997), 1647–1669.
- [41] S. Ishiwata, A Berry-Esseen type theorem on nilpotent covering graphs, *Canad. J. Math.* **56** (2004), no. 5, 963–982.
- [42] S. Ishiwata, Gradient estimate of the heat kernel on modified graphs, *Pot. Analysis* **27** (2007), no. 4, 335–351.
- [43] P. C. Kunstmann and L. Weis, *Maximal L_p regularity for parabolic problems, Fourier multiplier theorems and H^∞ -functional calculus*, Lect. Notes in Math. 1855. Springer-Verlag (2004).
- [44] H-Q Li, La transformation de Riesz sur les variétés coniques, *J. Funct. Anal.* **168** (1999), 145–238.
- [45] P. Li and S. T. Yau, On the parabolic kernel of the Schrödinger operator, *Acta Math.* **156** (1986), 153–201.
- [46] L. Liu, D. Yang and W. Yuan, Besov-type and Triebel-Lizorkin-type spaces associated with Heat kernels, Preprint 2013, arXiv:1309.1366.
- [47] Y. Meyer, *Wavelets and operators*, Cambridge Studies in advanced mathematics (1992).
- [48] A. Nagel, E. M. Stein and S. Wainger, Balls and metrics defined by vector fields I: Basic properties, *Acta Math.* **155** (1985), 103–147.
- [49] B. Qian, Hamilton type gradient estimate for the sub-elliptic operators, *Potential Anal.* (2015).
- [50] L. Saloff-Coste, Analyse sur les groupes de Lie à croissance polynomiale, *Ark. Mat.* **28** (1990), 315–331.
- [51] L. Saloff-Coste, A note on Poincaré, Sobolev, and Harnack inequalities, *Int. Math. Res. Not.* (1992), no. 2, 27–38.
- [52] E. M. Stein, *Topics in harmonic analysis related to the Littlewood-Paley theory*. Princeton University Press, 1970.
- [53] K.T. Sturm, Analysis on local Dirichlet spaces I. Recurrence, conservativeness and L^p -Liouville property, *J. Reine Angew. Math.* **456** (1994), 173–196.
- [54] K.T. Sturm, Analysis on local Dirichlet spaces II. Upper Gaussian estimates for the fundamental solutions of parabolic equations, *Osaka J. Math.* **32** (1995), no. 2, 275–312.
- [55] N. Th. Varopoulos, Analysis on Lie groups, *J. Funct. Anal.* **76** (1988), 346–410.
- [56] N. Th. Varopoulos, Small time Gaussian estimates of heat diffusion kernels. Part 1. The semigroup technique, *Bull. Sci. Math. Ser. 2* **113** (1989), 253–277.
- [57] N. Th. Varopoulos, L. Saloff-Coste and T. Coulhon, *Analysis and geometry on groups*. Cambridge University Press, 1992.
- [58] F.-Y. Wang and L. Yan, Gradient estimate on convex domains and applications, *Proc. Amer. Math. Soc.* **141** (2013), 1067–1081.
- [59] R. Zhu and X. Zhu, Three dimensional Navier-Stokes equation driven by space-time white noise, arXiv:1406.0047, (2014).
- [60] R. Zhu and X. Zhu, Approximating three-dimensional Navier-Stokes equations driven by space-time white noise, arXiv:1409.4864, (2014).

INSTITUT DE RECHERCHE MATHÉMATIQUES DE RENNES, 263 AVENUE DU GENERAL LECLERC, 35042 RENNES, FRANCE

E-mail address: ismael.bailleul@univ-rennes1.fr

CNRS - UNIVERSITÉ DE NANTES, LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, 2, RUE DE LA HOUSINIÈRE 44322 NANTES CEDEX 03, FRANCE

E-mail address: frederic.bernicot@univ-nantes.fr

CNRS - UNIVERSITÉ PARIS-SUD, LABORATOIRE DE MATHÉMATIQUES, UMR 8628, 91405 ORSAY, FRANCE

E-mail address: dorothee.frey@univ-nantes.fr