



**HAL**  
open science

# Approximate controllability via adiabatic techniques for the three-inputs controlled Schrödinger equation

Francesca Chittaro, Paolo Mason

► **To cite this version:**

Francesca Chittaro, Paolo Mason. Approximate controllability via adiabatic techniques for the three-inputs controlled Schrödinger equation. *SIAM Journal on Control and Optimization*, 2017, 55 (6), pp.4202-4226. 10.1137/15m1041419 . hal-01117828v2

**HAL Id: hal-01117828**

**<https://hal.science/hal-01117828v2>**

Submitted on 7 Mar 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1                    **APPROXIMATE CONTROLLABILITY VIA ADIABATIC**  
2                    **TECHNIQUES FOR THE THREE-INPUTS CONTROLLED**  
3                    **SCHRÖDINGER EQUATION\***

4                    FRANCESCA CARLOTTA CHITTARO <sup>†</sup> AND PAOLO MASON <sup>‡</sup>

5                    **Abstract.** We consider a system described by a controlled bilinear Schrödinger equation with  
6 three external inputs. We provide a constructive method to approximately steer the system from a  
7 given energy level to a superposition of energy levels corresponding to a given probability distribution.  
8 The method is based on adiabatic techniques and works if the spectrum of the Hamiltonian, as a  
9 function of the control parameters, admits conical intersections of eigenvalues. We provide sharp  
10 estimates of the relation between the error and the controllability time, and we show how to improve  
11 these estimates by selecting special control paths. As a byproduct of our results we show that conical  
12 intersections are stable with respect to general perturbations of the Hamiltonian and we also provide  
13 some results on the regularity of the eigenfamilies along paths locally around conical intersections.

14                    **Key words.** Schrödinger equation, quantum control, controllability, adiabatic methods, conical  
15 intersections

16                    **AMS subject classifications.** 93B05, 93C20, 81Q93

17                    **1. Introduction.** A typical issue in quantum control concerns the controllability  
18 of the bilinear Schrödinger equation

19 (1)                    
$$i \frac{d\psi}{dt} = \left( H_0 + \sum_{k=1}^m u_k(t) H_k \right) \psi(t),$$

20 where  $\psi$  belongs to the unit sphere of a (finite or infinite dimensional) complex separable  
21 Hilbert space  $\mathcal{H}$  and  $H_0, \dots, H_m$  are self-adjoint operators on  $\mathcal{H}$ . Here  $H_1, \dots, H_m$   
22 represent the action of external fields on the system, whose strength is given by the  
23 scalar-valued controls  $u_1, \dots, u_m$ , while  $H_0$  describes the uncontrolled dynamics of  
24 the system.

25                    The controllability problem aims at establishing whether, for every pair of states  
26  $\psi_0$  and  $\psi_1$  in the Hilbert sphere, there exist controls  $u_k(\cdot)$  and a time  $T$  such that the  
27 solution of (1) with initial condition  $\psi(0) = \psi_0$  satisfies  $\psi(T) = \psi_1$ .

28                    While the case where  $\mathcal{H}$  is a finite dimensional Hilbert space has been widely  
29 understood [3, 15], in the infinite dimensional case the answer is far from being  
30 given. In particular, negative results have been proved when  $\mathcal{H}$  is infinite-dimensional  
31 (see [4, 30]). Hence one has to look for weaker controllability properties as, for in-  
32 stance, approximate controllability (see for instance [7, 9, 13, 23, 25]), or controllability  
33 between subfamilies of states (in particular the eigenstates of  $H_0$ , which are  
34 the most relevant physical states) or in suitably regular subspaces of the space of  
35 square-integrable functions (see [5, 6]).

---

\*Submitted to the editors DATE. This work is partially based on the conference proceeding [11].

**Funding:** It has been supported by a public grant overseen by the French National Research Agency (ANR) as part of the “Investissement d’Avenir” program, through the iCODE Institute project funded by the IDEX Paris-Saclay, ANR-11-IDEX-0003-02, and by the project CARTT-IUT Toulon.

<sup>†</sup> Aix Marseille Université, CNRS, ENSAM, LSIS UMR 7296, 13397 Marseille, France, and Université de Toulon, CNRS, LSIS UMR 7296, 83957 La Garde, France ([francesca-carlotta.chittaro@univ-tln.fr](mailto:francesca-carlotta.chittaro@univ-tln.fr))

<sup>‡</sup> Laboratoire des Signaux et Systèmes, CNRS - CentraleSupélec - Univ. Paris-Sud, Université Paris-Saclay, 3, rue Joliot Curie, 91192, Gif-sur-Yvette, France ([paolo.mason@l2s.centralesupelec.fr](mailto:paolo.mason@l2s.centralesupelec.fr)).

36 Unlike most of the known controllability results, mainly obtained by means of  
 37 non-constructive arguments, the method proposed in this paper permits to explicitly  
 38 select control inputs steering the system from the initial state to an arbitrarily small  
 39 neighborhood of the given target state.

40 Adiabatic theory and conical intersections between eigenvalues constitute the  
 41 main tools of the control strategy we propose in this paper.

42 Roughly speaking, the adiabatic theorem (see [8, 24, 28]) states that the occupa-  
 43 tion probabilities associated with the energy levels of a time-dependent Hamiltonian  
 44  $H(\cdot)$  are approximately preserved along the evolution given by  $i\dot{\psi}(t) = H(t)\psi(t)$ , pro-  
 45 vided that  $H(\cdot)$  varies slowly enough. This result works whenever the energy levels  
 46 (i.e. the eigenvalues of  $H(\cdot)$ ) are isolated for every  $t$ . On the other hand, if two  
 47 eigenvalues intersect, and provided that  $H(\cdot)$  is smooth enough, the passage through  
 48 the intersections may determine (approximate) exchanges of the corresponding occu-  
 49 pation probabilities (see [28, Corollary 2.5] and [17]). For these reasons, adiabatic  
 50 methods are largely used in quantum control to induce population transfers (see for  
 51 instance the techniques known as Stimulated Raman Adiabatic Passage (STIRAP),  
 52 Stark-chirped rapid adiabatic passage (SCRAP)) and to prepare superposition states  
 53 [20]. The applications of adiabatic methods in quantum control, as a tool for ob-  
 54 taining controllability results, have already been exploited in previous papers (see for  
 55 instance [1, 10, 22, 33]). The general idea is to use slowly varying controls, taking  
 56 advantage of the adiabatic theorem, and “climb” the energy levels through the conical  
 57 intersections.

58 The method proposed in this paper is based on some ideas developed in [10] in  
 59 the case  $m = 2$  for self-adjoint Hamiltonians with real matrix elements. It exploits a  
 60 generalization of [28, Corollary 2.5] stating that it is possible to arbitrarily recombine  
 61 the probability weights associated with two subsequent energy levels by following  
 62 (slowly) a suitable control path passing through a conical intersection between them.  
 63 One of the main limits of the results of [10] consists in the fact that only Hamiltonians  
 64 with real matrix elements (with respect to some suitable basis of the Hilbert space) are  
 65 considered, and only bounded operators are admitted as control Hamiltonians. Many  
 66 important classes of physical systems are thus excluded: for instance, spin systems in  
 67 magnetic fields (described by Pauli matrices), and, in infinite dimension, Schrödinger  
 68 Hamiltonians containing external fields coupled with the momentum. The purpose  
 69 of this paper is to overcome this issue by adapting the control strategy introduced  
 70 in [10] to the general case of self-adjoint Hamiltonians, assuming that three controlled  
 71 Hamiltonians, relatively bounded with respect to the uncontrolled one, are employed.  
 72 Preliminary results in this sense were discussed in [11].

73 The control strategy applies whenever a part of the spectrum of the Hamiltonian  
 74 operator is uniformly separated from the rest of the spectrum (as a function of the  
 75 control parameters), is discrete and, within it, each pair of subsequent eigenvalues  
 76 intersect in a conical intersection. When there exists such a portion of the spectrum,  
 77 called *separated discrete spectrum*, this control strategy permits to attain (approx-  
 78 imately) a state having a prescribed distribution of probability (relative to the energy  
 79 levels of the separated discrete spectrum) starting from an eigenstate. In particular  
 80 this entails a controllability property, that we call *spread controllability*, which, al-  
 81 though weaker than the usual approximate controllability property, is more practical.  
 82 Indeed, the relative phases between pairs of components in the eigenbasis decompo-  
 83 sition are essentially uncontrollable since they evolve according to the gaps between  
 84 the corresponding energy levels. Furthermore, notice that this method allows us to  
 85 control the population inside some portion of the discrete spectrum, if well separated

86 from the rest, even in the presence of continuous spectrum, unlike many other classical  
87 methods.

88 Concerning the precision of the method, an application of the adiabatic theorem  
89 together with [28, Corollary 2.5] shows that the maximal error is of the order of the  
90 square root of the control speed. On the other hand the precision of the transfer may  
91 be remarkably improved if one follows some special paths in the space of controls;  
92 namely, such paths permit to attain a state with a prescribed probability distribution  
93 with an error of the order of the control speed. In practice, this means that in order to  
94 guarantee a given precision one may significantly reduce the duration of the process,  
95 whose extent constitutes one of the main disadvantages of the implementation of  
96 adiabatic techniques.

97 From a technical point of view, the choice of three instead of two controlled  
98 Hamiltonians leads to a different (and more involved) analysis of the properties of  
99 conical intersections and of the special paths introduced in the control algorithm.  
100 Also, many changes are due to the different regularity properties of the spectrum  
101 and of its related objects in the case of possibly unbounded controlled Hamiltoni-  
102 ans. These properties are carefully investigated (see Appendix A) by using tools from  
103 perturbation theory. Notice that the choice of three controls is quite natural when  
104 looking for eigenvalues intersections, since it is well-known, for Hermitian matrices or  
105 within spaces of self-adjoint operators satisfying particular transversality conditions,  
106 that the set of operators admitting multiple eigenvalues is a submanifold of codi-  
107 mension three (see e.g. [2, 29, 31]). Moreover conical intersections do not constitute  
108 a pathological phenomenon since all eigenvalue intersections are generically conical  
109 in the finite dimensional case and in some physically relevant infinite dimensional  
110 models (for reasons of space, this result will be presented in a future work). Conical  
111 intersections are also structurally stable with respect to variations of the Hamiltonian  
112 operator, as shown in Theorem 23. Concerning the relation between conical intersec-  
113 tions and controllability properties of the bilinear Schrödinger equation, let us finally  
114 mention [12].

115 The structure of the paper is the following. In Section 2 we introduce the notations  
116 used throughout the paper, the main assumptions and definitions, and we adapt the  
117 classical statement of the adiabatic theorem to our setting. In Section 3 we discuss  
118 some properties of conical intersections and related results that allow to propose  
119 the basic control strategy. In Section 4 we define special paths and, by means of a  
120 series of technical results, we show that they can be included in the control algorithm  
121 in order to improve its performance. As a byproduct, we get a structural stability  
122 result concerning conical intersections. Some numerical examples are provided in  
123 Section 5. In Section 6 we briefly mention some extensions of the control strategy  
124 and of the controllability results obtained earlier. Appendix A gathers the technical  
125 results concerning the regularity of the spectrum and of the spectral projections that  
126 are needed throughout the paper.

127 **2. Notations and preliminary results.** We start this section by introducing  
128 the notations that will be used in the rest of the paper.

129 For a set  $A \subset \mathbb{C}$  and  $z \in \mathbb{C}$ , we denote by  $d(z, A) = \inf_{x \in A} |z - x|$  the distance between  
130 the point  $z$  and the set  $A$ .

131 For a function  $f(\cdot)$  of a real parameter  $s$ , we use the following notation for its right  
132 and left limits at  $s_0$ :

$$133 \quad f(s_0^\pm) = \lim_{s \rightarrow s_0^\pm} f(s).$$

134 Whenever  $\gamma(s)$ ,  $s \in [s_1, s_2]$  is a curve on  $\mathbb{R}^3$  and  $Q(\cdot)$  is a function of  $\mathbf{v} \in \mathbb{R}^3$  then,  
 135 with abuse of notations, we denote by  $\dot{Q}(\gamma(r))$  the derivative of the composition  
 136  $Q(\gamma(\cdot))$  computed at  $r$ , that is  $\dot{Q}(\gamma(r)) := \frac{d}{ds}Q(\gamma(s))|_{s=r} = \frac{dQ}{d\mathbf{v}}(\gamma(r))\frac{d\gamma}{ds}(r)$ . Similarly,  
 137  $Q^{(l)}(\gamma(r)) := \frac{d^l}{ds^l}Q(\gamma(s))|_{s=r}$ .

138 The scalar product of two vectors  $\mathbf{w}_1, \mathbf{w}_2$  in an euclidean space  $\mathbb{R}^k$  is denoted by  
 139  $\mathbf{w}_1 \cdot \mathbf{w}_2$ , while the norm of  $\mathbf{w} \in \mathbb{R}^k$  is denoted by  $|\mathbf{w}|$ .

140 For a given vector  $\mathbf{v}$  or matrix  $A$  the respective transpose is denoted by  $\mathbf{v}^T$  and  $A^T$ .

141 The inverse of the transpose of an invertible square matrix  $A$  is denoted with  $A^{-T}$ .

142 Given a vector  $\mathbf{v} = (v_1, v_2, v_3) \in \mathbb{C}^3$ , we denote its complex conjugate  $(v_1^*, v_2^*, v_3^*)$  by  
 143  $\mathbf{v}^*$  and its real and imaginary parts respectively by

$$144 \quad \Re(\mathbf{v}) = (\Re(v_1), \Re(v_2), \Re(v_3)) \quad \Im(\mathbf{v}) = (\Im(v_1), \Im(v_2), \Im(v_3)).$$

145 The symbol  $\text{id}$  is used to denote the identity operator on a vector space which is  
 146 specified at each occurrence, whenever not clear from the context.

147 Given a linear operator  $A$  defined on a Hilbert space, its domain is denoted as  $\mathcal{D}(A)$ ,  
 148 the symbol  $\sigma(A)$  denotes the spectrum of  $A$ , while the resolvent set  $\rho(A)$  is the  
 149 complement of  $\sigma(A)$  in  $\mathbb{C}$ .

150 **2.1. General setting.** Let  $\mathcal{H}$  be a separable complex Hilbert space with scalar  
 151 product  $\langle \cdot, \cdot \rangle$  and norm  $\|\cdot\|$ ; let us introduce the following notion of relative bound-  
 152 edness between operators:

153 **DEFINITION 1** (*A-smallness and A-boundedness*). *Let  $A, B$  be two densely defined*  
 154 *operators with domains  $\mathcal{D}(A) \subset \mathcal{D}(B)$ . We say that  $B$  is  $A$ -bounded if there exist*  
 155  *$a, b > 0$  such that  $\|B\psi\| \leq a\|A\psi\| + b\|\psi\|$  for every  $\psi \in \mathcal{D}(A)$ .  $B$  is said to be*  
 156  *$A$ -small if for every  $\alpha > 0$  there exists  $\beta > 0$  such that  $\|B\psi\| \leq \alpha\|A\psi\| + \beta\|\psi\|$  for*  
 157 *every  $\psi \in \mathcal{D}(A)$ . (The latter notion is called infinitesimal smallness with respect to*  
 158  *$A$  in [26].)*

159 Given a self-adjoint operator  $A$  on  $\mathcal{H}$ , for every  $A$ -bounded operator  $B$  we define its  
 160 norm with respect to  $A$  as

$$161 \quad (2) \quad \|B\|_A = \sup_{\psi \in \mathcal{D}(A)} \frac{\|B\psi\|}{\|A\psi\| + \|\psi\|}.$$

162 This provides a norm in the space  $\mathcal{L}(\mathcal{D}(A), \mathcal{H})$  of continuous linear operators from  
 163  $\mathcal{D}(A)$  (endowed with the graph norm of  $A$ ) to  $\mathcal{H}$ .

164 We consider the Hamiltonian

$$165 \quad H(\mathbf{u}) = H_0 + u_1 H_1 + u_2 H_2 + u_3 H_3,$$

166 with  $\mathbf{u} = (u_1, u_2, u_3) \in \mathbb{R}^3$ , and where  $H_i$ ,  $i = 0, \dots, 3$  satisfy the following assump-  
 167 tion:

168 **(H0)**  $H_0$  is a self-adjoint operator on a separable complex Hilbert space  $\mathcal{H}$ , and  $H_i$   
 169 are  $H_0$ -small self-adjoint operators on  $\mathcal{H}$  for  $i = 1, 2, 3$ . Moreover,  $H_0$  is bounded  
 170 from below, that is there exists a constant  $C > 0$  such that  $\langle \psi, H_0 \psi \rangle > -C\|\psi\|^2$  for  
 171 every  $\psi \in \mathcal{D}(H_0)$ .

172 Under assumption **(H0)**, [26, Theorem X.12] guarantees that  $H(\mathbf{u})$  is self-adjoint  
 173 with domain  $\mathcal{D}(H_0)$  and bounded from below (uniformly for  $\mathbf{u}$  belonging to any com-  
 174 pact subset of  $\mathbb{R}^3$ ). Moreover, it is easy to see that for every  $\mathbf{u}$ ,  $H_0$  is  $H(\mathbf{u})$ -bounded,  
 175 and therefore  $H_i$  is  $H(\mathbf{u})$ -small, for every  $i = 1, 2, 3$ , with constants  $a, b$  (as in Defini-  
 176 tion 1) that depend continuously on  $\mathbf{u}$ .

177 Schrödinger Hamiltonians are typical Hamiltonian operators describing quantum  
 178 phenomena and can be represented in the form  $-\Delta + V$  on the Hilbert space  $L^2(\Omega)$ ,  
 179 where  $\Omega$  is a domain of  $\mathbb{R}^n$ ,  $\Delta$  is the Laplacian on  $\Omega$  (with Dirichlet or Neumann  
 180 boundary conditions) and  $V : \Omega \rightarrow \mathbb{R}$  has to be interpreted as a multiplicative operator  
 181 on  $L^2(\Omega)$ . In particular such Hamiltonian operators are unbounded operators. In  
 182 this context Hypothesis **(H0)** is thus intended to describe a Hamiltonian operator of  
 183 the previous form that can be controlled by means of three external inputs so that  
 184  $H_0 = -\Delta + V_0$  and  $H_i = V_i$  for some multiplicative operators  $V_i$ , for  $0 \leq i \leq 3$ .

185 Finite dimensional representations of quantum systems are also common, for in-  
 186 stance in the description of spin systems. In this case the Hamiltonian operator  $H(\mathbf{u})$   
 187 is a Hermitian matrix. Consider for instance the case of a spin-1/2 particle immersed  
 188 in a controlled magnetic field. In this case,  $H_i$  are the Pauli matrices, and the controls  
 189 are the components of the magnetic field.

190 The dynamics of the quantum systems we consider are described by the time-  
 191 dependent Schrödinger equation

$$192 \quad (3) \quad i \frac{d\psi}{dt} = H(\mathbf{u}(t))\psi(t).$$

193 Such an equation has mild solutions under hypothesis **(H0)**,  $\mathbf{u}(\cdot)$  piecewise  $\mathcal{C}^1$  and  
 194 with an initial condition in the domain of  $H_0$  (see e.g. [26, Theorem X.70] and [4]).

195 We are interested in controlling (3) inside some portion of the discrete spectrum  
 196 of  $H(\mathbf{u})$ . More precisely, we assume that (3) possesses a *separated discrete spectrum*,  
 197 according to the following definition.

198 **DEFINITION 2.** *A separated discrete spectrum is a pair  $(\omega, \Sigma)$  where  $\omega$  is a do-  
 199 main in  $\mathbb{R}^3$  and  $\Sigma$  a map defined on  $\omega$  that associates with each  $\mathbf{u} \in \omega$  a subset  
 200  $\Sigma(\mathbf{u})$  of the discrete spectrum of  $H(\mathbf{u})$  such that there exist two continuous functions  
 201  $f_1, f_2 : \omega \rightarrow \mathbb{R}$  satisfying*

- 202 •  $f_1(\mathbf{u}) < f_2(\mathbf{u})$  and  $\Sigma(\mathbf{u}) \subset [f_1(\mathbf{u}), f_2(\mathbf{u})] \quad \forall \mathbf{u} \in \omega.$
- 203 •  $\inf_{\mathbf{u} \in \omega} \inf_{\lambda \in \sigma(H(\mathbf{u})) \setminus \Sigma(\mathbf{u})} d(\lambda, [f_1(\mathbf{u}), f_2(\mathbf{u})]) > \Gamma$  for some  $\Gamma > 0.$

204 **REMARK 3.** *Thanks to Proposition 26 and Lemma 28, in order to guarantee the  
 205 existence of a separated discrete spectrum for the Hamiltonian  $H(\mathbf{u})$  satisfying **(H0)**,  
 206 it is enough that there exist some open interval  $I$  and some point  $\bar{\mathbf{u}}$  such that  $\sigma(H(\bar{\mathbf{u}})) \cap$   
 207  $I$  contains only a finite number of points. This in particular happens when  $H(\bar{\mathbf{u}})$  has  
 208 compact resolvent.*

209 **Notation** We label the eigenvalues belonging to a separated discrete spectrum  $\Sigma(\mathbf{u})$   
 210 in such a way that  $\Sigma(\mathbf{u}) = \{\lambda_{\hat{j}}(\mathbf{u}), \dots, \lambda_{\hat{j}+k}(\mathbf{u})\}$  for some non-negative integers  $\hat{j}, k$ ,  
 211 where  $\lambda_{\hat{j}}(\mathbf{u}) \leq \dots \leq \lambda_{\hat{j}+k}(\mathbf{u})$  are counted according to their multiplicity (note that  
 212 the separation of  $\Sigma$  from the rest of the spectrum guarantees that  $k$  is constant).  
 213 Moreover we denote by  $\phi_{\hat{j}}(\mathbf{u}), \dots, \phi_{\hat{j}+k}(\mathbf{u})$  an associated orthonormal family of eigen-  
 214 states. Notice that in this notation  $\lambda_{\hat{j}}$  does not necessarily correspond to the  $\hat{j}$ -th  
 215 energy level of the system.

216 Our techniques rely on the existence of conical intersections between the eigen-  
 217 values, which constitute a well studied phenomenon in molecular physics (see for  
 218 instance [8, 16, 17, 21, 32]). In this paper we will adopt the following definition,  
 219 consistent with the one already given in [10] for the two-inputs case.

220 **DEFINITION 4.** *Let  $H(\cdot)$  satisfy hypothesis **(H0)**. We say that  $\bar{\mathbf{u}} \in \mathbb{R}^3$  is a conical  
 221 intersection between two subsequent eigenvalues  $\lambda_j$  and  $\lambda_{j+1}$  if  $\lambda_j(\bar{\mathbf{u}}) = \lambda_{j+1}(\bar{\mathbf{u}})$  has*

222 *multiplicity two and there exists a constant  $c > 0$  such that for any unit vector  $\mathbf{v} \in \mathbb{R}^3$*   
 223 *and  $t > 0$  small enough we have that*

$$224 \quad \lambda_{j+1}(\bar{\mathbf{u}} + t\mathbf{v}) - \lambda_j(\bar{\mathbf{u}} + t\mathbf{v}) > ct.$$

225 See Section 5 for some examples of conical intersections in both the finite and infinite-  
 226 dimensional case.

227 To conclude this section, let us make some remarks on the regularity properties  
 228 and the asymptotic behavior of the eigenfamilies of  $H(\mathbf{u})$  in our setting. Notice  
 229 that in general the regularity properties of the Hamiltonian induce similar regularity  
 230 properties of the eigenfamilies (see e.g. Proposition 7 below). Moreover, it is well  
 231 known that the eigenvectors can be chosen analytic along straight lines  $\mathbf{u}(\cdot)$  possibly  
 232 passing through eigenvalues intersections (see [19],[27, Theorem XII.13]).

233 Let  $I$  be an interval and consider a  $\mathcal{C}^1$  curve  $\mathbf{u} : I \rightarrow \mathbb{R}^3$ . By direct computa-  
 234 tions we obtain that the following equations hold whenever  $\lambda_l(t) \neq \lambda_m(t)$  are simple  
 235 eigenvalues of  $H(\mathbf{u}(t))$  with corresponding eigenstates  $\phi_l(t), \phi_m(t)$ :

$$236 \quad (4) \quad \dot{\lambda}_l(t) = \langle \phi_l(t), (\dot{u}_1(t)H_1 + \dot{u}_2(t)H_2 + \dot{u}_3(t)H_3)\phi_l(t) \rangle,$$

$$237 \quad (5) \quad (\lambda_m(t) - \lambda_l(t)) \langle \phi_l(t), \dot{\phi}_m(t) \rangle = \langle \phi_l(t), (\dot{u}_1(t)H_1 + \dot{u}_2(t)H_2 + \dot{u}_3(t)H_3)\phi_m(t) \rangle.$$

239 Assume that  $\lambda_j(\bar{\mathbf{u}}) = \lambda_{j+1}(\bar{\mathbf{u}})$  and consider the half-line  $r_{\mathbf{v}}(t) = \bar{\mathbf{u}} + t\mathbf{v}$  with  $\mathbf{v} =$   
 240  $(v_1, v_2, v_3)$  unit vector and  $t \geq 0$ . Then, thanks to (5) and since each  $H_i$  is  $H_0$ -  
 241 bounded, we have

$$242 \quad \lim_{t \rightarrow 0^+} \langle \phi_j(r_{\mathbf{v}}(t)), (v_1H_1 + v_2H_2 + v_3H_3)\phi_{j+1}(r_{\mathbf{v}}(t)) \rangle$$

$$243 \quad (6) \quad = \langle \phi_j^{\mathbf{v}}, (v_1H_1 + v_2H_2 + v_3H_3)\phi_{j+1}^{\mathbf{v}} \rangle = 0,$$

245 where  $\phi_j^{\mathbf{v}}$  and  $\phi_{j+1}^{\mathbf{v}}$  are the limits of  $\phi_j(r_{\mathbf{v}}(t))$  and  $\phi_{j+1}(r_{\mathbf{v}}(t))$  as  $t$  tends to zero.

246 **2.2. The adiabatic theorem.** In this section we recall a classical formulation  
 247 of the time-adiabatic theorem ([8, 18, 24] adapted to our framework. For a general  
 248 overview see the monograph [28]).

249 Let  $H(\mathbf{u}) = H_0 + \sum_{i=1}^3 u_i H_i$  satisfy **(H0)** and have a separated discrete spectrum  
 250  $(\omega, \Sigma)$ . Assume that the map  $I = [\tau_0, \tau_f] \ni \tau \mapsto \mathbf{u}(\tau) = (u_1(\tau), u_2(\tau), u_3(\tau))$  belongs  
 251 to  $\mathcal{C}^2(I, \omega)$ . We introduce a small parameter  $\varepsilon > 0$  that controls the time scale, and  
 252 the slow Hamiltonian  $H(\mathbf{u}(\varepsilon t))$ ,  $t \in [\tau_0/\varepsilon, \tau_f/\varepsilon]$ . In these notations,  $\tau$  is a geometric  
 253 parameter used to describe the curve in the space of controls, while  $t$  is the actual  
 254 time of the evolution along the control path.

255 We denote by  $U^\varepsilon(t, t_0)$  the time evolution (from  $t_0 = \tau_0/\varepsilon$  to  $t = \tau/\varepsilon$ ) generated  
 256 by  $H(\mathbf{u}(\varepsilon t))$ , and with  $U_a^\varepsilon(t, t_0)$  the time evolution generated by the Hamiltonian  
 257  $H_a(\varepsilon t)$ , where  $H_a(\tau) = H(\mathbf{u}(\tau)) - i\varepsilon P_*(\mathbf{u}(\tau))\dot{P}_*(\mathbf{u}(\tau)) - i\varepsilon P_*^\perp(\mathbf{u}(\tau))\dot{P}_*^\perp(\mathbf{u}(\tau))$  is the  
 258 *adiabatic Hamiltonian*,  $P_*(\mathbf{u})$  denotes the spectral projection of  $H(\mathbf{u})$  on  $\Sigma(\mathbf{u})$ , and  
 259  $P_*^\perp(\mathbf{u}) = \text{id} - P_*(\mathbf{u})$ . A crucial property of the adiabatic Hamiltonian is that the  
 260 corresponding evolution decouples the dynamics relative to the subspace  $P_*(\mathbf{u})\mathcal{H}$  from  
 261 the remainder of the Hilbert space, in the sense that

$$262 \quad P_*(\mathbf{u}(t))U_a^\varepsilon(t, t_0) = U_a^\varepsilon(t, t_0)P_*(\mathbf{u}(t_0)).$$

263 **THEOREM 5 (Adiabatic Theorem).** *Assume that  $H(\mathbf{u}) = H_0 + \sum_{i=1}^3 u_i H_i$  sat-*  
 264 *isfies **(H0)** and has a separated discrete spectrum  $(\omega, \Sigma)$ . Let  $I \subset \mathbb{R}$  and  $\mathbf{u} : I \rightarrow \omega$*



265 be a  $\mathcal{C}^2$  curve. Then  $P_* \in \mathcal{C}^2(I, \mathcal{L}(\mathcal{H}))$  and there exists a constant  $C > 0$  such that  
 266 for all  $\tau_0, \tau \in I$ , and setting  $t_0 = \tau_0/\varepsilon, t = \tau/\varepsilon$ , one has  $\|U^\varepsilon(t, t_0) - U_a^\varepsilon(t, t_0)\| \leq$   
 267  $C\varepsilon(1 + \varepsilon|t - t_0|)$ . In particular

$$268 \quad \|P_*(\mathbf{u}(t))U^\varepsilon(t, t_0) - U^\varepsilon(t, t_0)P_*(\mathbf{u}(t_0))\| \leq 2C\varepsilon(1 + \varepsilon|t - t_0|).$$

269 Let us now assume that  $\Sigma = \{\lambda_j, \lambda_{j+1}\}$ ; we can take advantage of the adiabatic  
 270 theorem to decouple the dynamics associated with the band  $\Sigma$  from those associated  
 271 with the rest of the spectrum, in order to focus on the former.

272 Let  $\mathcal{W}(\tau)$  denote the subspace spanned by the eigenstates associated with  $\lambda_j(\mathbf{u}(\tau))$   
 273 and  $\lambda_{j+1}(\mathbf{u}(\tau))$ . Since  $\mathcal{W}(\tau)$  is two-dimensional for any  $\tau$ , it is possible to map it  
 274 isomorphically on  $\mathbb{C}^2$  and identify an effective Hamiltonian whose evolution is a rep-  
 275 resentation of  $U_a^\varepsilon(t, t_0)|_{\mathcal{W}(\varepsilon t_0)}$  on  $\mathbb{C}^2$ . In particular, if we can find a  $\mathcal{C}^1$  eigenstate basis  
 276  $\{\Phi_1(\mathbf{u}(\tau)), \Phi_2(\mathbf{u}(\tau))\}$  of  $\mathcal{W}(\tau)$  (associated with a reordering  $\{\Lambda_1(\mathbf{u}(\tau)), \Lambda_2(\mathbf{u}(\tau))\}$  of  
 277  $\{\lambda_j(\mathbf{u}(\tau)), \lambda_{j+1}(\mathbf{u}(\tau))\}$ ), then the isomorphism  $\mathcal{U}(\tau) : \mathcal{W}(\tau) \rightarrow \mathbb{C}^2$  is continuous. Rep-  
 278 resented in  $\mathbb{C}^2$ , the evolution  $U_a^\varepsilon(\cdot, t_0)|_{\mathcal{W}(\varepsilon t_0)}$  is governed by the Hamiltonian  $H_{\text{eff}}^\varepsilon(\varepsilon t)$ ,  
 279 where  $H_{\text{eff}}^\varepsilon(\cdot)$  is the *effective Hamiltonian*, whose form is

$$280 \quad (7) \quad H_{\text{eff}}^\varepsilon(\tau) = \begin{pmatrix} \Lambda_1(\mathbf{u}(\tau)) & 0 \\ 0 & \Lambda_2(\mathbf{u}(\tau)) \end{pmatrix} - i\varepsilon \begin{pmatrix} \langle \Phi_1(\mathbf{u}(\tau)), \dot{\Phi}_1(\mathbf{u}(\tau)) \rangle & \langle \Phi_2(\mathbf{u}(\tau)), \dot{\Phi}_1(\mathbf{u}(\tau)) \rangle \\ \langle \Phi_1(\mathbf{u}(\tau)), \dot{\Phi}_2(\mathbf{u}(\tau)) \rangle & \langle \Phi_2(\mathbf{u}(\tau)), \dot{\Phi}_2(\mathbf{u}(\tau)) \rangle \end{pmatrix},$$

281 with associated propagator  $U_{\text{eff}}^\varepsilon(t, t_0) = \mathcal{U}(\varepsilon t)U_a^\varepsilon(t, t_0)\mathcal{U}^*(\varepsilon t_0)$ .

282 Theorem 5 implies the following.

283 **THEOREM 6.** Assume that  $(\omega, \{\lambda_j, \lambda_{j+1}\})$  is a separated discrete spectrum and let  
 284  $\mathbf{u} : I \rightarrow \omega$  be a  $\mathcal{C}^2$  curve such that there exists a  $\mathcal{C}^1$ -varying basis of  $\mathcal{W}(\cdot)$  made of  
 285 eigenstates of  $H(\mathbf{u}(\cdot))$ . Then there exists a constant  $C$  such that for all  $\tau_0, \tau \in I$ , and  
 286 setting  $t_0 = \tau_0/\varepsilon, t = \tau/\varepsilon$ ,

$$287 \quad \|(U^\varepsilon(t, t_0) - \mathcal{U}^*(\varepsilon t)U_{\text{eff}}^\varepsilon(t, t_0)\mathcal{U}(\varepsilon t_0))|_{\mathcal{W}(\varepsilon t_0)}\| \leq C\varepsilon(1 + \varepsilon|t - t_0|).$$

### 288 3. Conical Intersections and general control strategy.

289 **3.1. Properties of conical intersections.** Let  $\lambda_j(\mathbf{u}), \lambda_{j+1}(\mathbf{u})$  denote two eigen-  
 290 values of  $H(\mathbf{u})$ , and assume that they are (locally) separated from the rest of the  
 291 spectrum. It is well known that the projection  $\mathbf{P}_{\mathbf{u}}$  associated with the sum of the  
 292 corresponding eigenspaces is smooth with respect to  $\mathbf{u}$ . More in general, the result  
 293 holds for any portion of the spectrum of  $H(\mathbf{u})$ , in presence of a gap (see e.g. [28]).

294 On the other hand, the projections  $P_j, P_{j+1}$ , associated respectively with  $\lambda_j$  and  
 295  $\lambda_{j+1}$ , are smooth with respect to  $\mathbf{u}$  at any point such that  $\lambda_j \neq \lambda_{j+1}$ , but they  
 296 are not necessarily continuous at the eigenvalues intersections. Nevertheless, along  
 297 regular curves passing through a conical intersection, it is possible to extend these  
 298 projections, obtaining operators whose regularity depends on the regularity of the  
 299 curve, as stated in the following result, proved in Appendix A.

300 **PROPOSITION 7.** Let  $\gamma : I \rightarrow \mathbb{R}^3, I = [-R, 0]$ , be a  $\mathcal{C}^k(I)$  curve such that  $\gamma(0) = \bar{\mathbf{u}}$   
 301 is a conical intersection between the eigenvalues  $\lambda_j$  and  $\lambda_{j+1}$  and  $\dot{\gamma}(t) \neq 0$  for every  
 302  $t \in I$ , and consider its  $k$ -jet at the origin  $\ell_k(t) = \gamma(0) + \sum_{j=1}^k \frac{1}{j!} t^j \frac{d^j}{dt^j} \gamma(t)|_{t=0}$ . Then  
 303  $P_j(\gamma(\cdot))$  is  $\mathcal{C}^k$  on  $[-R, 0]$ , it is  $\mathcal{C}^{k-1}$  at the singularity, and

$$304 \quad \lim_{t \rightarrow 0^-} \frac{d^l}{dt^l} P_j(\gamma(t)) = \lim_{t \rightarrow 0^-} \frac{d^l}{dt^l} P_j(\ell_k(t)), \quad l = 0, \dots, k-1,$$

305 where the limits above hold in the operator norm. The same result holds for  $P_{j+1}(\gamma(\cdot))$ .



306 Conical intersections have a characterization in terms of the non-degeneracy of a  
 307 particular matrix, which contains some geometric properties of the eigenspaces relative  
 308 to the intersecting eigenvalues, as shown below.

309 DEFINITION 8. We define the conicity matrix associated with two orthonormal  
 310 elements  $\psi_1, \psi_2 \in \mathcal{D}(H_0)$  as

$$311 \quad \mathcal{M}(\psi_1, \psi_2) = \begin{pmatrix} \langle \psi_1, H_1 \psi_2 \rangle & \langle \psi_1, H_1 \psi_2 \rangle^* & \langle \psi_2, H_1 \psi_2 \rangle - \langle \psi_1, H_1 \psi_1 \rangle \\ \langle \psi_1, H_2 \psi_2 \rangle & \langle \psi_1, H_2 \psi_2 \rangle^* & \langle \psi_2, H_2 \psi_2 \rangle - \langle \psi_1, H_2 \psi_1 \rangle \\ \langle \psi_1, H_3 \psi_2 \rangle & \langle \psi_1, H_3 \psi_2 \rangle^* & \langle \psi_2, H_3 \psi_2 \rangle - \langle \psi_1, H_3 \psi_1 \rangle \end{pmatrix}.$$

312 The following lemma can be proved by direct computation.

313 LEMMA 9. The quantity  $\det \mathcal{M}(\psi_1, \psi_2)$  is purely imaginary and the function  $(\psi_1, \psi_2) \mapsto$   
 314  $\det \mathcal{M}(\psi_1, \psi_2)$  is invariant under unitary transformation of the argument, that is if  
 315  $\begin{pmatrix} \widehat{\psi}_1 \\ \widehat{\psi}_2 \end{pmatrix} = \mathbb{U} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$  for a pair  $\psi_1, \psi_2$  of orthonormal elements of  $\mathcal{D}(H_0)$  and  $\mathbb{U} \in U(2)$ ,  
 316 then one has  $\det \mathcal{M}(\widehat{\psi}_1, \widehat{\psi}_2) = \det \mathcal{M}(\psi_1, \psi_2)$ .

317 As a consequence of the result here above, the determinant of  $\mathcal{M}(\psi_1, \psi_2)$  depends  
 318 only on the complex space spanned by  $\psi_1$  and  $\psi_2$ . Therefore, in a neighborhood of a  
 319 conical intersection between the levels  $\lambda_j, \lambda_{j+1}$  we can define the following function:

$$320 \quad (8) \quad F(\mathbf{u}) = \det \mathcal{M}(\psi_1, \psi_2)$$

321 where  $\{\psi_1, \psi_2\}$  is an orthonormal basis for the sum of eigenspaces relative to  $\lambda_j, \lambda_{j+1}$ .  
 322 In particular, outside the intersection we can take, for instance,  $\psi_1 = \phi_j$  and  $\psi_2 =$   
 323  $\phi_{j+1}$ . Thanks to the continuity of  $\mathbf{P}_{\mathbf{u}}$ , which follows from Proposition 26, we obtain  
 324 that  $F$  is continuous (see [10]).

325 The following result characterizes conical intersections in terms of the conicity  
 326 matrix.

327 PROPOSITION 10. Assume that  $(\omega, \{\lambda_j, \lambda_{j+1}\})$  is a separated discrete spectrum  
 328 with  $\lambda_j(\bar{\mathbf{u}}) = \lambda_{j+1}(\bar{\mathbf{u}})$ , for some  $\bar{\mathbf{u}} \in \omega$ . Let  $\{\psi_1, \psi_2\}$  be an orthonormal basis of the  
 329 eigenspace associated with the double eigenvalue. Then  $\bar{\mathbf{u}}$  is a conical intersection if  
 330 and only if  $\mathcal{M}(\psi_1, \psi_2)$  is nonsingular.

331 *Proof.* Define  $r_{\mathbf{v}}(t) = \bar{\mathbf{u}} + t\mathbf{v}$ , where  $\mathbf{v}$  is a unit vector in  $\mathbb{R}^3$ , and let  $\phi_j^{\mathbf{v}}, \phi_{j+1}^{\mathbf{v}}$  be  
 332 the limits of  $\phi_j(r_{\mathbf{v}}(t)), \phi_{j+1}(r_{\mathbf{v}}(t))$  as  $t \rightarrow 0^+$  (recall that the eigenfunctions  $\phi_j, \phi_{j+1}$   
 333 can be chosen analytic along  $r_{\mathbf{v}}$  for  $t \geq 0$ ). Assume that the intersection is not conical.  
 334 Then for every  $\varepsilon > 0$  there is a unit vector  $\mathbf{v}_\varepsilon = (v_1^\varepsilon, v_2^\varepsilon, v_3^\varepsilon)$  such that

$$335 \quad \left. \frac{d}{dt} \right|_{t=0^+} \left[ \lambda_{j+1}(r_{\mathbf{v}_\varepsilon}(t)) - \lambda_j(r_{\mathbf{v}_\varepsilon}(t)) \right] \leq \varepsilon.$$

336 By (4) and (6) we deduce that, if  $\mathbb{A}_\varepsilon$  is an orthogonal matrix having  $\mathbf{v}_\varepsilon$  as first row,  
 337 then the first row of the matrix  $\mathbb{A}_\varepsilon \mathcal{M}(\phi_j^{\mathbf{v}_\varepsilon}, \phi_{j+1}^{\mathbf{v}_\varepsilon})$  is equal to  $(0, 0, a)$  for some  $a$  whose  
 338 absolute value is smaller than  $\varepsilon$ . As a consequence

$$339 \quad |\det \mathcal{M}(\phi_j^{\mathbf{v}_\varepsilon}, \phi_{j+1}^{\mathbf{v}_\varepsilon})| = |\det (\mathbb{A}_\varepsilon \mathcal{M}(\phi_j^{\mathbf{v}_\varepsilon}, \phi_{j+1}^{\mathbf{v}_\varepsilon}))| \leq C\varepsilon(\alpha|\lambda_j(\mathbf{u})| + \beta)^2,$$

340 with  $C, \alpha$  and  $\beta$  suitable positive constants, where we have used the fact that  $H_i$  is  
 341  $H(\mathbf{u})$ -bounded for  $i = 1, 2, 3$ . By arbitrariness of  $\varepsilon$ , the conicity matrix is singular.  
 342 Let us now prove the converse statement: assume that  $\bar{\mathbf{u}}$  is a conical intersection

343 and, by contradiction, that  $\mathcal{M}(\psi_1, \psi_2)$  is singular for every orthonormal basis of the  
 344 eigenspace associated with the double eigenvalue. We introduce the matrix

$$345 \quad \widetilde{\mathcal{M}}(\psi_1, \psi_2) = \begin{pmatrix} \Re(\langle \psi_1, H_1 \psi_2 \rangle) & \Im(\langle \psi_1, H_1 \psi_2 \rangle) & (\langle \psi_2, H_1 \psi_2 \rangle - \langle \psi_1, H_1 \psi_1 \rangle) \\ \Re(\langle \psi_1, H_2 \psi_2 \rangle) & \Im(\langle \psi_1, H_2 \psi_2 \rangle) & (\langle \psi_2, H_2 \psi_2 \rangle - \langle \psi_1, H_2 \psi_1 \rangle) \\ \Re(\langle \psi_1, H_3 \psi_2 \rangle) & \Im(\langle \psi_1, H_3 \psi_2 \rangle) & (\langle \psi_2, H_3 \psi_2 \rangle - \langle \psi_1, H_3 \psi_1 \rangle) \end{pmatrix},$$

346 and we notice that  $\det \mathcal{M}(\psi_1, \psi_2) = -2i \det \widetilde{\mathcal{M}}(\psi_1, \psi_2)$  so that  $\mathcal{M}(\psi_1, \psi_2)$  is singular  
 347 if and only if  $\widetilde{\mathcal{M}}(\psi_1, \psi_2)$  is. The condition of conical intersection, together with (4)  
 348 and (6), implies that  $\mathbf{v}^T \widetilde{\mathcal{M}}(\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}}) = (0, 0, a)$  for some  $a \neq 0$ , so that the third  
 349 column of the matrix  $\widetilde{\mathcal{M}}(\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}})$  is never linearly dependent from the first two. In  
 350 particular the matrix  $\widetilde{\mathcal{M}}(\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}})$  is singular only if the first two columns of the  
 351 matrix are linearly dependent. Thus, up to multiplying  $\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}}$  by phase factors, we  
 352 can always assume that  $\langle \phi_j^{\mathbf{y}}, H_i \phi_{j+1}^{\mathbf{y}} \rangle \in \mathbb{R}$ ,  $i = 1, 2, 3$ .

353 Let us now fix a unit vector  $\mathbf{v} \in \mathbb{R}^3$  and let us call  $W$  the orthogonal complement  
 354 in  $\mathbb{R}^3$  of the vector  $(\langle \phi_j^{\mathbf{y}}, H_1 \phi_{j+1}^{\mathbf{y}} \rangle, \langle \phi_j^{\mathbf{y}}, H_2 \phi_{j+1}^{\mathbf{y}} \rangle, \langle \phi_j^{\mathbf{y}}, H_3 \phi_{j+1}^{\mathbf{y}} \rangle)$ . We have that  $\mathbf{v} \in W$   
 355 and  $\dim W \geq 2$ . By direct computations it is easy to prove that, for every  $\mathbf{w} \in W$ ,  
 356 the limit basis  $\{\phi_j^{\mathbf{w}}, \phi_{j+1}^{\mathbf{w}}\}$  is equal to  $\{\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}}\}$ , up to exchanges between the two  
 357 elements and up to phases.

358 For  $\mathbf{w} \in W$  let us consider the vector

$$359 \quad \Upsilon = \begin{pmatrix} \langle \phi_{j+1}^{\mathbf{w}}, H_1 \phi_{j+1}^{\mathbf{w}} \rangle - \langle \phi_j^{\mathbf{w}}, H_1 \phi_j^{\mathbf{w}} \rangle \\ \langle \phi_{j+1}^{\mathbf{w}}, H_2 \phi_{j+1}^{\mathbf{w}} \rangle - \langle \phi_j^{\mathbf{w}}, H_2 \phi_j^{\mathbf{w}} \rangle \\ \langle \phi_{j+1}^{\mathbf{w}}, H_3 \phi_{j+1}^{\mathbf{w}} \rangle - \langle \phi_j^{\mathbf{w}}, H_3 \phi_j^{\mathbf{w}} \rangle \end{pmatrix},$$

360 which corresponds to the third column of  $\widetilde{\mathcal{M}}(\phi_j^{\mathbf{w}}, \phi_{j+1}^{\mathbf{w}})$ . Notice that the definition of  
 361  $\Upsilon$ , up to a sign, does not depend on the choice of  $\mathbf{w} \in W$ , by previous remarks. Since  
 362  $\Upsilon^\perp$ , the orthogonal complement of  $\Upsilon$  in  $\mathbb{R}^3$ , has dimension 2 there exists a non-zero  
 363  $\tilde{\mathbf{w}} \in W \cap \Upsilon^\perp$ . By definition of  $\tilde{\mathbf{w}}$  we have that  $\tilde{\mathbf{w}} \cdot \Upsilon = 0$ .

364 We get a contradiction, thus the matrix  $\widetilde{\mathcal{M}}(\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}})$  must be nonsingular, and  
 365 therefore also  $\mathcal{M}(\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}})$  has to.  $\square$

366 A peculiarity of conical intersections is that, when approaching the singularity  
 367 from different directions, the eigenstates corresponding to the intersecting eigenvalues  
 368 have different limits. The following proposition provides the relation between these  
 369 limits.

370 **PROPOSITION 11.** *Let  $\tilde{\mathbf{u}}$  be a conical intersection between  $\lambda_j$  and  $\lambda_{j+1}$ . Let  $\mathbf{v}_0, \mathbf{v} \in$   
 371  $\mathbb{R}^3$  be two unit vectors, and call  $\phi_j^0, \phi_{j+1}^0$  the limits as  $t \rightarrow 0^+$  of the eigenstates  
 372  $\phi_j(r_0(t)), \phi_{j+1}(r_0(t))$  along a straight line  $r_0(t) = \tilde{\mathbf{u}} + t\mathbf{v}_0$ , and  $\phi_j^{\mathbf{y}}, \phi_{j+1}^{\mathbf{y}}$  the limit  
 373 basis along the straight line  $r_{\mathbf{v}}(t) = \tilde{\mathbf{u}} + t\mathbf{v}$ . Then, up to phases, the following relation  
 374 holds:*

$$375 \quad (9) \quad \begin{pmatrix} \phi_j^{\mathbf{y}} \\ \phi_{j+1}^{\mathbf{y}} \end{pmatrix} = \begin{pmatrix} \cos \Xi & e^{-i\beta} \sin \Xi \\ -e^{i\beta} \sin \Xi & \cos \Xi \end{pmatrix} \begin{pmatrix} \phi_j^0 \\ \phi_{j+1}^0 \end{pmatrix},$$

376 where the parameters  $\Xi = \Xi(\mathbf{v})$  and  $\beta = \beta(\mathbf{v})$  satisfy the following equations:

$$377 \quad (10) \quad \tan 2\Xi(\mathbf{v}) = (-1)^\xi \frac{2|\langle \phi_j^0, H_{\mathbf{v}} \phi_{j+1}^0 \rangle|}{\langle \phi_j^0, H_{\mathbf{v}} \phi_j^0 \rangle - \langle \phi_{j+1}^0, H_{\mathbf{v}} \phi_{j+1}^0 \rangle}$$

$$378 \quad (11) \quad \beta(\mathbf{v}) \stackrel{(\text{mod } 2\pi)}{=} \arg \langle \phi_j^0, H_{\mathbf{v}} \phi_{j+1}^0 \rangle + \xi\pi,$$

380 where  $H_{\mathbf{v}} = \sum_{i=1}^m H_i v_i$  and  $\xi = 0, 1$ .

381 *Proof.* First of all, we notice that all pairs of orthonormal eigenstates of  $H(\bar{\mathbf{u}})$   
 382 relative to the degenerate eigenvalue can be obtained through the action of the group  
 383  $U(2)$  on the pair  $(\phi_j^0, \phi_{j+1}^0)$ . However, if one takes into account the equivalence relation  
 384  $(e^{i\beta_1}\psi_1, e^{i\beta_2}\psi_2) \sim (\psi_1, \psi_2) \forall \beta_1, \beta_2 \in \mathbb{R}$ , it is enough to consider the transformations  
 385 of the form (9). If  $\mathbf{v} = \pm\mathbf{v}_0$ , then  $\langle \phi_j^0, H_{\mathbf{v}}\phi_{j+1}^0 \rangle = 0$  by (6), and we can assume  
 386  $\Xi(\mathbf{v}) = k\pi/2$  for some integer  $k$ , while  $\beta(\mathbf{v})$  can be any real number. If  $\mathbf{v}$  is not  
 387 parallel to  $\mathbf{v}_0$  then, as a consequence of Proposition 10,  $\mathbf{v}^T \mathcal{M}(\phi_j^0, \phi_{j+1}^0)$  is not parallel  
 388 to  $\mathbf{v}_0^T \mathcal{M}(\phi_j^0, \phi_{j+1}^0)$ , so that  $\langle \phi_j^0, H_{\mathbf{v}}\phi_{j+1}^0 \rangle \neq 0$ . The two cases defined by (10)-(11)  
 389 follow from  $\langle \phi_j^y, H_{\mathbf{v}}\phi_{j+1}^y \rangle = 0$ , once we replace  $\phi_j^y, \phi_{j+1}^y$  with the expression given  
 390 in (9).  $\square$

391 It can be seen that not all the solutions of (10)-(11) provide the correct trans-  
 392 formation (9), which, nevertheless, is easy to detect. The good solutions of (10)-(11)  
 393 constitute four branches which are continuous with respect to  $\mathbf{v}$ , and they can be  
 394 constructed as follows. Let  $\mathbf{w}(s)$ ,  $s \in [0, \bar{s}]$ , be a curve joining  $\mathbf{v}_0$  to  $\mathbf{v}$  such that  
 395  $\mathbf{w}(s) \notin \{\mathbf{v}_0, -\mathbf{v}_0\}$  for every  $s \in (0, \bar{s})$ ; for conical intersections, it is possible to as-  
 396 sociate with such a curve a continuous solution  $(\Xi(\mathbf{w}(s)), \beta(\mathbf{w}(s)))$  of (10)-(11) with  
 397  $\Xi(\mathbf{v}_0) = 0$  and compatible with (9). In particular, if we choose  $\Xi(\mathbf{v})$  satisfying (10)  
 398 with  $\xi = 0$ , it is easy to see that  $\Xi(\mathbf{w}(s)) \in [-\pi/2, 0]$  for  $s \in [0, \bar{s}]$ , from which  
 399 one deduces that the final value  $\Xi(\mathbf{v}) = \Xi(\mathbf{w}(\bar{s}))$  is independent of the chosen path  
 400 and continuously depends on  $\mathbf{v}$ . Moreover, it is easy to see that  $\Xi(-\mathbf{v}_0) = -\pi/2$ .  
 401 Similarly  $\beta(\mathbf{v}) = \beta(\mathbf{w}(\bar{s}))$  is independent of the chosen path and continuous outside  
 402  $\{\mathbf{v}_0, -\mathbf{v}_0\}$ . Note that the fact that  $\beta$  is discontinuous at  $-\mathbf{v}_0$  implies that the corre-  
 403 sponding limit basis  $(\phi_j^y, \phi_{j+1}^y)$  has a discontinuity at  $-\mathbf{v}_0$ , that is, its limit depends  
 404 on the path. We can repeat the same argument choosing  $\Xi(\mathbf{v})$  satisfying (10) with  
 405  $\xi = 1$  and with initial condition  $\Xi(\mathbf{v}_0) = 0$ . The other two continuous branches are  
 406 obtained choosing the initial condition  $\Xi(\mathbf{v}_0) = \pi$ .

407 **3.2. The basic control algorithm.** Let us consider the following controllability  
 408 problem.

409 *Let  $H(\cdot)$  satisfy (H0) and have a separated discrete spectrum  $(\omega, \Sigma)$  with  $\Sigma(\cdot) =$   
 410  $\{\lambda_0(\cdot), \dots, \lambda_k(\cdot)\}$ . Then, given  $\eta > 0$ ,  $\mathbf{u}^s, \mathbf{u}^f \in \omega$ ,  $j \in \{0, \dots, k\}$  and  $p \in [0, 1]^{k+1}$   
 411 such that  $\sum_{l=0}^k p_l^2 = 1$ , find  $T > 0$  and a path  $\mathbf{u} : [0, T] \rightarrow \omega$  with  $\mathbf{u}(0) = \mathbf{u}^s$  and  
 412  $\mathbf{u}(T) = \mathbf{u}^f$  such that*

$$413 \quad \|\psi(T) - \sum_{l=0}^k p_l e^{i\vartheta_l} \phi_l(\mathbf{u}^f)\| \leq \eta,$$

414 *where  $\psi(\cdot)$  is the solution of (3) with  $\psi(0) = \phi_j(\mathbf{u}^s)$ , and  $\vartheta_0, \dots, \vartheta_k \in \mathbb{R}$  are some  
 415 possibly unknown phases.*

416 If all levels are connected by means of conical intersections occurring at different  
 417 values of the control, the results obtained in the previous section provide the basic  
 418 elements to construct a family of control paths solving the problem here above. This  
 419 can be done by taking advantage of the following proposition, which describes the  
 420 spreading of occupation probabilities induced when a path in the space of controls  
 421 passes through a conical intersection.

422 **PROPOSITION 12.** *Let  $\bar{\mathbf{u}}$  be a conical intersection between the eigenvalues  $\lambda_j, \lambda_{j+1}$ .  
 423 Consider the curve  $\gamma : [0, 1] \rightarrow \omega$  defined by*

$$424 \quad \gamma(t) = \begin{cases} \bar{\mathbf{u}} + (t - \bar{\tau})\mathbf{w}_0 & t \in [0, \bar{\tau}] \\ \bar{\mathbf{u}} + (t - \bar{\tau})\mathbf{v} & t \in [\bar{\tau}, 1] \end{cases},$$

425 for some  $\bar{\tau} \in (0, 1)$  and some unit vectors  $\mathbf{w}_0, \mathbf{v}$ . Then there exists  $C > 0$  such that,  
 426 for any  $\varepsilon > 0$ ,

$$427 \quad \|\psi(1/\varepsilon) - \pi_1 e^{i\vartheta_1} \phi_j(\gamma(1)) - \pi_2 e^{i\vartheta_2} \phi_{j+1}(\gamma(1))\| \leq C\sqrt{\varepsilon}$$

428 where  $\vartheta_1, \vartheta_2 \in \mathbb{R}$ ,  $\psi(\cdot)$  is the solution of equation (3) with  $\psi(0) = \phi_j(\gamma(0))$  corre-  
 429 sponding to the control  $\mathbf{u} : [0, 1/\varepsilon] \rightarrow \omega$  defined by  $\mathbf{u}(t) = \gamma(\varepsilon t)$ ,

$$430 \quad \pi_1 = |\cos(\Xi(\mathbf{v}))|, \quad \pi_2 = |\sin(\Xi(\mathbf{v}))|,$$

431 and  $\Xi(\cdot)$  is the only solution of equation (10) with  $\xi = 0$  such that  $\Xi(\mathbf{v}) \in (-\pi/2, 0]$   
 432 for  $\mathbf{v} \neq -\mathbf{w}_0$ , and  $\Xi(-\mathbf{w}_0) = -\pi/2$ , where the limit basis in (10) is given by the  
 433 limits  $\phi_j(\gamma(\bar{\tau}^-)), \phi_{j+1}(\gamma(\bar{\tau}^-))$ , respectively.

434 *Proof.* We consider the Hamiltonian  $H(\mathbf{u}(t))$ ,  $t \in [0, 1/\varepsilon]$ . Since the control  
 435 function  $\mathbf{u}(\cdot)$  is not  $\mathcal{C}^2$  at the singularity, we cannot directly apply the adiabatic  
 436 theorem. Instead, we consider separately the evolution on the two subintervals (in  
 437 time  $t$ )  $[0, \bar{\tau}/\varepsilon]$  and  $[\bar{\tau}/\varepsilon, 1/\varepsilon]$ . Since the eigenstates  $\phi_j(\mathbf{u}(t)), \phi_{j+1}(\mathbf{u}(t))$  are piecewise  
 438  $\mathcal{C}^1$ , we can apply [28, Corollary 2.5] and obtain that there exists a phase  $\theta_1$  (depending  
 439 on  $\varepsilon$ ) such that

$$440 \quad \|\psi(\bar{\tau}/\varepsilon) - e^{i\theta_1} \phi_j(\gamma(\bar{\tau}^-))\| \leq C'\sqrt{\varepsilon},$$

441 for some constant  $C' > 0$ . By Proposition 11, this implies that

$$442 \quad \|\psi(\bar{\tau}/\varepsilon) - e^{i\theta_1} \left( \cos \Xi(\mathbf{v}) \phi_j(\gamma(\bar{\tau}^+)) - e^{-i\beta(\mathbf{v})} \sin \Xi(\mathbf{v}) \phi_{j+1}(\gamma(\bar{\tau}^+)) \right)\| \leq C'\sqrt{\varepsilon},$$

443 with  $\Xi(\mathbf{v})$  as in the statement of the proposition and  $\beta(\mathbf{v}) = \arg\langle \phi_j^0, H_{\mathbf{v}} \phi_{j+1}^0 \rangle$ . By  
 444 applying [28, Corollary 2.5] also in the time interval  $(\bar{\tau}/\varepsilon, 1/\varepsilon]$  we get the thesis.  $\square$

445 For control purposes, it is interesting to consider the problem of determining a  
 446 path that induces the desired transition in the case in which the initial probability is  
 447 concentrated in the first level and the final occupation probabilities  $\pi_1^2$  and  $\pi_2^2$  are pre-  
 448 scribed. For a given line reaching the conical intersection, the outward directions that  
 449 provide the required spreading of probability are given in the following proposition.  
 450 The proof easily follows from Proposition 11.

451 **PROPOSITION 13.** *Let  $\bar{\mathbf{u}}$  be a conical intersection between the eigenvalues  $\lambda_j, \lambda_{j+1}$ ,  
 452 and let  $\pi_1, \pi_2$  be positive constants such that  $\pi_1^2 + \pi_2^2 = 1$ . Consider the line  $r(t) =$   
 453  $\bar{\mathbf{u}} + (t - \bar{\tau})\mathbf{w}_0$ ,  $t \in [0, \bar{\tau}]$ , for a unit vector  $\mathbf{w}_0 \in \mathbb{R}^3$  and some  $\bar{\tau} \in (0, 1)$ , and set  
 454  $\phi_j^0 = \phi_j(r(\bar{\tau}^-))$  and  $\phi_{j+1}^0 = \phi_{j+1}(r(\bar{\tau}^-))$ .*

455 *Then the locus formed by the directions  $\mathbf{v} \in \mathbb{R}^3$  that give rise to transformation (9)  
 456 with  $\pi_1 = |\cos \Xi(\mathbf{v})|$  and  $\pi_2 = |\sin \Xi(\mathbf{v})|$  is given by the following expression whenever  
 457  $\pi_1 \notin \{0, 1\}$*

$$458 \quad \widetilde{\mathcal{M}}(\phi_j^0, \phi_{j+1}^0)^{-T}(\mathcal{K}),$$

459 where  $\mathcal{K} = \{(x, y, z) \in \mathbb{R}^3 : \sqrt{x^2 + y^2} = \frac{\pi_1 \pi_2}{\pi_1^2 - \pi_2^2} z\}$ . Otherwise, if  $\pi_1 = 0$  then  $\mathbf{v} = \mathbf{w}_0$   
 460 and if  $\pi_1 = 1$  then  $\mathbf{v} = -\mathbf{w}_0$ .

461 The controllability problem presented at the beginning of this section can be solved  
 462 taking advantage of the results shown above. The strategy consists in constructing  
 463 a piecewise  $\mathcal{C}^2$  path joining  $\mathbf{u}^s$  with  $\mathbf{u}^f$  that passes through the conical intersections  
 464  $\bar{\mathbf{u}}_j$  between the  $j$ -th and the  $(j + 1)$ -th levels,  $j = 0, \dots, k - 1$ , and avoids any  
 465 other degeneracy point. The tangent directions at the conical intersection are chosen  
 466 according to the probability weights  $p_j^2$ , as explained in Proposition 13.




FIG. 1. Construction of the path  $\gamma(\cdot)$ . The corners at the points  $\mathbf{u}_i$ , corresponding to conical intersections, are chosen in such a way that they induce the desired spreading.

467 For simplicity we assume that  $\psi(0) = \phi_0(\mathbf{u}^s)$ ; in the other cases a path can be  
 468 obtained similarly. We set  $\gamma(0) = \mathbf{u}^s$  and, for some  $0 < \tau_0 < 1$  we choose  $\gamma|_{[0, \tau_0]}$   
 469 in such a way that  $\gamma(\tau_0) = \bar{\mathbf{u}}_0$  and all the eigenvalues  $\lambda_l(\gamma(\tau))$  are simple for every  
 470  $l = 0, \dots, k$  and  $\tau \in [0, \tau_0)$ . Moreover,  $\gamma(\cdot)$  is chosen tangent to a segment in a  
 471 neighborhood of  $\bar{\mathbf{u}}_0$ . The rest of the path is then constructed recursively as follows.  
 472 Assume that the path has been defined up to time  $\tau_j$ , for some  $j = 0, \dots, k-2$ , with  
 473  $\gamma(\tau_j) = \bar{\mathbf{u}}_j$ , and that it is tangent to a segment of direction  $\mathbf{v}_j^-$  in a neighborhood of  
 474  $\bar{\mathbf{u}}_j$ . Then  $\gamma|_{[\tau_j, \tau_{j+1}]}$ , where  $\tau_{j+1} \in (\tau_j, 1)$ , is chosen to be tangent, in a neighborhood  
 475 of  $\bar{\mathbf{u}}_j$ , to a segment directed as  $\mathbf{v}_j^+$ , where  $\mathbf{v}_j^+$  is obtained applying Proposition 13 with  
 476  $\mathbf{w}_0 = \mathbf{v}_j^-$ ,  $\pi_1 = p_j / \sqrt{\sum_{l=j}^k p_l^2}$  and  $\pi_2 = \sqrt{1 - \pi_1^2}$ . Moreover  $\gamma|_{[\tau_j, \tau_{j+1}]}$  is such that  
 477  $\gamma(\tau_{j+1}) = \bar{\mathbf{u}}_{j+1}$  and is tangent to a straight line on a neighborhood of  $\bar{\mathbf{u}}_{j+1}$ . The last  
 478 arc defined on  $(\tau_{k-1}, 1]$  is simply constructed by joining  $\bar{\mathbf{u}}_{k-1}$  with  $\mathbf{u}^f$ , taking care  
 479 of choosing  $\gamma(\cdot)$  tangent to the outward direction obtained through Proposition 13,  
 480 in a neighborhood of  $\bar{\mathbf{u}}_{k-1}$ . To avoid highly non-homogeneous parameterizations, the  
 481 path  $\gamma(\cdot)$  can be reparameterized by arc-length. The geometric construction of the  
 482 path is represented in Figure 1.

483 Let us now reparameterize the time by setting  $t = \tau/\varepsilon$ , for some small positive  $\varepsilon$ .  
 484 When we are far from all the conical intersections it follows from Theorem 5 that the  
 485 evolution of  $H(\gamma(\varepsilon t))$  conserves the occupation probabilities relative to each energy  
 486 level  $\lambda_l$ ,  $l = 0, \dots, k$ , with an approximation of order  $\varepsilon$ . Similarly, in a neighborhood  
 487 of the conical intersection between  $\lambda_j$  and  $\lambda_{j+1}$ ,  $j = 0, \dots, k-1$ , the evolution of  
 488  $H(\gamma(\varepsilon t))$  conserves the occupation probabilities relative to each non-intersecting en-  
 489 ergy level with an approximation of order  $\varepsilon$ . The occupation probabilities relative to  
 490 the  $j$ -th and  $(j+1)$ -th levels are instead estimated by combining Theorem 5 with  
 491 the estimate provided by Proposition 12.

492 Summing up, we obtain the estimate

$$493 \quad \|\psi(1/\varepsilon) - \sum_{l=0}^k p_l e^{i\vartheta_l} \phi_l(\mathbf{u}^f)\| \leq C\sqrt{\varepsilon},$$

494 for some  $\vartheta_0, \dots, \vartheta_k \in \mathbb{R}$  and some  $C > 0$  depending on the path  $\gamma(\cdot)$  and on the gaps  
 495 in the spectrum.

496 **4. An improvement of the efficiency of the algorithm: the non-mixing**  
 497 **field.** The problem of reducing transition times is quite important in quantum control,  
 498 in particular for the need of preventing decoherence. In this section we show how  
 499 to construct special paths in the space of controls that permit to speed up the process:

indeed, the error accumulated when following these paths is inversely proportional to the total time of the transition, while the basic strategy described in Proposition 12 guarantees an error scaling with the inverse of the square root of the total time, so that shorter times are sufficient to guarantee a prescribed accuracy.

Let us consider a pair  $\{\lambda_j, \lambda_{j+1}\}$  of eigenvalues separated from the rest of the spectrum in a certain open set  $\omega \subset \mathbb{R}^3$ , according to Definition 2, and intersecting only at  $\bar{\mathbf{u}} \in \omega$  where they form a conical intersection. We are interested in the dynamics inside the subspace  $\mathbf{P}_{\bar{\mathbf{u}}}\mathcal{H}$ , where  $\mathbf{P}_{\bar{\mathbf{u}}}$  denotes the projection associated with the two levels  $\{\lambda_j(\mathbf{u}), \lambda_{j+1}(\mathbf{u})\}$  for  $\mathbf{u} \in \omega$ . To improve the algorithm described in the previous section, the idea is to cancel the off-diagonal terms in the effective Hamiltonian (7), which are responsible of the error of order  $\sqrt{\varepsilon}$  in the estimates given in [28, Corollary 2.5] and in Proposition 12. In order to do that, we choose some special trajectories in  $\omega$  along which the term  $\langle \phi_j, \dot{\phi}_{j+1} \rangle$  is null.

We denote the first column of the conicity matrix  $\mathcal{M}(\psi_1, \psi_2)$  by  $\mathbf{m}(\psi_1, \psi_2)$  and its components  $\langle \psi_1, H_i \psi_2 \rangle$  as  $m_i$ , and we define the vector

$$(12) \quad X(\psi_1, \psi_2) = \Im \mathbf{m}(\psi_1, \psi_2) \times \Re \mathbf{m}(\psi_1, \psi_2) = \Im (m_2 m_3^*, m_3 m_1^*, m_1 m_2^*)^T$$

where  $\times$  denotes the cross product.

REMARK 14. *Let us remark that the vector  $X(\psi_1, \psi_2)$  is invariant under phase changes in the argument, that is  $X(\psi_1, \psi_2) = X(e^{i\beta_1} \psi_1, e^{i\beta_2} \psi_2)$ . Notice however that  $X(\psi_1, \psi_2) = -X(\psi_2, \psi_1)$ .*

DEFINITION 15. *Given a conical intersection  $\bar{\mathbf{u}}$ , the vector field*

$$\mathcal{X}_P(\mathbf{u}) = X(\phi_j(\mathbf{u}), \phi_{j+1}(\mathbf{u})),$$

*defined in  $\omega \setminus \{\bar{\mathbf{u}}\}$ , is called the non-mixing field associated with  $\bar{\mathbf{u}}$ .*

The non-mixing field is smooth in its domain of definition. From (5) and (12), we have  $\langle \phi_j, \dot{\phi}_{j+1} \rangle = 0$  along its integral curves. Moreover, a simple computation leads to

$$(13) \quad X(\psi_1, \psi_2) \cdot \begin{pmatrix} \langle \psi_2, H_1 \psi_2 \rangle - \langle \psi_1, H_1 \psi_1 \rangle \\ \langle \psi_2, H_2 \psi_2 \rangle - \langle \psi_1, H_2 \psi_1 \rangle \\ \langle \psi_2, H_3 \psi_2 \rangle - \langle \psi_1, H_3 \psi_1 \rangle \end{pmatrix} = \frac{1}{2i} \det \mathcal{M}(\psi_1, \psi_2),$$

which implies the following result.

PROPOSITION 16. *Let  $\bar{\mathbf{u}}$  be a conical intersection and  $\gamma(\cdot) \subset \omega \setminus \{\bar{\mathbf{u}}\}$  be an integral curve of the non-mixing field. Then*

$$\frac{d}{dt} [\lambda_{j+1}(\gamma(t)) - \lambda_j(\gamma(t))] = \frac{1}{2i} F(\gamma(t)),$$

*where  $F(\cdot)$  is defined in (8). In particular, all the integral curves of the non-mixing field starting from a punctured neighborhood of the conical intersection reach it in finite time (up to a time reversal).*

Without loss of generality, we can assume that  $\bar{\mathbf{u}} = 0$ . Denote every  $\mathbf{u} \in \mathbb{R}^3$  by  $\mathbf{u} = \rho \mathbf{v}$ , where  $\rho$  is its module and  $\mathbf{v}$  the versor. Let us call  $\mu(\rho, \mathbf{v}) = \mathbf{m}(\phi_j(\rho, \mathbf{v}), \phi_{j+1}(\rho, \mathbf{v}))$ , where  $\phi_j(\rho, \mathbf{v}), \phi_{j+1}(\rho, \mathbf{v})$  denotes a choice of the eigenstates relative respectively to  $\lambda_j(\rho \mathbf{v})$  and  $\lambda_{j+1}(\rho \mathbf{v})$  (extended by continuity for  $\rho = 0$ ). We remark that  $\mu(\rho, \mathbf{v})$  is defined up to a phase. Finally, set  $\mathcal{X}_\mu(\rho, \mathbf{v}) = \mathcal{X}_P(\rho \mathbf{v})$ . Notice that, for different values of  $\mathbf{v}$ ,  $\mathcal{X}_\mu(\rho, \mathbf{v})$  has a priori different limits as  $\rho$  tends to 0.

The following estimates hold.

542 LEMMA 17. Assume that  $\bar{\mathbf{u}} = 0$  is a conical intersection. Then the inequality  
 543  $|\mu(\rho, \mathbf{v}) \cdot \mathbf{v}| \leq C\rho$  holds in a neighborhood of 0 for some constant  $C > 0$  uniform with  
 544 respect to  $\mathbf{v}$ .

545 *Proof.* Up to shifting  $H_0$  by a multiple of the identity we may assume  $\lambda_j(0) =$   
 546  $\lambda_{j+1}(0) = 0$ . Then

$$547 \quad \rho\mu(\rho, \mathbf{v}) \cdot \mathbf{v} = \rho \langle \phi_j(\rho, \mathbf{v}), (v_1 H_1 + v_2 H_2 + v_3 H_3) \phi_{j+1}(\rho, \mathbf{v}) \rangle$$

$$548 \quad = -\langle \phi_j(\rho, \mathbf{v}), H_0 \phi_{j+1}(\rho, \mathbf{v}) \rangle = -\langle \phi_j(\rho, \mathbf{v}) - \mathbf{P}_0 \phi_j(\rho, \mathbf{v}), H_0 \phi_{j+1}(\rho, \mathbf{v}) \rangle.$$

550 Assumption **(H0)** implies that

$$551 \quad \|H_0 \phi_{j+1}(\rho, \mathbf{v})\| \leq |\lambda_{j+1}(\rho \mathbf{v})| + \rho \sum_{i=1}^3 \|H_i \phi_{j+1}(\rho, \mathbf{v})\| \leq c\rho,$$

552 for some  $c > 0$ , locally around the intersection. By smoothness of the projection, we  
 553 get that

$$554 \quad |\langle \phi_j(\rho, \mathbf{v}) - \mathbf{P}_0 \phi_j(\rho, \mathbf{v}), H_0 \phi_{j+1}(\rho, \mathbf{v}) \rangle| \leq C\rho^2,$$

555 for a suitable  $C > 0$ , hence we get the thesis.  $\square$

556 We are now ready to prove the following result, which provides some information  
 557 on the behavior of the trajectories of the non-mixing field.

558 PROPOSITION 18. With the notations introduced above and for  $\rho$  small enough,  
 559 there exist three constants  $c_1, c_2, c_3 > 0$  such that  $c_1 \leq |\dot{\rho}| \leq c_2$  and  $|\dot{\mathbf{v}}| \leq c_3$  along the  
 560 trajectories of the non-mixing field.

561 *Proof.* Direct computations lead to the equations

$$562 \quad \dot{\rho} = \mathcal{X}_\mu \cdot \mathbf{v}, \quad \dot{\mathbf{v}} = \frac{1}{\rho} (\mathcal{X}_\mu - (\mathcal{X}_\mu \cdot \mathbf{v}) \mathbf{v}).$$

563 The upper bound for  $|\dot{\rho}|$  comes easily from the  $H(\mathbf{u})$ -boundedness of  $H_i$  for every  $i$ .

564 From (13) we get that  $|\mathcal{X}_\mu| \geq c$  for some  $c > 0$  in a neighborhood of the singularity,  
 565 which implies that the sinus of the angle between  $\Im \mathbf{m}(\mu(\rho, \mathbf{v}))$  and  $\Re \mathbf{e}(\mu(\rho, \mathbf{v}))$  is  
 566 uniformly far from zero. As a consequence of this fact and of Lemma 17, for any unit  
 567 vector  $\mathbf{z}$  in the plane spanned by  $\Im \mathbf{m}(\mu(\rho, \mathbf{v}))$  and  $\Re \mathbf{e}(\mu(\rho, \mathbf{v}))$  one has  $|\mathbf{z} \cdot \mathbf{v}| \leq \hat{c}\rho$  for  
 568 some  $\hat{c} > 0$  and  $\rho$  small enough. The orthogonal projection of  $\mathbf{v}$  on that plane may  
 569 be written as  $(\mathbf{v} \cdot \hat{\mathbf{v}})\hat{\mathbf{v}}$ , where  $\hat{\mathbf{v}}$  is a unit vector. Then

$$570 \quad |\mathcal{X}_\mu \cdot \mathbf{v}| = |\mathcal{X}_\mu \cdot (\mathbf{v} - (\mathbf{v} \cdot \hat{\mathbf{v}})\hat{\mathbf{v}})| = \sqrt{1 - (\mathbf{v} \cdot \hat{\mathbf{v}})^2} |\mathcal{X}_\mu| \geq (1 - \hat{c}^2 \rho^2) |\mathcal{X}_\mu|$$

571 and

$$572 \quad |\mathcal{X}_\mu - (\mathcal{X}_\mu \cdot \mathbf{v}) \mathbf{v}|^2 = |\mathcal{X}_\mu|^2 - (\mathcal{X}_\mu \cdot \mathbf{v})^2 \leq 2\hat{c}^2 \rho^2 |\mathcal{X}_\mu|^2.$$

573 The thesis follows.  $\square$

574 The following proposition is a generalization of [10, Proposition 5.9] in the three  
 575 dimensional case. The proof follows the same lines, thanks to Proposition 18, and is  
 576 thus omitted.

577 PROPOSITION 19. For every unit vector  $\mathbf{v}$  in  $\mathbb{R}^3$  there exists an integral curve  
 578  $\gamma : [-\eta, 0] \rightarrow \omega$  of  $\mathcal{X}_P$  with  $\gamma(0) = 0$ ,  $\eta > 0$ , such that  $\lim_{t \rightarrow 0^-} \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} = \mathbf{v}$ .



579 Thanks to Proposition 18, the integral curves of the non-mixing field are  $\mathcal{C}^1$  up to  
 580 the singularity included. In particular, they satisfy the hypotheses of Proposition 7  
 581 with  $k = 1$ , so that the projections  $P_j(\mathbf{u})$  and  $P_{j+1}(\mathbf{u})$  on the eigenspaces relative to  
 582 the intersecting eigenvalues are  $\mathcal{C}^1$  along the integral curves of the non-mixing field  
 583 outside the singularity, and can be continuously extended at the singularity. On the  
 584 other hand,  $\mathbf{P}_\mathbf{u}$  is  $\mathcal{C}^1$  along such curves, singularity included.

585 We remark moreover that, if  $\gamma : [t_0, t_1] \rightarrow \mathbb{R}^3$  is an integral curve of the non-mixing  
 586 field such that  $\lambda_j(\gamma(t)) \neq \lambda_{j+1}(\gamma(t))$  for  $t \in [t_0, t_1]$ , by definition of the non-mixing  
 587 field, it holds

$$588 \quad (14) \quad P_j(\gamma(t))\dot{P}_{j+1}(\gamma(t)) = 0 \quad P_{j+1}(\gamma(t))\dot{P}_j(\gamma(t)) = 0 \quad \forall t \in [t_0, t_1].$$

589 We have the following.

590 PROPOSITION 20. *Along every integral curve of the non-mixing field, there is a*  
 591 *choice of the eigenstates relative to the intersecting eigenvalues which is  $\mathcal{C}^1$  up to the*  
 592 *singularity included.*

593 *Proof.* Let  $\gamma : [-T, 0] \rightarrow \mathbb{R}^3$  be an integral curve of  $\mathcal{X}_P$  such that  $\gamma(0) = \bar{\mathbf{u}}$  is  
 594 a conical intersection between  $\lambda_j$  and  $\lambda_{j+1}$ . Outside the singularity, the eigenstates  
 595 are well defined, up to a phase. To fix the phase, we set  $\psi_j(t) = \frac{P_j(\gamma(t))\hat{\psi}}{\|P_j(\gamma(t))\hat{\psi}\|}$ , for  
 596 some  $\hat{\psi} \in \mathcal{H}$  satisfying  $P_j(\gamma(t))\hat{\psi} \neq 0$  on  $[0, T]$  (up to reducing  $T$ ). Thus  $\psi_j(t)$  is a  
 597 normalized eigenstate of  $H(\gamma(t))$  relative to  $\lambda_j(\gamma(t))$ . In order to prove that  $\psi_j(t)$  is  
 598  $\mathcal{C}^1$  it is enough to prove that  $P_j(\gamma(t))$  is. Since  $P_j(\gamma(t)) + P_{j+1}(\gamma(t)) = \mathbf{P}_{\gamma(t)}$  for  $t \in$   
 599  $[-T, 0)$ , and by (14), we get that  $\mathbf{P}_{\gamma(t)}\dot{P}_j(\gamma(t)) = P_j(\gamma(t))\dot{P}_j(\gamma(t)) = P_j(\gamma(t))\dot{\mathbf{P}}_{\gamma(t)}$ .  
 600 Therefore

$$601 \quad \dot{P}_j(\gamma(t)) = \dot{\mathbf{P}}_{\gamma(t)}P_j(\gamma(t)) + \mathbf{P}_{\gamma(t)}\dot{P}_j(\gamma(t)) = \dot{\mathbf{P}}_{\gamma(t)}P_j(\gamma(t)) + P_j(\gamma(t))\dot{\mathbf{P}}_{\gamma(t)},$$

602 where the right hand side has limit for  $t \rightarrow 0^-$ .

603 We can repeat the same procedure to show that there is a choice for  $\psi_{j+1}(t)$  such  
 604 that  $\dot{\psi}_{j+1}(t)$  has limit for  $t \rightarrow 0^-$ .  $\square$

605 A suitable improvement of the regularity properties provided by Proposition 20  
 606 is actually needed in order to apply Theorem 6 along the integral curves of the non-  
 607 mixing field, around conical intersections. This is achieved by the next result.

608 PROPOSITION 21. *In a neighborhood of a conical intersection, the integral curves*  
 609 *of its associated non-mixing field are  $\mathcal{C}^\infty$  up to the singularity included. Moreover, we*  
 610 *can choose  $\mathcal{C}^\infty$  eigenstates  $\psi_j, \psi_{j+1}$  along such a curve, up to the singularity included.*

611 *Proof.* Let  $\gamma : [-T, 0] \rightarrow \mathbb{R}^3$  be an integral curve of  $\mathcal{X}_P$  such that  $\gamma(0) = \bar{\mathbf{u}}$  and  
 612  $|\dot{\gamma}(t)| > 0$  for every  $t \in [-T, 0]$  (this is true up to choosing  $T$  sufficiently small), and  
 613 define the eigenstates  $\psi_j(\cdot), \psi_{j+1}(\cdot)$  as in Proposition 20.

614 For simplicity, we denote by  $\lambda_j(t), \lambda_{j+1}(t)$  and  $P_j(t), P_{j+1}(t)$  the  $j$ -th and  $(j+1)$ -th  
 615 eigenvalues and the corresponding spectral projections at  $\gamma(t)$ . We prove by induction  
 616 that, for every positive integer  $h$ ,  $\gamma(\cdot), \lambda_l(\cdot), P_l(\cdot)$  and  $H_i P_l(\cdot)$  are  $\mathcal{C}^h$  on  $[-T, 0]$ , where  
 617  $l = j, j+1$  and  $i = 1, 2, 3$ . The proposition then follows immediately.

618 For every non-negative integer  $h$ , whenever the  $h$ -th order derivatives of  $P_l(\cdot)$  and  
 619  $H_i P_l(\cdot)$  are well defined, the following inequality holds

$$620 \quad \|H_i(P_l^{(h)}(s) - P_l^{(h)}(t))\| \leq \|H_i\|_{H(\gamma(t))} (\|H(\gamma(s))P_l^{(h)}(s) - H(\gamma(t))P_l^{(h)}(t)\| \\
 621 \quad + \sum_i |\gamma_i(s) - \gamma_i(t)| \|H_i P_l^{(h)}(s)\| + \|P_l^{(h)}(s) - P_l^{(h)}(t)\|). \\
 622$$

623 In particular, thanks to Proposition 20 and since  $H_i$  is  $H(\gamma(s))$ -bounded (uniformly  
 624 with respect to  $s$ ), this yields the continuity of  $H_i P_l(\cdot)$  on  $[-T, 0]$  and the initialization  
 625 step of the induction. Let us now take  $h > 0$ ; we prove that, if  $\lambda_l(\cdot), P_l(\cdot)$  are  $\mathcal{C}^h$  and  
 626  $H_i P_l(\cdot)$  is  $\mathcal{C}^{h-1}$ , then  $H_i P_l^{(h)}(\cdot)$  is continuous. Indeed, by differentiating  $h$  times  
 627 the equality  $H(\gamma(\cdot))P_l(\cdot) = \lambda_l(\cdot)P_l(\cdot)$  and exploiting the regularity assumptions on  
 628  $\lambda_l(\cdot), P_l(\cdot)$  and  $H_i P_l(\cdot)$  we get that  $H(\gamma(\cdot))P_l^{(h)}(\cdot)$  is continuous in  $[-T, 0]$ . Since  
 629  $H_i$  is  $H(\gamma(s))$ -bounded we deduce that  $\|H_i P_l^{(h)}(s)\|$  is uniformly bounded on  $[-T, 0]$ .  
 630 Then, the continuity of  $H_i P_l^{(h)}(\cdot)$  follows from the above inequality.

631 Assume now that  $\gamma(\cdot), \lambda_l(\cdot), P_l(\cdot)$  and  $H_i P_l(\cdot)$  have been proved to be  $\mathcal{C}^h$ , for  
 632  $h \leq k - 1$ . Then, by definition of  $\mathcal{X}_{\mathcal{P}}$  it turns out that  $\gamma(\cdot)$  is  $\mathcal{C}^k$  in  $[-T, 0]$  and,  
 633 applying (4), we obtain that  $\lambda_l(\cdot)$  is  $\mathcal{C}^k$  in  $[-T, 0]$  too. Let us show that  $P_l(\cdot)$  is  $\mathcal{C}^k$ .  
 634 Note that  $P_l(\cdot) = \mathbf{P}_{\gamma(\cdot)} P_l(\cdot)$ , where  $\mathbf{P}_{\gamma(\cdot)}$  is  $\mathcal{C}^k$  in  $[-T, 0]$ , because of the regularity of  
 635  $\gamma(\cdot)$ . Then, by inductive hypothesis,  $P_l(\cdot)$  is  $\mathcal{C}^k$  if and only if  $\mathbf{P}_{\gamma(\cdot)} P_l^{(k)}(\cdot)$  is continuous  
 636 in  $[-T, 0]$ . We can develop  $\mathbf{P}_{\gamma(t)} P_j^{(k)}(\gamma(t))$  as

$$637 \quad \mathbf{P}_{\gamma(t)} P_j^{(k)}(\gamma(t)) = P_j(t) \mathbf{P}_{\gamma(t)}^{(k)} - P_j(t) P_{j+1}^{(k)}(t) + P_{j+1}(t) P_j^{(k)}(t).$$

638 The first term in the right-hand side is continuous in  $[-T, 0]$ . By (14), it follows that  
 639  $\frac{d^{k-1}}{dt^{k-1}} \left( P_j(t) \dot{P}_{j+1}(t) \right) \equiv 0$ . Then, by inductive hypothesis, we get that  $P_j(t) P_{j+1}^{(k)}(t)$   
 640 is continuous in  $[-T, 0]$ . By a symmetric argument,  $P_{j+1}(t) \dot{P}_j^{(k)}(t)$  is continuous  
 641 in  $[-T, 0]$ . We deduce that  $\mathbf{P}_{\gamma(t)} P_j^{(k)}(\gamma(t))$  is continuous in  $[-T, 0]$ , and so is  $P_j^{(k)}(\cdot)$   
 642 (and, by symmetry,  $P_{j+1}^{(k)}(\cdot)$ ). Since, as shown above,  $H_i P_l^{(k)}(\cdot)$  is continuous in  $[-T, 0]$   
 643 for  $l = j, j + 1$  and  $i = 1, 2, 3$ , the proof of the proposition is concluded.  $\square$

644 We are now ready to state the main result of this section, namely we show how  
 645 the curves tangent to the non-mixing field allow to improve the performances of the  
 646 control algorithm presented in Section 3.2.

647 **THEOREM 22.** *Let  $H(\mathbf{u}) = H_0 + u_1 H_1 + u_2 H_2 + u_3 H_3$  satisfy hypotheses **(H0)**.  
 648 Assume that  $(\omega, \{\lambda_0, \dots, \lambda_k\})$  is a separated discrete spectrum for  $H(\mathbf{u})$  and that there  
 649 exist conical intersections  $\bar{\mathbf{u}}_j \in \omega$ ,  $j = 0, \dots, k - 1$ , between the eigenvalues  $\lambda_j, \lambda_{j+1}$ ,  
 650 with  $\lambda_l(\bar{\mathbf{u}}_j)$  simple if  $l \neq j, j + 1$ . Then, for every  $\mathbf{u}^s$  and  $\mathbf{u}^f$  such that the eigenvalues  
 651  $\lambda_l$ ,  $l = 0, \dots, k$  are non degenerate at  $\mathbf{u}^s$  and  $\mathbf{u}^f$ , for every  $\bar{\phi} \in \{\phi_0(\mathbf{u}^s), \dots, \phi_k(\mathbf{u}^s)\}$ ,  
 652 and  $p \in [0, 1]^{k+1}$  such that  $\sum_{j=0}^k p_j^2 = 1$ , there exist  $C > 0$  and a continuous control  
 653  $\gamma(\cdot) : [0, 1] \rightarrow \mathbb{R}^m$  with  $\gamma(0) = \mathbf{u}^s$  and  $\gamma(1) = \mathbf{u}^f$ , such that for every  $\varepsilon > 0$*

$$654 \quad (15) \quad \|\psi(1/\varepsilon) - \sum_{j=0}^k p_j e^{i\vartheta_j} \phi_j(\mathbf{u}^f)\| \leq C\varepsilon,$$

655 where  $\psi(\cdot)$  is the solution of (3) with  $\psi(0) = \bar{\phi}$ ,  $\mathbf{u}(t) = \gamma(\varepsilon t)$ , and  $\vartheta_0, \dots, \vartheta_k \in \mathbb{R}$  are  
 656 some phases depending on  $\varepsilon$  and  $\gamma$ .

657 *Proof.* The strategy is analogous to the one presented in Section 3.2 and is based  
 658 on the construction of a suitable piecewise smooth path joining  $\mathbf{u}^s$  with  $\mathbf{u}^f$  that  
 659 passes through all the conical intersection; in particular, we assume that the path  
 660  $\gamma : [0, 1] \rightarrow \omega$  satisfies  $\gamma(0) = \mathbf{u}^s$ ,  $\gamma(1) = \mathbf{u}^f$  and  $\gamma(\tau_j) = \bar{\mathbf{u}}_j$ ,  $j = 0, \dots, k - 1$ , for  
 661 some  $0 < \tau_0 < \dots < \tau_{k-1} < 1$ . The only difference concerns the construction of  
 662 the path in the neighborhoods of the conical intersections: indeed, in these regions  
 663 the path is chosen to be tangent to the non-mixing field. At the intersection, the

664 inner and outer directions are selected according to Proposition 13, as explained  
 665 in Section 3.2, and the existence of corresponding trajectories tangent to the non-  
 666 mixing field is guaranteed by Proposition 19. Far from all the conical intersections the  
 667 evolution of  $H(\gamma(\varepsilon t))$  conserves the occupation probabilities relative to each energy  
 668 level  $\lambda_l$ ,  $l = 0, \dots, k$ , with an approximation of order  $\varepsilon$ . Let us now estimate the  
 669 probability distribution obtained after the passage through the conical intersection  
 670  $\bar{\mathbf{u}}_0$ . Since the path is tangent to the non-mixing field, we can apply Theorem 6 in order  
 671 to study the evolution inside the space  $\mathbf{P}_{\mathbf{u}(t)}\mathcal{H}$ , where  $\mathbf{P}_{\mathbf{u}(t)}$  is the spectral projection  
 672 associated with the two levels  $\{\lambda_0(\mathbf{u}), \lambda_1(\mathbf{u})\}$ . For  $\tau$  in a left neighborhood of  $\tau_0$ , the  
 673 effective Hamiltonian and its associated evolution operator  $U_{\text{eff}}^\varepsilon$  are diagonal, and this  
 674 implies the existence of a phase  $\theta_0$  (depending on  $\varepsilon$ ) and of a positive constant  $C_0$   
 675 such that  $\|\psi(\tau_0/\varepsilon) - e^{i\theta_0}\phi_0(\gamma(\tau_0^-))\| \leq C_0\varepsilon$ , where  $\psi(\cdot)$  is the solution of equation (3)  
 676 with  $\psi(0) = \phi_0(\gamma(0))$  corresponding to the control  $\mathbf{u}(t) = \gamma(\varepsilon t)$ , defined on  $[0, 1/\varepsilon]$ .  
 677 By Proposition 11, this implies that

$$678 \quad \|\psi(\tau_0/\varepsilon) - e^{i\theta_0} \left( \cos \Xi(\mathbf{v}) \phi_0(\gamma(\tau_0^+)) - e^{-i\beta(\mathbf{v})} \sin \Xi(\mathbf{v}) \phi_1(\gamma(\tau_0^+)) \right)\| \leq C_0\varepsilon,$$

679 with  $\Xi(\mathbf{v})$  and  $\beta(\mathbf{v})$  satisfying equations (10)-(11), and  $\mathbf{v}$  is the outer direction.

680 Since the effective Hamiltonian is diagonal also for  $\tau$  belonging to a right neigh-  
 681 borhood of  $\tau_0$ , we conclude that there exist two phases  $\alpha_0$  and  $\alpha_1$  (depending on  $\tau$   
 682 and  $\varepsilon$ ) and a positive constant  $\hat{C}_0$  such that

$$683 \quad \|\psi(\tau/\varepsilon) - e^{i\alpha_0} p_0 \phi_0(\gamma(\tau)) - e^{i\alpha_1} \sqrt{1 - p_0^2} \phi_1(\gamma(\tau))\| \leq \hat{C}_0\varepsilon.$$

684 Since analogous estimates hold for the passages through any other conical intersection  
 685 and outside the corresponding neighborhoods the theorem is proved.  $\square$

686 To conclude this section, we present below a result providing some information  
 687 on the structural stability of conical intersections based on the properties of the non-  
 688 mixing fields.

689 **THEOREM 23.** *Assume that  $H(\mathbf{u}) = H_0 + u_1H_1 + u_2H_2 + u_3H_3$  satisfies **(H0)**  
 690 and admits a separated discrete spectrum  $(\omega, \{\lambda_j, \lambda_{j+1}\})$ . Let  $\bar{\mathbf{u}} \in \omega$  be a conical  
 691 intersection for  $H(\mathbf{u})$  between the eigenvalues  $\lambda_j$  and  $\lambda_{j+1}$ . Then for every  $\varepsilon > 0$   
 692 there exists  $\delta > 0$  such that, if  $\hat{H}(\mathbf{u}) = \hat{H}_0 + u_1\hat{H}_1 + u_2\hat{H}_2 + u_3\hat{H}_3$  satisfies **(H0)** and*

$$693 \quad \sum_{i=0}^3 \|\hat{H}_i - H_i\|_{H_0} \leq \delta,$$

694 then the operator  $\hat{H}(\mathbf{u})$  has a separated discrete spectrum  $(\omega, \{\hat{\lambda}_j, \hat{\lambda}_{j+1}\})$  and it pos-  
 695 sesses a conical intersection of eigenvalues at  $\hat{\mathbf{u}}$ , with  $|\bar{\mathbf{u}} - \hat{\mathbf{u}}| \leq \varepsilon$ .

696 *Proof.* First of all, by equivalence of all norms  $\|\cdot\|_{H(\mathbf{u})}$ , without loss of generality  
 697 we can assume that  $\bar{\mathbf{u}} = 0$ . We notice that our assumptions guarantee that in a  
 698 neighborhood of the conical intersection the eigenvalues  $\lambda_j$  and  $\lambda_{j+1}$  are well sepa-  
 699 rated from the rest of the spectrum. Continuous dependence of the eigenvalues with  
 700 respect to perturbations of the Hamiltonian (see Lemma 28) ensures that, if  $\delta$  is small,  
 701 then  $\hat{H}(\cdot)$  admits two eigenvalues  $\hat{\lambda}_j, \hat{\lambda}_{j+1}$  close to  $\lambda_j, \lambda_{j+1}$ . Moreover  $\{\hat{\lambda}_j, \hat{\lambda}_{j+1}\}$  is  
 702 separated from the rest of the spectrum, locally around  $\bar{\mathbf{u}}$ . From the conicity of the  
 703 intersection between  $\lambda_j$  and  $\lambda_{j+1}$ , there exists  $\varepsilon > 0$  small enough such that  $|F(\mathbf{u})| \geq c$   
 704 for some  $c > 0$  on  $B(\bar{\mathbf{u}}, \varepsilon)$  and, moreover, by Proposition 18 the vector field  $\mathcal{X}_P$  (up

705 to the sign) points inside the ball  $B(\bar{\mathbf{u}}, \varepsilon)$  at every point of its boundary. If  $\delta$  is small  
 706 enough then  $\hat{\lambda}_j \neq \hat{\lambda}_{j+1}$  on  $\partial B(\bar{\mathbf{u}}, \varepsilon)$  and the gap between the two eigenvalues can be  
 707 assumed to be of order  $\varepsilon$ . Therefore we can define the conicity matrix  $\hat{\mathcal{M}}$  associated  
 708 with  $\hat{H}(\cdot)$  and the function  $\hat{F}(\mathbf{u}) = \det \hat{\mathcal{M}}(\psi_j(\mathbf{u}), \psi_{j+1}(\mathbf{u}))$ , where  $\{\psi_j(\mathbf{u}), \psi_{j+1}(\mathbf{u})\}$   
 709 is an orthonormal basis for the sum of eigenspaces relative to  $\{\hat{\lambda}_j(\mathbf{u}), \hat{\lambda}_{j+1}(\mathbf{u})\}$ . Since  
 710 the conicity matrix depends continuously on the control operators, with respect to  
 711 the norm  $\|\cdot\|_{H_0}$ , and as a consequence of Lemma 9 and Proposition 26, we can take  
 712  $\delta$  small enough such that  $|\hat{F}(\mathbf{u})| \geq c/2$  on  $B(\bar{\mathbf{u}}, \varepsilon)$ . This allows us to define, whenever  
 713  $\hat{\lambda}_j \neq \hat{\lambda}_{j+1}$ , the non-mixing field  $\hat{\mathcal{X}}_{\mathcal{P}}$  associated with  $\hat{H}(\cdot)$  and corresponding to the  
 714 band  $\{\hat{\lambda}_j, \hat{\lambda}_{j+1}\}$ ; thanks to Proposition 16, up to a time reversal the time derivative  
 715 of  $\hat{\lambda}_{j+1} - \hat{\lambda}_j$  along the integral curves of  $\hat{\mathcal{X}}_{\mathcal{P}}$  is smaller than  $-c/4$ . By Corollary 27  
 716 if  $\delta$  is small enough, then  $\hat{\mathcal{X}}_{\mathcal{P}}$  points inside  $B(\bar{\mathbf{u}}, \varepsilon)$  at every point of  $\partial B(\bar{\mathbf{u}}, \varepsilon)$ . Any  
 717 trajectory  $\hat{\gamma}(\cdot)$  of  $\hat{\mathcal{X}}_{\mathcal{P}}$  starting from  $B(\bar{\mathbf{u}}, \varepsilon)$  remains inside  $B(\bar{\mathbf{u}}, \varepsilon)$  in its interval of  
 718 definition and reaches in finite time a point  $\hat{\mathbf{u}}$  corresponding to a double eigenvalue  
 719  $\hat{\lambda}_j(\hat{\mathbf{u}}) = \hat{\lambda}_{j+1}(\hat{\mathbf{u}})$ . The conclusion follows from Proposition 10.  $\square$

720 **5. Examples.** Let us first consider a finite dimensional example; let  $H(\mathbf{u})$  be  
 721 the Hamiltonian in  $iu(3)$  defined by

$$722 \quad (16) \quad H_0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad H_1 = \begin{pmatrix} 1 & i & 0 \\ -i & 0 & 1 \\ 0 & 1 & -1 \end{pmatrix}, \quad H_2 = \begin{pmatrix} 0 & 0 & i \\ 0 & 1 & 0 \\ -i & 0 & 0 \end{pmatrix}, \quad H_3 = \begin{pmatrix} -1 & 1 & -1 \\ 1 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}.$$

723 The Hamiltonian  $H(\cdot)$  admits a double eigenvalue at  $\mathbf{u} = 0$  corresponding to the  
 724 two lowest levels. A simple computation leads to  $\det \mathcal{M}(e_1, e_2) = -2i$  where  $e_1 =$   
 725  $(1, 0, 0)^T, e_2 = (0, 1, 0)^T$  form a basis of the double eigenspace at 0. Thus the eigen-  
 726 value intersection is conical. Moreover, at the point  $\mathbf{u} = (1/2, -1/4, 0)$ , the second and  
 727 third eigenvalues degenerate, and the intersection is conical: indeed, the determinant  
 728 of the conicity matrix corresponding to these two levels is equal to  $4i/3$ .

729 Figure 2 describes the behavior of the trajectories of the non-mixing field relative  
 730 to the first two eigenvalues of  $H(\cdot)$ . Consistently with the results shown above, the  
 731 flow corresponding to the non-mixing vector field allows to identify two conical inter-  
 732 sections, one of them being the origin, among the first two levels. In particular, the  
 trajectories converge or diverge from them, locally.




FIG. 2. Trajectories of the non-mixing field relative to the two lowest eigenvalues for the Hamiltonian corresponding to (16)

733

734 Let us consider the case in which the initial value of the control is  $\mathbf{u}_0 = (-1, 0, 0)$ .  
 735 We want to approximately send an initial state  $\psi$  concentrated in the lowest energy  
 736 level into a state whose probability distribution with respect to the three states is

737 (1/3, 1/6, 1/2). To do that, we first pass through the conical intersection between the  
 738 two lowest levels, leaving a probability of 1/3 on the lowest one and sending 2/3 of  
 739 probability to the other; afterwards, we pass through the conical intersection between  
 740 the two highest levels in order to send 1/2 of the total probability to the third level;  
 741 finally, we come back to the initial point.

742 Figure 3, on the left, shows the chosen path, which follows the non-mixing field in a  
 743 neighborhood of the conical intersection. On the right the distribution of probability  
 744 is depicted. The simulations were run on arc-length parametrized curves, with a  
 745 parameter  $\varepsilon = 0.001$ .




FIG. 3. Control path and probability distributions for the Hamiltonian corresponding to (16). We use black for the probability of being in the second energy state, and the lighter shade of gray for the probability of being in the highest one. The starting point is labeled as  $O$  and the conical intersections  $(0, 0, 0)$  and  $(1/2, -1/4, 0)$  respectively as  $P_1, P_2$ .

746 Consider now the Hamiltonian  $H(\mathbf{u}) = -\Delta + u_1V_1 + u_2V_2 - iu_3(\nabla\mathbf{A} + \mathbf{A}\nabla)$ ,  
 747 where

$$748 \quad V_1(x) = x_2^2 + x_3^2, \quad V_2(x) = x_2x_3, \quad A = (0, -x_3/2, x_2/2)^T,$$

$$749 \quad x = (x_1, x_2, x_3) \in \Omega = (0, 1) \times (0, \sqrt{3}) \times (0, \sqrt{5})$$

751 with zero Dirichlet boundary conditions. We recall that the third controlled operator  
 752 acts on the elements of its domain as follows  $-i(\nabla\mathbf{A} + \mathbf{A}\nabla)\psi = -i\mathbf{A}\cdot\nabla\psi - i\operatorname{div}(\mathbf{A}\psi)$ .  
 753 This operator has the form of a vector potential coupled with the momentum. By  
 754 classical results (see [26, 27]), it is easy to check that  $H(\mathbf{u})$  satisfies hypothesis **(H0)**  
 755 and has a purely discrete spectrum with a finite number of eigenvalues in each compact  
 756 subset of  $\mathbb{R}$  (this, together with Lemma 28, guarantees that  $H(\mathbf{u})$  possesses a separated  
 757 discrete spectrum).

758 We claim that  $H(0)$  (representing the potential well in  $\Omega$ ) admits conical inter-  
 759 sections of eigenvalues. The eigenvalues and the eigenfunctions of  $H(0)$  take the form

$$760 \quad \lambda_{j_1, j_2, j_3} = \pi^2 \left( j_1^2 + \frac{j_2^2}{3} + \frac{j_3^2}{5} \right), \quad \psi_{j_1, j_2, j_3}(x) = \frac{2\sqrt{2}}{\sqrt[4]{15}} \sin(j_1\pi x_1) \sin\left(\frac{j_2\pi x_2}{\sqrt{3}}\right) \sin\left(\frac{j_3\pi x_3}{\sqrt{5}}\right)$$

761 where  $j_1, j_2, j_3$  are strictly positive integers. In particular it is easy to check that  
 762 the fourth and the fifth lowest eigenvalues take the same value  $\lambda_{1,1,3} = \lambda_{1,2,2}$  at the  
 763 origin, and, since the determinant of the conicity matrix is approximately  $0.029i$ , we  
 764 deduce that the intersection is conical.

765 Let us consider an initial control  $\mathbf{u}_0 = (1, 0, 0)$  and an initial state concentrated  
 766 on the fourth energy level of  $H(\mathbf{u}_0)$ . Our aim is to induce an approximate transition

767 to a state distributed between the fourth and fifth level of  $H(\mathbf{u}_0)$  with probability  $2/3$   
 768 and  $1/3$ , respectively. We follow arc-length parametrized curves with a parameter  $\varepsilon =$   
 769  $0.003$ , and we compare the transition obtained following control paths with a different  
 770 behavior around the singularity. Namely, we consider a path which is locally an  
 771 integral curve of the non-mixing field (top-left in Figure 4) and one which is piecewise  
 772 affine in the sense of Proposition 11 (bottom-left in Figure 4). The corresponding  
 773 probability distributions as functions of the time are shown in Figure 4, on the right  
 774 side. Both paths induce quite precisely the desired transition, although for long times  
 775 the accuracy degrades. We remark that the most significant difference between the  
 776 two strategies is given by the fact that the second one induces a less accurate spread  
 777 of probability at the passage through the conical intersection, as expected.

778 Note that, to compute the evolution of the controlled system, we truncated the  
 779 infinite dimensional system to the lowest 20 and to the lowest 30 energy levels. These  
 780 two choices did not exhibit significant differences: this suggests that the truncation  
 781 up to the first 20 energy levels well describes the behavior of the infinite dimensional  
 782 system. Figure 4 is obtained choosing this truncation.




FIG. 4. Control paths and probability distributions for  $H(\mathbf{u}) = -\Delta + u_1V_1 + u_2V_2 - iu_3(\nabla\mathbf{A} + \mathbf{A}\nabla)$ . We reproduce in gray the probability of being in the fourth energy level and in black the probability of being in the fifth one. The starting point is labeled as  $O$  and the conical intersection as  $P$ .

783 **6. Final remarks.** The integral curves of the non-mixing field are not the only  
 784 paths guaranteeing the estimate (15): suitable approximations (in the sense precised  
 785 just below) of these curves also ensure an adiabatic approximation of order  $\varepsilon$ . Indeed,  
 786 let us consider an arc-length parametrized curve  $\gamma^P(\tau)$ , tangent to the non-mixing  
 787 field, that reaches a conical intersection between  $\lambda_j$  and  $\lambda_{j+1}$  at time  $\tau = 0$ , and  
 788 let  $\gamma$  be a  $C^3$  curve such that  $|\gamma(\tau) - \gamma^P(\tau)| \leq C\tau^3$ , for  $\tau$  small enough and some  
 789 positive constant  $C$ . Let  $t = \tau/\varepsilon$  and consider the effective Hamiltonians evaluated  
 790 along the two curves. It is then easy to see, by simple computations and because of  
 791 Proposition 7, that the difference between the two effective Hamiltonians is less or  
 792 equal than  $C'\varepsilon^2t$  for some  $C' > 0$ . This term, integrated over a time interval of order

793  $1/\varepsilon$ , gives a difference of order  $\varepsilon$ .

794 An interesting controllability problem alternative to the one introduced in Sec-  
 795 tion 3.2 aims at sending (approximately) an initial state  $\psi^s = \sum_{j=0}^k c_j \phi_j(\mathbf{u}^s)$  to a  
 796 final state concentrated in a single energy level. This problem may appear completely  
 797 equivalent to the previous one, but is actually more delicate. Indeed, a natural way to  
 798 induce the desired transition would be to run backward in time a path constructed as  
 799 in Section 3.2. However, a simple computation shows that, at each passage through  
 800 a conical intersection, the components corresponding to the intersecting eigenvalues  
 801 recombine in a concentrated state only if their relative phase coincides with the one  
 802 induced by the unitary transformation of the limit basis (that is, the phase  $\beta(\mathbf{v})$  of  
 803 Proposition 11). On the other hand, the computation of dynamical phases, coming  
 804 from the integration of the energy on intervals whose length is of order  $1/\varepsilon$ , is very  
 805 sensitive to changes in the speed  $\varepsilon$ . This is in principle possible, but it compromises  
 806 the constructiveness of the algorithm.

807 Similarly, by means of non-constructive arguments (for instance, by exploiting  
 808 the rational independence of the gaps between the eigenvalues underlined in [12,  
 809 Lemma 14]) one can obtain the following controllability property, stronger than the  
 810 one considered in Section 3.2.

811 *Under assumption (H0), assuming that the Hamiltonian possesses a separated*  
 812 *discrete spectrum in which all energy levels are connected through conical intersections,*  
 813 *and for any given initial and target states  $\psi^s, \psi^f$  distributed in  $\Sigma$  and  $\eta > 0$ , there*  
 814 *exists a control input steering the system from  $\psi^s$  to a final state whose distance from*  
 815  *$\psi^f$  is less than  $\eta$ .*

816 It is opinion of the authors that all the results here above still hold in a non-linear  
 817 sufficiently smooth setting, that is for Hamiltonians of the form  $H(\mathbf{u})$  whose deriva-  
 818 tives with respect to the parameter  $\mathbf{u}$  are  $H(0)$ -small up to a suitable order and under  
 819 hypothesis (H1). This case is interesting, since it covers relevant physical models,  
 820 such as those described by Hamiltonians with controlled electromagnetic potentials.  
 821 Preliminary results in this sense have been obtained in [14].

## 822 Appendix A. Regularity properties and proof of Proposition 7.

823 Let  $\mathcal{H}$  be a complex separable Hilbert space; all operators in the following are  
 824 assumed to be operators on  $\mathcal{H}$ . In this section we derive some regularity results on  
 825 the eigenvalues and the eigenstates of self-adjoint operators with respect to the norm  
 826 defined in (2) (see e.g. [19] for similar regularity properties). Such results will be used  
 827 in particular to prove Proposition 7.

828 The resolvent of  $A$  in  $\zeta \in \rho(A)$  is denoted by  $R(A, \zeta) = (A - \zeta \text{id})^{-1}$ ; we recall  
 829 that it is a bounded linear operator that maps  $\mathcal{H}$  into  $\mathcal{D}(A)$ , and that, given two self-  
 830 adjoint operators  $A_1, A_2$  with the same domain, their resolvents satisfy the Second  
 831 Resolvent Identity

$$832 \quad (17) \quad R(A_2, \zeta) - R(A_1, \zeta) = R(A_1, \zeta)(A_1 - A_2)R(A_2, \zeta).$$

833 The identity

$$834 \quad (18) \quad \|(X - \zeta \text{id})^{-1}\| = d(\zeta, \sigma(X))^{-1}$$

835 holds for self-adjoint operators (see e.g. [19]).

836 Let us state the following technical lemma, which will be largely used in the  
 837 following. Its proof easily comes from the definition of  $\|\cdot\|_A$  and is thus omitted.



838 LEMMA 24. Let  $A, B$  be self-adjoint operators with  $B$   $A$ -bounded and  $\zeta \in \rho(A)$ .  
 839 Then the following inequality holds:

$$840 \quad (19) \quad \|BR(A, \zeta)\| \leq \left(1 + (|\zeta| + 1) \|R(A, \zeta)\|\right) \|B\|_A.$$

841 The following result shows that the resolvent set for a self-adjoint operator  $A$   
 842 possesses some continuity properties with respect to small perturbation in the space  
 843  $\mathcal{L}(\mathcal{D}(A), \mathcal{H})$ .

844 LEMMA 25. Let  $A_1$  be a self-adjoint operator. Let  $\mathcal{Z} \subset \rho(A_1)$  be a compact set.  
 845 There exists a  $\delta > 0$  such that, if a self-adjoint operator  $A_2$  satisfies  $\|A_1 - A_2\|_{A_1} \leq \delta$ ,  
 846 then  $A_1$  and  $A_2$  have the same domain,  $\mathcal{Z} \subset \rho(A_2)$  and the inequality

$$847 \quad (20) \quad \|R(A_2, \zeta) - R(A_1, \zeta)\| \leq C \|A_1 - A_2\|_{A_1}$$

848 holds true on  $\mathcal{Z}$  for some constant  $C$  depending on  $\mathcal{Z}$  and  $A_1$ .

849 *Proof.* To ensure that  $\mathcal{D}(A_1) = \mathcal{D}(A_2)$  it is clearly enough to assume  $\delta < 1$ . To  
 850 conclude the proof we proceed as follows. As a consequence of (18),  $\|R(A_1, \zeta)\|$  is  
 851 uniformly bounded on  $\mathcal{Z}$ . This, together with (19), implies that  $\|(A_2 - A_1)R(A_1, \zeta)\| <$   
 852  $C' \|A_1 - A_2\|_{A_1}$  for some  $C' > 0$ . Since

$$853 \quad R(A_2, \zeta) = R(A_1, \zeta) (\text{id} + (A_2 - A_1)R(A_1, \zeta))^{-1}$$

854 whenever the right-hand side is well defined, we deduce that  $\mathcal{Z} \subset \rho(A_2)$  and that  
 855  $\|R(A_2, \zeta)\| \leq 2\|R(A_1, \zeta)\|$  on  $\mathcal{Z}$ , provided that  $\delta$  is small enough. Finally, by apply-  
 856 ing (17), we get (20).  $\square$

857 Let  $\Sigma \subset \sigma(A)$  be constituted by a finite number of eigenvalues of the self-adjoint  
 858 operator  $A$ . For every positively-oriented closed path  $\Gamma \subset \mathbb{C}$  encircling  $\Sigma$ , and not  
 859 encircling any other element in  $\sigma(A)$ , the projection  $P$  onto the sum of the eigenspaces  
 860 relative to  $\Sigma$  is given by

$$861 \quad (21) \quad P = -(2\pi i)^{-1} \oint_{\Gamma} R(A, \zeta) d\zeta.$$

862 PROPOSITION 26. Let  $A_1$  be a self-adjoint operator. Assume that  $\Sigma_1 = \sigma(A_1) \cap$   
 863  $(\zeta_1, \zeta_2)$  is constituted by  $k$  eigenvalues, counted with multiplicity, and that  $\zeta_1, \zeta_2 \notin$   
 864  $\sigma(A_1)$ . Then for every  $\epsilon > 0$  there exists a  $\delta > 0$ , depending on  $\zeta_1, \zeta_2$  and  $A_1$ , such  
 865 that if  $A_2$  is self-adjoint and  $\|A_1 - A_2\|_{A_1} \leq \delta$ , then

- 866 *i)*  $\Sigma_2 = \sigma(A_2) \cap (\zeta_1, \zeta_2)$  is constituted by  $k$  eigenvalues, counted with multiplicity;  
 867 *ii)* Calling  $P_{\Sigma_1}^{A_1}$  the spectral projection onto the sum of eigenspaces of  $A_1$  relative  
 868 to  $\Sigma_1$  and  $P_{\Sigma_2}^{A_2}$  the spectral projection onto the sum of eigenspaces of  $A_2$   
 869 relative to  $\Sigma_2$ , it holds  $\|P_{\Sigma_1}^{A_1} - P_{\Sigma_2}^{A_2}\| \leq \epsilon$ .

870 *Proof.* Let  $\Gamma$  be the circle in the complex plane centered on the real axis and pass-  
 871 ing through  $\zeta_1$  and  $\zeta_2$ , and consider the projection  $P_{\Sigma_2}^{A_2} = -(2\pi i)^{-1} \oint_{\Gamma} R(A_2, \zeta) d\zeta$ .  
 872 From (20) we obtain that

$$873 \quad \left\|P_{\Sigma_1}^{A_1} - P_{\Sigma_2}^{A_2}\right\| \leq (2\pi)^{-1} \oint_{\Gamma} \|R(A_1, \zeta) - R(A_2, \zeta)\| d\zeta \leq \hat{C}\delta$$

874 for some  $\hat{C} > 0$  (depending on  $\Gamma$  and  $A_1$ ). In particular,  $\|P_{\Sigma_1}^{A_1} - P_{\Sigma_2}^{A_2}\| < 1$  for  $\delta$  small  
 875 enough, which easily implies that  $\dim \text{Range}(P_{\Sigma_2}^{A_2}) = \dim \text{Range}(P_{\Sigma_1}^{A_1}) = k$  (see [27,  
 876 page 14]), therefore  $\Sigma_2$  contains exactly  $k$  eigenvalues, counted with multiplicity, i.e.  
 877 *i)* holds true. By possibly choosing a smaller  $\delta$ , *ii)* also holds true.  $\square$

878 The following result immediately follows from Proposition 26.

879 **COROLLARY 27.** *Let  $\lambda$  be a simple eigenvalue of a self-adjoint operator  $A_1$ . Then*  
 880 *for every  $\epsilon > 0$  there exists  $\delta > 0$  such that any self-adjoint operator  $A_2$  satisfying*  
 881  *$\|A_1 - A_2\|_{A_1} \leq \delta$  admits a unique eigenvalue  $\mu$  with  $|\lambda - \mu| < \epsilon$ . Moreover, there*  
 882 *exists a choice  $\phi_\lambda^{A_1}, \phi_\mu^{A_2}$  for the corresponding eigenstates such that  $\|\phi_\lambda^{A_1} - \phi_\mu^{A_2}\| \leq \epsilon$ .*

883 The next result provides an estimate concerning regularity properties of the eigen-  
 884 values.

885 **LEMMA 28.** *Let  $A_1$  be a self-adjoint operator such that  $\sigma(A_1) \cap I$  is discrete and*  
 886 *without finite accumulation points for some open, possibly unbounded, interval  $I$ . If*  
 887  *$\delta > 0$  is small enough and  $A_2$  is a self-adjoint operator satisfying  $\|A_2 - A_1\|_{A_1} \leq \delta$ ,*  
 888 *then the eigenvalues of  $A_2$  contained in  $I$  are close to those of  $A_1$ , in the following*  
 889 *sense. Up to appropriately indexing on a subset of  $\mathbb{Z}$  the eigenvalues (counted with*  
 890 *multiplicity) in  $\sigma(A_j) \cap I$ , for  $j = 1, 2$ , and denoting them with  $\mu_i(A_j)$  we have*  
 891  *$|\mu_i(A_1) - \mu_i(A_2)| \leq \epsilon(1 + |\mu_i(A_1)|)$ , where  $\epsilon = e^{\frac{\delta}{1-\delta}} - 1$ .*

892 *Proof.* Let  $A_1$  satisfy the hypotheses of the lemma, and let  $A_2$  be a self-adjoint  
 893 operator with  $\|A_1 - A_2\|_{A_1} \leq \delta$ , where without loss of generality we assume that  
 894  $\delta < 1$ ; define  $A(t) = A_1 + t(A_2 - A_1)$ , for  $t \in [0, 1]$ . Let  $\lambda_i(t)$  be the analytic branch  
 895 of the eigenvalues of  $A(t)$  emanating from  $\lambda_i(0) = \mu_i(A_1)$ , and denote by  $\phi_i(t)$  a  
 896 corresponding analytic eigenstate. By hypothesis

$$897 \quad \|(A_2 - A_1)\phi_i(t)\| \leq \delta\|A_1\phi_i(t)\| + \delta \leq \delta|\lambda_i(t)| + \delta t\|(A_2 - A_1)\phi_i(t)\| + \delta,$$

898 which implies that  $\|(A_2 - A_1)\phi_i(t)\| \leq \frac{\delta}{1-\delta}(|\lambda_i(t)| + 1)$ . From

$$899 \quad |\dot{\lambda}_i(t)| = |\langle \phi_i(t), (A_2 - A_1)\phi_i(t) \rangle| \leq \|(A_2 - A_1)\phi_i(t)\|$$

900 we easily get the thesis.  $\square$

901 Consider a control-dependent Hamiltonian  $H(\mathbf{u})$  satisfying assumption **(H0)**. It  
 902 is easy to see that for any  $\mathbf{u}_1, \mathbf{u}_2$  the norms  $\|\cdot\|_{H(\mathbf{u}_1)}, \|\cdot\|_{H(\mathbf{u}_2)}$  are equivalent, thanks to  
 903 the  $H_0$ -smallness of the control Hamiltonians. From Lemma 28, and the equivalence  
 904 of the norms  $\|\cdot\|_{H(\mathbf{u})}$ , the eigenvalues  $\lambda_i(\cdot)$  of  $H(\cdot)$  are locally Lipschitz, and the  
 905 corresponding Lipschitz constants locally depend on the magnitude of  $\lambda_i(\cdot)$ . Moreover,  
 906 we remark the following fact: let  $\bar{\mathbf{u}}$  be a conical intersection between the eigenvalues  
 907  $\lambda_j$  and  $\lambda_{j+1}$ , that satisfy a gap condition, according to Definition 2. By the definition  
 908 of conical intersection and the Lipschitz continuity of the eigenvalues we can conclude  
 909 that there exist a suitably small neighborhood  $U$  of  $\bar{\mathbf{u}}$  and two constants  $C_1 > 0$  and  
 910  $C_2 > 0$  such that

$$911 \quad (22) \quad \lambda_{j+1}(\mathbf{u}) - \lambda_j(\mathbf{u}) \geq C_1|\mathbf{u} - \bar{\mathbf{u}}| \quad \forall \mathbf{u} \in U$$

912 and

$$913 \quad (23) \quad |\lambda_i(\mathbf{u}) - \lambda_i(\mathbf{u}')| \leq C_2|\mathbf{u} - \mathbf{u}'| \quad \forall \mathbf{u}, \mathbf{u}' \in U, \quad i = j, j + 1.$$

914 We are now ready to prove Proposition 7.

915 *Proof of Proposition 7.* For simplicity, throughout this proof we will write  $R(\mathbf{u}, \xi)$   
 916 to denote the resolvent  $R(H(\mathbf{u}), \xi)$ . The property that the projections  $P_j(\gamma(t))$ ,  $J =$   
 917  $j, j + 1$ , are  $\mathcal{C}^k$  at any  $t \in [-R, 0)$  whenever  $\gamma(\cdot)$  is  $\mathcal{C}^k$  has been shown e.g. in [28]. It  
 918 comes from (21) and a recursive application of the identity

$$919 \quad (24) \quad \frac{d}{dt}R(f(t), \xi) = R(f(t), \xi) \left( \frac{d}{dt}H(f(t)) \right) R(f(t), \xi),$$

920 valid for any differentiable path  $f$  taking values in  $\mathbb{R}^3$ , together with (19).

921 It remains to study the regularity of the projections at the singularity. We first  
 922 consider the case  $k = 1$ . Without loss of generality we assume  $|\dot{\gamma}(0)| = 1$ . Let  
 923  $\rho = C_1/4$ , where  $C_1$  is as in (22), and for every  $t \in [-R, 0)$  consider the circle  
 924  $\Gamma_t \subset \mathbb{C}$  of radius  $\rho t$  centered at  $\lambda_j(\gamma(t))$ . There exists  $0 < T \leq R$  such that for every  
 925  $t \in [-T, 0)$

$$926 \quad |\lambda_{j+1}(\gamma(t)) - \lambda_j(\gamma(t))| \geq \frac{3}{4}C_1 t = 3\rho t$$

927 so that  $|\lambda_{j+1}(\gamma(t)) - \zeta| \geq 2\rho t$  for every  $\zeta \in \Gamma_t$ . Thus  $d(\zeta, \sigma(H(\gamma(t)))) = \rho t$  and, by (23)  
 928 and the definition of  $\ell_1(\cdot)$ ,  $d(\zeta, \sigma(H(\ell_1(t)))) \geq \rho t/2$ , up to reducing  $T$ . Therefore,  
 929 from (18), for  $\zeta \in \Gamma_t$  it holds

$$930 \quad (25) \quad \|R(\gamma(t), \zeta)\| = \frac{1}{\rho t}, \quad \|R(\ell_1(t), \zeta)\| \leq \frac{2}{\rho t}.$$

931 It is left to prove that  $\oint_{\Gamma_t} (R(\gamma(t), \zeta) - R(\ell_1(t), \zeta)) d\zeta$  tends to 0. Estimate (19) gives

$$932 \quad \|(H(\ell_1(t)) - H(\gamma(t)))R(\gamma(t), \zeta)\| \leq C \left| \frac{\ell_1(t) - \gamma(t)}{t} \right|$$

933 for some  $C > 0$ , which, together with (17) and (25) yields the thesis.

934 Let us now tackle the general case; the proof follows similar arguments. We  
 935 define the circuit  $\Gamma_\tau$  as above, and we notice that for every fixed  $\tau \in (-T, 0)$  there is  
 936 a neighborhood  $I_\tau$  of  $\tau$  such that (25) can be replaced by the similar estimate

$$937 \quad (26) \quad \|R(\gamma(t), \zeta)\| \leq \frac{2}{\rho t}, \quad \|R(\ell_k(t), \zeta)\| \leq \frac{2}{\rho t},$$

938 holding for every  $\zeta \in \Gamma_\tau$  and  $t \in I_\tau$ . By applying (17) we get, for every  $t \in I_\tau$  and  
 939  $l \leq k - 1$ ,

$$940 \quad \frac{d^l}{dt^l} P_j(\gamma(t)) - \frac{d^l}{dt^l} P_j(\ell(t)) = -(2\pi i)^{-1} \oint_{\Gamma_\tau} \frac{d^l}{dt^l} \left[ R(\ell_k(t), \zeta) (H(\ell_k(t)) - H(\gamma(t))) R(\gamma(t), \zeta) \right] d\zeta.$$

941 The proof can then be easily completed by applying recursively the identity (24)  
 942 together with the estimates (26) and (19), and by exploiting the regularity of  $\gamma(\cdot)$  and  
 943 the definition of  $\ell_k(\cdot)$ .  $\square$

944 *Acknowledgements.* The authors would like to thank Ugo Boscain for fruitful discus-  
 945 sions.

946

## REFERENCES

- 947 [1] R. Adami and U. Boscain. Controllability of the Schrödinger equation via intersection of eigen-  
 948 values. In *Proceedings of the 44th IEEE Conference on Decision and Control, December*  
 949 *12-15*, pages 1080–1085, 2005.
- 950 [2] A. A. Agrachev. Spaces of symmetric operators with multiple ground states. *Funct. Anal.*  
 951 *Appl.*, 45(4):241–251, 2011.
- 952 [3] F. Albertini and D. D’Alessandro. Notions of controllability for bilinear multilevel quantum  
 953 systems. *IEEE Trans. Automat. Control*, 48(8):1399–1403, 2003.
- 954 [4] J. M. Ball, J. E. Marsden, and M. Slemrod. Controllability for distributed bilinear systems.  
 955 *SIAM J. Control Optim.*, 20(4):575–597, 1982.
- 956 [5] K. Beauchard and J.-M. Coron. Controllability of a quantum particle in a moving potential  
 957 well. *J. Funct. Anal.*, 232(2):328–389, 2006.

- 958 [6] K. Beauchard and C. Laurent. Local controllability of 1D linear and nonlinear Schrödinger  
959 equations with bilinear control. *J. Math. Pures Appl. (9)*, 94(5):520–554, 2010.
- 960 [7] A. M. Bloch, R. W. Brockett, and C. Rangan. Finite controllability of infinite-dimensional  
961 quantum systems. *IEEE Trans. Automat. Control*, 55(8):1797–1805, 2010.
- 962 [8] M. Born and V. Fock. Beweis des adiabatsatzes. *Zeitschrift für Physik A Hadrons and  
963 Nuclei*, 51(3–4):165–180, 1928.
- 964 [9] U. Boscain, M. Caponigro, T. Chambrion, and M. Sigalotti. A weak spectral condition for  
965 the controllability of the bilinear Schrödinger equation with application to the control of  
966 a rotating planar molecule. *Communications in Mathematical Physics*, 2(311):423–455,  
967 2012.
- 968 [10] U. Boscain, F. Chittaro, P. Mason, and M. Sigalotti. Adiabatic control of the Schrödinger  
969 equation via conical intersections of the eigenvalues. *IEEE Trans. Automat. Control*,  
970 57(8):1970–1983, 2012.
- 971 [11] U. Boscain, F. Chittaro, P. Mason, and M. Sigalotti. Controllability of the Schrödinger equation  
972 via adiabatic methods and conical intersections of the eigenvalues. In *Proceedings of the  
973 51st IEEE Conference on Decision and Control, December 10-13*, pages 3044–3049, 2012.
- 974 [12] U. Boscain, J.-P. Gauthier, F. Rossi, and M. Sigalotti. Approximate controllability, exact con-  
975 trollability, and conical eigenvalue intersections for quantum mechanical systems. *Communi-  
976 cations in Mathematical Physics*, 333(3):1225–1239, 2015.
- 977 [13] T. Chambrion, P. Mason, M. Sigalotti, and U. Boscain. Controllability of the discrete-spectrum  
978 Schrödinger equation driven by an external field. *Ann. Inst. H. Poincaré Anal. Non  
979 Linéaire*, 26(1):329–349, 2009.
- 980 [14] F. Chittaro and P. Mason. Approximate controllability by adiabatic methods of the Schrödinger  
981 equation with nonlinear hamiltonian. In *Proceedings of the 54th IEEE Conference on  
982 Decision and Control, December 15-18*, pages 7771–7776, 2015.
- 983 [15] D. D’Alessandro. *Introduction to quantum control and dynamics*. Applied Mathematics and  
984 Nonlinear Science Series. Boca Raton, FL: Chapman, Hall/CRC., 2008.
- 985 [16] C. Fermanian Kammerer and C. Lasser. Propagation through generic level crossings : a surface  
986 hopping semigroup. *SIAM J. of Math. Anal.*, 140(1):103–133, 2008.
- 987 [17] G. A. Hagedorn. Molecular propagation through electron energy level crossings. *Mem. Amer.  
988 Math. Soc.*, 111(536):vi+130, 1994.
- 989 [18] T. Kato. On the adiabatic theorems of quantum mechanics. *Phys. Soc. Japan*, 5:435–439, 1950.
- 990 [19] T. Kato. *Perturbation theory for linear operators*. Die Grundlehren der mathematischen Wis-  
991 senschaften, Band 132. Springer-Verlag New York, Inc., New York, 1966.
- 992 [20] M. Lapert, S. Guérin, and D. Sugny. Field-free quantum cogwheel by shaping of rotational  
993 wave packets. *Phys. Rev. A*, 83:013403 (5 pages), 2011.
- 994 [21] C. Lasser and S. Teufel. Propagation through conical crossings: an asymptotic semigroup.  
995 *Comm. Pure Appl. Math.*, 58(9):1188–1230, 2005.
- 996 [22] Z. Leghtas, A. Sarlette, and P. Rouchon. Adiabatic passage and ensemble control of quantum  
997 systems. *Journal of Physics B*, 44(15), 2011.
- 998 [23] M. Mirrahimi. Lyapunov control of a quantum particle in a decaying potential. *Ann. Inst. H.  
999 Poincaré Anal. Non Linéaire*, 26(5):1743–1765, 2009.
- 1000 [24] G. Nenciu. On the adiabatic theorem of quantum mechanics. *J. Phys A*, 13:15–18, 1980.
- 1001 [25] V. Nersisyan. Global approximate controllability for Schrödinger equation in higher Sobolev  
1002 norms and applications. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 27(3):901–915, 2010.
- 1003 [26] M. Reed and B. Simon. *Methods of modern mathematical physics. II. Fourier analysis, self-  
1004 adjointness*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1975.
- 1005 [27] M. Reed and B. Simon. *Methods of modern mathematical physics. IV. Analysis of operators*.  
1006 Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978.
- 1007 [28] S. Teufel. *Adiabatic perturbation theory in quantum dynamics*, volume 1821 of *Lecture Notes  
1008 in Mathematics*. Springer-Verlag, Berlin, 2003.
- 1009 [29] M. Teytel. How rare are multiple eigenvalues? *Comm. Pure Appl. Math.*, 52(8):917–934, 1999.
- 1010 [30] G. Turinici. On the controllability of bilinear quantum systems. In M. Defranceschi and  
1011 C. Le Bris, editors, *Mathematical models and methods for ab initio Quantum Chemistry*,  
1012 volume 74 of *Lecture Notes in Chemistry*. Springer, 2000.
- 1013 [31] J. Von Neumann and E. Wigner. Über das Verhalten von Eigenwerten bei adiabatischen  
1014 Prozessen. *Physikalische Zeitschrift*, 30:467–470, 1929.
- 1015 [32] D. R. Yarkony. Diabolical conical intersections. *Reviews of Modern Physics*, 68:985–1013, 1996.
- 1016 [33] L. Yatsenko, S. Guérin, and H. Jauslin. Topology of adiabatic passage. *Phys. Rev. A*, 65:043407  
1017 (7 pages), 2002.