

The GR3D model, a tool to explore the Global Repositioning Dynamicsof Diadromous fish Distribution

Thibault Rougier, Hilaire Drouineau, N. Dumoulin, T. Faure, G. Deffuant,

Eric Rochard, Patrick Lambert

▶ To cite this version:

Thibault Rougier, Hilaire Drouineau, N. Dumoulin, T. Faure, G. Deffuant, et al.. The GR3D model, a tool to explore the Global Repositioning Dynamicsof Diadromous fish Distribution. Ecological Modelling, 2014, 283, pp.31-44. 10.1016/j.ecolmodel.2014.03.019. hal-01118691

HAL Id: hal-01118691 https://hal.science/hal-01118691

Submitted on 19 Feb2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 The GR3D model, a tool to explore the Global Repositioning Dynamics of

2 Diadromous fish Distribution

- 3 Thibaud ROUGIER^a, Hilaire DROUINEAU^a, Nicolas DUMOULIN^b, Thierry FAURE^b,
- 4 Guillaume DEFFUANT^b, Eric ROCHARD^a, Patrick LAMBERT^a
- 5
- ⁶ ^aIrstea, EABX, Unité de recherches Ecosystèmes Aquatiques et Changements Globaux, 50
- 7 avenue de Verdun, Gazinet Cestas, F-33612 Cestas, France

- 9
- 10 ^bIrstea, LISC, Laboratoire d'Ingénierie des Systèmes Complexes, 9 avenue Blaise Pascal –
- 11 CS 20085, 63178 Aubière, France
- 12
- 13
- 13
- 14

^{8 &}lt;u>thibaud.rougier@irstea.fr</u>

15 ABSTRACT

16 Within the context of ongoing environmental changes, the life history of diadromous fish 17 represents a real potential for exploring and colonizing new environments due to high 18 potential dispersal abilities. The use of dynamic approaches to assess how these species 19 will respond to climate change is a challenging issue and mechanistic models able to 20 incorporate biological and evolutionary processes are a promising tool. To this end we 21 developed an individual-based model, called GR3D (Global Repositioning Dynamics for 22 Diadromous fish Distribution), combining climatic requirements and population dynamics 23 with an explicit dispersal process to evaluate potential evolution of their distribution area 24 in the context of climatic change. This paper describes thoroughly the model structure and 25 presents an exploratory test case where the repositioning of a virtual allis shad (Alosa alosa 26 L.) population between two river catchments under a scenario of temperature increase was 27 assessed. The global sensitivity analysis showed that landscape structure and parameters 28 related to sea lifespan and to survival at sea were crucial to determine the success of 29 colonization. These results were consistent with the ecology of this species. The integration 30 of climatic factors directly into the processes and the explicit dispersal process make 31 GR3D an original and relevant tool to assess the repositioning dynamics and persistence of 32 diadromous fish facing climate change.

33

34 Keywords: Climate change, Diadromous fish, Dispersal, Distribution, Individual-based
35 model, Population dynamics

36

37

38 **1. Introduction**

39 Organisms have two ways to survive to the current major biodiversity threats such as 40 habitat loss and fragmentation, land use changes and more recently climate change. They 41 may adapt to new conditions on site through phenotypic plasticity or evolution, but this 42 might not be possible within the modification ranges or the timescales imposed by climate 43 change (Gienapp et al., 2008; Visser, 2008). Alternatively, species may shift their range, 44 searching for new suitable habitats. Regarding this last option, many studies have now 45 highlighted changes in distribution and community structure in various species (Thomas 46 and Lennon, 1999; Walther et al., 2002; Parmesan and Yohe, 2003; MacKenzie et al., 47 2007; Nicolas et al., 2011). The success of these relocations or repositioning are contingent 48 upon the dispersal ability of the studied species and the availability of new suitable habitats 49 (Gaston and Blackburn, 2002; Thomas et al., 2004).

50 Many studies using statistical approaches and species distribution models (Guisan 51 and Zimmermann, 2000) have analysed the geographical distribution of species or 52 community with respect to their present environment and used the results to predict 53 expected distribution areas under climate change scenarios. This type of analysis has been 54 carried out for many taxa such as plants (Midgley et al., 2002; Thuiller, 2003; 55 Zimmermann et al., 2009), reptiles and amphibians (Segurado and Araújo, 2004; Araújo et 56 al., 2006), birds (Huntley et al., 2006), mammals (Thuiller et al., 2006), insects (Heikkinen 57 et al., 2007; Barrows et al., 2008), fish (Buisson, 2009) or diadromous fish (Lassalle, 2008). However, such projections do not consider populations' abilities to adapt to 58 59 changing environmental conditions (Pulliam, 2000; Guisan and Thuiller, 2005; Jackson et 60 al., 2009).

Despite a call by the scientific community in the last decade for developing a new
generation of models (Guisan and Thuiller, 2005; Keith et al., 2008; Thuiller et al., 2008;

63 Franklin, 2010) combining climatic suitability, population dynamics and dispersal 64 (Huntley et al., 2010), few studies have developed mechanistic or semi-mechanistic models 65 to deal with population dynamics and dispersal limitations of species under climate change 66 scenarios (but see Brooker et al., (2007), Keith et al., (2008), Engler and Guisan (2009), Midgley et al., (2010), Hein et al. (2011), Fennell et al., (2012)). Mechanistic models aim 67 68 to incorporate the processes by which a species disperses through an environment (Merow 69 et al., 2011) and, unlike correlative models, allows one to compare potential and realized 70 distributions (Franklin, 2010; Gallien et al., 2010; Merow et al., 2011). Such models would 71 be a key point for guiding management and conservation of species in a period of rapid 72 environmental changes (Kinnison and Hairston, 2007) especially in fragmented and 73 heterogeneous environments. However, the development of these sorts of models is 74 generally limited by the lack of knowledge about population dynamics and on processes 75 involved in dispersal of the studied species.

76 The dispersal process is often a critical issue in modelling species movements 77 (Travis et al., 2012). Dispersal is generally defined as (non-returning) movements of 78 individuals away from where they are born, from where they have once reproduced to 79 another breeding site, or between social groups with potentially impacts on gene flow 80 (Clobert et al., 2001; Ronce, 2007; Clobert et al., 2009). It is described as a three-stage 81 process with departure (= emigration), transience (= transfer), and settlement as elementary 82 components (Clobert et al., 2009; Travis et al., 2012). Travis et al., (2012) recently 83 proposed an eco-evolutionary framework for modelling dispersal. However, it is not 84 straightforward to transpose these definitions of dispersal and to define dispersers for 85 diadromous fish according to the diversity and complexity of their life cycles.

Biadromous fishes rely on freshwater, estuarine and marine ecosystems to complete
their life cycles (McDowall, 2009; Quinn et al., 2009). Anadromous species achieve most

88 of their growth at sea and migrate to freshwater to reproduce either in their natal river 89 (natal homing behavior) or in other rivers (straying) while catadromous species spend most 90 of their life in freshwaters and migrate to sea in order to reproduce (McDowall, 1988). 91 Diadromous life history strategies have evolved in phylogenetically diverse fish groups 92 (McDowall, 1997). This original life history characteristic has enabled them to adjust their 93 distribution to cope with environmental changes (McDowall, 1997; McDowall, 2009) and 94 could allow them to do the same in future changing environments (Lassalle et al., 2009). 95 Since diadromous species have a real potential to explore and colonize new environments 96 with a part of their life cycle at sea, a diadromous strategy is an adaptive asset compared to 97 a holobiotic life history strategy in a context of global change. In this paper, we will only 98 focus on anadromous species, for which we assume that dispersers are strayed spawners 99 (i.e. strayers) not making natal homing behavior (i.e. not returning to their natal stream for 100 spawning). However, processes involved in the determination of strayers as well as the 101 behavior of the strayers remain unknown for diadromous species. As dispersal is closely 102 linked to individual behavior, individual-based-models (IBMs) appear to be a promising 103 approach to address the question of dispersal and repositioning dynamics of diadromous 104 fishes. IBMs also offers a flexible and easy way to simulate different behaviors.

105 Despite their complexity often seen to hamper interpretation of their results, the use 106 of IBMs in ecological modelling has increased exponentially (Grimm, 1999) from the 107 seminal works of Huston et al. (1988). Moreover, recent progresses have been made to 108 allow complex IBMs to be better communicated and described (Grimm et al., 2006; 109 Grimm et al., 2010). Many IBMs have already involved fish populations (Van Winkle et 110 al., 1993; Van Winkle et al., 1998; Grimm, 1999; Charles et al., 2008) and more especially 111 diadromous fish such as chinook salmon, Oncorhynchus tshawytscha (Jager et al., 1997), 112 European eel, Anguilla anguilla (Lambert and Rochard, 2007) or more recently Sockeve

Salmon, Onchorhynchus nerka (Reed et al., 2011), brown trout, Salmo trutta (Frank and 113 114 Baret, 2013) and Atlantic salmon, Salmo salar (Piou and Prévost, 2012, 2013). 115 Furthermore, IBMs are powerful tools to represent biological and habitat mechanisms, 116 individual behavior and variability (Grimm, 1999). They also allow analysing potential 117 climate change effects on natural systems (Reed et al., 2011; Piou and Prévost, 2013) as 118 environmental effects can be directly incorporated in processes. Although complexity is an 119 essential feature of IBMs and can result in unforeseen emergent properties, they may 120 appear over-parameterized if compared to much more parsimonious statistical models. 121 Furthermore, a number of population dynamics parameters are often imprecise and poorly 122 estimated. Hence, it is important to carry out sensitivity analyses (Saltelli, 2004) of this 123 kind of model. Global sensitivity analysis methods are now well known to advance the 124 comprehension and exploration of system modelled (Faivre et al., 2013). In such an 125 analysis, parameters are varied simultaneously allowing identification of interactions by 126 statistical analysis of the simulation results. These methods have already been used in 127 industry (Kleijnen, 1998), ecology (Cariboni et al., 2007) or fishery science (Drouineau et 128 al., 2006; Drouineau et al., 2008; Lehuta et al., 2010).

129 For diadromous species, developing a model combining climatic requirements and 130 population dynamics with explicit dispersal processes is a real challenge and will be a 131 major step forward for guiding management policies. If successful, it could help to 132 determine in which catchment and for which species restoration and restocking actions will 133 be relevant. Moreover, the development of this kind of model for diadromous fish does not 134 appear to be limited by the knowledge of their ecology and their population dynamics 135 which are now well documented (at least for the continental phase) for several species such 136 as Atlantic salmon Salmo salar (Peterson et al., 1977; Bagliniere and Maisse, 1985; Fleming, 1996; Friedland, 1998; Hutchings and Jones, 1998; Forseth et al., 2001; Jonsson 137

and Jonsson, 2009; Buoro et al., 2010), European eel *Anguilla anguilla* (Gatto and Rossi,
139 1979; Vollestad and Jonsson, 1988; De Leo and Gatto, 1995; Dekker, 2000; Lambert,
2005; Lambert and Rochard, 2007; Bevacqua et al., 2011), European sturgeon *Acipenser*141 *sturio* (Rochard and Lambert, 2011) or allis shad *Alosa alosa* (Lambert et al., 2001; Acolas
142 et al., 2006; Lassalle et al., 2008b; Mota and Antunes, 2011; Rougier et al., 2012).

143 In this context, we developed an exploratory simulation model (GR3D: Global 144 Repositioning Dynamics for Diadromous fish Distribution) integrating the whole 145 population dynamics of diadromous fish (reproduction, growth, survival, maturation, 146 downstream and upstream migrations). The GR3D model is individual-based, stochastic 147 and spatially explicit. To test climate change effect on population dynamics, we introduced 148 temperature variations and their effects on demographic traits, such as growth, 149 reproduction and survival. We introduced individual variability through growth, survival, 150 reproduction and migration.

Models of high complexity require a substantial exploration phase to avoid misinterpretation of simulation results (Grimm and Railsback, 2005). The more complex a model is, the more its calibration, and the extrapolation and analysis of its results are difficult (Levins, 1966; Caswell and John, 1992). From this consideration, testing a complex model with simple exploratory cases is a reasonable and rational approach (Faivre et al., 2013).

Here, we present the GR3D model and show how this generic model may be relevant to study the repositioning dynamics and persistence of diadromous fish in a context of climate change. To illustrate the potential of the model, we applied GR3D in a virtual exploratory study where we assess the repositioning of a virtual anadromous allis shad population between two river catchments under a scenario of temperature increase.

- 162 We run the associated sensitivity analysis in order to identify parameters influencing model
- 163 outputs.
- 164
- 165 **2. Materials and methods**
- 166 2.1. Model description

167 The GR3D model has been developed in Java using the "SimAquaLife" framework 168 (Dumoulin, 2007) which is an individual-based, process-oriented toolkit for aquatic life 169 simulation.

The description of the GR3D model follows the ODD protocol (Overview, Design concepts, Details) for describing individual- and agent-based models (Grimm and Railsback, 2005; Grimm et al., 2006; Grimm et al., 2010).

173 2.1.1. Purpose

The general purpose of GR3D is to evaluate diadromous fish local persistence (defined as the probability for a species to maintain a population in a specific river catchment) (Jager et al., 2013), global persistence (defined as the probability for a species to maintain a population in at least one catchment) and concomitant potential evolution of their distribution area, in the context of climatic change at a European scale combining population dynamics, repositioning behaviors through dispersal processes and climatic requirements.

- 181
- 182 2.1.2. State variables and scales

Temporal scales: GR3D simulates a seasonal time step with distinct processes and
 scheduling in each of these steps.

Entities and spatial scales: Three types of entities compose the GR3D model: one for
fish agents and two for environment elements. GR3D is spatially explicit with a continental

187 compartment and a sea compartment. The continental compartment is split up in "River
188 Basins". Each "River Basin" (RB) is characterized by a name, a position (latitude and
189 longitude at the mouth), a catchment area (km²) and seasonal mean water temperatures at
190 the mouth (Table 1).

According to the studied species, the sea compartment is either split into "Sea Basins" (SB) associated with one of the river basins, or composed of a unique "sea basin" connected to all the river basins. Each SB is characterized by a name and seasonal mean water temperatures (Table 1).

Temperatures in compartments are updated at each time step according to a virtual
scenario of water temperature increase or according to a real projection of climate change.
Actually, GR3D is designed to work either with virtual environments or real environments
such as the Atlantic coast.

199 *State variables of diadromous fish individuals:* Each diadromous fish is described 200 by an ID, a gender, a maturation stage (mature or immature), an age, a body length, a 201 location (a river basin or a sea basin), a birth place and a number of reproductions.

202

203 2.1.3. Process overview and scheduling

GR3D is developed to cover the entire life cycle of any diadromous fish species trying to be the more generic as possible and taking into account the differences between anadromous and catadromous species. Hence, we structured the model in six submodels consistent with the life cycle events and processes of any diadromous species (reproduction, growth, survival, downstream migration, maturation, and upstream migration including dispersal). As the present test case deals with an anadromous species (see section 2.2.), we only presented the computational order of life cycle events and

211 processes in a conceptual diagram for an anadromous species with allis shad as an212 illustrative case (Fig. 1).

213

214 2.1.4. Design concepts

215 Basic principles: GR3D is a population dynamic model that explicitly includes 216 environmental effects in processes and that incorporates an original dispersal process. 217 During the upstream migration of anadromous fish, we distinguished two kinds of 218 individuals: individuals having a homing behavior (i.e. returning to their natal river to 219 spawn) or having a straying behavior (i.e. not returning to their natal stream). In the 220 absence of information on marine dispersion behavior, we assumed that the choice of a 221 river basin by an individual is the result of three components: 1) a propensity to adopt a 222 homing behavior (characteristic of the species), 2) the accessibility of the river basin, and 223 3) its attractiveness.

Interactions: Indirect interactions are included in the reproduction process (see Section 2.1.7.1) through a demographic Allee effect (Allee, 1931; Stephens et al., 1999) and through a density-dependant survival of eggs and larvae.

227 *Stochasticity:* Demographic stochasticity was incorporated for growth, survival,
 228 reproduction and dispersal.

Observation: Graphical outputs from "SimAquaLife" interface directly show population-level variables (declared as observable in GR3D) as well as spatial distribution of the individuals. As example, the number of spawners (distinguishing those exhibiting natal homing behavior from those exhibiting straying) and juveniles at each reproduction for each river basin can be recorded as well as the year of the first non-null reproduction and the last year without reproduction.

236 2.1.5. Initialization

237	The environment, the start and the number of year of simulations, the abundance, the age
238	and the initial distribution of the individuals are initialised according the application case.
239	Population parameters are initialised according to the studied species (see section 2.2.).
240	
241	2.1.6. Input
242	In the current version of GR3D, environmental conditions changed over time via
243	temperature, which changed over space and season. The model is designed to integrate any
244	kind of seasonal temperature temporal series. In a same way, to define the environment, an
245	input file containing river basins characteristics (cf. § 2.1.2) can be used.
246	
247	2.1.7. Submodels
248	GR3D contains 42 parameters, which are listed in Table 2 according to the submodel they

are involved in. According to the studied species, their values are obtained either from observations (literature) or are reasonable guessed (i.e. we have only an idea of what would be a realistic value of the parameter) or are calibrated. GR3D is based on six submodels representing fundamental biological processes.

253

254 2.1.7.1. Reproduction:

Reproduction occurs yearly during the reproduction season (defined as a population parameter, Table 2) in each river basin when spawners are present. We assumed that numbers of recruits R_j produced by S_j spawners in a spawning basin *j* follows a Beverton & Holt stock-recruitment relationship of parameters α_j and β_j (BH S-R) (Beverton and Holt, 1957). However we modified the traditional BH S-R:

260 We included an Allee effect (Eq. (1)) to take into account difficulties to settle a 261 population with limited numbers of fish in new habitats. Depensation strength is 262 modeled as a function of river basin watershed area through two species specific 263 parameters: η a parameter linking the basin watershed area WA_i to its carrying 264 capacity (the higher η is, the higher the stock level to reach 95% of the asymptotic recruitment) and θ which controls the Allee effect strength (a high θ corresponds to a 265 266 strong depensation, i.e. the stock level to produce half of the asymptotic recruitment is 267 close to the stock level producing 95% of the asymptotic recruitment).

268
$$R_{j} = \frac{\alpha_{j}S_{j} \frac{1}{1 + e^{\left(-\ln(19)\frac{S_{j} - \eta/\theta \cdot WA_{j}}{\eta \cdot WA_{j} - \eta/\theta \cdot WA_{j}}\right)}}}{\beta_{j} + S_{j} \frac{1}{1 + e^{\left(-\ln(19)\frac{S_{j} - \eta/\theta \cdot WA_{j}}{\eta \cdot WA_{j} - \eta/\theta \cdot WA_{j}}\right)}}$$
(1)

269

270 By comparison to the traditional BH S-R, we considered that the non-density dependent mortality from egg to recruit (duration Δt_{rec} , Table 2) is a function of 271 temperature, consequently parameters α_i and β_i depend on temperature (Eq. (2) and 272 Eq. (3)). Egg survival is optimal ($surv_{optRep}$, Table 2) at T_{opt} and there is no recruitment 273 274 when temperature is out of the range T_{min} - T_{max} (Eq. (4), Table 2). The density 275 dependent mortality of the BH S-R is assumed to be dependent on the basin surface 276 (through a population parameter λ , Table 2) to take into account resource limitations 277 in small basins (Eq. (5)).

278
$$\alpha_{j} = \frac{b_{j} e^{-b_{j} \Delta t_{rec}}}{c_{j} \left(1 - e^{-b_{j} \Delta t_{rec}}\right)}, \qquad (2)$$

279
$$\beta_j = \frac{b_j}{ac_j \left(1 - e^{-b_j \Delta t_{rec}}\right)},$$
 (3)

280 The parameter *a* is linked to the fecundity of the species (Table 2),

281
$$b_{j} = -\frac{1}{\Delta t_{rec}} \ln \left[surv_{optRep} \frac{\left(T_{j} - T_{minRep}\right) \left(T_{j} - T_{maxRep}\right)}{\left(T_{j} - T_{minRep}\right) \left(T_{j} - T_{maxRep}\right) - \left(T_{j} - T_{optRep}\right)^{2}} \right], \quad (4)$$

 $c_j = \frac{\lambda}{Surf_j}$.

(5)

282 and

283

284

We introduce stochasticity in this process by drawing the value of the effective recruitment $R_{Eff,j}$ in a lognormal distribution of standard deviation σ_{rep} around the stock-recruitment relationship (Table 2).

After the reproduction, the population parameter Sp_{sp} (Table 2) defined the survival probability of the spawners to take into account the iteroparity of the studied species.

290

291 2.1.7.2. Downstream migration.

This process occurs at different life stage and season according to the studied species. The age and the season at which this migration occurs are defined as population parameters (Table 2). In this process, individuals migrate from their river location to a determined sea compartment according to the studied species.

296

297 2.1.7.3. Growth.

The population parameter L_{ini} determines the initial length of individuals at age Δt_{rec} (i.e. after reproduction).We then use a von Bertalanffy growth function (von Bertalanffy, 1938) derived from Fabens (1965) for modelling of the growth process. Each season, each individual length is updated according to its previous length. In order to introduce individual variability and to avoid negative growth, we use a lognormal distribution to determine the growth increment

$$\Delta L = LogN(\mu_{\Delta L}, \sigma_{\Delta L}), \qquad (6)$$

305 with $\mu_{\Delta L}$ the mean and $\sigma_{\Delta L}$ the standard deviation of the variable's natural logarithm 306 (Table 2). So, $\mu_{\Delta L}$ is expressed

307
$$\mu_{\Delta L} = \log\left(\left(L_{\infty} - TL_{t}\right) \times \left(1 - \exp^{-\kappa \Delta t}\right)\right) - \frac{\sigma_{\Delta L}^{2}}{2}, \qquad (7)$$

where TL_t is the total length of a fish at time unit t, L_{∞} is a population parameter 308 309 corresponding to the asymptotic length of fish (Table 2) and κ is the growth coefficient. 310 Since temperature is known to affect growth rate (Brown et al., 2004), Rosso's function 311 (Rosso et al., 1995) is used to correlate the growth coefficient to temperature and introduce 312 seasonal variability (Mallet et al., 1999; Dion and Hughes, 2004; Kielbassa et al., 2010; 313 Bal et al., 2011). According to this function, the growth coefficient κ and temperature are 314 linked by a bell-shaped relationship. In fact, κ is equal to 0 if the temperature is lower or equal to the minimal growth temperature $T_{minGrow}$; it increases with temperature up to an 315 optimum $\kappa_{optGrow}$ at the optimal growth temperature $T_{optGrow}$. If temperature continues to 316 317 increase, the growth coefficient κ rapidly decreases down to 0 at the maximal growth 318 temperature $T_{maxGrow}$. The relationship is expressed

319

320
$$\kappa = \kappa_{optGrow} \frac{(T - T_{minGrow})(T - T_{maxGrow})}{(T - T_{minGrow})(T - T_{maxGrow}) - (T - T_{optGrow})^{2}}.$$
 (8)

321

322 Parameters
$$T_{minGrow}$$
, $T_{optGrow}$, $T_{maxGrow}$ and $\kappa_{optGrow}$ are population parameters (Table 2).

323

324 2.1.7.4. Survival.

At each time step, the survival of each individual is assessed depending on its location andstage.

For individuals at sea, the seasonal survival probability, Sp_{sea} , is calculated using the annual mortality coefficient at sea, Z_{sea} , defined as a population parameter (Table 2) and a survival equation $Sp_{sea} = \exp^{-Z_{sea} \times \Delta t_{season}}$, (9)

331 where Δt_{season} is the duration of a season (i.e. 0.25 year⁻¹). We assumed that Z_{sea}

incorporates all sources of individual mortality at sea.

333 For individuals in river, the seasonal survival probability, Sp_{riv} , is calculated using two

annual mortality coefficients M_{riv} (natural mortality coefficient) and H_{riv} (human-induced

- 335 mortality coefficient) defined as population parameters (Table 2) and a survival equation
- $336 Sp_{riv} = \exp^{-(M_{riv} + H_{riv}) \times \Delta t_{season}}. (10)$

337 The natural mortality coefficient M_{riv} is assumed to be dependent on the river temperature

338 T_i and was computed as follows

339
$$M_{riv} = \frac{-\ln\left(surv_{optRiv} \frac{\left(T_{j} - T_{minSurvRiv}\right)\left(T_{j} - T_{maxSurvRiv}\right)}{\left(T_{j} - T_{minSurvRiv}\right)\left(T_{j} - T_{maxSurvRiv}\right) - \left(T_{j} - T_{optsurvRiv}\right)^{2}\right)}{\Delta t_{season}}, \quad (11)$$

340 where $T_{minSurvRiv}$, $T_{optSurvRiv}$, $T_{maxSurvRiv}$ and $surv_{optRiv}$ were population parameters (Table 2). 341 We assumed that the other mortality coefficient H_{riv} incorporates the other sources of 342 mortality in river (fishery, pollution, dams...).

- 343
- 344 2.1.7.5. *Maturation*.

In the current version of GR3D, individual maturation is not explicit and did not involve energy allocation processes. An individual becomes mature as soon as its length reached the length at first maturity L_{mat} (defined as a population parameter, Table 2) and its maturation stage changed from *immature* to *mature*.

350 2.1.7.6. Upstream migration.

This submodel is essential in the model since we assumed that dispersal is linked to this migration. This process occurs at different life stages and seasons depending on the studied species. The age and the season at which the downstream migration occurs are defined as population parameters (Table 2). It has been designed as a three-stage process with emigration, movement and settlement phases.

356 *1. The emigration phase:* in this phase, individuals have a probability to adopt a homing (357 p_{hom}) or a straying behavior $(1-p_{hom})$ which is assumed to be a characteristic of the 358 species. p_{hom} is defined as a population parameter (Table 2).

359

360 2. The movement phase: Individuals that do not become strayers simply migrate in their 361 natal river (homing behavior). For strayers, the probability to migrate in each river basin is 362 assumed to be a function of its accessibility and its attractiveness. We assumed that 363 accessibility depends on the dispersal distance $D_{j-birthPlace_i}$ between basin j and the birth 364 place of an individual *i*, and on the total length TL_i of the individual *i*. The basin 365 attractiveness is assumed to be a function of its watershed area WA_i as a proxy of the river 366 flow. Then, the "weight" of each basin relative to an individual *i* is computed using a logit 367 function

$$\log it\left(w_{i}^{j}\right) = \alpha_{const} + \alpha_{dist}\left(\frac{D_{j-birthPlace_{i}} - \overline{D_{j-birthPlace}}}{\sigma_{j-birthPlace}}\right), \quad (12)$$
$$+ \alpha_{TL}\left(\frac{TL_{i} - \overline{TL}}{\sigma_{TL}}\right) + \alpha_{WA}\left(\frac{WA_{j} - \overline{WA}}{\sigma_{WA}}\right)$$

369 where α_{const} , α_{dist} , α_{TL} and α_{WA} are four population parameters (Table 2). In order to have 370 comparable parameters, we used standard cores values of each factor influencing the function and we defined $\overline{D_{j-birthPlace}}$, $\sigma_{j-birthPlace}$, \overline{TL} , σ_{TL} , \overline{WA} and σ_{WA} as population parameters (Table 2).

We assumed that individuals may also not find any basin so we introduce a virtual "death basin" with a fixed weight $w^{deathBasin}$. Then w_i^j are standardized so that their sum equals 1, providing probabilities to choose each river basin (including death basin). The choice is

- then modelled by a simple multinomial process.
- 377 *3. The settlement phase:* Individuals enter in the selected destination, survive if conditions
 378 are suitable and reproduce if they find mating requirements.
- 379

380 *2.2. The test case*

381 2.2.1. Description of the studies species

382 Allis shad (Alosa alosa L.) is an anadromous clupeid spawning in the higher middle 383 watercourse of rivers. Fish migrate to sea during their first year, where they grow and 384 return to freshwater to spawn at between 3 and 6 years (Bagliniere and Elie, 2000; 385 Baglinière et al., 2003). Currently, populations of allis shad exist along the northeastern 386 Atlantic coasts in some large rivers of France (Loire, Gironde-Garonne-Dordogne, and Adour) and Portugal (Minho and Lima) (Lassalle et al., 2008b) and despite some 387 388 protective measures, this species appears to have been in serious decline for a number of 389 years (Limburg and Waldman, 2009; Rougier et al., 2012). Biology and ecology of allis 390 shad have therefore received a great deal of attention in the last 30 years (Mennesson-391 Boisneau and Boisneau, 1990; Prouzet et al., 1994; Bagliniere and Elie, 2000; Acolas et 392 al., 2004; Acolas et al., 2006) and several studies also dealt with its population dynamics 393 (Martin Vandembulcke, 1999; Lambert et al., 2001; Rougier et al., 2012).

394

395 *2.2.2. Environment, time and initialization*

An environment with two rivers basins (RB1 and RB2) and two sea basins (SB1 and SB2) highly associated to the river basins (i.e. RB1 with SB1 and RB2 with SB2) was used for the test case. Our experiment mimics a situation where temperature increase opened the opportunity for a stable allis shad population in RB1 to expand into a new suitable river RB2. Two environmental parameters were defined to determine the initial configuration of the environment: *WA_RB2* corresponding to the watershed area of the uninhabited river basin RB2 and *intDist* describing the distance between RB1 and RB2.

Each simulation starts in summer and lasts 200 years (i.e. 800 time steps). During 403 404 the first fifty years of simulation, a virtual population of allis shad was created in RB1 by 405 introducing there 500000 juveniles at the first time step. From the year 50 to the year 150, 406 we introduced in the two basins a linear water temperature increase of 3°C in 100 years 407 (i.e. water temperature increase by 0.0075°C at each time step) and RB2 becomes suitable 408 for allis shad reproduction. At the end of the temperature increase, only RB2 is suitable for 409 allis shad for the last 50 years of simulations. Temperatures in RB1 and RB2 were 410 initialized with the same values except for the spring temperature as we used it to control 411 the suitability of each river during the simulation. In order to work with realistic values, we 412 used daily estimates of Gironde estuarine water temperature (°C) provided by EDF 413 (Electricité De France) between 1991 and 2009 to initialize the winter, summer and 414 autumn temperatures in RB1 and RB2 as the Gironde basin is located at the center of the 415 allis shad distribution area (Baglinière, 2000; Lassalle, 2008). Spring temperature of RB1 was set to 3°C below $T_{maxSurvRiv}$ and set to T_{minRep} in RB2. The watershed area of RB1 was 416 set to 40000 km². The environmental parameters WA RB2 and *intDist* were included in the 417 418 global sensitivity analysis to assess the influence of the environment configuration on 419 model outputs (see section 2.2.4.).

420 The seasonal mean water temperatures of SB1 and SB2 were calculated as the 421 mean between 12°C and the seasonal mean water temperatures of RB1 and RB2, 422 respectively.

Initial length of each individual is set at 2 centimetres. The individual state variables were initialized as follows for each new individual: gender was assigned according to a sex ratio of 1:1; stage was set to "immature", the number of reproduction was set to 0; and the birth place was set according to their birth location.

427

428 2.2.3. Model parameterisation

For this test case with allis shad, 20 population dynamics parameters were obtained from
literature, 18 were parameterized as reasonable guesses (based upon expertise), and 4 were
estimated using off-line calibration (Table 2).

For the off-line calibration, we used a previous work (Rougier et al., 2012) about the population dynamics of allis shad in the Gironde basin for the parameterization of the reproduction process (i.e. parameter η , θ , *surv*_{optRep} and λ of the BH S-R relationship, Table 2) and used their estimates of $\alpha_{Gironde}$, $\beta_{Gironde}$ and η . We assumed that the recruitment was a number of juveniles in estuary and set Δt_{rec} to 0.33 year (4 months) as it corresponds to the average duration of the juvenile downstream migration towards the sea (Lochet et al., 2008).

We assumed dispersal of allis shad was only dependent of the distance between the suitable habitats. Hence, α_{TL} and α_{WA} were set to 0. We set $\sigma_{j-birthPlace}$ to 978 km (using the EuroDiad 2.0 database of European river basins from Lassalle (2008)). We set $\overline{D_{j-birthPlace}}$ to 300 km as we assumed that this value was close to the maximal distance of dispersal for allis shad. Then we assumed a weight of 0.95 and 0.05 for a basin located at a 444 distance of 10 km and 300 km respectively. In consequence, α_{const} and α_{dist} were set to 445 -2.9 and 19.7 respectively (Table 2).

- 446
- 447 2.2.4. Sensitivity analysis

For these simulations, we mainly focused on the success of the colonization of RB2. Hence, we recorded the following 2 output variables: (i) *fs* was a qualitative variable describing the final state of RB2 using the value of the coefficient of variation (CV) of the recruitment in RB2 during the last ten years of simulations, i.e. that it took either the value 0 if the CV was null, or the value 1 if the CV was higher than a threshold arbitrarily set to 5%, or the value 2 if the CV was positive and lower than the threshold; (ii) *mlr* was the mean of the recruitment in RB2 during the last ten years of simulations.

455 We defined two variables using the fs output variable. fsP indicated the 456 presence/absence of fish in RB2 (i.e. fs = 0 for absence and fs > 0 for presence). 457 Considering only positive values of fs, we defined fsS as a variable describing a kind of 458 stability of the RB2 population (i.e. relatively unstable for $f_s = 1$ because it means that 459 there is high variation in the recruitment values in the ten last year of simulation, and stable for fs = 2 because it means that recruitment values are stable in the ten last years of 460 461 simulation). Hence, we used these two variables to analyze the influence of parameters on 462 both the presence/absence of fish in RB2 and on the stability of the population in RB2. mlr 463 was a metric indicating the abundance of fish in RB2. As the null values described absence 464 of fish in RB2, we only used positive values of *mlr* to analyze the influence of parameters 465 on abundance in RB2.

In order to screen non-influential and influential parameters on GR3D model outputs, we conducted a global sensitivity analyses by varying both the model and the environmental parameters. We identified 20 uncertain population dynamics parameters of

469 the GR3D model to incorporate in the sensitivity analysis and we added 2 environmental 470 parameters in the analysis (WA RB2 the watershed area of RB2 and intDist the distance 471 between RB1 and RB2; Table 3). Considering a maximal range of variation of 20% often 472 used in sensitivity analysis (Drouineau et al., 2006; Lehuta et al., 2010), we set a low and a 473 high value for each population parameter based on our own expertise and knowledge of 474 allis shad population dynamics (Table 3). Concerning environmental parameters, we considered three modalities for WA_RB2 and four for intDist (Table 3). As a complete 475 476 design would have taken too much time since it would involve more than 12 million simulations without replicates $(2^{20} \times 3 \times 4)$, we first used an experimental design to 477 478 identify the most influential parameters on model outputs. Regarding the number of factors 479 and modalities, the most appropriate experimental design are D-optimal designs 480 (Drouineau et al., 2006; Faivre et al., 2013). In this type of design, the estimations of the 481 effects are partially correlated (i.e. that contrary to factorial design, confusion between 482 effects estimates may exist, but an algorithm is used to find the set of experiments), 483 maximizing the determinant of the information matrix (XX' with X, the matrix of the 484 different combinations of modalities for each experiment) of the design, equivalent to 485 maximizing the efficiency of the estimation (Droesbeke et al., 1997; Drouineau et al., 486 2006). We use the AlgDesign R package (Wheeler, 2011) to generate our D-Optimal 487 design and, finally, the resulting experimental design consisted of 1300 simulations that we 488 replicated 10 times. Only main effects were estimated with this design using a logistic 489 regression model for *fsP* and *fsS* and analysis of variance (ANOVA) for *mlr*. For each 490 model output response y, first order indices (SI) of a parameter x_i measuring the importance 491 of the deviance (or variance for *mlr* output) of the y mean conditionally on factor x_i , were 492 considered as a sensitivity index (Saltelli, 2004; Saltelli et al., 2008; Faivre et al., 2013) 493 and were calculated as follows

494
$$SI(x_i) = \frac{Dev[E(y | x_i)]}{Dev(y)}.$$
 (18)

495 Focusing on the ten highest SI of the 3 model outputs, we selected the most influential 496 parameters. For computation time reasons, we were not able to select more than 13 497 parameters. Then with the most influential selected parameters, we used a complete 498 sampling with 10 replicates in order to estimate main effects and first order interactions 499 effects on the sensitivity of model outputs. We used the same models as those used with 500 the experimental design to estimate SI and global sensitivity indices (TSI). TSI, that 501 measured the mean of the y deviance (or variance for *mlr* output) on the uncertain 502 parameters different from x_i (Saltelli, 2004; Saltelli et al., 2008; Faivre et al., 2013), were 503 calculated for each parameter as follows

504
$$TSI(x_i) = \frac{E\left[Dev(y \mid x_j, j \neq i)\right]}{Dev(y)}.$$
 (19)

505 TSI corresponds also to a measure of the sum of the main effect of a parameter x_i and the 506 effects of interactions of this parameter with all the other uncertain parameters (Saltelli, 507 2004; Saltelli et al., 2008; Faivre et al., 2013). Both indices SI and TSI are between 0 and 1 508 but TSI is higher than SI as it includes interaction effects. Estimating the impact of first 509 order interaction was necessary to assess the combined effect of uncertainties on two 510 parameters, which may be different from the sum of the main effects of the two 511 parameters. We expressed SI and TSI in percentage and we assumed the more distant from 512 0 they were, the more impact they have on model results.

513 Statistical analyses of model outputs were carried out using R (R Core Team, 514 2013). To run all our simulations of the sensitivity analysis, we used OpenMOLE which is 515 a workflow engine specifically tailored for the distributed exploration of simulation models 516 (Reuillon et al., 2013).

518 **3. Results**

519 We present in Fig 2 an illustrative example of model output where we followed the 520 dynamics over time in RB1 and RB2 of the recruitment for a same set of parameters by 521 just varying the value of the distance between RB1 and RB2.

522 *3.1.* Selection of the most influential parameters

523 Among the 13000 simulations with the D-Optimal design, we observed failure of 524 colonization of RB2 in 48% of simulations (i.e. fs = 0). Thirteen percent of simulations led 525 to an unstable population (i.e. fs = 1) and 39% led to stable situations in RB2 (i.e. fs = 2). 526 The logistic regression applied to *fsP* and *fsS* explained 78% and 70% of the null deviance 527 respectively and the analysis of variance of *mlr* explained 60% of the null variance (Table 528 4). Focusing on the ten highest SI of the three model outputs, we identified 13 different 529 parameters meaning that model outputs are relatively sensitive to the same parameter (Fig. 530 3). It was especially true for *fsP* and *fsS* which were very sensitive to the distance between 531 RB1 and RB2 that explained around 50% of the deviance for these two variables (Fig. 3). These two variables were also sensitive to parameters of the growth ($\kappa_{optGrow}$), maturation 532 533 (L_{mat}) , and survival processes (Z_{sea}) which are highly linked to the duration of the 534 individual life at sea (Fig. 3). So, the success of colonization and the stability of the 535 population were, not surprisingly, negatively linked to *intDist*, but also to parameters that 536 increased the duration of the life at sea. mlr was mainly sensitive to the watershed area of 537 RB2 (Fig. 3) with a positive correlation between *mlr* and *WA RB2* as this parameter is 538 involved in the reproduction process to determine the production capacity of RB2. mlr was 539 also sensitive to the other parameters of the reproduction process (Surv_{optRep}, λ and T_{optRep}) 540 and to the environmental parameter *intDist*. To select the most influential parameters, we 541 kept 12 of the 13 parameters identified in the ten highest SI of the three model outputs. 542 Actually, as the outputs were not very sensitive to $T_{maxSurvRiv}$ which was a parameter used to

control the suitability of RB1 in our test case, we did not select this parameter and rather selected the parameter $w^{deathBasin}$ involved in dispersal. Among the 13 selected parameters, four concerned the reproduction process, two concerned the growth, survival and upstream migration (i.e. dispersal) processes, one concerned the maturation process and the last two were environmental parameters (Table 3). Consequently, in the second step of the sensitivity analysis, the complete sampling consisted in 24576 simulations that we replicated 10 times.

550 3.2. Analysis of interaction effects

551 Among the 245760 simulations, we observed 49% of unsuccessful colonization, 11% of 552 simulations led to an unstable population in RB2 and 40% led to a stable situation. 553 Although we considered fewer parameters than with the D-Optimal design, the explained 554 deviance and variance of the outputs analysis increased in this second step of the 555 sensitivity analysis because we took first order interactions effects into account. We 556 explained 90% and 85% of the deviance for fsP and fsS respectively and 79% of the 557 variance for *mlr* (Table 4). The output variables were globally sensitive to the same 558 parameters than those observed with the D-Optimal design (Fig. 4). This sensitivity 559 analysis confirmed that the model outputs *fsP* and *fsS* were particularly sensitive to the 560 environmental parameter *intDist* and to parameters increasing the duration of the life at sea $(\kappa_{antGrow}, L_{mat}, \text{ and } Z_{sea})$ and that mlr was sensitive to the two environmental parameters and 561 562 to parameters of the reproduction process. Additionally, we observed a great importance of 563 the interaction effects for two parameters: the homing probability p_{hom} and the parameter 564 θ of the reproduction process which is linked to the strength of the Allee effect (Fig. 4). 565 This was especially true for the *fsS* output where the main effect of the homing probability 566 and the parameter θ were very low (0.02% and 0.03% respectively) while the interaction 567 effects of these factors with other parameters explained 6.3% and 4.9% of the null

deviance respectively (Fig. 4). Interaction effects of the homing probability were particularly high with the distance between RB1 and RB2 and the optimal growth coefficient $\kappa_{optGrow}$. For the parameter θ , interaction effects were high with the length at first maturity, the optimal growth coefficient and the annual mortality coefficient at sea. We also observed a high interaction between the weight of the death basin and the distance between RB1 and RB2 to explain the stability of the population in RB2.

To summarize, the success and the stability of the colonization of RB2 are very sensitive to the landscape structure represented through the distance between RB1 and RB2 and to parameters related to the duration of the life at sea and the survival condition in this compartment. Interaction effects mainly highlighted that the role of the homing and the Allee effect should be considered with attention. Concerning the abundance of the recruitment in RB2, it is sensitive to parameters of the reproduction process and to the watershed area of RB2.

581

582 **4. Discussion**

583 We developed, implemented and tested an individual-based model combining population 584 dynamics, dispersal and temperature requirements. GR3D has been designed to assess the 585 persistence of diadromous fish and the evolution of their distribution area in a context of 586 global warming. To our knowledge, this is the first attempt to incorporate dispersal 587 abilities of diadromous fish within a dynamic approach at the scale of their distribution 588 range. This article focuses on the individual-based model description and its exploration 589 with a basic virtual test case. We deliberately chose to run our simulations in a simple, 590 virtual environment in order to clearly disentangle the effect of the environment and the 591 effect of uncertainties in population dynamics on model outputs using a global sensitivity 592 analysis. We demonstrated the sensitivity of the outputs to some key population dynamics

593 parameters and to environmental parameters which were consistent with our knowledge of 594 this species. We considered our results are satisfactory enough to address more precisely, 595 in further studies, the issue of allis shad persistence in a context of global change at the real 596 scale of its distribution range. GR3D has been designed as a flexible and generic model 597 and can represent an integrative tool to study persistence of diadromous fish whose 598 population dynamics is well documented. After discussing the innovations and the 599 structure of the model, we come back to the understandings brought by our test case and its 600 associated sensitivity analysis.

601

602 4.1. Innovations and model structure

603 The GR3D model incorporates an explicit dynamic model of diadromous fish dispersal 604 process at large scale. This is particularly relevant in a context of climate change since 605 Brooker et al. (2007) has demonstrated that dispersal ability of individuals interacts with 606 the rate of climate change to determine range shifting dynamics. To model dispersal in 607 GR3D, we took both landscape structure and individual dispersal abilities into account. We 608 defined a generic accessibility model for available habitats (i.e. river basins) that depends 609 on the dispersal distance and the size of the individuals (as it represents a proxy of their 610 swimming capacities). Dispersal distance has already been recognized as an important 611 factor in dispersal processes and is often represented through dispersal kernels in dispersal 612 modelling approaches (Chesson and Lee, 2005; Slone, 2011; Bocedi et al., 2012). 613 Moreover the individual size has been recognized as closely linked to dispersal potential of 614 individuals with potential impacts on species' range shifts through phenotype dependent 615 dispersal processes (McCauley and Mabry, 2011). Furthermore, incorporating the size of 616 individuals in a context of climate change is relevant as shifts in body size and its 617 consequences in dispersal (and also in a lot of other life history parameters) have been

618 identified as the third major response to climate change (Gardner et al., 2011). In our 619 dispersal process, we also modelled the attractiveness of the available habitats as a function of their watershed area as we assumed it was a proxy of the river flow. River flow has 620 621 already been showed as being the vector of attractive factors for diadromous species such 622 as larval pheromone by the sea lamprey (*Petromyzon marinus*) (Vrieze and Sorensen, 623 2001) or natural stream odor by salmon (Barinaga, 1999). The function we used for 624 dispersal modelling can then be parameterized and calibrated according to the studied 625 species and the knowledge of its population dynamics. This process has been designed in 626 order to integrate easily other effects that could potentially influence dispersal of a species.

627 Moreover, although our test case dealt with an anadromous species, the GR3D 628 model could also be adapted to catadromous species assuming for these species that 629 dispersers are strayed juveniles that do not use the same growing area (river catchment or 630 estuary) than their parents and have to choose consequently another river basin for their 631 freshwater life. However, the high heterogeneity of diadromous fish species life histories 632 (McDowall, 1988; McDowall, 1997) imply that further specific and local adjustments 633 might be required for different species. For instance, in its current version, GR3D is not 634 able to deal with the diversity and complexity of Atlantic salmon life cycle. A large 635 amount of literature is available for this species describing its population dynamics in 636 many catchments (Mangel, 1994; Rivot et al., 2004; Reed et al., 2011). Recently, Piou and 637 Prevost (2012) developed and parameterized IBASAM, a complex individual-based model 638 to simulate population dynamics of Atlantic salmon in the Scorff River (Brittany, France). 639 Despite the high complexity of this model, it does not incorporate all phenotypes of 640 Atlantic salmon life histories and further developments would also be needed according to 641 the Atlantic salmon population studies. The IBASAM model, described as a tool for the 642 investigation of potential climate change effects on Atlantic salmon population structure,

did not consider individual dispersal and so potential modification of distribution range in
response to environmental change. Consequently, the GR3D and IBASAM models might
be complementary dynamic approaches in order to study Atlantic salmon response to
climate change.

647 As GR3D is a complex model integrating numerous parameters, our study dealt 648 also with the question of the degree of complexity that should be incorporated in modeling 649 approach as the more the complexity is high in a model, the more the model is able to 650 reproduce the observed reality but the more it is difficult to calibrate, to extrapolate and to 651 analyze the model results (Levins, 1966; Caswell and John, 1992). Some authors consider 652 that simple models are the best way to lead to generality in ecology (Holling, 1966) while 653 other authors argue that this point of view may be an obstacle to the progress of ecological 654 research and demonstrate that in some cases at least, complex models are desirable, 655 general, and can be linked with simple models (Lassalle, 2008; Evans et al., 2013). In the 656 GR3D model, several simplifications were made in the structure of the model. First, in its 657 current version, GR3D does not deal with genetic and evolution issues although future 658 works have already been envisaged in this aim. Second, we did not represent explicit 659 movements of individuals within compartments (e.g. with a random walk process 660 (O'Sullivan and Perry, 2009)) and we assumed homogeneous conditions within a 661 compartment. Third, despite the expected impact of climate change on precipitation and 662 therefore river discharge, we did not incorporate the effect of discharge on population 663 dynamics processes as we were not able to link the effect of this parameter on life history 664 traits, largely because models linking precipitation to river flow are complex and focus 665 mostly at local scales (Milly et al., 2005; Qi et al., 2009; Beyene et al., 2010). Finally, 666 GR3D is a generic mono-species model and thus does not take into account potential 667 interactions between or among species. There are thus opportunities for further

668 improvements of the model. However, these improvements would undoubtedly lead to a 669 significant increase in model complexity and consequently of the computation time. In our 670 opinion, our mechanistic representation is realistic enough to address the issue of 671 diadromous fish dispersal and their persistence under climate change at a large scale. 672 Consequently, GR3D appears as a great step forward in management of diadromous fish 673 species. It could be used as an integrative tool to assess potential impact of climate change 674 on endangered diadromous fish species. For instance, we might be able to identify for an 675 endangered population if demographic rescue from neighboring populations will be likely 676 or if restocking program would be relevant and necessary (Seddon et al., 2009; Loss et al., 677 2011). The GR3D model is able to generate predictions of change in the distribution of 678 species but also predictions of population abundances under defined climate change 679 scenario.

680

681 4.2. Model parameterization and sensitivity

682 GR3D was parameterized to reproduce dispersal and population dynamics of a virtual allis 683 shad population between two river catchments in a context of temperature increase. After 684 50 years of simulations to initialise an abundant and stable population in RB1, temperature 685 started to increase and RB2 became gradually suitable while suitability of RB1 gradually 686 decreases. This scenario was a way to mimic a situation where a population threatened by 687 temperature warming had an opportunity to colonize a new suitable habitat. Consequently 688 for analyzing model results, we mainly focused on the colonization of previously 689 uninhabited river basin RB2. With a more complex environment, it would have been 690 difficult to clearly disentangle the effect of the environment and the effect of uncertainties 691 in population dynamics on model outcomes. We chose therefore this simple test case.

692 GR3D is a complex model incorporating a high number of parameters. 693 Consequently, several analyses were necessary to further understand the relative role of 694 different parameters in the model outputs. Such analyses are essential for complex models 695 to avoid misinterpretation of simulation results (Grimm and Railsback, 2005). From this 696 consideration, the fast computation time of the GR3D model is a real advantage. Actually, 697 in our simple test case, the model dealt with millions of individuals but each simulation of 698 200 years took less than one second with a mean duration of simulations around 0.3 699 second. We tested the model with more complex virtual environment (e.g. with 25 river 700 basins) and the duration of simulation was always below one minute.

In the test case, we assumed that dispersal depended only on the dispersal distance. We did not incorporate an effect of the individual length as we considered only one population and as the length of allis shad spawners (i.e. potential dispersers for this species) has always been observed as relatively homogeneous within a population (Bagliniere and Elie, 2000). We also ignored the effect of the watershed area in the dispersal function, as only one river was available for dispersers.

707 Analyzing the success of the colonization of RB2 and the stability of the population 708 in RB2, our sensitivity analyses showed the importance of dispersal distance and of 709 parameters related to the duration of the life at sea ($\kappa_{outGrow}$, L_{mat}) and the survival condition 710 in this compartment (Z_{sea}) . These results were consistent with our expectation and 711 reproduced the expected dynamics of the model. We populated the environment initially in 712 order to have an abundant and stable population in RB1. Consequently, a high number of 713 dispersers were able to colonize the uninhabited river and were only limited by the distance 714 they have to cross. The duration of the life at sea is highly linked to the resilience of 715 diadromous species (i.e. the more the individuals stay at sea, the more the mortality of 716 individuals is high and the resilience of the studied species is low) and so unsurprisingly

717 influences dispersal and stability of population as well as the survival conditions at sea. 718 The analysis of interactions effects showed the importance of the homing (p_{hom}) and of the 719 Allee effect (parameter θ). As the probability of homing determines the number of 720 dispersers, it was also not surprising to obtain a high interaction effect with the dispersal 721 distance. In a same way, we introduced an Allee effect in the reproduction dynamics in 722 order to take into account potential difficulties to establish a population in an uninhabited 723 river basin such as mate limitation and reproductive facilitation problems or a decrease of 724 juvenile mortality with decreasing stock size (Berec et al., 2007). It explains consequently 725 the high effect of the interactions of this parameter. Concerning the abundance of the RB2 726 population, the most influential parameters were, as expected, parameters linked to the 727 success of the reproduction. Actually, parameters $surv_{optRep}$, λ and T_{optRep} defined the 728 survival of individuals between egg and recruit stages and the watershed area of RB2 were 729 directly linked to its production capacity.

Furthermore, model outcomes were not very sensitive to thermal parameters. This result might be explained by a high thermal tolerance of the allis shad species and by the use of the Gironde basin temperature to initialize temperature of RB1 and RB2 (except the spring temperature that we modified to control the suitability of the two rivers). The Gironde system is located in the middle of the allis shad distribution range and, although the Gironde allis shad population is endangered, this might not be because of an environmental change (Rougier et al., 2012).

The results of this test case are a good way to illustrate the dynamics included in the GR3D model. This step was necessary before applying GR3D to a more complex case study to avoid misinterpretation on the model results. Simulation results of the GR3D model might also be compared with results of a species distribution model applied to the same species by Lassalle (2008). We could quantify the divergence in prediction of future

742 potential distribution between static and dynamic approaches as has already be done for 743 two virtual plant species by Engler and Guisan (2009), who showed significant differences 744 between the two kind of approach. Recently, Estes et al. (2013) also compared mechanistic 745 and empirical model projections for the suitability of a crop species and concluded that 746 both approaches might be complementary. Lassalle et al. (2008a) described the 1900 747 observed distribution of European diadromous fish species. Using this 1900 distribution in 748 GR3D as an input defining the initial distribution of individuals, and then, simulate 749 population dynamics from 1900 to today might be a way to validate the GR3D model. We 750 could thus compare projected distributions of GR3D simulation results with the current 751 distribution of European diadromous fish species to analyze whether species have 752 responded in the way that the GR3D model suggests (Kerr and Dobrowski, 2013).

753

754 Acknowledgments

We wish to thank G. Lassalle, F. Daverat and N. Séon-Massin for fruitful discussions helping in developing the model. Sincere thanks are due to Karin Limburg for English corrections and some valuable comments and suggestions. This study was supported by grants from the French National Research Institute of Science and Technology for Environment and Agriculture (Irstea) and the French National Agency for Water and Aquatic Environments (ONEMA).

761

762 **References**

Acolas, M.L., Bégout Anras, M.L., Véron, V., Jourdan, H., Sabatié, M.R., Baglinière, J.L.,

764 2004. An assessment of the upstream migration and reproductive behaviour of allis

shad (*Alosa alosa* L.) using acoustic tracking. ICES J. Mar. Sci. 61, 1291-1304.

- Acolas, M.L., Véron, V., Jourdan, H., Bégout, M.L., Sabatié, M.R., Baglinière, J.L., 2006.
- 767 Upstream migration and reproductive patterns of a population of allis shad in a small
- river (L'Aulne, Brittany, France). ICES J. Mar. Sci. 63, 476-484.
- Allee, W.C., 1931. Animal aggregations: a study in general sociology. University of
 Chigaco Press, Chicago 431 pp.
- 771 Aprahamian, M.W., Bagliniere, J.L., Sabatié, M.R., Alexandrino, P., Aprahamian, C.D.,

2002. Synopsis of biological data on *Alosa alosa* and *Alosa fallax* spp., Warrington.

- Araújo, M.B., Thuiller, W., Pearson, R.G., 2006. Climate warming and the decline of
 amphibians and reptiles in Europe. J. Biogeogr. 33, 1712-1728.
- Baglinière, J.L., 2000. Le genre *Alosa* sp., in: Baglinière, J.L., Elie, P. (eds.), Les aloses
 (*Alosa alosa* et *Alosa fallax* spp.). Cemagref Editions INRA Editions, Paris, pp. 330.
- Bagliniere, J.L., Elie, P., 2000. Les aloses (*Alosa alosa* et *Alosa fallax* spp.) : écobiologie
 et variabilité des populations. Cemagref, Inra Éditions, Paris 275 pp.
- Bagliniere, J.L., Maisse, G., 1985. Precocious maturation and smoltification in wild
 Atlantic salmon in the Armorican massif, France. Aquaculture 45, 249-263.
- 782 Baglinière, J.L., Sabatié, R., Rochard, E., Alexandrino, P., Aprahamian, M.W., 2003. The
- allis shad *Alosa alosa*: biology, ecology, range, and status of populations. Am. Fish.
 Soc. Symp. 35, 85-102.
- Bal, G., Rivot, E., Prévost, E., Piou, C., Baglinière, J.L., 2011. Effect of water temperature
 and density of juvenile salmonids on growth of young-of-the-year Atlantic salmon *Salmo salar*. J. Fish Biol. 78, 1002-1022.
- Barinaga, M., 1999. Salmon follow watery odors home. Science 286, 705-706.

- 789 Barrows, C.W., Preston, K.L., Rotenberry, J.T., Allen, M.F., 2008. Using occurrence
- records to model historic distributions and estimate habitat losses for twopsammophilic lizards. Biol. Conserv. 141, 1885-1893.
- Berec, L., Angulo, E., Courchamp, F., 2007. Multiple Allee effects and population
 management. Trends Ecol. Evol. 22, 185-191.
- Bevacqua, D., Andrello, M., Melià, P., Vincenzi, S., De Leo, G.A., Crivelli, A.J., 2011.
- 795 Density-dependent and inter-specific interactions affecting European eel settlement796 in freshwater habitats. Hydrobiologia 671, 259-265.
- Beverton, R.J.H., Holt, S.J., 1957. On the dynamics of exploited fish populations. Her
 majesty's stationery Office, London 553 pp.
- Beyene, T., Lettenmaier, D.P., Kabat, P., 2010. Hydrologic impacts of climate change on
 the Nile River Basin: Implications of the 2007 IPCC scenarios. Clim. Change 100,
 433-461.
- Bocedi, G., Pe'er, G., Heikkinen, R.K., Matsinos, Y., Travis, J.M.J., 2012. Projecting
 species' range expansion dynamics: sources of systematic biases when scaling up
 patterns and processes. Methods in Ecology and Evolution 3, 1008-1018.
- Brooker, R.W., Travis, J.M.J., Clark, E.J., Dytham, C., 2007. Modelling species' range
 shifts in a changing climate: The impacts of biotic interactions, dispersal distance and
 the rate of climate change. J. Theor. Biol. 245, 59-65.
- Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M., West, G.B., 2004. Toward a
 metabolic theory of ecology. Ecology 85, 1771-1789.
- 810 Buisson, L., 2009. Poissons des rivières françaises et changement climatique : impacts sur
- 811 la distribution des espèces et incertitudes des projections. Thèse de Doctorat. Institut
- 812 National Polytechnique de Toulouse, Toulouse, p. 282.

- 813 Buoro, M., Prévost, E., Gimenez, O., 2010. Investigating evolutionary trade-offs in wild
- 814 populations of Atlantic salmon (*Salmo salar*): Incorporating detection probabilities
- and individual heterogeneity. Evolution 64, 2629-2642.
- 816 Cariboni, J., Gatelli, D., Liska, R., Saltelli, A., 2007. The role of sensitivity analysis in
 817 ecological modelling. Ecol. Model. 203, 167-182.
- 818 Cassou-Leins, J.J., Cassou-Leins, F., Boisneau, P., Baglinière, J.L., 2000. La reproduction,
- 819 in: Baglinières, J.L., Elie, P. (eds.), Les aloses (*Alosa alosa* et *Alosa fallax* spp.).
 820 Cemagref Editions INRA Editions, Paris, pp. 73-92.
- Caswell, H., John, A.M., 1992. From the individual to the population in demographic
 models, in: DeAngelis, D.L., Gross, L.J. (eds.), Individual-based models and
 approaches in ecology: populations, communities and ecosystems. Chapman et Hall,
 New York., pp. 36-61.
- 825 Charles, S., Subtil, F., Kielbassa, J., Pont, D., 2008. An individual-based model to describe
 826 a bullhead population dynamics including temperature variations. Ecol. Model. 215,
 827 377-392.
- 828 Chesson, P., Lee, C.T., 2005. Families of discrete kernels for modeling dispersal. Theor.
 829 Popul. Biol. 67, 241-256.
- 830 Clobert, J., Danchin, E., Dhondt, A.A., Nichols, J.D., 2001. Dispersal. Oxford University
 831 Press, New York 452 pp.
- Clobert, J., Le Galliard, J.-F., Cote, J., Meylan, S., Massot, M., 2009. Informed dispersal,
 heterogeneity in animal dispersal syndromes and the dynamics of spatially structured
 populations. Ecol. Lett. 12, 197-209.
- Bassing De Leo, G.A., Gatto, M., 1995. A size and age-structured model of the European eel
 (Anguilla anguilla L.). Can. J. Fish. Aquat. Sci. 52, 1351-1367.

- B37 Dekker, W., 2000. A Procrustean assessment of the European eel stock. ICES J. Mar. Sci.
 S38 57, 938-947.
- Bion, C.A., Hughes, N.F., 2004. Testing the ability of a temperature-based model to
 predict the growth of age-0 arctic grayling. Trans. Am. Fish. Soc. 133, 1047-1050.
- B41 Droesbeke, J.J., Fine, J., Saporta, G., 1997. Plans d'expériences: applications à l'entreprise.
 B42 Editions Technip, Paris.
- B43 Drouineau, H., Mahévas, S., Bertignac, M., Fertin, A., 2008. Assessing the impact of
 discretisation assumptions in a length-structured population growth model. Fisheries
 Research 91, 160-167.
- Beliaeff, B., 2006. Assessing the impact of
 different management options using ISIS-Fish: The French *Merluccius merluccius* -*Nephrops norvegicus* mixed fishery of the Bay of Biscay. Aquat. Living Resour. 19,
- 849 15-29.
- B50 Dumoulin, N., 2007. SimAquaLife : un cadriciel pour la modélisation de la dynamique
 spatiale d'organismes aquatiques utilisant les métadonnées pour la production
 automatique de fonctionnalités. Techniques et Sciences Informatiques 26.
- Engler, R., Guisan, A., 2009. MigClim: Predicting plant distribution and dispersal in a
 changing climate. Divers. Distrib. 15, 590-601.
- Estes, L.D., Bradley, B.A., Beukes, H., Hole, D.G., Lau, M., Oppenheimer, M.G., Schulze,
 R., Tadross, M.A., Turner, W.R., 2013. Comparing mechanistic and empirical model
 projections of crop suitability and productivity: implications for ecological
- forecasting. Glob. Ecol. Biogeogr. 22, 1007-1018.
- Evans, M.R., Grimm, V., Johst, K., Knuuttila, T., de Langhe, R., Lessells, C.M., Merz, M.,
 O'Malley, M.A., Orzack, S.H., Weisberg, M., Wilkinson, D.J., Wolkenhauer, O.,

- 861 Benton, T.G., 2013. Do simple models lead to generality in ecology? Trends Ecol.
- 862 Evol. 28, 578-583.
- Fabens, A.J., 1965. Properties and fitting of the Von Bertalanffy growth curve. Growth 29,
 265-289.
- Faivre, R., Iooss, B., Mahévas, S., Makowski, D., Monod, H., 2013. Analyse de sensibilité
 et exploration de modèles: application aux sciences de la nature et de
 l'environnement, Editions Quae ed, Versailles 352 pp.
- Fennell, M., Murphy, J.E., Armstrong, C., Gallagher, T., Osborne, B., 2012. Plant Spread
 Simulator: A model for simulating large-scale directed dispersal processes across
 heterogeneous environments. Ecol. Model. 230, 1-10.
- Fleming, I.A., 1996. Reproductive strategies of Atlantic salmon: Ecology and evolution.
 Rev. Fish Biol. Fish. 6, 379-416.
- Forseth, T., Hurley, M.A., Jensen, A.J., Elliott, J.M., 2001. Functional models for growth
 and food consumption of Atlantic salmon parr, *Salmo salar*, from a Norwegian river.
 Freshw. Biol. 46, 173-186.
- 876 Frank, B.M., Baret, P.V., 2013. Simulating brown trout demogenetics in a river/nursery
- brook system: The individual-based model DemGenTrout. Ecol. Model. 248, 184-202.
- Franklin, J., 2010. Moving beyond static species distribution models in support of
 conservation biogeography. Divers. Distrib. 16, 321-330.
- Friedland, K.D., 1998. Ocean climate influences on critical Atlantic salmon (*Salmo salar*)
 life history events. Can. J. Fish. Aquat. Sci. 55, 119-130.
- Gallien, L., Münkemüller, T., Albert, C.H., Boulangeat, I., Thuiller, W., 2010. Predicting
 potential distributions of invasive species: where to go from here? Divers. Distrib.
 16, 331-342.

- 886 Gardner, J.L., Peters, A., Kearney, M.R., Joseph, L., Heinsohn, R., 2011. Declining body
- size: a third universal response to warming? Trends Ecol. Evol. 26, 285-291.
- Gaston, K.J., Blackburn, T.M., 2002. Large-scale dynamics in colonization and extinction
 for breeding birds in Britain. J. Anim. Ecol. 71, 390-399.
- Gatto, M., Rossi, R., 1979. A method for estimating mortalities and abundances of the
 Valli Di Comacchio eels. Memorie dell'Istituto Italiano di Idrobiologia suppl. 37,
 107-114.
- Gienapp, P., Teplitsky, C., Alho, J.S., Mills, J.A., MerilÄ, J., 2008. Climate change and
 evolution: disentangling environmental and genetic responses. Mol. Ecol. 17, 167178.
- Grimm, V., 1999. Ten years of individual-based modelling in ecology: what have welearned and what could we learn in the future? Ecol. Model. 115, 129-148.
- 898 Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V., Giske, J., Goss-Custard, J.,
- Grand, T., Heinz, S.K., Huse, G., 2006. A standard protocol for describing
 individual-based and agent-based models. Ecol. Model. 198, 115-126.
- 901 Grimm, V., Berger, U., DeAngelis, D.L., Polhill, J.G., Giske, J., Railsback, S.F., 2010. The
- 902 ODD protocol: A review and first update. Ecol. Model. 221, 2760-2768.
- 903 Grimm, V., Railsback, S., 2005. Individual-based modeling and ecology. Princeton
 904 University Press 413 pp.
- Guisan, A., Thuiller, W., 2005. Predicting species distribution: offering more than simple
 habitat models. Ecol. Lett. 8, 993-1009.
- 907 Guisan, A., Zimmermann, N.E., 2000. Predictive habitat distribution models in ecology.
 908 Ecol. Model. 135, 147-186.

- 909 Heikkinen, R.K., Luoto, M., Kuussaari, M., Toivonen, T., 2007. Modelling the spatial
- 910 distribution of a threatened butterfly: Impacts of scale and statistical technique.
- 911 Landsc. Urban Plann. 79, 347-357.
- Hein, C.L., Öhlund, G., Englund, G., 2011. Dispersal through stream networks: modelling
 climate-driven range expansions of fishes. Divers. Distrib., no-no.
- Holling, C.S., 1966. The strategy of building models of complex ecological systems, in:

915 Watt, K.E.F. (ed.), Systems Analysis in Ecology. Academic Press, pp. 195-214.

- 916 Huntley, B., Barnard, P., Altwegg, R., Chambers, L., Coetzee, B.W.T., Gibson, L.,
- 917 Hockey, P.A.R., Hole, D.G., Midgley, G.F., Underhill, L.G., Willis, S.G., 2010.
- 918 Beyond bioclimatic envelopes: dynamic species' range and abundance modelling in
- 919 the context of climatic change. Ecography 33, 621-626.
- 920 Huntley, B., Collingham, Y.C., Green, R.E., Hilton, G.M., Rahbek, C., Willis, S.G., 2006.
- 921 Potential impacts of climatic change upon geographical distributions of birds. Ibis922 148, 8-28.
- Huston, M., DeAngelis, D., Post, W., 1988. New computers models unify ecological
 theory. Bioscience 38, 682-692.
- Hutchings, J.A., Jones, M.E.B., 1998. Life history variation and growth rate thresholds for
 maturity in Atlantic salmon, *Salmo salar*. Can. J. Fish. Aquat. Sci. 55, 22-47.
- Jackson, S.T., Betancourt, J.L., Booth, R.K., Gray, S.T., 2009. Ecology and the ratchet of
 events: Climate variability, niche dimensions, and species distributions. Proceedings
- 929 of the National Academy of Sciences 106, 19685-19692.
- 930 Jager, H.I., Cardwell, H.E., Sale, M.J., Bevelhimer, M.S., Coutant, C.C., Van Winkle, W.,
- 931 1997. Modelling the linkages between flow management and salmon recruitment in932 rivers. Ecol. Model. 103, 171-191.

- Jager, H.I., Peterson, D.L., Farrae, D., Bevelhimer, M.S., 2013. A population model to
 assess influences on the viability of the Shortnose sturgeon population in the
 Ogeechee river, Georgia. Trans. Am. Fish. Soc. 142, 731-746.
- Jonsson, B., Jonsson, N., 2009. A review of the likely effects of climate change on
 anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with
 particular reference to water temperature and flow. J. Fish Biol. 75, 2381-2447.
- 939 Keith, D.A., Akçakaya, H.R., Thuiller, W., Midgley, G.F., Pearson, R.G., Phillips, S.J.,
- Regan, H.M., Araújo, M.B., Rebelo, T.G., 2008. Predicting extinction risks under
 climate change: coupling stochastic population models with dynamic bioclimatic
 habitat models. Biology Letters 4, 560-563.
- Kerr, J.T., Dobrowski, S.Z., 2013. Predicting the impacts of global change on species,
 communities and ecosystems: it takes time. Glob. Ecol. Biogeogr. 22, 261-263.
- Kielbassa, J., Delignette-Muller, M.L., Pont, D., Charles, S., 2010. Application of a
 temperature-dependent von Bertalanffy growth model to bullhead (*Cottus gobio*).
 Ecol. Model. 221, 2475-2481.
- Kinnison, M.T., Hairston, N.G., 2007. Eco-evolutionary conservation biology:
 contemporary evolution and the dynamics of persistence. Funct. Ecol. 21, 444-454.
- Kleijnen, J.P.C., 1998. Experimental design for sensitivity analysis, optimization, and
 validation of simulation models, in: Banks, J. (ed.), Handbook of simulation.
 Principles, Methodology, Advances, Applications and Practise. John Wiley, New
- 953 York, pp. 173-224.
- Kottelat, M., Freyhof, J., 2007. Handbook of European freshwater fishes. Kottelat, Cornol,
 Switzerland and Freyhof, Berlin, Germany 646 pp.

- Lambert, P., 2005. Exploration multiscalaire des paradigmes de la dynamique de la
 population d'anguilles européennes à l'aide d'outils de simulation. Thèse de Doctorat.
 Université Bordeaux I, Bordeaux, p. 224.
- Lambert, P., Martin-Vandembulcke, D., Rochard, E., Bellariva, J.L., Castelnaud, G., 2001.
- 960 Age à la migration de reproduction des géniteurs de trois cohortes de grandes aloses
- 961 (Alosa alosa) dans le bassin versant de la Garonne (France) la (France). Bull. Fr.
- 962 Peche Piscic. 362/363, 973-987.
- Lambert, P., Rochard, E., 2007. Identification of the inland population dynamics of the
 European eel using a pattern-oriented modelling. Ecol. Model. 206, 166-178.
- 965 LANUV, 2010. The reintroduction of the allis shad (*Alosa alosa*) to the Rhine System -

966 Life project - LANUV-Fachbericht 28. Recklinghausen.

- Lassalle, G., 2008. Impacts des changements globaux sur la distribution des poissons
 migrateurs amphihalins, une approche par modélisation à l'échelle continentale.
 Thèse de Doctorat. Université Bordeaux I, Bordeaux, p. 244.
- Lassalle, G., Béguer, M., Beaulaton, L., Rochard, E., 2008a. Diadromous fish conservation
 plans need to consider global warming issues: An approach using biogeographical
 models. Biol. Conserv. 141, 1105-1118.
- Lassalle, G., Béguer, M., Beaulaton, L., Rochard, E., 2009. Learning from the past to
 predict the future: responses of European diadromous fish to climate change, in:
 Haro, A.J., Smith, K.L., Rulifson, R.A., Moffitt, C.M., Klauda, R.J., Dadswell, M.J.,
 Cunjak, R.A., Cooper, J.E., Beal, K.L., Avery, T.S. (eds.), Challenges for
 diadromous fishes in a dynamic global environment, vol. Symposium 69. American
 Fisheries Society, Bethesda, Maryland, pp. 175-193.

- Lassalle, G., Trancart, T., Lambert, P., Rochard, E., 2008b. Latitudinal variations in age
 and size at maturity among allis shad *Alosa alosa* populations. J. Fish Biol. 73, 17991809.
- Lehuta, S., Mahévas, S., Petitgas, P., Pelletier, D., 2010. Combining sensitivity and
 uncertainty analysis to evaluate the impact of management measures with ISIS–Fish:
 marine protected areas for the Bay of Biscay anchovy (*Engraulis encrasicolus*)
 fishery. ICES Journal of Marine Science: Journal du Conseil 67, 1063-1075.
- Levins, R., 1966. The strategy of model building in population ecology. Am. Sci. 54, 421–
 431.
- 288 Limburg, K.E., Waldman, J.R., 2009. Dramatic declines in north Atlantic diadromous
 289 fishes. Bioscience 59, 955-965.
- Lochet, A., Jatteau, P., Tomàs, J., Rochard, E., 2008. Retrospective approach to
 investigating the early life history of a diadromous fish: Allis shad *Alosa alosa* (L.)

in the Gironde-Garonne-Dordogne watershed. J. Fish Biol. 72, 946-960.

- Loss, S.R., Terwilliger, L.A., Peterson, A.C., 2011. Assisted colonization: Integrating
 conservation strategies in the face of climate change. Biol. Conserv. 144, 92-100.
- 995 MacKenzie, B.R., Gislason, H., Möllmann, C., Köster, F.W., 2007. Impact of 21st century
- 996 climate change on the Baltic Sea fish community and fisheries. Global Change997 Biology 13, 1348-1367.
- Mallet, J.P., Charles, S., Persat, H., Auger, P., 1999. Growth modelling in accordance with
 daily water temperature in European grayling (*Thymallus thymallus* L.). Can. J. Fish.
 Aquat. Sci. 56, 994-1000.
- Mangel, M., 1994. Climate change and salmonid life history variation. Deep-Sea Research
 Part II: Topical Studies in Oceanography 41, 75-106.

- 1003 Martin Vandembulcke, D., 1999. Dynamique de population de la grande alose (Alosa
- 1004 *alosa*, L. 1758) dans le bassin versant Gironde-Garonne-Dordogne (France): analyse
- 1005 et prévision par modélisation. Thèse de Doctorat. Ecole Nationale Polytechnique,
- 1006 Toulouse, p. 155.
- McCauley, S.J., Mabry, K.E., 2011. Climate change, body size, and phenotype dependentdispersal. Trends Ecol. Evol. 26, 554-555.
- McDowall, R.M., 1988. Diadromy in fishes: migration between freshwater and marineenvironments. Croom Helm, London 308 pp.
- McDowall, R.M., 1997. The evolution of diadromy in fishes (revisited) and its place in
 phylogenetic analysis. Rev. Fish Biol. Fish. 7, 443-462.
- McDowall, R.M., 2009. Making the best of two worlds: diadromy in the evolution,
 ecology, and conservation of aquatic organisms, in: Haro, A.J., Smith, K.L.,
- 1015 Rulifson, R.A., Moffitt, C.M., Klauda, R.J., Dadswell, M.J., Cunjak, R.A., Cooper,
- 1016J.E., Beal, K.L., Avery, T.S. (eds.), Challenges for diadromous fishes in a dynamic1017global environment. American Fisheries Society, Symposium 69, Berthesda,
- 1018 Maryland, pp. 1-22.
- 1019 Mennesson-Boisneau, C., Aprahamian, M.W., Sabatié, M.R., Cassou-Leins, J.J., 2000a.
- 1020 Caractéristiques des adultes, in: Baglinière, J.L., Elie, P. (eds.), Les aloses (*Alosa*1021 *alosa* et *Alosa fallax* spp.). Cemagref Editions INRA Editions, Paris, pp. 33-54.
- Mennesson-Boisneau, C., Aprahamian, M.W., Sabatié, M.R., Cassou-Leins, J.J., 2000b.
 Remontée migratoire des adultes, in: Baglinière, J.L., Elie, P. (eds.), Les aloses
- 1024 (*Alosa alosa* et *Alosa fallax* spp.). Cemagref Editions INRA Editions, Paris, pp. 551025 72.

- 1026 Mennesson-Boisneau, C., Boisneau, P., 1990. Migration, répartition, reproduction,
- 1027 caractéristiques biologiques et taxonomie des aloses (Alosa sp) dans le bassin de la
- 1028 Loire. Université Rennes I et Paris XII, p. 143.

- 1029 Merow, C., Lafleur, N., Silander Jr, J.A., Wilson, A.M., Rubega, M., 2011. Developing
- 1030 dynamic mechanistic species distribution models: Predicting bird-mediated spread of 1031 invasive plants across northeastern North America. Am. Nat. 178, 30-43.
- Midgley, G.F., Davies, I.D., Albert, C.H., Altwegg, R., Hannah, L., Hughes, G.O., 1032
- O'Halloran, L.R., Seo, C., Thorne, J.H., Thuiller, W., 2010. BioMove an integrated 1034 platform simulating the dynamic response of species to environmental change. 1035 Ecography 33, 612-616.
- 1036 Midgley, G.F., Hannah, L., Millar, D., Rutherford, M.C., Powrie, L.W., 2002. Assessing the vulnerability of species richness to anthropogenic climate change in a 1037 1038 biodiversity hotspot. Glob. Ecol. Biogeogr. 11, 445-451.
- 1039 Milly, P.C.D., Dunne, K.A., Vecchia, A.V., 2005. Global pattern of trends in streamflow 1040 and water availability in a changing climate. Nature 438, 347-350.
- 1041 Mota, M., Antunes, C., 2011. First report on the status of Allis shad (Alosa alosa) in the 1042 Minho River (Northwestern Iberian Peninsula). J. Appl. Ichthyol. 27, 56-59.
- 1043 Nicolas, D., Chaalali, A., Drouineau, H., Lobry, J., Uriarte, A., Borja, A., Boët, P., 2011.
- 1044 Impact of global warming on European tidal estuaries: some evidence of northward 1045 migration of estuarine fish species. Reg Environ Change 11, 639-649.
- 1046 O'Sullivan, D., Perry, G.L.W., 2009. A discrete space model for continuous space dispersal 1047 processes. Ecological Informatics 4, 57-68.
- 1048 Parmesan, C., Yohe, G., 2003. A globally coherent fingerprint of climate change impacts 1049 across natural systems. Nature 421, 37-42.

- Peterson, R.H., Spinney, H.C.E., Sreedharan, A., 1977. Development of atlantic salmon
 (*Salmo salar*) eggs and alevins under varied temperature regimes. J. Fish. Res. Board
 Can. 34, 31-43.
- Piou, C., Prévost, E., 2012. A demo-genetic individual-based model for Atlantic salmon
 populations: Model structure, parameterization and sensitivity. Ecol. Model. 231, 37-
- 1055 52.
- Piou, C., Prévost, E., 2013. Contrasting effects of climate change in continental vs. oceanic
 environments on population persistence and microevolution of Atlantic salmon.
 Global Change Biology 19, 711-723.
- Prouzet, P., Martinet, J.P., Badia, J., 1994. Biological characteristics and catch variation of
 allis shad (*Alosa alosa*) from commercial catches in the Adour River (Pyrenees
 atlantiques, France). Aquat. Living Resour. 7, 1-10.
- Pulliam, H.R., 2000. On the relationship between niche and distribution. Ecol. Lett. 3, 349-361.
- Qi, S., Sun, G., Wang, Y., McNulty, S.G., Myers, J.A.M., 2009. Streamflow response to
 climate and landuse changes in a coastal watershed in North Carolina. Transactions
 of the ASABE 52, 739-749.
- Quinn, T.P., Doctor, K., Kendall, N., Rich, H.B., 2009. Diadromy and the life history of
 sockeye salmon: nature, nurture and the hand of man, in: Haro, A.J., Smith, K.L.,
 Rulifson, R.A., Moffitt, C.M., Klauda, R.J., Dadswell, M.J., Cunjak, R.A., Cooper,
 J.E., Beal, K.L., Avery, T.S. (eds.), Challenges for diadromous fishes in a dynamic
- 1071 global environment. American Fisheries Society, Symposium 69, Berthesda,1072 Maryland, pp. 23-42.
- 1073 R Core Team, 2013. R: A language and environment for statistical computing.

- 1074 Reed, T.E., Schindler, D.E., Hague, M.J., Patterson, D.A., Meir, E., Waples, R.S., Hinch,
- 1075 S.G., 2011. Time to evolve? Potential evolutionary responses of Fraser river sockeye
- salmon to climate change and effects on persistence. PLoS ONE 6, e20380.
- 1077 Reuillon, R., Leclaire, M., Rey-Coyrehourcq, S., 2013. OpenMOLE, a workflow engine
 1078 specifically tailored for the distributed exploration of simulation models. Future
 1079 Generation Computer Systems 29, 1981-1990.
- Rivot, E., Prevost, E., Parent, E., Bagliniere, J.L., 2004. A Bayesian state-space modelling
 framework for fitting a salmon stage-structured population dynamic model to
 multiple time series of field data. Ecol. Model. 179, 463-485.
- Rochard, E., Lambert, P., 2011. Chapter 30 Modelling the future of stocked fish. Biology
 and conservation of the Atlantic European sturgeon Acipenser sturio L., 1758, 417424.
- Ronce, O., 2007. How does it feel to be like a rolling stone? Ten questions about dispersal
 evolution. Annual Review of Ecology, Evolution, and Systematics 38, 231-253.
- 1088 Rosso, L., Lobry, J.R., Bajard, S., Flandrois, J.P., 1995. Convenient model to describe the
- 1089 combined effects of temperature and pH on microbial growth. Appl. Environ.1090 Microbiol. 61, 610-616.
- Rougier, T., Lambert, P., Drouineau, H., Girardin, M., Castelnaud, G., Carry, L.,
 Aprahamian, M., Rivot, E., Rochard, E., 2012. Collapse of allis shad, *Alosa alosa*, in
 the Gironde system (southwest France): environmental change, fishing mortality, or
- 1094Allee effect? ICES Journal of Marine Science: Journal du Conseil 69, 1802-1811.
- Saltelli, A., 2004. What is sensitivity analysis?, in: Saltelli, A., Chan, K., Scott, E.M.
 (eds.), Sensitivity Analysis. Probability and Statistics Series. John Wiley, New york.
 475 pp, pp. 3-13.

- 1098 Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., Saisana,
- 1099 Tarantola, S., 2008. Global sensitivity analysis: the primer. Wiley, New York.
- 1100 Seddon, P.J., Armstrong, D.P., Soorae, P., Launay, F., Walker, S., Ruiz-Miranda, C.R.,
- 1101 Molur, S., Koldewey, H., Kleiman, D.G., 2009. The risks of assisted colonization.
- 1102 Conserv. Biol. 23, 788-789.
- Segurado, P., Araújo, M.B., 2004. An evaluation of methods for modelling species
 distributions. J. Biogeogr. 31, 1555-1568.
- Slone, D.H., 2011. Increasing accuracy of dispersal kernels in grid-based populationmodels. Ecol. Model. 222, 573-579.
- Stephens, P.A., Sutherland, W.J., Freckleton, R.P., 1999. What is the Allee effect? Oikos87, 185-190.
- 1109 Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham,
- 1110 Y.C., Erasmus, B.F.N., Ferreira De Siqueira, M., Grainger, A., Hannah, L., Hughes,
- 1111 L., Huntley, B., Van Jaarsveld, A.S., Midgley, G.F., Miles, L., Ortega-Huerta, M.A.,
- 1112 Peterson, A.T., Phillips, O.L., Williams, S.E., 2004. Extinction risk from climate
- 1113 change. Nature 427, 145-148.
- 1114 Thomas, C.D., Lennon, J.J., 1999. Birds extend their ranges northwards. Nature 399, 213.
- Thuiller, W., 2003. BIOMOD optimizing predictions of species distributions and
 projecting potential future shifts under global change. Global Change Biology 9,
 1117 1353-1362.
- 1118 Thuiller, W., Albert, C., Araújo, M.B., Berry, P.M., Cabeza, M., Guisan, A., Hickler, T.,
- 1119 Midgley, G.F., Paterson, J., Schurr, F.M., Sykes, M.T., Zimmermann, N.E., 2008.
- 1120 Predicting global change impacts on plant species' distributions: Future challenges.
- 1121 Perspectives in Plant Ecology, Evolution and Systematics 9, 137-152.

- Thuiller, W., Broennimann, O., Hughes, G., Alkemade, J.R.M., Midgley, G.F., Corsi, F.,
 2006. Vulnerability of African mammals to anthropogenic climate change under
- 1124 conservative land transformation assumptions. Global Change Biology 12, 424-440.
- Tomas, J., Augagneur, S., Rochard, E., 2005. Discrimination of the natal origin of youngof-the-year Allis shad (*Alosa alosa*) in the Garonne-Dordogne basin (south-west
 France) using otolith chemistry. Ecol. Freshw. Fish. 14, 185-190.
- Travis, J.M.J., Mustin, K., Bartoń, K.A., Benton, T.G., Clobert, J., Delgado, M.M.,
 Dytham, C., Hovestadt, T., Palmer, S.C.F., Van Dyck, H., Bonte, D., 2012.
 Modelling dispersal: an eco-evolutionary framework incorporating emigration,
 movement, settlement behaviour and the multiple costs involved. Methods in
- 1132 Ecology and Evolution 3, 628-641.
- 1133 Van Winkle, W., Jager, H.I., Railsback, S.F., Holcomb, B.D., Studley, T.K., Baldrige, J.E.,
 1134 1998. Individual-based model of sympatric populations of brown and rainbow trout
 1135 for instream flow assessment: model description and calibration. Ecol. Model. 110,
- 1136 175-207.
- 1137 Van Winkle, W., Rose, K.A., Chambers, R.C., 1993. Individual-based approach to fish
 population dynamics: An overview. Trans. Am. Fish. Soc. 122, 397-403.
- Visser, M.E., 2008. Keeping up with a warming world; assessing the rate of adaptation to
 climate change. Proceedings of the Royal Society B: Biological Sciences 275, 649659.
- Vollestad, L.A., Jonsson, B., 1988. A 13-year study of the population dynamics and
 growth of the European eel *Anguilla anguilla* in a Norwegian river: Evidence for
 density-dependent mortality, and development of a model for predicting yield. J.
 Anim. Ecol. 57, 983-997.

- 1146 von Bertalanffy, L., 1938. A quantitative theory of organic growth. Hum. Biol. 10, 181-
- 1147 213.
- 1148 Vrieze, L.A., Sorensen, P.W., 2001. Laboratory assessment of the role of a larval 1149

pheromone and natural stream odor in spawning stream localization by migratory sea

- 1150 lamprey (Petromyzon marinus). Can. J. Fish. Aquat. Sci. 58, 2374-2385.
- 1151 Walther, G.R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T.J.C., Fromentin,
- 1152 J.M., Hoegh-Guldberg, O., Bairlein, F., 2002. Ecological responses to recent climate 1153 change. Nature 416, 389-395.
- 1154 Wheeler, B., 2011. AlgDesign: Algorithmic Experimental Design. R package version 1.1-
- 1155 7. http://CRAN.R-project.org/package=AlgDesign.
- 1156 Zimmermann, N.E., Yoccoz, N.G., Edwards Jr, T.C., Meier, E.S., Thuiller, W., Guisan, A.,
- 1157 Schmatz, D.R., Pearman, P.B., 2009. Climatic extremes improve predictions of
- 1158 spatial patterns of tree species. Proc. Natl. Acad. Sci. U. S. A. 106, 19723-19728.
- 1159
- 1160
- 1161

List of tables :

Table1. List of agents intervening in the GR3D individual-based model, with their state variables and corresponding status or measure unit.

Agent	State Variable	Descritpion	Status/unit of measure
Sea basin	name	Name ID	SB and a number
	seasonal temperature	Water temperature	Numeric (°C)
River basin	name	Name ID	RB and a number
	longitude	Longitude at the mouth	Numeric (°)
	latitude	Latitude at the mouth	Numeric(°)
	watershed area	Watershed area of the river basin	Numeric (km ²)
	seasonal temperature	Water temperature	Numeric (°C)
Diadromous fish	ID	Fish identification	Numeric (-)
	gender	Sex	M, F
	stage	Stage	Mature, immature
	age	Age	Numeric (-)
	body-length	Body length	Numeric (cm)
	location	Current location	Name of the compartment
	birth place	Birth place	Name of the birth compartment
	number of reproduction	Number of reproduction	Numeric (-)

Table 2. GR3D parameters description and selected nominal values for the test case with allis

shad.

Parameter name	Description	Value	Reference or remarks
Reproduction			
repSeason	Season of the reproduction	Spring	(Mennesson-Boisneau et al., 2000b)
Δt_{rec}	Assumed age of juvenile produced by the reproduction (Year)	0.33	As we assumed that recruitment were juveniles in estuary, we used the study from Lochet et al. (2008)
η	Parameter to relate $S_{95,j}$ and the surface of a spawning place (Ind/km ²)	2.4	Offline calibration using the study from Rougier et al. (2012)
θ	Ratio between $S_{95,j}$ and $S_{50,j}$ in each spawning place	1.9	Offline calibration using the study from Rougier et al. (2012)
a	Parameter of the S-R relationship linked to the fecundity of the studied species (Eggs/individual)	135000	(Cassou-Leins et al., 2000; LANUV, 2010)
SURV _{optRep}	Optimal survival rate of an individual from eggs to the age Δt_{rec}	1.7*10 ⁻³	Offline calibration using the study from Rougier et al. (2012)
$T_{minRep}, T_{optRep}, T_{maxRep}$	Water temperature (°C) regulating survival of an individual from eggs to the age Δt_{rec}	14, 20, 26	(Cassou-Leins et al., 2000; Kottelat and Freyhof, 2007)
λ	Parameter to relate c_j and the surface of a spawning place	4.1*10 ⁻⁴	Offline calibration using the study from Rougier et al. (2012)
$\sigma_{\scriptscriptstyle rep}$	Standard deviation of log- normal distribution of the recruitment	0.2	Reasonable guessed
Sp_{sp}	Survival probability of spawners after reproduction (i.e. iteroparous rate)	0.1	(Mennesson-Boisneau et al., 2000b)
Downstream migration			
downMigAge	Age of individual when it runs toward the sea (Year)	0.33	(Lochet et al., 2008); In the test case, this migration concerns also spawners which survive after reproduction but, as mature individuals, they migrate automatically at sea the season following the reproduction In the test case, both juveniles and
downMigSeason	Season of the run of individuals toward the sea	summer	spawners which survive after reproduction migrate in summer (Cassou-Leins et al., 2000)
Growth			
L_{ini}	Initial length of juvenile in estuary (cm)	2	As we considered juveniles in estuary, we used the study from Lochet et al. (2008)
$\sigma_{_{\Delta\!L}}$	Standard deviation of log- normal distribution of the	0.2	Reasonable guessed

L_{∞}	growth increment Asymptotic length of the individual (cm)	60	(Mennesson-Boisneau et al., 2000a)
$T_{minGrow}, T_{optGrow},$ $T_{maxGrow}$	Water temperature (°C) regulating the growth	3, 17, 26	Reasonable guessed
$\kappa_{optGrow}$	Optimal growth coefficient (cm/season)	0.3	(Bagliniere and Elie, 2000; Aprahamian et al., 2002)
Survival			
Z_{sea}	Annual mortality coefficient at sea (Year ⁻¹)	0.4	Reasonable guessed based upon Rougier et al. (2012)
H _{riv}	from natural) coefficient in river (Year ⁻¹)	0	Reasonable guessed to 0 to limit the complexity of the test case
$T_{minSurvRiv}$, $T_{optSurvRiv}$, $T_{maxSurvRiv}$	Water temperature (°C) regulating survival of individual in river	10, 20, 23	In the test case, it concerns only spawners during the reproduction run and this phase has been shown highly dependent of water temperature (Cassou-Leins et al., 2000; Kottelat and Freyhof, 2007)
SURV _{optRiv}	Optimal natural survival rate of individuals in river (Year ⁻¹)	1	Natural mortality was assumed to be negligible at optimal conditions during the reproduction run as it concerns a short period of 3 months (Cassou- Leins et al., 2000)
Maturation			
	Length at first maturity (cm)	40	(Cassou-Leins et al., 2000; Lassalle et al., 2008b)
Upstream migration			In the test area, this migration
upMigAge	Age of individual when it runs toward the river (Year)	-	concerns only mature individuals and is not age-specific consequently
upMigSeason	Season of the return of spawners in river for spawning	spring	All mature individuals at sea migrate in river at this season
p_{hom}	Probability to do natal homing behavior	0.75	(Tomas et al., 2005)
$lpha_{_{const}},\ lpha_{_{dist}},\ lpha_{_{TL}}, \ lpha_{_{W\!A}}$	Parameters of the logit function used to determine the weight of each accessible basin for dispersers	-2.9, 19.7, 0, 0	Reasonable guessed assuming a weight of 0.95 and 0.05 for a basin located at a distance of 10km and 300km respectively and assuming no effect of individual size and surface of basins on dispersal
$\overline{D_{j-birthPlace}}$, $\sigma_{j-birthPlace}$, \overline{TL} , σ_{TL} , \overline{WA} , σ_{WA}	Mean and standard deviation used for standard core values in the logit function	300, 978,-,-,-,-	Reasonable guessed assuming that 300 km was close to the maximal distance of allis shad dispersal and from EuroDiad 2.0 database (Lassalle, 2008). Parameters linked to fish length and surface basin were not relevant in the test case.
W ^{deathBasin}	Weigth of the death basin used to introduced a mortality of dispersers	0.4	Reasonable guessed

-

Table 3. Uncertain population dynamics parameters of the GR3D model and environmental

Parameter name	Tested values	Selected for complete sampling (Yes or No)
Reproduction SM1		
η	[2.2; 2.6]	Ν
heta	[1.8; 2]	Y
SURV _{optRep}	$[1*10^{-3}; 2*10^{-3}]$	Y
T_{minRep}	[13.3; 14.6]	Ν
T_{optRep}	[19.3; 20.6]	Y
T_{maxRep}	[25.3; 26.6]	Ν
λ	[3*10 ⁻⁴ ; 5*10 ⁻⁴]	Y
$\sigma_{_{rep}}$	[0.1; 0.3]	Ν
Growth SM2		
$\sigma_{_{\Delta\!L}}$	[0.1; 0.3]	Ν
$T_{minGrow}$	[2.3; 3.6]	Ν
$T_{optGrow}$	[16.3; 17.6]	Y
$T_{maxGrow}$	[25.3; 26.6]	Ν
$\kappa_{optGrow}$	[0.2; 0.4]	Y
Survival SM3		
Z_{sea}	[0.3; 0.5]	Y
$T_{minSurvRiv}$	[9.3; 10.6]	Ν
$T_{optSurvRiv}$	[19.3; 20.6]	Y
T _{maxSurvRiv}	[22.3; 23.6]	Ν
Maturation SM5		
L _{mat}	[36; 44]	Y
Anadromous migration S	SM6	
p_{hom}	[0.6; 0.9]	Y
W ^{deathBasin}	[0.2; 0.6]	Ν
Environmental paramete	ers	
WA_RB2	[20000; 40000; 80000]	Y
intDist	[100; 300; 600; 900]	Ŷ

parameters tested in the sensitivity analysis of the test case.

Table 4. Percentage of explained deviance (for *fsP* and *fsS*) and explained variance (for *mlr*)

by the model applied to outputs variables according to the experimental.

	Output variable		
Design	fsP	fsS	mlr
D-Optimal design	78%	70%	60%
Complete sampling			
Without interaction effects	81%	69%	54%
With interaction effects	90%	85%	79%

List of figures :

Fig. 1. Conceptual diagram of the life cycle of anadromous species (adapted for allis shad) represented in the GR3D individual-based model.



Fig. 2. Illustrative example of simulation results over 200 years showing the dynamics over time of the recruitment in RB1 and RB2 using the nominal values of the model parameters and with (A) a distance of 300 km between RB1 and RB2 and a success of the colonization of RB2 (fs = 2), (B) a distance of 600 km between RB1 and RB2 and very low recruitments in RB2 at the end of simulation (fs = 1), and (C) a distance of 900 km between RB1 and RB2 and no colonization of RB2 (fs = 0).



Fig. 3. Ten highest SI values (in %; the higher the percentage, the more sensitive the model is to the considered parameter) for the *fsP* (A), *.fsS* (B) and *mlr* (C) output variables calculated from logistic regression for *fsP* and *fsS* and ANOVA for *mlr* with the D-Optimal experimental design (theta and lambda correspond to the parameters θ and λ of the reproduction process respectively).



Fig. 4. TSI values (in %; the higher the percentage, the more sensitive the model is to the considered parameter) for the *fsP* (A), *.fsS* (B) and *mlr* (C) output variables calculated from logistic regression on *fsP* and *fsS* and ANOVA on *mlr* with the complete sampling design (theta, lambda, weightDB correspond to the parameters θ , λ and $w^{deathBasin}$ respectively).

