



HAL
open science

Continuous fields of properly infinite C^* -algebras

Etienne Blanchard

► **To cite this version:**

| Etienne Blanchard. Continuous fields of properly infinite C^* -algebras. 2015. hal-01123514v5

HAL Id: hal-01123514

<https://hal.science/hal-01123514v5>

Preprint submitted on 15 Apr 2015 (v5), last revised 17 Apr 2015 (v7)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONTINUOUS FIELDS OF PROPERLY INFINITE C*-ALGEBRAS

ETIENNE BLANCHARD

ABSTRACT. Any unital separable continuous $C(X)$ -algebra with properly infinite fibres is properly infinite as soon as the compact Hausdorff space X has finite topological dimension. We study conditions under which this is still the case in the infinite dimensional case.

1. INTRODUCTION

One of the basic C*-algebras studied in the classification programme launched by G. Elliott ([Ell94]) of nuclear C*-algebras through K-theoretical invariants is the Cuntz C*-algebra \mathcal{O}_∞ generated by infinitely many isometries with pairwise orthogonal ranges ([Cun77]). This C*-algebra is pretty rigid in so far as it is a strongly self-absorbing C*-algebra ([TW07]): Any separable unital continuous $C(X)$ -algebra A the fibres of which are isomorphic to the same strongly self-absorbing C*-algebra D is a trivial $C(X)$ -algebra provided the compact Hausdorff base space X has finite topological dimension. Indeed, the strongly self-absorbing C*-algebra D tensorially absorbs the Jiang-Su algebra \mathcal{Z} ([Win09]). Hence, this C*-algebra D is K_1 -injective ([Rør04]) and the $C(X)$ -algebra A satisfies $A \cong D \otimes C(X)$ ([DW08]).

But I. Hirshberg, M. Rørdam and W. Winter have built a non-trivial unital continuous C*-bundle over the infinite dimensional compact product $\prod_{n=0}^\infty S^2$ such that all its fibres are isomorphic to the strongly self-absorbing UHF algebra of type 2^∞ ([HRW07, Example 4.7]). More recently, M. Dădărlat has constructed in [Dăd09, §3] for all pair (Γ_0, Γ_1) of countable abelian torsion groups a unital separable continuous $C(X)$ -algebra A such that

- the base space X is the compact Hilbert cube $X = B_\infty$ of infinite dimension,
- all the fibres A_x ($x \in B_\infty$) are isomorphic to the strongly self-absorbing Cuntz C*-algebra \mathcal{O}_2 generated by two isometries s_1, s_2 satisfying $1_{\mathcal{O}_2} = s_1 s_1^* + s_2 s_2^*$,
- $K_i(A) \cong C(Y_0, \Gamma_i)$ for $i = 0, 1$, where $Y_0 \subset [0, 1]$ is the canonical Cantor set.

These K-theoretical conditions imply that the $C(B_\infty)$ -algebra A is not a trivial one. But these arguments do not work anymore when the strongly self-absorbing algebra D is the Cuntz algebra \mathcal{O}_∞ ([Cun77]), in so far as $K_0(\mathcal{O}_\infty) = \mathbb{Z}$ is a torsion free group.

We describe in this article the link between several notions of proper infiniteness for $C(X)$ -algebras which appeared during the recent years ([KR00],[BRR08], [CEI08], [RR11]). We then study whether certain unital continuous $C(X)$ -algebras with fibres \mathcal{O}_∞ , especially the Pimsner-Toeplitz algebra ([Pim95]) of Hilbert $C(X)$ -modules with

2010 *Mathematics Subject Classification.* Primary: 46L80; Secondary: 46L06, 46L35.

Key words and phrases. C*-algebra, Classification, Proper Infiniteness.

fibres of dimension greater than 2 and with compact base space X of infinite topological dimension, are properly infinite C^* -algebras.

I especially thank E. Kirchberg and a referee for a few inspiring remarks.

2. A FEW NOTATIONS

We present in this section the main notations which are used in this article. We denote by $\mathbb{N} = \{0, 1, 2, \dots\}$ the set of positive integers and we denote by $[S]$ the closed linear span of a subset S in a Banach space.

Definition 2.1. ([Dix69], [Kas88], [Blan97]) *Let X be a compact Hausdorff space and let $C(X)$ be the C^* -algebra of continuous function on X .*

- *A unital $C(X)$ -algebra is a unital C^* -algebra A endowed with a unital morphism of C^* -algebra from $C(X)$ to the centre of A .*
- *For all closed subset $F \subset X$ and all element $a \in A$, one denotes by $a|_F$ the image of a in the quotient $A|_F := A/C_0(X \setminus F) \cdot A$. If $x \in X$ is a point in X , one calls fibre at x the quotient $A_x := A|_{\{x\}}$ and one write a_x for $a|_{\{x\}}$.*
- *The $C(X)$ -algebra A is said to be continuous if the upper semicontinuous map $x \in X \mapsto \|a_x\| \in \mathbb{R}_+$ is continuous for all $a \in A$.*

Remarks 2.2. a) ([Cun81], [BRR08]) For all integer $n \geq 2$, the C^* -algebra $\mathcal{T}_n := \mathcal{T}(\mathbb{C}^n)$ is the universal unital C^* -algebra generated by n isometries s_1, \dots, s_n satisfying the relation

$$s_1 s_1^* + \dots + s_n s_n^* \leq 1. \quad (2.1)$$

b) A unital C^* -algebra A is said to be *properly infinite* if and only if one the following equivalent conditions holds true ([Cun77], [Rør03, Proposition 2.1]):

- the C^* -algebra A contains two isometries with mutually orthogonal range projections, *i.e.* A unittally contains a copy of \mathcal{T}_2 ,
- the C^* -algebra A contains a unital copy of the simple Cuntz C^* -algebra \mathcal{O}_∞ generated by infinitely many isometries with pairwise orthogonal ranges.

c) If A is a C^* -algebra and E is a Hilbert A -module, one denotes by $\mathcal{L}(E)$ the set of adjointable A -linear operators acting on E ([Kas88]).

3. GLOBAL PROPER INFINITENESS

Proposition 2.5 of [BRR08] and section 6 of [Blan13] entail the following stable proper infiniteness for continuous $C(X)$ -algebras with properly infinite fibres.

Proposition 3.1. *Let X be a second countable perfect compact Hausdorff space, *i.e.* without any isolated point, and let A be a separable unital continuous $C(X)$ -algebra with properly infinite fibres.*

1) *There exist:*

- (a) *a finite integer $n \geq 1$,*
- (b) *a covering $X = \overset{\circ}{F}_1 \cup \dots \cup \overset{\circ}{F}_n$ by the interiors of closed balls F_1, \dots, F_n ,*

(c) *unital embeddings of C*-algebra* $\sigma_k : \mathcal{O}_\infty \hookrightarrow A|_{F_k}$ ($1 \leq k \leq n$).

2) *The tensor product* $M_p(\mathbb{C}) \otimes A$ *is properly infinite for all large enough integer* p .

Proof. 1) For all point $x \in X$, the semiprojectivity of the C*-subalgebra $\mathcal{O}_\infty \hookrightarrow A_x$ ([Blac04, Theorem 3.2]) entails that there are a closed neighbourhood $F \subset X$ of the point x and a unital embedding $\mathcal{O}_\infty \otimes C(F) \hookrightarrow A|_F$ of $C(F)$ -algebra. The compactness of the topological space X enables to conclude.

2) Proposition 2.7 of [BRR08] entails that the C*-algebra $M_{2^{n-1}}(A)$ is properly infinite. Proposition 2.1 of [Rør97] implies that $M_p(A)$ for all integer $p \geq 2^{n-1}$. \square

Remark 3.2. If X is a second countable compact Hausdorff space and A is a separable unital continuous $C(X)$ -algebra, then $\tilde{X} := X \times [0, 1]$ is a perfect compact space, $\tilde{A} := A \otimes C([0, 1])$ is a unital continuous $C(\tilde{X})$ -algebra and every morphism of unital C*-algebra $\mathcal{O}_\infty \rightarrow \tilde{A}$ induces a unital *-homomorphism $\mathcal{O}_\infty \rightarrow A$ by composition with the projection map $\tilde{A} \rightarrow A$ coming from the injection $x \in \tilde{X} \mapsto (x, 0) \in \tilde{X}$.

The proper infiniteness of the tensor product $M_p(\mathbb{C}) \otimes A$ does not always imply that the C*-algebra A is properly infinite ([HR98]). Indeed, there exists a unital C*-algebra A which is not properly infinite, but such that the tensor product $M_2(\mathbb{C}) \otimes A$ is a properly infinite C*-algebra ([Rør03, Proposition 4.5]). The following corollary nevertheless holds true.

Corollary 3.3. *Let* j_0, j_1 *denote the two canonical unital embeddings of the Cuntz extension* \mathcal{T}_2 *in the full unital free product* $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$ *and let* $\tilde{u} \in \mathcal{U}(\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2)$ *be a* K_1 -*trivial unitary satisfying* $j_1(s_1) = \tilde{u} \cdot j_0(s_1)$ ([BRR08, Lemma 2.4]).

The following assertions are equivalent:

- (a) *The full unital free product* $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$ *is* K_1 -*injective.*
- (b) *The unitary* \tilde{u} *belongs to the connected component* $\mathcal{U}_0(\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2)$ *of* $1_{\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2}$.
- (c) *Every separable unital continuous* $C(X)$ -*algebra* A *with properly infinite fibres is a properly infinite C*-algebra.*

Proof. (a) \Rightarrow (b) A unital C*-algebra A is called K_1 -injective if and only if all K_1 -trivial unitaries $v \in \mathcal{U}(A)$ are homotopic to the unit 1_A in $\mathcal{U}(A)$ (see e.g. [Roh09]). Thus, (b) is a special case of (a) since $K_1(\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2) = \{1\}$ (see e.g. [Blan10, Lemma 4.4]).

(b) \Rightarrow (c) Let A be a separable unital continuous $C(X)$ -algebra with properly infinite fibres. Take a finite covering $X = \overset{\circ}{F}_1 \cup \dots \cup \overset{\circ}{F}_n$ such that there exist unital embeddings $\sigma_k : \mathcal{T}_2 \rightarrow A|_{F_k}$ for all $1 \leq k \leq n$. Set $G_k := F_1 \cup \dots \cup F_k \subset X$ and let us construct by induction isometries $w_k \in A|_{G_k}$ such that the two projections $w_k w_k^*$ and $1|_{G_k} - w_k w_k^*$ are properly infinite and full in the restriction $A|_{G_k}$:

- If $k = 1$, the isometry $w_1 := \sigma_1(s_1)$ has the requested properties.
- If $k \in \{1, \dots, n-1\}$ and the isometry $w_k \in A|_{G_k}$ is already constructed, then Lemma 2.4 of [BRR08] implies that there exists a morphism of unital C*-algebra $\pi_k :$

$\mathcal{T}_2 *_C \mathcal{T}_2 \rightarrow A|_{G_k \cap F_{k+1}}$ satisfying

$$\begin{aligned} - \quad & \pi_k(j_0(s_1)) = w_{k|G_k \cap F_{k+1}}, \\ - \quad & \pi_k(j_1(s_1)) = \sigma_{k+1}(s_1)|_{G_k \cap F_{k+1}} = \pi_k(\tilde{u}) \cdot w_{k|G_k \cap F_{k+1}}. \end{aligned} \quad (3.1)$$

If the unitary \tilde{u} belongs to the connected component $\mathcal{U}_0(\mathcal{T}_2 *_C \mathcal{T}_2)$, then $\pi_k(\tilde{u})$ is homotopic to $1_{A|_{G_k \cap F_{k+1}}} = \pi_k(1_{\mathcal{T}_2 *_C \mathcal{T}_2})$ in $\mathcal{U}(A|_{G_k \cap F_{k+1}})$, so that $\pi_k(\tilde{u})$ admits a unitary lifting z_{k+1} in $\mathcal{U}_0(A|_{F_{k+1}})$ (see *e.g.* [LLR00, Lemma 2.1.7]). The only isometry $w_{k+1} \in A|_{G_{k+1}}$ satisfying the two constraints

$$\begin{aligned} - \quad & w_{k+1|G_k} = w_k, \\ - \quad & w_{k+1|F_{k+1}} = (z_{k+1})^* \cdot \sigma_{k+1}(s_1) \end{aligned} \quad (3.2)$$

verifies that the two projections $w_{k+1}w_{k+1}^*$ and $1_{|G_{k+1}} - w_{k+1}w_{k+1}^*$ are properly infinite and full in $A|_{G_{k+1}}$.

The proper infiniteness of the projection $w_n w_n^*$ in $A|_{G_n} = A$ implies that the unit $1_A = w_n^* w_n = w_n^* \cdot w_n w_n^* \cdot w_n$ is also a properly infinite projection in A , *i.e.* the C^* -algebra A is properly infinite.

(c) \Rightarrow (a) The C^* -algebra $\mathcal{D} := \{f \in C([0, 1], \mathcal{T}_2 *_C \mathcal{T}_2); f(0) \in j_0(\mathcal{T}_2) \text{ and } f(1) \in j_1(\mathcal{T}_2)\}$ is a unital continuous $C([0, 1])$ -algebra the fibres of which are all properly infinite. Thus, condition (c) implies that the C^* -algebra \mathcal{D} is properly infinite, a statement which is equivalent to the K_1 -injectivity of $\mathcal{T}_2 *_C \mathcal{T}_2$ ([Blan10, Proposition 4.2]). \square

4. A QUESTION OF PROPER INFINITENESS

We describe in this section the link between the different notions of proper infiniteness which have been introduced during the last decades.

Recall first the link between properly infinite C^* -algebras and properly infinite Hilbert C^* -modules studied by K. T. Coward, G. Elliott and C. Ivanescu.

Proposition 4.1. ([CEI08]) *Let A be a stable C^* -algebra and let $a \in A_+$ be a positive element. The following assertions are equivalent:*

- (a) *a is properly infinite in A , *i.e.* $a \oplus a \precsim a$ in $\mathcal{K} \otimes A$ ([KR00, definition 3.2]).*
- (b) *There is an embedding of Hilbert A -module $\ell^2(\mathbb{N}) \otimes A \hookrightarrow [aA]$ ([CEI08]).*

Proof. (a) \Rightarrow (b) If there exists an infinite sequence $\{d_i\}$ in A such that $a = d_i^* d_i \geq \sum_{j \in \mathbb{N}} d_j d_j^*$ for all $i \in \mathbb{N}$, then $[aA] \supset \sum_j [d_j A] \cong \ell^2(\mathbb{N}) \otimes A$.

(b) \Rightarrow (a) One has embeddings of Hilbert A -modules $[aA] \oplus [aA] \subset \ell_2(\mathbb{N}) \otimes A \subset [aA]$. \square

The following holds for continuous deformations of properly infinite C^* -algebras.

Proposition 4.2. *Let X be a second countable compact Hausdorff space, let H be a separable Hilbert $C(X)$ -module with non-zero fibres and let $a \in \mathcal{K}(H)$ be a strictly positive compact contraction. Consider the following assertions:*

- (a) *All the Hilbert spaces H_x are properly infinite, *i.e.* the operator a_x is properly infinite in the C^* -algebra $\mathcal{K}(H)_x \cong \mathcal{K}(H_x)$ for all $x \in X$.*
- (b) *H is a properly infinite Hilbert $C(X)$ -module, *i.e.* a is properly infinite in $\mathcal{K}(H)$.*

(c) The multiplier C^* -algebra $\mathcal{L}(H) = \mathcal{M}(\mathcal{K}(H))$ is properly infinite.

Then (c) \Rightarrow (b) \Rightarrow (a). But (a) $\not\Rightarrow$ (b) and (b) $\not\Rightarrow$ (c).

Proof. (c) \Rightarrow (b) If $\sigma : \mathcal{T}_2 = C^* \langle s_1, s_2 \rangle \rightarrow \mathcal{L}(H)$ is a unital $*$ -homomorphism, then the two elements $d_1 = \sigma(s_1) \cdot a^{1/2}$ and $d_2 = \sigma(s_2) \cdot a^{1/2}$ satisfy $d_i^* d_j = \delta_{i=j} \cdot a$ in $\mathcal{K}(H)$.

(b) \Rightarrow (a) The relations $c_i^* c_j = \delta_{i=j} \cdot a$ between 3 operators c_1, c_2, a in a $C(X)$ -algebra D entails that $(c_i)_x^* (c_j)_x = \delta_{i=j} \cdot a_x$ in the quotient $D_x = D/C_0(X \setminus \{x\}) \cdot D$ for all $x \in X$.

(a) $\not\Rightarrow$ (b) Let $B_3 = \{(x_1, x_2, x_3) \in \mathbb{R}^3; x_1^2 + x_2^2 + x_3^2 \leq 1\}$ be the unit ball of dimension 3, let B_3^+, B_3^- be the two open semi-disks $B_3^+ = \{(x_1, x_2, x_3) \in B_3; x_3 > -\frac{1}{2}\}$, $B_3^- = \{(x_1, x_2, x_3) \in B_3; x_3 < \frac{1}{2}\}$ and let $S_2 = \{(x_1, x_2, x_3) \in B_3; x_1^2 + x_2^2 + x_3^2 = 1\} \subset B_3$ be the unit sphere of dimension 2.

The self-adjoint operator $f \in C(B_3) \otimes M_2(\mathbb{C}) \cong C(B_3, M_2(\mathbb{C}))$ given by

$$f(x_1, x_2, x_3) = \frac{1}{2} \cdot \begin{pmatrix} 1 + x_3 & x_1 - ix_2 \\ x_1 + ix_2 & 1 - x_3 \end{pmatrix} \quad (4.1)$$

is a positive contraction since each self-adjoint matrix $f(x_1, x_2, x_3) \in M_2(\mathbb{C})$ satisfies

$$f(x_1, x_2, x_3)^2 = f(x_1, x_2, x_3) + \frac{1}{4}(x_1^2 + x_2^2 + x_3^2 - 1) \cdot 1_{M_2(\mathbb{C})}, \quad (4.2)$$

i.e. $(f(x_1, x_2, x_3) - \frac{1}{2} \cdot 1_{M_2(\mathbb{C})})^2 = \frac{x_1^2 + x_2^2 + x_3^2}{4} \cdot 1_{M_2(\mathbb{C})} \leq (\frac{1}{2} \cdot 1_{M_2(\mathbb{C})})^2$.

The nontrivial Hilbert $C(B_3)$ -module $F := [f \cdot \begin{pmatrix} C(B_3) \\ C(B_3) \end{pmatrix}]$ satisfies the two isomorphisms of Hilbert $C(B_3)$ -module:

$$\begin{aligned} - & F \cdot C_0(B_3^+) \cong C_0(B_3^+) \oplus C_0(B_3^+ \setminus S_2 \cap B_3^+) \\ - & F \cdot C_0(B_3^-) \cong C_0(B_3^-) \oplus C_0(B_3^- \setminus S_2 \cap B_3^-). \end{aligned} \quad (4.3)$$

The set $B_\infty := \{x \in \ell^2(\mathbb{N}); \sum_p |x_p|^2 \leq 1\}$ is a metric compact space called the *complex Hilbert cube* when equipped with the distance $d((x_p), (y_p)) = \sum_p 2^{-p-2} |x_p - y_p|$. Denote by E_{DD} the non-trivial Hilbert $C(B_\infty)$ -module with fibres $\ell^2(\mathbb{N})$ constructed by J. Dixmier and A. Douady ([DD63, §17], [BK04a, Proposition 3.6]).

Finally, consider the product $X := B_\infty \times B_3$ and the Hilbert $C(X)$ -module

$$H := E_{DD} \otimes C(B_3) \oplus C(B_\infty) \otimes F. \quad (4.4)$$

The two Hilbert $C(X)$ -submodules $H \cdot C_0(B_\infty \times B_3^+)$ and $H \cdot C_0(B_\infty \times B_3^-)$ are properly infinite, *i.e.* there exist embeddings of Hilbert $C(X)$ -module $\ell^2(\mathbb{N}) \otimes C_0(B_\infty \times B_3^+) \hookrightarrow H \cdot C_0(B_\infty \times B_3^+)$ and $\ell^2(\mathbb{N}) \otimes C_0(B_\infty \times B_3^-) \hookrightarrow H \cdot C_0(B_\infty \times B_3^-)$. Hence, all the fibres of the Hilbert $C(X)$ -module H are properly infinite Hilbert spaces, *i.e.* $\ell^2(\mathbb{N}) \hookrightarrow H_x$ for all point x in the compact space $X = B_\infty \times B_3^+ \cup B_\infty \times B_3^-$ ([CEI08]). But the Hilbert $C(X)$ -module H is not properly infinite *i.e.* $\ell^2(\mathbb{N}) \otimes C(X) \not\hookrightarrow H$ ([RR11, Example 9.11]). The equality $C(B_3) = C_0(B_3^+) + C_0(B_3^-)$ only implies that $\ell^2(\mathbb{N}) \otimes C(X) \hookrightarrow H \oplus H$.

(b) $\not\Rightarrow$ (c) There exists a continuous field \tilde{H} of Hilbert spaces over the compact space $Y := B_\infty \times (B_3)^\infty$ such that $\tilde{H} = [a \cdot \tilde{H}]$ for some properly infinite contraction $a \in \mathcal{K}(\tilde{H})$ and the C^* -algebra $\mathcal{L}(\tilde{H})$ is not properly infinite. Indeed, let $\eta \in \ell^\infty(B_\infty, \ell^2(\mathbb{N}) \oplus \mathbb{C})$

be the section $x \mapsto x \oplus \sqrt{1 - \|x\|^2}$, let \ddot{F} be the closed Hilbert $C(B_\infty)$ -module $\ddot{F} := [C(B_\infty, \ell^2(\mathbb{N}) \oplus 0) + C(B_\infty) \cdot \eta]$, let $\theta_{\eta, \eta} \in \mathcal{L}(\ddot{F})$ be the projection $\zeta \mapsto \eta \langle \eta, \zeta \rangle$ and let $E_{DD} = (1 - \theta_{\eta, \eta}) \cdot \ddot{F}$ be the Hilbert $C(B_\infty)$ -submodule built in [DD63]. Define also the sequence of contractions $\tilde{f} = (\tilde{f}_n)$ in $\ell^\infty\left(\mathbb{N}, M_2(C((B_3)^\infty))\right)$ by

$$x_n = (x_{n,k}) \in (B_3)^\infty \longmapsto \tilde{f}_n(x_n) := f(x_{n,n}) \in M_2(\mathbb{C}). \quad (4.5)$$

The Hilbert $C(Y)$ -module

$$\tilde{H} := C(Y) \oplus E_{DD} \otimes C((B_3)^\infty) \oplus C(B_\infty) \otimes [\tilde{f} \cdot \ell^2\left(\mathbb{N}, \left(\begin{smallmatrix} C((B_3)^\infty) \\ C((B_3)^\infty) \end{smallmatrix}\right)\right)] \quad (4.6)$$

has the desired properties ([RR11, Example 9.13]). \square

Question 4.3. – What can be written when the operator a in Proposition 4.2 is a projection in a unital continuous $C(X)$ -algebra?

– Is the full unital free product $\mathcal{T}_2 *_C \mathcal{T}_2$ a properly infinite C^* -algebra which is not K_1 -injective? (see the equivalence (a) \Leftrightarrow (c) in Corollary 3.3)

5. THE PIMSNER-TOEPLITZ ALGEBRA OF A HILBERT $C(X)$ -MODULE

We look in this section at the proper infiniteness of the unital continuous $C(X)$ -algebras with fibres \mathcal{O}_∞ corresponding to the Pimsner-Toeplitz $C(X)$ -algebras of Hilbert $C(X)$ -modules with infinite dimension fibres.

Definition 5.1. ([Pim95]) *Let X be a compact Hausdorff space and let E be a full Hilbert $C(X)$ -module, i.e. without any zero fibre.*

a) *The full Fock Hilbert $C(X)$ -module $\mathcal{F}(E)$ of E is the direct sum*

$$\mathcal{F}(E) := \bigoplus_{m \in \mathbb{N}} E^{(\otimes_{C(X)})^m}, \quad (5.1)$$

where $E^{(\otimes_{C(X)})^m} := \begin{cases} C(X) & \text{if } m = 0, \\ E \otimes_{C(X)} \dots \otimes_{C(X)} E \text{ (} m \text{ terms)} & \text{if } m \geq 1. \end{cases}$

b) *The Pimsner-Toeplitz $C(X)$ -algebra $\mathcal{T}(E)$ of E is the unital subalgebra of the $C(X)$ -algebra $\mathcal{L}(\mathcal{F}(E))$ of adjointable $C(X)$ -linear operators acting on $\mathcal{F}(E)$ generated by the creation operators $\ell(\zeta)$ ($\zeta \in E$), where*

$$\begin{aligned} - \ell(\zeta) (f \cdot \hat{1}_{C(X)}) &:= f \cdot \zeta = \zeta \cdot f & \text{for } f \in C(X) & \text{and} \\ - \ell(\zeta) (\zeta_1 \otimes \dots \otimes \zeta_k) &:= \zeta \otimes \zeta_1 \otimes \dots \otimes \zeta_k & \text{for } \zeta_1, \dots, \zeta_k \in E & \text{if } k \geq 1. \end{aligned} \quad (5.2)$$

c) *Let $(C^*(\mathbb{Z}), \Delta)$ be the abelian compact quantum group generated by a unitary \mathbf{u} with spectrum the unit circle and with coproduct $\Delta(\mathbf{u}) = \mathbf{u} \otimes \mathbf{u}$. Then, there is a unique coaction α of the Hopf C^* -algebra $(C^*(\mathbb{Z}), \Delta)$ on the Pimsner-Toeplitz $C(X)$ -algebra $\mathcal{T}(E)$ such that $\alpha(\ell(\zeta)) = \ell(\zeta) \otimes \mathbf{u}$ for all $\zeta \in E$, i.e.*

$$\begin{aligned} \alpha : \mathcal{T}(E) &\rightarrow \mathcal{T}(E) \otimes C^*(\mathbb{Z}) = C(\mathbb{T}, \mathcal{T}(E)) \\ \ell(\zeta) &\mapsto \ell(\zeta) \otimes \mathbf{u} = (z \mapsto \ell(z\zeta)) \end{aligned} \quad (5.3)$$

The fixed point $C(X)$ -subalgebra $\mathcal{T}(E)^\alpha = \{a \in \mathcal{T}(E); \alpha(a) = a \otimes 1\}$ under this coaction is the closed linear span

$$\mathcal{T}(E)^\alpha = \left[C(X) \cdot 1_{\mathcal{T}(E)} + \sum_{k \geq 1} \ell(E)^k \cdot (\ell(E)^k)^* \right]. \quad (5.4)$$

Besides, the projection $P \in \mathcal{L}(\mathcal{F}(E))$ onto the submodule E induces a quotient morphism of $C(X)$ -algebra $a \in \mathcal{T}(E)^\alpha \mapsto \bar{\mathbf{q}}(a) := P \cdot a \cdot P \in \mathcal{K}(E) + C(X) \cdot 1_{\mathcal{L}(E)} \subset \mathcal{L}(E)$.

Proposition 5.2. *Let X be a second countable compact Hausdorff perfect space and let E be a separable Hilbert $C(X)$ -module with infinite dimensional fibres.*

1) *There exist a covering $X = \overset{\circ}{F}_1 \cup \dots \cup \overset{\circ}{F}_m$ by the interiors of closed subsets F_1, \dots, F_m and m norm 1 sections ζ_1, \dots, ζ_m in E such that $\mathcal{T}(E) = C^* \langle \mathcal{T}(E)^\alpha, \ell(\zeta_1), \dots, \ell(\zeta_m) \rangle$, $(\ell(\zeta_k) \ell(\zeta_k)^*)_{|_{F_k}}$ and $(1 - \ell(\zeta_k) \ell(\zeta_k)^*)_{|_{F_k}}$ are properly infinite projections in $\mathcal{T}(E)_{|_{F_k}}$ for all index $k \in \{1, \dots, m\}$.*

2) *Set $G_k := F_1 \cup \dots \cup F_k$ for all integer $k \in \{1, \dots, m\}$ and $\bar{G}_l := G_l \cap F_{l+1}$ for all integer $l \in \{1, \dots, m-1\}$. If $\xi(l) \in E_{|_{\bar{G}_l}}$ is a section such that $\|\xi(l)_y\| = 1$ for all $y \in \bar{G}_l$, then there is a unitary $w_l \in \mathcal{T}(E)_{|_{\bar{G}_l}}$ such that*

- (a) $w_l \cdot \ell(\xi(l))_{|_{\bar{G}_l}} = \ell(\zeta_{l+1})_{|_{\bar{G}_l}}$,
- (b) $w_l \oplus 1_{|_{\bar{G}_l}}$ is homotopic to $1_{|_{\bar{G}_l}} \oplus 1_{|_{\bar{G}_l}}$ among the unitaries in $M_2(\mathcal{T}(E)_{|_{\bar{G}_l}})$.

3) *If for all K_1 -trivial unitary $w_k \in \mathcal{T}(E)_{|_{\bar{G}_k}}$ there is a unitary $z_{k+1} \in \mathcal{T}(E)^\alpha_{|_{F_{k+1}}}$ such that $(z_{k+1})_{|_{\bar{G}_k}} = w_k$ ($1 \leq k \leq m-1$), then there is a section $\xi \in E$ satisfying*

$$\forall x \in X, \quad \|\xi_x\| = 1, \quad (5.5)$$

so that Lemma 6.1 of [Blan13] implies that the C^* -algebra $\mathcal{T}(E)$ is properly infinite.

Proof. 1) Given a point $x \in X$ and a unit vector $\zeta \in E_x$, let ξ_1, ξ_2, ξ_3 be three norm 1 sections in E such that $(\xi_1)_x = \zeta$ and the matrix $a = [\langle \xi_i, \xi_j \rangle] \in M_3(C(X))$ satisfies $a_x = 1_3 \in M_3(\mathbb{C})$. Let $F \subset X$ be a closed neighbourhood of x such that $\|a_y - 1_3\| \leq 1/2$ for all $y \in F$. Define the sections ξ'_1, ξ'_2, ξ'_3 in $E_{|_F}$ by

$$\begin{pmatrix} (\xi'_1)_y \\ (\xi'_2)_y \\ (\xi'_3)_y \end{pmatrix} = \begin{pmatrix} (\xi_1)_y \\ (\xi_2)_y \\ (\xi_3)_y \end{pmatrix} \cdot (a_y^* a_y)^{-1/2} \quad \text{for all } y \in F. \quad (5.6)$$

One has $[\langle \xi'_i, \xi'_j \rangle] = (a_{|_F}^* a_{|_F})^{-1/2} \cdot (a_{|_F}^* a_{|_F}) \cdot (a_{|_F}^* a_{|_F})^{-1/2} = 1$ in $M_3(C(F))$. Hence, $\ell(\xi'_1) \ell(\xi'_1)^*$ and $q := 1_{|_F} - \ell(\xi'_1) \ell(\xi'_1)^*$ are properly infinite projections in $\mathcal{T}(E)_{|_F}$ since

$$\begin{aligned} - \quad 1_{|_F} - q &= \ell(\xi'_1) \ell(\xi'_1)^* \geq \ell(\xi'_1) \ell(\xi'_2) \ell(\xi'_2)^* \ell(\xi'_1)^* + \ell(\xi'_1) \ell(\xi'_3) \ell(\xi'_3)^* \ell(\xi'_1)^* \\ - \quad q &\geq \ell(\xi'_2) \ell(\xi'_2)^* + \ell(\xi'_3) \ell(\xi'_3)^* \geq \ell(\xi'_2) q^2 \ell(\xi'_2)^* + \ell(\xi'_3) q^2 \ell(\xi'_3)^*, \end{aligned} \quad (5.7)$$

so that there exist unital $*$ -homomorphisms from \mathcal{T}_2 to $(1-q) \cdot \mathcal{T}(E)_{|_F} \cdot (1-q)$ and $q \cdot \mathcal{T}(E)_{|_F} \cdot q$ given by $s_i \mapsto \ell(\xi'_1) \ell(\xi'_{1+i}) \ell(\xi'_1)^*$ and $s_i \mapsto \ell(\xi'_{1+i}) q$ for $i = 1, 2$.

The compactness of the space X enables to end the proof of this first assertion.

2) Let $v_l \in \mathcal{T}(E)_{|_{\bar{G}_l}}$ be the partial isometry $v_l := \ell(\zeta_{l+1})_{|_{\bar{G}_l}} \cdot \ell(\xi(l))_{|_{\bar{G}_l}}^*$. There exists by Lemma 2.4 of [BRR08] a K_1 -trivial unitary w_l in the properly infinite unital C^* -algebra $\mathcal{T}(E)_{|_{\bar{G}_l}}$ which has the two requested properties (a) and (b).

3) One constructs inductively the restrictions $\xi|_{G_k}$ in $E|_{G_k}$. Set $\xi|_{G_1} := \zeta_1$ and assume $\xi|_{G_k}$ already constructed. As $\ell(\xi|_{G_k})|_{\bar{G}_k} = z_{k+1}^* \cdot \ell(\zeta_{k+1})|_{\bar{G}_k}$, the only extension $\xi|_{G_{k+1}} \in E|_{G_{k+1}}$ such that $(\xi|_{G_{k+1}})|_{G_k} = \xi|_{G_k}$ and $(\xi|_{G_{k+1}})|_{F_{k+1}} = \bar{\mathbf{q}}(z_{k+1})^* \cdot (\zeta_{k+1})|_{F_{k+1}}$ satisfies $\|(\xi|_{G_{k+1}})_x\| = 1$ for all point $x \in G_{k+1}$. \square

Remarks 5.3. a) The nontrivial separable Hilbert $C(B_\infty)$ -module E_{DD} constructed by J. Dixmier and A. Douady ([DD63]) has infinite dimensional fibres and every section $\zeta \in E_{DD}$ satisfies $\zeta_x = 0$ for at least one point $x \in B_\infty$. Thus, it cannot satisfy the assumptions for the assertion 3) of Proposition 5.2. There are some $k \in \{1, \dots, m-1\}$ and a unitary $a_{k+1} \in \mathcal{U}_0(M_2(\mathcal{T}(E_{DD})|_{F_{k+1}}))$ such that

$$(a_{k+1})|_{\bar{G}_k} = w_k \oplus 1|_{\bar{G}_k} \quad (5.8)$$

and either $a_{k+1} \notin \mathcal{T}(E_{DD})|_{F_{k+1}} \oplus C(F_{k+1})$ or $\alpha(a_{k+1}) \neq a_{k+1} \otimes 1$.

b) If each of the K_1 -trivial unitaries w_l introduced in assertion 2) of Proposition 5.2 satisfies $\alpha(w_l) = w_l \otimes 1$ and $w_l \sim_h 1|_{\bar{G}_l}$ in $\mathcal{U}(\mathcal{T}(E)|_{\bar{G}_l})^\alpha$, then there exist by [LLR00, Lemma 2.1.7] $m-1$ unitaries $z_{l+1} \in \mathcal{T}(E)|_{F_{l+1}}^\alpha$ such that $(z_{l+1})|_{\bar{G}_l} = w_l$, so that there exists a section $\xi \in E$ with $\xi_x \neq 0$ for all $x \in X$.

c) Let A be a separable unital continuous $C(B_\infty)$ -algebra with fibres isomorphic to \mathcal{O}_2 such that $K_i(A) \cong C(Y_0, \Gamma_i)$ for $i = 0, 1$, where (Γ_0, Γ_1) is a pair of countable abelian torsion groups ([Dăd09, §3]). Let φ be a continuous field of faithful states on A . Then the $C(B_\infty)$ -algebra $A' \subset \mathcal{L}(L^2(A, \varphi))$ generated by $\pi_\varphi(A)$ and the algebra of compact operators $\mathcal{K}(L^2(A, \varphi))$ is a continuous $C(B_\infty)$ -algebra since both the ideal $\mathcal{K}(L^2(A, \varphi))$ and the quotient $A \cong A'/\mathcal{K}(L^2(A, \varphi))$ are continuous (see *e.g.* [Blan09, Lemma 4.2]). All the fibres of A' are isomorphic to the Cuntz extension \mathcal{T}_2 . But A' is not a trivial $C(B_\infty)$ -algebra since $K_0(A') = C(Y_0, \Gamma_0) \oplus \mathbb{Z}$ and $K_1(A') = C(Y_0, \Gamma_1)$.

Question 5.4. The Pimsner-Toeplitz algebra $\mathcal{T}(E_{DD})$ is locally purely infinite ([BK04b, Definition 1.3]) since all its simple quotients are isomorphic to the Cuntz algebra \mathcal{O}_∞ ([BK04b, Proposition 5.1]). But is $\mathcal{T}(E_{DD})$ properly infinite?

REFERENCES

- [Blac04] B. Blackadar, *Semiprojectivity in simple C^* -algebras*, Adv. Stud. Pure Math. **38** (2004), 1–17.
- [Blan97] E. Blanchard, *Subtriviality of continuous fields of nuclear C^* -algebras*, with an appendix by E. Kirchberg, J. Reine Angew. Math. **489** (1997), 133–149.
- [Blan09] E. Blanchard, *Amalgamated free products of C^* -bundles*, Proc. Edinburgh Math. Soc. **52** (2009), 23–36.
- [Blan10] E. Blanchard, *K_1 -injectivity for properly infinite C^* -algebras*, Clay Math. Proc. **11** (2010), 49–54.
- [Blan13] E. Blanchard, *Continuous fields with fibres \mathcal{O}_∞* , Math. Scand. **115** (2014), 189–205.
- [BK04a] E. Blanchard, E. Kirchberg, *Global Glimm halving for C^* -bundles*, J. Op. Th. **52** (2004), 385–420.
- [BK04b] E. Blanchard, E. Kirchberg, *Non-simple purely infinite C^* -algebras: the Hausdorff case*, J. Funct. Anal. **207** (2004), 461–513.
- [BRR08] E. Blanchard, R. Rohde, M. Rørdam, *Properly infinite $C(X)$ -algebras and K_1 -injectivity*, J. Noncommut. Geom. **2** (2008), 263–282.
- [CEI08] K. T. Coward, G. Elliott, C. Ivanescu, *The Cuntz semigroup as an invariant for C^* -algebras*, J. Reine Angew. Math. **623** (2008), 161–193.

- [Cun77] J. Cuntz, *Simple C^* -Algebras Generated by Isometries*, Commun. Math. Phys. **57** (1977), 173–185.
- [Cun81] J. Cuntz, *K -theory for certain C^* -algebras*, Ann. of Math. **113** (1981), 181–197.
- [DD63] J. Dixmier, A. Douady, *Champs continus d'espaces hilbertiens et de C^* -algèbres*, Bull. Soc. Math. France **91** (1963), 227–284.
- [Dix69] J. Dixmier, *Les C^* -algèbres et leurs représentations*, Gauthiers-Villars Paris (1969).
- [Dăd09] M. Dădărlat, *Fiberwise KK -equivalence of continuous fields of C^* -algebras*, J. K-Theory **3** (2009), 205–219.
- [DW08] M. Dădărlat, W. Winter, *Trivialization of $C(X)$ -algebras with strongly self-absorbing fibres*, Bull. Soc. Math. France **136** (2008), 575–606.
- [Ell94] G. Elliott, *The classification problem for amenable C^* -algebras*, Proc. Internat. Congress of Mathematicians (Zurich, Switzerland, 1994), Birkhauser Verlag, Basel (1995), 922–932.
- [HRW07] I. Hirshberg, M. Rørdam, W. Winter, *$C_0(X)$ -algebras, stability and strongly self-absorbing C^* -algebras*, Math. Ann. **339** (2007), 695–732.
- [HR98] J. Hjeltnborg, M. Rørdam, *On stability of C^* -algebras*. J. Funct. Anal. **155** (1998), 153–170.
- [Kas88] G.G. Kasparov, *Equivariant KK -theory and the Novikov conjecture*, Invent. Math. **91** (1988), 147–201.
- [KR00] E. Kirchberg, M. Rørdam, *Non-simple purely infinite C^* -algebras*, Amer. J. Math. **122** (2000), 637–666.
- [LLR00] F. Larsen, N. J. Laustsen, M. Rørdam, *An Introduction to K -theory for C^* -algebras*, London Mathematical Society Student Texts **49** (2000) CUP, Cambridge.
- [Pim95] M. V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbb{Z}* , Free probability theory, Fields Inst. Commun. **12** (1997), 189–212.
- [RR11] L. Robert, M. Rørdam, *Divisibility properties for C^* -algebras*, Proc. London Math. Soc. **106** (2013), 1330–1370.
- [Roh09] R. Rohde, *K_1 -injectivity of C^* -algebras*, Ph.D. thesis at Odense (2009)
- [Rør97] M. Rørdam, *Stability of C^* -algebras is not a stable property.*, Doc. Math. J. **2** (1997), 375–386.
- [Rør03] M. Rørdam, *A simple C^* -algebra with a finite and an infinite projection*, Acta Math. **191** (2003), 1091–1142.
- [Rør04] M. Rørdam, *The stable and the real ranks of \mathcal{Z} -absorbing C^* -algebras*, Internat. J. Math. **15** (2004), 1065–1084.
- [TW07] A. Toms, W. Winter, *Strongly self-absorbing C^* -algebras*, Trans. Amer. Math. Soc. **359** (2007), 3999–4029.
- [Win09] W. Winter, *Strongly self-absorbing C^* -algebras are \mathcal{Z} -stable*, J. Noncommut. Geom. **5** (2011), 253–264.

Etienne.Blanchard@imj-prg.fr

IMJ-PRG, UP7D - Campus des Grands Moulins, Case 7012
F-75205 Paris Cedex 13