


# From Sparse Regression to Sparse Multiple Correspondence Analysis

Gilbert Saporta

CEDRIC, CNAM, Paris

[gilbert.saporta@cnam.fr](mailto:gilbert.saporta@cnam.fr)

- Joint work with Anne Bernard, Ph.D student funded by the R&D department of Chanel cosmetic company
- Industrial context and motivation:
  - Relate gene expression data to skin aging measures



–  $n=500$ ,  $p= 800\,000$  SNP's, 15 000 genes

# Outline

1. Introduction
2. Regularized regression
3. Sparse regression
4. Sparse PCA
5. Sparse MCA
6. Conclusion and perspectives

# 1. Introduction

- High dimensional data:  $p \gg n$ 
  - Gene expression data
  - Chemometrics
  - etc.
- Several solutions for regression problems with **all** variables; but interpretation is difficult
- Sparse methods: provide combinations of **few** variables

- This talk:
  - a survey of sparse methods for supervised (regression) and unsupervised (PCA) problems
  - New propositions in the unsupervised case when variables belong to disjoint groups or blocks:
    - Group sparse PCA
    - Sparse multiple correspondence analysis

## 2. Regularized regression

- No OLS solution when  $p > n$
- A special case of multicollinearity
- Usual regularized regression techniques:
  - Component based: PCR, PLS
  - Ridge



## 2.1 Principal components regression

- First papers: Kendall, Hotelling (1957), Malinvaud (1964)
- At most  $n$  components when  $p \gg n$
- Select  $q$  components and regress  $y$  upon them
  - Orthogonal components  sum of univariate regressions
  - Back to original variables:

**C** components matrix n,q      **U** loadings matrix p,q

$$\mathbf{C} = \mathbf{X}\mathbf{U}$$

$$\hat{\mathbf{y}} = \mathbf{C}\hat{\boldsymbol{\alpha}} = \alpha_1 \mathbf{c}_1 + \dots + \alpha_q \mathbf{c}_q = \mathbf{X}\mathbf{U}\hat{\boldsymbol{\alpha}} = \mathbf{X}\hat{\boldsymbol{\beta}} \quad \hat{\boldsymbol{\beta}} = \mathbf{U}\hat{\boldsymbol{\alpha}}$$

- Principal components unrelated to the response variable y:
  - Ranking the components
    - Not according to their eigenvalues
    - but according to  $r^2(y; c_j)$
- Choice of q
  - crossvalidation

## 2.2 PLS regression

- Proposed by H. and S.Wold (1960's)
- Close to PCR: projection onto a set of orthogonal combinations of predictors
- PLS components optimised to be predictive of both X and y variables
- Tucker's criterium:  $\max \text{cov}^2(y ; Xw)$

- Trade-off between maximizing correlation between  $t=Xw$  and  $y$  (OLS) and maximizing variance of  $t$  (PCA) :

$$\text{cov}^2(y ; Xw) = r^2(y ; Xw) V(Xw) V(y)$$

- Easy solution:
  - $w_j$  proportional to  $\text{cov}(y; x_j)$
  - No surprising signs..
- Further components by iteration on residuals
- Stopping rule: cross-validation

## 2.3 Ridge regression

- Hoerl & Kennard (1970)

$$\hat{\beta}_R = (\mathbf{X}'\mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}'\mathbf{y}$$

- Several interpretations
  - Tikhonov regularization

$$\min \|\mathbf{y} - \mathbf{X}\beta\|^2 \quad \text{with } \|\beta\|^2 \leq c^2$$

$$\text{or } \min \left( \|\mathbf{y} - \mathbf{X}\beta\|^2 + \lambda \|\beta\|^2 \right)$$

- Bayesian regression

- Gaussian prior for  $\beta$   $N(\mathbf{0}; \psi^2 \mathbf{I})$
- Gaussian distribution  $\mathbf{Y}/\beta$   $N(\mathbf{X}\beta; \sigma^2 \mathbf{I})$

Maximum a posteriori or posterior expectation :


$$\hat{\beta} = \left( \mathbf{X}'\mathbf{X} + \frac{\sigma^2}{\psi^2} \mathbf{I} \right)^{-1} \mathbf{X}'\mathbf{y}$$

Gives an interpretation for k

- Choice of k :
  - cross-validation

- Shrinkage properties (Hastie et al. , 2009)
  - PCR discards low variance directions
  - PLS shrinks low variance directions but inflates high variance directions
  - Ridge shrinks all principal directions but shrinks more low variance directions
- Lost properties:
  - Bias, scale invariance  
need standardised data

# 3. Sparse regression


- Keeping all predictors is a drawback for high dimensional data: combinations of too many variables cannot be interpreted
- Sparse methods simultaneously shrink coefficients and select variables, hence better predictions

## 3.1 Lasso and elastic-net

- Lasso (Tibshirani, 1996) imposes a  $L_1$  constraint on the coefficients  $\sum_{j=1}^p |b_j| < c$   
$$\hat{\beta}_{lasso} = \arg \min_{\beta} \left( \|\mathbf{y} - \mathbf{X}\beta\|^2 + \lambda \sum_{j=1}^p |\beta_j| \right)$$
- Lasso continuously shrinks the coefficients towards zero
- Convex optimisation; no explicit solution

- Constraints and log-priors
  - Like ridge regression, the Lasso is a bayesian regression but with an exponential prior

$$f(\beta_j) = \frac{\lambda}{2} \exp(-\lambda |\beta_j|)$$



- $|\beta_j|$  is proportional to the log-prior

- Finding the optimal parameter
  - Cross validation if optimal prediction is needed
  - BIC when the sparsity is the main concern


$$\lambda_{opt} = \arg \min_{\lambda} \left( \frac{\|y - X \hat{\beta}(\lambda)\|^2}{n\sigma^2} + \frac{\log(n)}{n} \hat{df}(\lambda) \right)$$

a good unbiased estimate of df is the number of nonzero coefficients . (Zou et al., 2007)

- A more general form:

$$\hat{\boldsymbol{\beta}}_{lasso} = \arg \min_{\boldsymbol{\beta}} \left( \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|^2 + \lambda \sum_{j=1}^p |\beta_j|^q \right)$$

- $q=2$  ridge;  $q=1$  Lasso;  $q=0$  subset selection (counts the number of variables)
- $q>1$  do not provide null coefficients (derivability)




**FIGURE 3.12.** Contours of constant value of  $\sum_j |\beta_j|^q$  for given values of  $q$ .

- Lasso produces a sparse model but the number of selected variables cannot exceed the number of units
- **Elastic net:** combine ridge penalty and lasso penalty to select more predictors than the number of observations (Zou & Hastie, 2005)

$$\hat{\beta}_{en} = \arg \min_{\beta} \left( \|y - X\beta\|^2 + \lambda_2 \|\beta\|^2 + \lambda_1 \|\beta\|_1 \right)$$

## 3.2 Group-lasso

- X matrix divided into J sub-matrices  $X_j$  of  $p_j$  variables
- **Group Lasso:** extension of Lasso for selecting groups of variables  
(Yuan & Lin, 2007):



$$\hat{\boldsymbol{\beta}}_{GL} = \arg \min_{\boldsymbol{\beta}} \left\| \mathbf{y} - \sum_{j=1}^J \mathbf{X}_j \boldsymbol{\beta}_j \right\|^2 + \lambda \sum_{j=1}^J \sqrt{p_j} \|\boldsymbol{\beta}_j\|$$

If  $p_j=1$  for all  $j$ , group Lasso = Lasso


- Drawback: no sparsity within groups
- A solution: **sparse group lasso** (Simon et al. , 2012)

$$\min_{\beta} \left( \left\| \mathbf{y} - \sum_{j=1}^J \mathbf{X}_j \beta_j \right\|^2 + \lambda_1 \sum_{j=1}^J \|\beta_j\| + \lambda_2 \sum_{j=1}^J \sum_{i=1}^{p_j} |\beta_{ij}| \right)$$

- Two tuning parameters: grid search

### 3.3 Other sparse regression methods

- SCAD penalty (Fan & Li, 2001)
  - « smoothly clipped absolute deviation »
  - Non-convex
- Sparse PLS
  - Several extensions
    - Chun & Keles (2010)
    - Le Cao et al. (2008)



# 4. Sparse PCA

- In PCA, each PC is a linear combination of **all** the original variables : difficult to interpret the results
- **Challenge of SPCA:** obtain components easily interpretable (lot of zero loadings in principal factors)
- **Principle of SPCA:** modify PCA imposing lasso/elastic-net constraints to construct modified PCs with sparse loadings
- **Warning:** Sparse PCA does not provide a global selection of variables but a selection **dimension by dimension** : different from the regression context (Lasso, Elastic Net, ...)

## 4.1 First attempts:

- **Simple PCA**
  - by Vines (2000) : integer loadings
  - Rousson, V. and Gasser, T. (2004) : loadings (+ , 0, -)
- **SCoTLASS** (Simplified Component Technique – Lasso) by Jolliffe & al. (2003) : extra  $L_1$  constraints

$$\max \mathbf{u}' \mathbf{V} \mathbf{u} \text{ with } \|\mathbf{u}\|^2 = \mathbf{u}' \mathbf{u} = 1 \text{ and } \sum_{j=1}^p |u_j| \leq t$$

## SCotLass properties:

$t \geq \sqrt{p}$  usual PCA

$t < 1$  no solution

$t = 1$  only one nonzero coefficient

$1 < t < \sqrt{p}$

- Non convex problem

## 4.2 S-PCA by Zou et al (2006)

Let the SVD of  $\mathbf{X}$  be  $\mathbf{X} = \mathbf{UDV}'$  with  $\mathbf{Z} = \mathbf{UD}$  the principal components

Ridge regression:

$$\hat{\boldsymbol{\beta}}_{ridge} = \arg \min_{\boldsymbol{\beta}} \|\mathbf{Z} - \mathbf{X}\boldsymbol{\beta}\|^2 + \lambda \|\boldsymbol{\beta}\|^2$$

$$\mathbf{X}'\mathbf{X} = \mathbf{V}\mathbf{D}^2\mathbf{V}' \text{ with } \mathbf{V}'\mathbf{V} = \mathbf{I}$$

$$\hat{\boldsymbol{\beta}}_{i,ridge} = (\mathbf{X}'\mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}' (\mathbf{X}\mathbf{v}_i) = \mathbf{v}_i \frac{d_{ii}^2}{d_{ii}^2 + \lambda} \quad \longrightarrow \quad \tilde{\mathbf{v}} = \mathbf{v}_i$$

**Loadings can be recovered by regressing (ridge regression) PCs on the p variables**

→ PCA can be written as a **regression-type optimization problem**

Sparse PCA add a new penalty to produce sparse loadings:

$$\hat{\beta} = \arg \min_{\beta} \|Z - X\beta\|^2 + \lambda \|\beta\|^2 + \lambda_1 \|\beta\|_1$$

Lasso  
penalty

$\hat{v}_i = \frac{\hat{\beta}}{\|\hat{\beta}\|}$  is an approximation to  $v_i$ , and  $X\hat{v}_i$  the  $i^{\text{th}}$  approximated component

→ Produces sparse loadings with zero coefficients to facilitate interpretation

Alternated algorithm between elastic net and SVD

## 4.3 S-PCA via regularized SVD

- Shen & Huang (2008) : starts from the SVD with a smooth penalty (L1, SCAD, etc.)

$$\mathbf{X}^{(k)} = \sum_{j=1}^k d_j \mathbf{u}_j \mathbf{v}_j'$$


$$\min_{\mathbf{u}, \mathbf{v}} \|\mathbf{X} - \mathbf{u}\mathbf{v}'\|^2 + \sum_{j=1}^p h_\lambda(|v_j|)$$

- Loss of orthogonality
  - SCotLass: orthogonal loadings but correlated components
  - S-PCA: neither loadings, nor components are orthogonal
  - Necessity of adjusting the % of explained variance

## 4.4 Group Sparse PCA

Data matrix X divided into J groups  $X_j$  of  $p_j$  variables

**Group Sparse PCA:** compromise between SPCA and group Lasso



**Goal:** select groups of continuous variables (zero coefficients to entire blocks of variables)

**Principle:** replace the penalty function in the SPCA algorithm

$$\hat{\beta} = \arg \min_{\beta} \|\mathbf{Z} - \mathbf{X}\beta\|^2 + \lambda \|\beta\|^2 + \lambda_1 \|\beta\|_1$$

by that defined in the group Lasso

$$\hat{\beta}_{GL} = \arg \min_{\beta} \left\| \mathbf{Z} - \sum_{j=1}^J \mathbf{X}_j \beta_j \right\|^2 + \lambda \sum_{j=1}^J \sqrt{p_j} \|\beta_j\|$$

ECDA 2013, Luxemburg

## Group Sparse PCA Algorithm

- ① Apply the SVD to  $\mathbf{X}$  and obtain the best rank-one approximation of  $\mathbf{X}$  as  $\delta \tilde{\mathbf{u}} \tilde{\mathbf{v}}^T$ . Set  $\tilde{\mathbf{v}}_{old} = \tilde{\mathbf{v}} = [\tilde{\mathbf{v}}_1, \dots, \tilde{\mathbf{v}}_K]$  and  $\tilde{\mathbf{u}}_{old} = \delta \tilde{\mathbf{u}}$ .
- ② Update:
  - a)  $\tilde{\mathbf{v}}_{new} = [h_\lambda(\mathbf{X}_{[1]}^T \tilde{\mathbf{u}}_{old}), \dots, h_\lambda(\mathbf{X}_{[K]}^T \tilde{\mathbf{u}}_{old})]$
  - b)  $\tilde{\mathbf{u}}_{new} = \mathbf{X} \tilde{\mathbf{v}}_{new} / \|\mathbf{X} \tilde{\mathbf{v}}_{new}\|$
- ③ Repeat Step 2 replacing  $\tilde{\mathbf{u}}_{old}$  and  $\tilde{\mathbf{v}}_{old}$  by  $\tilde{\mathbf{u}}_{new}$  and  $\tilde{\mathbf{v}}_{new}$  until convergence.

Sparse loadings  $\mathbf{v}_i$  ( $i > 1$ ) are obtained via rank-one approximation of residual matrices.

→ **Choice of  $\lambda$**  using cross-validation or an ad-hoc approach.

# 5.Sparse MCA

Original table

X <sub>j</sub>
1
p <sub>j</sub>
:
:
3

In MCA:

Selection of **1 column** in the original table  
(categorical variable X<sub>j</sub>)

=

Selection of **a block of p<sub>j</sub> indicator variables**  
in the complete disjunctive table

Complete disjunctive table

X <sub>j1</sub>	...	X <sub>j<sub>pj</sub></sub>
1		0
0		1
:		:
:		:
0		0

**Sparse MCA :** select categorical variables, not categories

**Principle:** a straightforward extension of Group Sparse PCA for groups of indicator variables, with the chi-square metric . Uses s-PCA r-SVD algorithm.

Let  $F$  be the  $n \times q$  disjunctive table divided by the number of units

$$\mathbf{r} = F\mathbf{1}_q \quad \mathbf{c} = F^T\mathbf{1}_n \quad \mathbf{D}_r = \text{diag}(\mathbf{r}) \quad \mathbf{D}_c = \text{diag}(\mathbf{c})$$

And  $\tilde{F}$  be the matrix of standardised residuals:

$$\tilde{F} = \mathbf{D}_r^{-\frac{1}{2}} (F - \mathbf{r}\mathbf{c}^T) \mathbf{D}_c^{-\frac{1}{2}}$$


Singular Value Decomposition  $\tilde{F} = \mathbf{U}\Lambda\mathbf{V}^T$

Properties	MCA	Sparse MCA
Uncorrelated Components	TRUE	FALSE
Orthogonal loadings	TRUE	FALSE
Barycentric property	TRUE	PARTLY TRUE
% of inertia	$\frac{\lambda_j}{tot} \times 100$	$\ \tilde{\mathbf{Z}}_{j,1,\dots,j-1}\ ^2$
Total inertia	$\frac{1}{p} \sum_{j=1}^p p_j - 1$	$\sum_{j=1}^k \ \tilde{\mathbf{Z}}_{j,1,\dots,j-1}\ ^2$

$\tilde{\mathbf{Z}}_{j,1,\dots,j-1}$  are the residuals after adjusting  $\tilde{\mathbf{Z}}_j$  for  $\tilde{\mathbf{Z}}_{1,\dots,j-1}$  (regression projection)

## Toy example: Dogs

$X_1$ Size	...	$X_6$ Aggressiveness
large (L)		aggressive (A)
medium (M)		aggressive (A)
:		:
small (S)	...	nonaggressive (N)



$K_1$ Size			...	$K_6$ Aggressiveness	
S.	M.	L.		A	N
0	0	1		1	0
0	1	0		1	0
:	:	:	...	:	:
1	0	0		0	1

### Data:

$n=27$  breeds of dogs

$p=6$  variables


$q=16$  (total number of columns)

$X : 27 \times 6$  matrix of categorical variables

$K : 27 \times 16$  complete disjunctive table  $\rightarrow K=(K_1, \dots, K_6)$

**1 block  
= 1  $K_j$  matrix**

## Toy example: Dogs



$\lambda = 0:25$  is a compromise between the number of variables selected and the % of variance lost.


## Toy example: Comparison of the loadings

Variable	MCA				Sparse MCA			
	Comp1	Comp2	Comp3	Comp4	Comp1	Comp2	Comp3	Comp4
large	-0.361	0.071	-0.005	0.060	-0.389	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
medium	0.280	0.287	0.300	-0.055	0.226	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
small	0.291	-0.400	-0.293	-0.041	0.390	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
lightweight	0.316	-0.389	-0.193	-0.081	0.368	-0.256	<b>0.000</b>	<b>0.000</b>
heavy	-0.047	0.390	-0.133	0.088	-0.075	0.451	<b>0.000</b>	<b>0.000</b>
very heavy	-0.294	-0.215	0.458	-0.055	-0.305	-0.479	<b>0.000</b>	<b>0.000</b>
slow	0.059	-0.383	0.296	0.133	<b>0.000</b>	-0.561	<b>0.000</b>	<b>0.000</b>
fast	0.224	0.256	0.057	-0.299	<b>0.000</b>	0.282	<b>0.000</b>	<b>0.000</b>
veryfast	-0.303	0.156	-0.391	0.168	<b>0.000</b>	0.328	<b>0.000</b>	<b>0.000</b>
unintelligent	0.173	0.157	0.356	0.236	<b>0.000</b>	<b>0.000</b>	0.693	-0.693
avg intelligent	-0.145	-0.309	-0.168	0.125	<b>0.000</b>	<b>0.000</b>	-0.327	0.327
very intelligent	-0.086	0.125	-0.330	-0.491	<b>0.000</b>	<b>0.000</b>	-0.642	0.642
unloving	-0.366	-0.084	0.030	0.087	-0.462	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
veryaffectionate	0.353	0.081	-0.029	-0.084	0.445	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
agressive	-0.170	-0.096	0.162	-0.515	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
non agressive	0.164	0.093	-0.156	0.497	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
Nb non-zero loadings	16	16	16	16	8	6	3	3
Adjusted variance (%)	28.19	22.80	13.45	9.55	23.03	17.40	10.20	9.50

# Application on genetic data

## Single Nucleotide Polymorphisms

SNP1=X <sub>1</sub>	...	SNP537=X <sub>537</sub>
AA		AB
AB		BB
:		:
:		:
AA		AA
BB		AA



SNP1=D <sub>[1]</sub>			...	SNP537=D <sub>[537]</sub>		
AA	AB	BB		AA	AB	BB
1	0	0		0	1	0
0	1	0		0	0	1
:	:	:		:	:	:
:	:	:		:	:	:
1	0	0		1	0	0
0	0	1		1	0	0

### Data:

n=502 individuals

p=537 SNPs (among more than 800 000 of the original data base, 15000 genes)

q=1554 (total number of columns)

X : 502 x 537matrix of qualitative variables

K : 502 x 1554 complete disjunctive table → K=(K<sub>1</sub>, ..., K<sub>1554</sub>)

**1 block**


=

**1 SNP = 1 K<sub>j</sub> matrix**


# Application on genetic data

## Single Nucleotide Polymorphisms

Cumulative % of variance  
depending on lambda





Nb of non-zero loadings  
depending on lambda



# Application on genetic data

## Single Nucleotide Polymorphisms



$\lambda = 0.005$ : CPEV = 0.32% and 174 columns selected on Comp 1

# Application on genetic data

## Comparison of the loadings

SNPs	MCA		SMCA	
	Comp1	Comp2	Comp1	Comp2
SNP1.AA	-0.078	0.040	-0.092	0.102
SNP1.AG	-0.014	-0.027	-0.022	-0.053
SNP1.GG	0.150	-0.002	0.132	-0.003
SNP2.AA	-0.082	0.041	-0.118	<b>0.000</b>
SNP2.AG	-0.021	-0.025	-0.020	<b>0.000</b>
SNP2.GG	-0.081	0.040	-0.001	<b>0.000</b>
SNP3.CC	-0.004	0.050	<b>0.000</b>	<b>0.000</b>
SNP3.CG	0.016	0.021	<b>0.000</b>	<b>0.000</b>
SNP3.GG	-0.037	-0.325	<b>0.000</b>	<b>0.000</b>
SNP4.AA	0.149	-0.003	0.050	<b>0.000</b>
SNP4.AG	-0.016	-0.025	-0.002	<b>0.000</b>
SNP4.GG	-0.081	0.040	-0.100	<b>0.000</b>
...	...	...	...	...
Nb non-zero loadings	1554	1554	172	108
Variance (%)	1.14	0.63	0.32	0.16
Cumulative variance (%)	1.14	1.77	0.32	0.48

# 6. Conclusions and perspectives

- Sparse techniques provide elegant and efficient solutions to problems posed by high-dimensional data:
  - A new generation of data analysis methods with few restrictive hypothesis
- Very powerful in a context of variable selection in high dimension issues:
  - reduce noise as well as computation time.

- 2 new methods in a unsupervised multiblock data context: **Group Sparse PCA** for continuous variables, and **Sparse MCA** for categorical variables
  - Both methods produce sparse loadings structures that makes easier the interpretation and the comprehension of the results
  - Possibility of selecting superblocks (genes)
- **Research in progress:**
  - Extension of Sparse MCA to select groups and predictors within a group (sparsity within groups)
    - sparsity at both group and individual feature levels
    - compromise between Sparse MCA and sparse group lasso developed by Simon et al. (2012).

# **Thanks for your attention**

# References

- Chun, H. and Keles, S. (2010), Sparse partial least squares for simultaneous dimension reduction and variable selection, *Journal of the Royal Statistical Society - Series B*, Vol. 72, pp. 3–25.
- Fan, J. and Li, R. Variable Selection via Nonconcave Penalized Likelihood and its Oracle Properties. *Journal of American Statistical Association*, **96**, 1348-1360, 2001
- Hastie T., Tibshirani R., Friedman J. (2009) *The elements of statistical learning*, 2nd edition, Springer, 2009
- Jolliffe, I.T. , Trendafilov, N.T. and Uddin, M. (2003) A modified principal component technique based on the LASSO. *Journal of Computational and Graphical Statistics*, **12**, 531–547,
- Rousson, V. , Gasser, T. (2004), Simple component analysis. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **53**, 539-555
- Shen, H. and Huang, J. Z. (2008). Sparse principal component analysis via regularized low rank matrix approximation. *Journal of Multivariate Analysis*, 99:1015-1034.

- Simon, N., Friedman, J., Hastie, T., and Tibshirani, R. (2012) A Sparse-Group Lasso. *Journal of Computational and Graphical Statistics*,
- Tenenhaus M. (1998) *La régression PLS*, Technip
- Tibshirani, R. (1996) Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society, Series B*, **58**, 267-288
- Vines, S.K., (2000) Simple principal components, *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **49**, 441-451
- Yuan, M., Lin, Y. (2007) Model selection and estimation in regression with grouped variables. *Journal of the Royal Statistical Society, Series B*, **68**, 49-67,
- Zou, H., Hastie , T. (2005) Regularization and variable selection via the elastic net. *Journal of Computational and Graphical Statistics*, **67**, 301-320,
- Zou, H., Hastie, T. and Tibshirani, R. (2006) Sparse Principal Component Analysis. *Journal of Computational and Graphical Statistics*, **15**, 265-286.
- H. Zou, T. Hastie, R. Tibshirani, (2007), On the “degrees of freedom” of the lasso, *The Annals of Statistics*, 35, 5, 2173–2192.