

## Correction: GPR50 is the mammalian ortholog of Mel1c: Evidence of rapid evolution in mammals

Laurence Dufourny, A. Levasseur, Martine Migaud, I. Callebaut, P.

Pontarotti, Benoit Malpaux, Philippe Monget

## ▶ To cite this version:

Laurence Dufourny, A. Levasseur, Martine Migaud, I. Callebaut, P. Pontarotti, et al.. Correction: GPR50 is the mammalian ortholog of Mel1c: Evidence of rapid evolution in mammals. BMC Evolutionary Biology, 2012, 12 (Article 28), pp.1-3. 10.1186/1471-2148-8-105 . hal-01129620

## HAL Id: hal-01129620 https://hal.science/hal-01129620

Submitted on 28 Mar 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

### CORRECTION




**Open Access** 

# Correction: GPR50 is the mammalian ortholog of Mel1c: Evidence of rapid evolution in mammals

Laurence Dufourny<sup>1+</sup>, Anthony Levasseur<sup>2+</sup>, Martine Migaud<sup>1</sup>, Isabelle Callebaut<sup>3</sup>, Pierre Pontarotti<sup>2</sup>, Benoit Malpaux<sup>1\*</sup> and Philippe Monget<sup>1\*</sup>

#### Correction

The core of the text as well as the legends remain the same as in our article [1] since the changes provided here do not change any of the outcomes of the former study including the results and our related conclusions. However, new figure files are provided since some arrows and stars have been previously misplaced in figures three A (figure 1) and four (figure 2), and the for-



**Figure 1** Phylogenetic analysis of the GPR50/M11/M12/Me11c genes. (A) Overall phylogenetic tree showing 3 groups of genes: GPR50, M11/ MT2, and Mel1c genes and the animal orders where each branch is expressed. The trees (npl) are the fusion of three phylogenetic trees built based on Neighbour joining, maximum Parsimony and maximum Likelihood (see "Materials and Methods" section for further details). The italic letters correspond to the name given to the branches for the likelihood ratio tests (B) Phylogenetic tree of GPR50 genes. Please note that only mammalian species appear in the tree. (C) Phylogenetic tree of MT1 genes (D) Phylogenetic tree of MT2 genes. (E) Phylogenetic tree of Mel1c genes that do seem to appear only in non mammalian species. Bootstrap values are reported for each npl method.

\* Correspondence: malpaux@tours.inra.fr; monget@tours.inra.fr

<sup>1</sup>Physiologie de la Reproduction et des Comportements, UMR 6175 INRA-

CNRS-Université François Rabelais de Tours-Haras Nationaux, 37380 Nouzilly, France


Full list of author information is available at the end of the article




© 2012 Dufourny et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

<sup>+</sup> Contributed equally

GPR50 locus (unpublished data). p.d.: predicted gene. Chrm: chromosome.





**Figure 3 Sequence alignment of human MT1, MT2 and GPR50 with bovine rhodopsin** (pdb 1F88). Sequence identities are reported white on a black background, whereas sequence similarities are boxed (A). The positions of the transmembrane helices, as observed in the bovine rhodopsin structure, are reported above its sequence. Arrows indicate the positions of the amino acids that, in GPR50, evolved under positive selection. Stars indicate amino acids which have been shown to play a key role for melatonin binding in MT1 (dark blue), MT2 (light blue) or both (red). A ribbon representation of the GPR50 3D structure model is represented (B), with transmembrane helices colored according to the sequence alignment. Amino acids evolving under positive selection and amino acids important for melatonin binding in MT1/MT2 are shown according to the colors reported in the sequence alignment.

mer figure six (figure 3) showed the GPR50 mouse sequence instead of the human sequence as stated in the legend [1].

#### Author details

<sup>1</sup>Physiologie de la Reproduction et des Comportements, UMR 6175 INRA-CNRS-Université François Rabelais de Tours-Haras Nationaux, 37380 Nouzilly, France. <sup>2</sup>EA3781: Evolution Biologique - Laboratoire de PhyloGénomique -Université de Provence - Marseille - France. <sup>3</sup>Biologie Structurale, IMPMC, UMR7590, CNRS, Université Paris 6 et 7, 75015 Paris, France.

#### Received: 12 January 2012 Accepted: 7 March 2012 Published: 7 March 2012

#### Reference

 Dufourny L, Levasseur A, Migaud M, Callebaut I, Pontarotti P, Malpaux B, Monget P: GPR50 is the mammalian ortholog of Mel1c: evidence of rapid evolution in mammals. *BMC Evol Biol* 2008, 8:105.

#### doi:10.1186/1471-2148-12-28

**Cite this article as:** Dufourny *et al.*: **Correction: GPR50** is the mammalian ortholog of Mel1c: Evidence of rapid evolution in mammals. *BMC Evolutionary Biology* 2012 **12**:28.

# Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit