



HARMONIC FUNCTIONS ON MANIFOLDS WHOSE LARGE SPHERE ARE SMALL

Gilles Carron

► To cite this version:

Gilles Carron. HARMONIC FUNCTIONS ON MANIFOLDS WHOSE LARGE SPHERE ARE SMALL. Annales Mathématiques Blaise Pascal, 2016, 23 (2), pp. 249-261. 10.5802/ambp.362 . hal-01132922

HAL Id: hal-01132922

<https://hal.science/hal-01132922>

Submitted on 18 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HARMONIC FUNCTIONS ON MANIFOLDS WHOSE LARGE SPHERE ARE SMALL .

GILLES CARRON

ABSTRACT. We study the growth of harmonic functions on complete Riemannian manifolds where the extrinsic diameter of geodesic spheres is sublinear. It is an generalization of a result of A. Kazue. We also get a Cheng and Yau estimates for the gradient of harmonic functions.

RÉSUMÉ : On étudie la croissance des fonctions harmoniques sur les variétés riemanniennes complètes dont le diamètre des grandes sphères géodésiques croît sous linéairement. Il s'agit de généralisation de travaux de A. Kazue. Nous obtenons aussi une estimée de type Cheng-Yau pour le gradient des fonctions harmoniques.

1. INTRODUCTION

When (M, g) is a complete Riemannian manifold with non negative Ricci curvature, S-Y. Cheng and S-T. Yau have proven that any harmonic function $h: M \rightarrow \mathbb{R}$ satisfies the gradient estimate [4] :

$$\sup_{z \in B(x, R)} |dh|(z) \leq \frac{C(n)}{R} \sup_{z \in B(x, 2R)} |h(z)|.$$

This result implies that such a manifold can not carry non constant harmonic function $h: M \rightarrow \mathbb{R}$ with sublinear growth :

$$|h(x)| = o(d(o, x)) \text{ , } d(o, x) \rightarrow +\infty .$$

A celebrated conjecture of S-T. Yau predicted the finite dimensionality of the space of harmonic functions with polynomial growth on a complete Riemannian manifold with non negative Ricci curvature :

$$\mathcal{H}_\nu(M, g) = \{h \in \mathcal{C}^2(M) , \Delta_g h = 0, |h(x)| = \mathcal{O}(d^\nu(o, x))\} .$$

This conjecture has been proven by T. Colding and B. Minicozzi in a much more general setting.

We say that a complete Riemannian manifold (M^n, g) satisfies the *doubling* condition if there is a constant ϑ such that for any $x \in M$ and radius $R > 0$:

$$\text{vol } B(x, 2R) \leq \vartheta \text{vol } B(x, R).$$

If $B \subset M$ is a geodesic ball, we will use the notation $r(B)$ for the radius of B and κB for the ball concentric to B and with radius $\kappa r(B)$. And if f is an integrable function on a subset $\Omega \subset M$, we will note f_Ω its mean over Ω :

$$f_\Omega = \frac{1}{\text{vol } \Omega} \int_\Omega f.$$

We say that a complete Riemannian manifold (M^n, g) satisfies the scale (L^2) Poincaré inequality if there is a constant μ such that for any ball $B \subset M$ and any function $\varphi \in C^1(2B)$:

$$\|\varphi - \varphi_B\|_{L^2(B)}^2 \leq \mu r^2(B) \|d\varphi\|_{L^2(2B)}^2.$$

Theorem. [5] *If (M, g) is a complete Riemannian manifold that is doubling and that satisfies the scale Poincaré inequality then for any ν , the space of harmonic function of polynomial growth of order ν has finite dimension:*

$$\dim \mathcal{H}_\nu(M, g) < +\infty.$$

It is well known that a complete Riemannian manifold with non negative Ricci curvature is doubling and satisfies the scale Poincaré inequality, hence the Yau's conjecture is true.

The proof is quantitative and gives a precise estimation of the dimension of $\dim \mathcal{H}_\nu(M, g)$. In fact, the condition on the Poincaré inequality can be weakened and the result holds on a doubling manifold (M, g) that satisfies the mean value estimation [6, 10] : for any harmonic function defined over a geodesic ball $3B$:

$$\sup_{x \in B} |h(x)| \leq \frac{C}{\text{vol } 2B} \int_{2B} |h|.$$

An example of Riemannian manifold satisfying the above condition are Riemannian manifold (M, g) that outside a compact set (M, g) is isometric to the warped product

$$([1, \infty) \times \Sigma, (dr)^2 + r^{2\gamma} h)$$

where (Σ, h) is a closed connected manifold and $\gamma \in (0, 1]$. But when $\gamma \in (0, 1)$, a direct analysis, separation of variables, shows that any harmonic function h satisfying for some $\epsilon > 0$:

$$h(x) = \mathcal{O}\left(e^{Cr^{1-\gamma-\epsilon}}\right)$$

is necessary constant. In particular, a harmonic function with polynomial growth is constant. In [8, 9], A. Kasue has shown that this was a general result for manifold whose Ricci curvature satisfies a quadratic decay lower bound and whose geodesic spheres have sublinear growth (see also [11] for a related results):

Theorem. *If (M, g) is complete Riemannian manifold with a based point o whose Ricci curvature satisfies a quadratic decay lower bound:*

$$\text{Ricci} \geq -\frac{\kappa^2}{d^2(o, x)} g,$$

and whose geodesic sphere have sublinear growth:

$$\text{diam } \partial B(o, R) = o(R), \quad R \rightarrow +\infty$$

then any harmonic function with polynomial growth is constant.

Following A. Grigor'yan and L. Saloff-Coste [7], we say that a ball $B(x, r)$ is remote (from a fixed point o) if

$$3r \leq d(o, x).$$

Our first main result is a refinement of A. Kasue's result when the hypothesis of the Ricci curvature is replaced by a scale Poincaré inequality for remote ball : There is a constant μ such that all remote balls $B = B(x, r)$ satisfy a scale Poincaré inequality :

$$\forall \varphi \in \mathcal{C}^1(2B) : \|\varphi - \varphi_B\|_{L^2(B)}^2 \leq \mu r^2 \|d\varphi\|_{L^2(2B)}^2$$

Theorem A. *Let (M, g) be a complete Riemannian manifold whose remote balls satisfy the scale Poincaré inequality and assume that geodesic spheres have sub-linear growth:*

$$\text{diam } \partial B(o, R) = o(R) , \quad R \rightarrow +\infty.$$

If $h : M \rightarrow \mathbb{R}$ is a harmonic function such that for $I_R := \int_{B(o, R)} h^2 :$

$$\liminf_{R \rightarrow +\infty} \log(I_R) \frac{\text{diam } \partial B(o, R)}{R} = 0$$

then h is constant.

For instance, on such a manifold, a harmonic function $h : M \rightarrow \mathbb{R}$ satisfying :

$$|h(x)| \leq C d(o, x)^\nu (\text{vol } B(o, d(o, x)))^{-\frac{1}{2}}$$

is constant. Moreover if the diameter of geodesic sphere satisfies

$$\text{diam } \partial B(o, R) \leq C R^\gamma ,$$

for some $\gamma \in (0, 1)$ then if $h : M \rightarrow \mathbb{R}$ is a harmonic function such that for some positive constant C and ϵ :

$$|h(x)| \leq C e^{d(o, x)^{1-\gamma-\epsilon}} \text{vol } B(o, d(o, x))$$

then h is constant.

A by product of the proof will imply that on the class of manifold considered by A. Kasue, the doubling condition implies an estimate *à la* Cheng-Yau for the gradient of harmonic function:

Theorem B. *Let (M^n, g) be a complete Riemannian manifold that is doubling and whose Ricci curvature satisfies a quadratic decay lower bound. Assume that the diameter of geodesic sphere has a sublinear growth*

$$\text{diam } \partial B(o, R) = \sup_{x, y \in \partial B(o, R)} d(x, y) = o(R) ,$$

then there is a constant C such that for any geodesic ball $B \subset M$ and any harmonic function $h : 3B \rightarrow \mathbb{R}$

$$\sup_{x \in B} |dh|^2(x) \leq \frac{C}{\text{vol } 2B} \int_{2B} |dh|^2.$$

This result has consequences for the boundness of the Riesz transform. When (M^n, g) is a complete Riemannian manifold with infinite volume, the Green formula and the spectral theorem yield the equality:

$$\forall f \in \mathcal{C}_0^\infty(M) , \quad \int_M |df|_g^2 \text{dvol}_g = \langle \Delta f, f \rangle_{L^2} = \int_M \left| \Delta^{\frac{1}{2}} f \right|^2 \text{dvol}_g .$$

Hence the Riesz transform

$$R := d\Delta^{-\frac{1}{2}} : L^2(M) \rightarrow L^2(T^*M)$$

is a bounded operator. It is well known [12] that on a Euclidean space, the Riesz transform has a bounded extension $R : L^p(\mathbb{R}^n) \rightarrow L^p(T^*\mathbb{R}^n)$ for every $p \in (1, +\infty)$. Also according to D. Bakry, the same is true on manifolds with non-negative Ricci curvature [2]. As it was noticed in [3, section 5], in the setting of the Theorem B, the analysis of A. Grigor'yan and L. Saloff-Coste [7] implies a scale L^1 -Poincaré inequality: there is a constant C such that any balls $B = B(x, r)$ satisfies :

$$\forall \varphi \in \mathcal{C}^1(2B) : \|\varphi - \varphi_B\|_{L^1(B)} \leq Cr^2 \|d\varphi\|_{L^1(2B)}.$$

And according to the analysis of P. Auscher and T. Coulhon [1] (see also the explanations in [3, section 5]), the Theorem B implies :

Corollary C. *Under the assumption of Theorem B, the Riesz transform is bounded on L^p for every $p \in (1, +\infty)$.*

Acknowledgements. *I thank Hans-Joachim Hein : this project had begun by a very fruitful discussion where we proved together the key lemma (2.1). I'm partially supported by the grants ACG: ANR-10-BLAN 0105 and GTO : ANR-12-BS01-0004.*

2. ABSENCE OF HARMONIC FUNCTIONS

Recall that when (M, g) is a complete Riemannian manifold and $o \in M$, we say that a geodesic ball $B(x, r)$ is *remote* (from o) if

$$3r \leq d(o, x).$$

We define ρ the radius function by $\rho(t) = \inf_{x \in \partial B(o, t)} \max_{y \in \partial B(o, t)} d(x, y)$, we have

$$\rho(t) \leq \text{diam } \partial B(o, t) \leq 2\rho(t).$$

2.1. An inequality.

Lemma 2.1. *Let (M, g) be a complete Riemannian manifold whose all remote balls $B = B(x, r)$ satisfy a scale Poincaré inequality :*

$$\forall \varphi \in \mathcal{C}^1(2B) : \|\varphi - \varphi_B\|_{L^2(B)}^2 \leq \mu r^2(B) \|d\varphi\|_{L^2(2B)}^2.$$

Then there are constants $C > 0$ and $\kappa \in (0, 1)$ depending only on μ such that if

$$\forall r \in [R, 2R] : \rho(r) \leq \varepsilon r \text{ with } \varepsilon \in (0, 1/12)$$

and if h is a harmonic function on $B(o, 2R)$ then

$$\int_{B(o, R)} |dh|^2 \leq C \kappa^{\frac{1}{\varepsilon}} \int_{B(o, 2R)} |dh|^2.$$

Proof. Let $r \in [R + 4\varepsilon R, 2R - 4\varepsilon R]$, our hypothesis implies that there is some $x \in \partial B(o, r)$ such that

$$B(o, r + \varepsilon R) \setminus B(o, r) \subset B(x, \varepsilon R + \varepsilon r).$$

Let $h: B(o, 2R) \rightarrow \mathbb{R}$ be a harmonic function and $c \in \mathbb{R}$ a real number. We use the Lipschitz function :

$$\chi(x) = \begin{cases} 1 & \text{on } B(o, r) \\ \frac{r + \varepsilon R - d(o, x)}{\varepsilon R} & \text{on } B(o, r + \varepsilon R) \setminus B(o, r) \\ 0 & \text{outside } B(o, r + \varepsilon R) \end{cases}$$

Then integrating by part and using the fact that h is harmonic we get

$$\int_M \chi^2 |d(h - c)|^2 + 2\chi(h - c) \langle d\chi, d(h - c) \rangle = \int_M \langle d((h - c)\chi^2), d(h - c) \rangle = 0$$

So that we have :

$$\begin{aligned} \int_M |d(\chi(h - c))|^2 &= \int_M \chi^2 |d(h - c)|^2 + 2\chi(h - c) \langle d\chi, d(h - c) \rangle + (h - c)^2 |d\chi|^2 \\ &= \int_{B(o, r + \varepsilon R)} (h - c)^2 |d\chi|^2, \end{aligned}$$

and hence

$$\begin{aligned} \int_{B(o, r)} |dh|^2 &\leq \int_{B(o, r + \varepsilon R)} |d(\chi(h - c))|^2 = \int_{B(o, r + \varepsilon R)} (h - c)^2 |d\chi|^2 \\ &\leq \frac{1}{\varepsilon^2 R^2} \int_{B(o, r + \varepsilon R) \setminus B(o, r)} (h - c)^2 \\ &\leq \frac{1}{\varepsilon^2 R^2} \int_{B(x, \varepsilon R + \varepsilon r)} (h - c)^2. \end{aligned}$$

The hypothesis that $\varepsilon \leq 1/12$ implies that the ball $B(x, \varepsilon R + \varepsilon r)$ is remote, hence if we choose

$$c = h_{B(x, \varepsilon(R + r))} = \frac{1}{\text{vol } B(x, \varepsilon(R + r))} \int_{B(x, \varepsilon(R + r))} h$$

then the Poincaré inequality and the fact that $r + R \leq 3R$ imply :

$$\int_{B(o, r)} |dh|^2 \leq 9\mu \int_{B(x, 6\varepsilon R)} |dh|^2.$$

But we have :

$$B(x, 6\varepsilon R) \subset B(o, r + 6\varepsilon R) \setminus B(o, r - 6\varepsilon R),$$

hence we get

$$\int_{B(o, r - 6\varepsilon R)} |dh|^2 \leq 9\mu \int_{B(o, r + 6\varepsilon R) \setminus B(o, r - 6\varepsilon R)} |dh|^2.$$

And for all $r \in [R, R - 12\varepsilon R]$ we get :

$$\int_{B(o,r)} |dh|^2 \leq \frac{9\mu}{1+9\mu} \int_{B(o,r+12\varepsilon R)} |dh|^2.$$

We iterate this inequality and get

$$\int_{B(o,R)} |dh|^2 \leq \left(\frac{9\mu}{1+9\mu} \right)^N \int_{B(o,2R)} |dh|^2$$

provide that $N12\varepsilon R \leq R$; hence the result with $C = 1 + \frac{1}{9\mu}$ and

$$\kappa = \left(\frac{9\mu}{1+9\mu} \right)^{\frac{1}{12}}.$$

□

2.2. Harmonic function with polynomial growth. We can now prove the following extension of Kasue's results :

Theorem 2.2. *Let (M, g) be a complete Riemannian manifold whose all remote balls $B = B(x, r)$ satisfy a scale Poincaré inequality :*

$$\forall \varphi \in C^1(2B) : \|\varphi - \varphi_B\|_{L^2(B)}^2 \leq \mu r^2(B) \|d\varphi\|_{L^2(2B)}^2$$

Assume that balls anchored at o have polynomial growth :

$$\text{vol } B(o, R) \leq CR^\mu$$

and that geodesic spheres have sublinear diameter growth :

$$\lim_{t \rightarrow +\infty} \frac{\rho(t)}{t} = 0$$

then any harmonic function on (M, g) with polynomial growth is constant.

Proof. Let $h : M \rightarrow \mathbb{R}$ be a harmonic function with polynomial growth :

$$h(x) \leq C(1 + d(o, x))^\nu.$$

We will defined

$$E_R = \int_{B(o,R)} |dh|^2 \text{ and } \epsilon(r) = \sup_{t \geq r} \frac{\rho(t)}{t}.$$

We remark first that using the cut off function ξ defined by

$$\xi(x) = \begin{cases} 1 & \text{on } B(o, R) \\ \frac{2R-d(o,x)}{R} & \text{on } B(o, 2R) \setminus B(o, R) \\ 0 & \text{outside } B(o, 2R) \end{cases}$$

We obtain

$$(1) \quad E_R \leq \int_{B(o,2R)} |d(\xi h)|^2 = \int_{B(o,2R)} |h|^2 |d\xi|^2 \leq CR^{2\nu+\mu-2}.$$

If we iterate the inequality obtained in Lemma 2.1, we get for all R such that $\epsilon(R) \leq 1/12$:

$$E_R \leq C^\ell \kappa^{\sum_{j=0}^{\ell-1} \frac{1}{\epsilon(2^j R)}} E_{2^\ell R} .$$

Using the estimation (1), we get

$$(2) \quad E_R \leq C(R) e^{\ell \left(\frac{\log \kappa}{\ell} \sum_{j=0}^{\ell-1} \frac{1}{\epsilon(2^j R)} + \log(2)(2\nu + \mu - 2) + \log C \right)} .$$

But the Cesaro theorem convergence implies that :

$$\lim_{\ell \rightarrow +\infty} \frac{1}{\ell} \sum_{j=0}^{\ell-1} \frac{1}{\epsilon(2^j R)} = +\infty$$

hence if we let $\ell \rightarrow +\infty$ in the inequality (2) we get $E_R = 0$ and this for all sufficiently large R , hence h is constant. \square

2.3. Extension. A slight variation of the arguments yields the following extension:

Theorem 2.3. *Let (M, g) be a complete Riemannian manifold whose all remote balls $B = B(x, r)$ satisfies a scale Poincaré inequality :*

$$\forall \varphi \in C^1(2B) : \|\varphi - \varphi_B\|_{L^2(B)}^2 \leq \mu r^2(B) \|d\varphi\|_{L^2(2B)}^2$$

Assume that the geodesic spheres have sublinear diameter growth :

$$\lim_{t \rightarrow +\infty} \frac{\rho(t)}{t} = 0 \quad \text{and} \quad \text{let } \epsilon(r) = \sup_{t \geq r} \frac{\rho(t)}{t} .$$

Let $h : M \rightarrow \mathbb{R}$ be a harmonic function and assume that $I_R = \int_{B(o, R)} h^2$ satisfy

$$\log I(R) = o \left(\int_1^{R/4} \frac{dt}{t\epsilon(t)} \right)$$

then h is constant.

Proof. Indeed, the above argumentation shows that if R is large enough then

$$E_R \leq M(\ell, R) I(2^{\ell+1} R) 4^{-\ell} R^{-2}$$

where

$$\begin{aligned} \log(M(\ell, R)) &= \log \left(C^\ell \kappa^{\sum_{j=0}^{\ell-1} \frac{1}{\epsilon(2^j R)}} \right) \\ &= \ell \log C + \log \kappa \left(\sum_{j=0}^{\ell-1} \frac{1}{\epsilon(2^j R)} \right) . \end{aligned}$$

But

$$\sum_{j=0}^{\ell-1} \frac{1}{\epsilon(2^j R)} \geq \frac{1}{\log 2} \sum_{j=0}^{\ell-1} \int_{2^{j-1} R}^{2^j R} \frac{dt}{t\epsilon(t)} \geq \frac{1}{\log 2} \int_{R/2}^{2^{\ell-1} R} \frac{dt}{t\epsilon(t)} .$$

Hence we get the inequality :

$$\log E_R \leq \log I(2^{\ell+1}R) - \ell \log(4) + \ell \log C + \frac{\log \kappa}{\log 2} \int_{R/2}^{2^{\ell-1}R} \frac{dt}{t\epsilon(t)} - 2 \log R.$$

It is then easy to conclude. \square

3. LIPSCHITZ REGULARITY OF HARMONIC FUNCTIONS

We are going to prove that a Lipschitz regularity for harmonic function analogous to the the Cheng-Yau gradient inequality :

Theorem 3.1. *Let (M^n, g) be a complete Riemannian manifold that satisfy the doubling condition : there is a constant ϑ such that for any $x \in M$ and radius $R > 0$:*

$$\text{vol } B(x, 2R) \leq \vartheta \text{vol } B(x, R)$$

and assume moreover that the Ricci curvature satisfies a quadratic decay lower bound

$$\text{Ricci} \geq -\frac{\kappa^2}{r^2(x)}g,$$

where for a fixed point $o \in M$: $r(x) := d(o, x)$.

Assume that the diameter of geodesic sphere growth slowly

$$\text{diam } \partial B(o, R) = \sup_{x, y \in \partial B(o, R)} d(x, y) = o(R)$$

then there is a constant C such that for any geodesic ball $B \subset M$ and any harmonic function $h : 3B \rightarrow \mathbb{R}$

$$\sup_{x \in B} |dh|^2(x) \leq \frac{C}{\text{vol } 2B} \int_{2B} |dh|^2.$$

Proof. According to [3, Proposition 5.3], we need only to show that there is a constant C such that if $R > 0$ and if $h : B(o, 2R) \rightarrow \mathbb{R}$ is a harmonic function then for any $s \leq \sigma \leq R$:

$$(3) \quad \frac{1}{\text{vol } B(o, s)} \int_{B(o, s)} |dh|^2 \leq \frac{C}{\text{vol } B(o, \sigma)} \int_{B(o, \sigma)} |dh|^2.$$

According to the Lemma 2.1, for all $\eta > 0$, there is a $R_0 > 0$ such that for all $R \geq R_0$, then

$$\int_{B(o, R)} |dh|^2 \leq \eta \int_{B(o, 2R)} |dh|^2.$$

Hence for all $R \geq R_0$:

$$\frac{1}{\text{vol } B(o, R)} \int_{B(o, R)} |dh|^2 \leq \eta \vartheta \frac{1}{\text{vol } B(o, 2R)} \int_{B(o, 2R)} |dh|^2.$$

Choose $\eta = \vartheta^{-1}$, then we get that for all $R_0 \leq s \leq \sigma \leq R$:

$$\frac{1}{\text{vol } B(o, s)} \int_{B(o, s)} |dh|^2 \leq \frac{\vartheta}{\text{vol } B(o, \sigma)} \int_{B(o, \sigma)} |dh|^2.$$

The Ricci curvature being bounded on $B(o, 3R_0)$, the Cheng and Yau gradient estimate yields a constant B such that for all $x \in B(o, R_0)$:

$$|dh|^2(x) \leq \frac{B}{\text{vol } B(o, 2R_0)} \int_{B(o, 2R_0)} |dh|^2$$

Hence the estimate (3) holds with $C = \max\{B, \vartheta\}$. \square

REFERENCES

- [1] P. Auscher, T. Coulhon, Riesz transform on manifolds and Poincaré inequalities, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **4** (2005), no. 3, 531–555.
- [2] D. Bakry : Etude des Transformations de Riesz dans les variétés Riemanniennes à courbure de Ricci minorée. *Séminaire de probabilités XIX*, 351:145–174, 1983-1984. Lecture Notes in Mathematics, 1123.
- [3] G. Carron, Riesz transform on manifolds with quadratic curvature decay , preprint arXiv:1403.6278.
- [4] S.Y. Cheng, S.T. Yau : Differential equations on Riemannian manifolds and their geometric applications, *Comm. Pure Appl. Math.* **28** (1975), 333–354.
- [5] T-H. Colding, W-P. Minicozzi : Harmonic functions on manifolds. *Ann. of Math. (2)* **146** (1997), no. 3, 725–747.
- [6] T-H. Colding, W-P. Minicozzi : Liouville theorems for harmonic sections and applications. *Commun. Pure Appl. Math.* **51** (1998), 113–138.
- [7] A. Grigor'yan, L. Saloff-Coste : Stability results for Harnack inequalities, *Ann. Inst. Fourier (Grenoble)* **55** (2005), 825–890.
- [8] A. Kasue : Harmonic functions of polynomial growth on complete Riemannian manifolds I, *Proc. Symp. Pure Math.*, Vol. 54, part II, (1993), 281–290.
- [9] A. Kasue, Harmonic functions of polynomial growth on complete Riemannian manifolds II, *J. Math. Soc. Japan*, **47** (1995), 37–65.
- [10] P. Li : Harmonic sections of polynomial growth. *Math. Res. Lett.* **4**(1) (1997) 35–44.
- [11] C. Sormani : Harmonic functions on manifolds with nonnegative Ricci curvature and linear volume growth. *Pacific J. Math.* **192** (2000), no. 1, 183–189.
- [12] E.M. Stein : *Singular Integrals and Differentiability of Functions*, Princeton University Press, Princeton (1970).
- [13] S.T. Yau : Some function-theoretic properties of complete Riemannian manifold and their applications to geometry, *Indiana Univ. Math. J.* **25** (1976), no. 7, 659–670.

LABORATOIRE DE MATHÉMATIQUES JEAN LERAY (UMR 6629), UNIVERSITÉ DE NANTES,
2, RUE DE LA HOUSSINIÈRE, B.P. 92208, 44322 NANTES CEDEX 3, FRANCE
E-mail address: Gilles.Carron@univ-nantes.fr