

Understanding protected area resilience: a multi-scale socioal-ecological approach

G. S. Cumming, Craig Allen, Natalie Ban, Duan Biggs, Harry C. Biggs, David H.M. Cumming, Alta de Vos, Graham Epstein, Michel Etienne, Kristine Maciejewski, et al.

▶ To cite this version:

G. S. Cumming, Craig Allen, Natalie Ban, Duan Biggs, Harry C. Biggs, et al.. Understanding protected area resilience: a multi-scale socioal-ecological approach. Ecological Applications, 2014, 25 (2), pp.299-319. 10.1890/13-2113.1. hal-01134583v1

HAL Id: hal-01134583 https://hal.science/hal-01134583v1

Submitted on 23 Mar 2015 (v1), last revised 22 Sep 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ECOLOGICAL SOCIETY OF AMERICA

Ecology/Ecological Monographs/Ecological Applications

PREPRINT

This preprint is a PDF of a manuscript that has been accepted for publication in an ESA journal. It is the final version that was uploaded and approved by the author(s). While the paper has been through the usual rigorous peer review process of ESA journals, it has not been copy-edited, nor have the graphics and tables been modified for final publication. Also note that the paper may refer to online Appendices and/or Supplements that are not yet available. We have posted this preliminary version of the manuscript online in the interest of making the scientific findings available for distribution and citation as quickly as possible following acceptance. However, readers should be aware that the final, published version will look different from this version and may also have some differences in content.

The doi for this manuscript and the correct format for citing the paper are given at the top of the online (html) abstract.

Once the final published version of this paper is posted online, it will replace the preliminary version at the specified doi.

UNDERSTANDING PROTECTED AREA RESILIENCE: A MULTI-SCALE,

2	SOCIAL-ECOLOGICAL APPROACH
3	
4	Graeme S. Cumming ¹ , Craig R. Allen ² , Natalie C. Ban ³ , Duan Biggs ⁴ , Harry C. Biggs ⁵ ,
5	DAVID H. M. CUMMING ¹ , ALTA DE VOS ¹ , GRAHAM EPSTEIN ⁶ , MICHEL ETIENNE ⁷ , KRISTINE
6	Maciejewski ¹ , Raphaël Mathevet ⁸ , Christine Moore ¹ , Mateja Nenadovic ⁹ , Michael
7	SCHOON ¹⁰
8	
9	¹ Percy FitzPatrick Institute, DST/NRF Centre of Excellence, University of Cape Town, Rondebosch, Cape Town
10	7701, South Africa. E-mail graeme.cumming@uct.ac.za
11	² U.S. Geological Survey, Nebraska Cooperative Fish and Wildlife Research Unit, School of Natural Resources,
12	University of Nebraska – Lincoln, 68583
13	³ School of Environmental Studies, University of Victoria, PO Box 1700 STN CSC, Victoria BC V8W 2Y2, Canada
14	⁴ Australian Research Council Centre of Excellence for Environmental Decisions, Centre for Biodiversity and
15	Conservation Science, University of Queensland, Brisbane, Queensland 4072, Australia
16	⁵ Conservation Services Division, South African National Parks, Private Bag X402, Skukuza 1350, South Africa
17	⁶ The Vincent and Elinor Ostrom Workshop in Political Theory and Policy Analysis, Indiana University, 513 North
18	Park Ave, Bloomington, IN, 47408, USA
19	⁷ INRA Ecodevelopment Unit, Site Agroparc 84914, Avignon Cedex 9, France
20	⁸ UMR 5175 Centre d'Ecologie Fonctionnelle et Evolutive, CNRS, 1919 route de Mende, 34293 Montpellier Cedex
21	5, France
22	⁹ Duke University Marine Laboratory, Nicholas School of the Environment, Duke University
23	135 Duke Marine Lab Rd, Beaufort, NC 28516, USA
24	¹⁰ School of Sustainability, Arizona State University, PO Box 875502, Tempe, AZ 85257, USA
25	

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

Abstract. Protected Areas (PAs) remain central to the conservation of biodiversity. Classical PAs were conceived as areas that would be set aside to maintain a natural state with minimal human influence. However, global environmental change and growing cross-scale anthropogenic influences mean that PAs can no longer be thought of as 'ecological islands' that function independently of the broader social-ecological system in which they are located. For PAs to be resilient (and to contribute to broader social-ecological resilience) they must be able to adapt to changing social and ecological conditions over time in a way that supports the long-term persistence of populations, communities, and ecosystems of conservation concern. We extend Ostrom's social-ecological systems framework to consider the long-term persistence of PAs, as a form of land-use embedded in social-ecological systems, with important cross-scale feedbacks. Most notably we highlight the cross-scale influences and feedbacks on PAs that exist from the local to the global scale, contextualizing PAs within multi-scale social-ecological 'functional landscapes'. Such functional landscapes are integral to understand and manage individual PAs for long-term sustainability. We illustrate our conceptual contribution with three case studies that highlight cross-scale feedbacks and social-ecological interactions in the functioning of PAs and in relation to regional resilience. Our analysis suggests that while ecological, economic and social processes are often directly relevant to PAs at finer scales, at broader scales the dominant processes that shape and alter PA resilience are primarily social and economic. Key words: conservation, cross-scale, resilience, social-ecological system, national park, reserve,

47

spatial resilience

48

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

49 Introduction

Protected areas (PAs) remain one of conservation biology's most important approaches for ensuring that representative examples of ecological populations, communities, and ecosystems are maintained for current and future generations. Historically, most PAs were created as places that would remain 'natural' (Brandon et al. 1998). Over time, as the original focus of conservation biology on rare and endangered species has expanded into a more general awareness of the relevance of ecosystems (and the services they provide) for human wellbeing, our understanding of PAs and their objectives has changed. PAs now range from strict PAs, in which no harvesting of fauna or flora occurs and human visitation is restricted, to multiple use areas in which sustainable use of natural resources is the norm (Table 1; Dudley (2008)). PAs can no longer be viewed as purely ecological islands (Janzen 1983). Instead, as we come to better understand (1) the driving roles of regional processes (i.e., those that occur over broader extents than most PAs) in the composition of ecological communities and their spatial and temporal population dynamics; (2) the complex political and economic influences that underpin PA establishment and maintenance; (3) the role of PAs as providers of benefits for local communities and society within a broader landscape context; and (4) the potential costs of PAs, including opportunity costs, it is becoming increasingly clear that PAs are social-ecological systems that both respond to and influence a wide range of ecological, social, and political processes.

TABLE 1 HERE

PAs are human constructs in which institutions are used to try to achieve ecological and social goals. Human activities in most PAs are limited so that recognized natural, ecological and/or cultural values for some social groups are maintained (Table 1; Dudley 2008). In order to meet ecological goals, conservationists have strived to influence PA location, pattern, management and governance. The creation of state-owned or public conservation areas is usually driven by the ecological consciousness and political will of the participants (Mathevet and Mauchamp 2005) but must also confront a variety of ecological and political constraints. Defining the formal boundaries of protected areas is impossible without support from external institutions such as national and international policies, laws, and agreements. This means that the creation and maintenance of PAs is heavily dependent on their compatibility with institutions in the broader social and economic system. Each PA has social and ecosystem characteristics, often including stated management goals, that influence (and are influenced by) governance, affecting economic outputs and social outcomes in the social-ecological system (Ostrom 2009).

PAs are vulnerable to political change (Agrawal 2005, Clement 2010), economic fluctuations, and ecological change. Understanding what makes PAs resilient to both ecological and socioeconomic change is therefore important for conservation. We view resilience as being comprised of (1) the ability of a system to remain within the same regime (system state characterized by key processes) following a perturbation; and (2) the capacity of a system to adapt to change and persist through times of change (Carpenter et al. 2001, Lundy and Montgomery 2010). Resilience may also be viewed as the maintenance by a system of a continuous identity in space and time (Cumming and Collier 2005). Resilience itself is not a normative concept, and the resilience of some social-ecological states (e.g., poverty traps) may

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

be negative from a conservation perspective; we focus here on 'positive' resilience, in the sense of resilience that helps PAs to achieve conservation goals. PAs must change and adapt to changing environmental conditions through time (Lee and Jetz 2008), while seeking to maintain their cultural and social roles as important elements of their identity. The core of their identity, however, lies in the fact that they support, or at least are intended to support, the long-term persistence of populations, species and communities of a wide range of organisms, as well as related abiotic ecosystem elements and processes (Jax 2010) and ecosystem services. If PAs are to be resilient in social, economic and ecological terms, their physical location and boundaries, as well as their management and governance, must be politically viable well into the future (Folke et al. 1996, Adger et al. 2005) lest they become 'paper parks', are made smaller (e.g., the extent of Etosha National Park, in Namibia, is currently about a quarter of what it was in 1907), or are de-gazetted altogether. Management of decision-making processes is therefore at least as important for PA resilience as management of the biophysical system, suggesting that conservation science is necessarily interdisciplinary (Mathevet and Mauchamp 2005). Furthermore, PAs influence the regions in which they are embedded, and are in turn influenced by the broader context of those regions. Clearly, the maintenance and possibly enhancement of PA resilience – in a social-ecological context – is a key goal for conservation biology. The social-ecological nature of PAs has already received considerable recognition within both the peer-reviewed literature and cutting-edge conservation practice (Berkes et al. 2003, Fischer et al. 2009, Strickland-Munro et al. 2010, Ban et al. 2013). Despite the existence of a solid body of inter- and trans-disciplinary work on PAs, however, many gaps remain. Here we focus on three particular areas that require further development: (1) the relationships between a socialecological perspective on PAs and research from other fields on social-ecological systems and their resilience; (2) scale and the analysis of cross-scale influences and feedbacks on PAs; and (3) assessment of the resilience of PAs. Although many scholars have also argued for greater attention to issues of power in studies of environmental governance (Blaikie 2006, Jentoft 2007, Clement and Amezaga 2013), we do not explicitly focus on this topic in this paper. Nonetheless the close relationship between power and the rules, norms and conventions (i.e., institutions) of human societies means that power is rarely far removed from the discussion.

PAS AND SOCIAL-ECOLOGICAL SYSTEMS FRAMEWORKS

The study of social-ecological systems (SESs) has led to a wide range of frameworks, theories and models that aim to structure inquiry and explain or predict the dynamic outputs of complex adaptive systems. We use 'system' to refer to a cohesive, temporally continuous entity that consists of key elements, interactions, and a local environment (Cumming and Collier 2005). SESs are systems that include social, economic and ecological elements as well as the interactions between them. The concept of an SES is useful for PA management because it explicitly implies that the manager, other stakeholders, and related institutions are part of a cohesive whole, the system. This in turn suggests that approaches that incorporate these elements into dynamic models of system interactions - rather than treating them as immutable external influences on ecosystems - may identify opportunities to enhance the resilience of systems that would otherwise be overlooked. Moreover, PAs do not exist in a vacuum and interact with, contain, and/or are nested within other SESs.

Frameworks are underlying sets of ideas that serve to connect and make sense of different concepts (Pickett et al. 2007). They are used to aid the investigation of complex phenomena by

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

necessary.

identifying, organizing and simplifying relevant factors, and are generally compatible with multiple theories and models (Pickett et al. 2007, Schlager 2007, McGinnis 2011). Frameworks that have been explicitly developed for understanding social-ecological systems include, amongst many others, resilience (Holling 1973, Resilience Alliance 2007a, b), robustness (Anderies et al. 2004), vulnerability (Turner et al. 2003, Adger 2006), Self-organized Holarchic Open systems (SOHO) (Kay and Boyle 2008, Waltner-Toews et al. 2008), and sustainability science (Kates et al. 2001). All of these approaches have the potential to provide a unified approach for the study of SESs across multiple methods and disciplines (Ostrom 2007, 2009, Poteete et al. 2010); and all are potentially relevant as a platform to better understand the dynamics of PAs. Different frameworks have, however, tended to focus on different elements of the same problem, and no single existing framework can be considered fully comprehensive (Cumming 2011 pp. 37-40). In the context of PAs there is a strong need to bring key ideas from different frameworks together into a more comprehensive body of theory. We propose an approach that combines elements of resilience analysis (e.g., Holling 1973, Resilience Alliance 2007b) and the closely related SES framework of Ostrom (2007, 2009), while extending them in several directions. Before we consider how these frameworks can be applied to PAs, a brief summary of each set of ideas and their main strengths and weaknesses is

159

160

161

162

The Resilience Alliance workbooks (Resilience Alliance 2007a, b, 2010) attempt to operationalize key resilience concepts by posing a series of questions to strategically define and assess SESs. Within this broader framework, a nested framework (adapted from Chapin et al.

(2006)) offers a protocol to structure interacting, cross-scale social-ecological components, processes, institutions, and feedbacks. The workbooks use the adaptive cycle and panarchy models (Kenward et al. 2001, Gunderson and Holling 2002), and the adaptive governance and social-network literatures, to facilitate an understanding of system dynamics and interactions, assess governance and offer insights about potential actions. What they lack in a unified underpinning theory (Cumming 2011) is compensated for by their firm grounding in a rich empirical literature spanning many case-studies and multiple disciplines (reviewed in Haider et al. (2012)). In providing a practical way to structure multiple resilience perspectives in complex, dynamic SESs, the framework offers an approach to understand issues of scale in SESs, including PAs, and proposes a novel approach to natural resource management (Walker et al. 2009, Strickland-Munro et al. 2010, Haider et al. 2012).

The resilience approach has, however, been criticized for being difficult to operationalize (Strickland-Munro et al. 2010, Cumming 2011, Holt et al. 2012). Practical problems in applying resilience thinking have resulted in a relatively low number of directly comparable case-study examples. Practitioners have also lamented its lack of guidance for delineating system boundaries, developing tools to navigate a transition to desirable futures, and describing governance structures (Strickland-Munro et al. 2010, Haider et al. 2012, Holt et al. 2012). These criticisms are particularly relevant for PAs, where implicit geographic or ecologically-relevant boundaries (e.g., catchment edges) may not line up with PA boundaries (Mitchell 2011), and where identifying social thresholds and variables, articulating governance choices and incorporating relations of power (Strickland-Munro et al. 2010, Armitage et al. 2012) may be particularly important in defining elements that may contribute to or erode a system's resilience

(Walker et al. 2009).

Ostrom's SES framework (Figure 1) provides a useful complement to resilience approaches. It has its origins in institutional studies of the commons that made significant contributions towards a game theoretic understanding of environmental governance (Ostrom 1990, Ostrom et al. 1994). It provides researchers an analytical tool with which to capture, organize and analyze a diverse set of social and ecological variables that are considered relevant for a particular aspect of a system (Ostrom 2007, Poteete et al. 2009). In total, Ostrom's SES framework includes over fifty potentially influential classes of variables that are ordered within a multilevel classificatory system. The four core components (resource systems, resource units, actors and governance systems) are organized as a partially decomposable system (Simon 1991) where each of the potentially influential variables can be further unpacked to capture subclasses and cumulatively integrate knowledge concerning their effects on sustainability. There are two additional components that allow for linkages across levels of governance, or between systems, and an additional two components that are used to evaluate SES interactions and outcomes.

FIGURE 1 HERE

Ostrom's SES framework has been criticized on several fronts that generally point to two main issues. First, its origin in institutional analysis neglects alternative social scientific perspectives. Most notable among these omissions are the power-laden theories of political ecology that view environmental degradation as a direct consequence of imbalances of power between influential policymakers (e.g., national governments) and their associates (e.g., local elites and businesses)

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

and marginalized small-scale users (e.g., subsistence farmers and pastoralists) (Peet and Watts 1993, Robbins 2004). Second, the ecological aspects of the framework and their interactions remain underdeveloped (Berkes and Ross 2013). A particularly problematic issue for ecologists seeking to apply Ostrom's SES framework is its lack of clear definitions concerning resource units and resource systems. For example, resource units have been operationalized at multiple levels of biological organization, including species and communities (Gutierrez et al. 2011), water and land (Ostrom 2011) and even landscapes for tourism (Blanco 2011). While it could be argued that each of these studies presents an internally consistent application of the framework, it is unclear whether syntheses between such disparate case studies are feasible or if the findings necessarily apply to the broader population of SESs. Third, while dynamic and multi-scale analysis is technically possible, nearly all applications of the framework and its institutional analysis precursor focus on a single focal action situation (e.g., resolution of a natural resource management problem by multiple stakeholders) that occurs once only and in a single location (McGinnis 2011). Moreover, until recent modifications to the SES framework were introduced by Epstein et al. (2013), the framework was poorly equipped to analyze biophysical processes and diagnose ecological contributions to social-ecological outcomes. However, even with these changes, Epstein et al.'s (2013) analysis of the successful remediation of Lake Washington simply transforms inherently dynamic internal phosphorus loading processes into several static one-way relationships with the dependent variable. Although sufficient for their analysis, the failure to account for dynamic linkages within and across scales remains a major weakness of the SES framework. In fact most applications of the framework have a general tendency to focus on a single scale or level of governance, on a single resource, and to treat the problem as if all resources and actors were at the same focal scale.

As analytical approaches for understanding (and hence, better managing) PAs, both resilience approaches and Ostrom's SES framework have much to recommend them. Our objective in this article is to extend them to better integrate social-ecological feedbacks and cross-scale effects that often dominate the dynamics of PAs and other social-ecological systems.

EXTENDING EXISTING FRAMEWORKS TO INCLUDE SCALE AND CROSS-SCALE FEEDBACKS

The obvious tension between ecological and social demands in many PAs suggests that analysis of the resilience of PAs requires a hierarchical, cross-scale and multi-level framework in which different scales and institutional levels are connected by a set of interactions between different actors, resources and processes. Examples of interactions include the movements of actors and resources (e.g., tourism, water flows out of PAs to downstream communities) as well as the interplay of rules and information across scales. Holling (2001) suggested that complex system behaviours, such as those that we observe in PAs, arise from the interactions of processes that occur at a minimum of three different spatial and temporal extents; and furthermore, that in many cases, shifts between different system states are driven by changes in the slower variables (e.g., buildup of Phosphorus in a shallow lake, or loss of trust in human society) rather than the faster variables (e.g., trophic interactions or law enforcement).

It is important to recognize that PAs, which are institutions (in Ostrom's sense) rather than biophysical entities, have been created at a variety of different spatial scales and institutional levels. While PAs in the IUCN categories I-IV are often single tenure units, those in categories V and VI (such as Biosphere reserves and Transfrontier conservation areas) usually include

multiple, nested tenure units that are governed by different rules. For example, the current rulesin-use in the Greater Limpopo Transfrontier Conservation Area or the Causse Méjan (both of which are discussed in more detail later in this paper) differ between farms, core conservation areas, hunting areas, and designated buffer zones. Similarly, while larger areas may be expected to change more slowly because of the buffering effect of larger ecological populations, this is not inevitable; political change that has an influence at a national extent, for example, can happen swiftly. In heterogeneous landscapes, different tenure units at the same spatial scale may also interact (e.g., mines and conservation areas).

263

264

265

266

267

268

269

270

271

272

273

274

262

255

256

257

258

259

260

261

We propose a system description that includes five hierarchical levels of institutional organization. These in turn are related to five hierarchical spatial scales of analysis, with some flexibility, depending on the system that is under analysis (Figure 2). The first institutional level is defined as the *sub-tenure unit*. It refers to patches of habitat (or any other fine-scale, discrete ecological units that are managed differently), and/or specific human use areas, that (1) fall within a single tenure unit; (2) have a single management policy; and (3) exist at smaller spatial extents than the boundaries of the tenure unit. For example, different rules about making fires or leaving your vehicle may apply at picnic sites or bird hides (blinds) within a PA; and different habitats in a PA may have different management needs. Sub-tenure units will always, by definition, have a smaller spatial extent than a PA. They relate most closely to the patch scale of analysis, which reflects the grain and extent of habitat heterogeneity within the PA.

275

276

277

The second institutional level is that of a *single tenure unit*. Single tenure units belong to a single owner or organization. They may have the same extent or a smaller extent than that of a PA,

depending on the diversity of tenure types and human use zones occurring within the PA. Single tenure units define one or several scales of analysis that might (for example) correspond to the extent of a traditional game park, or to those of core ecological and farming areas respectively within a Biosphere reserve.

282

283

284

285

286

287

288

289

290

291

292

293

294

278

279

280

281

The third institutional level, the *proximate institutional context*, includes multiple tenure units as well as the institutions and organizations that are responsible for coordinating the interactions (where these occur) between tenure units. Depending on the nature of the study system, the proximate context might define a spatial scale that is only slightly larger than the PA, or a larger region that contains a network of PAs that are managed with a shared objective. For example, provincial parks in the Western Cape of South Africa form a network that is overseen by a regional conservation organization, Cape Nature; the proximate institutional context for any single provincial park includes Cape Nature, and related ecological scales of analysis include surrounding PAs and unprotected dispersal corridors that connect PAs. The proximate institutional context also includes institutions that relate to the governance of resources around PAs, particularly where (as in the case of water laws, for example) they relate directly to ecological flows (e.g., water, invasive species) that might enter the PA from surrounding areas.

295

296

297

298

299

300

Proximate institutions in turn sit within (or sometimes straddle) a national institutional context, the fourth level, which typically consists of the institutions of a single nation-state (e.g., its constitution and related governance structures). This institutional level aligns with a national extent of analysis. However, sometimes, as in the case of transboundary conservation areas, PAs may include as many as three or four nations, creating an international institutional context that

is the fifth and final institutional level. This fifth level includes international power relations and the global economy. International contexts are aligned with the broadest scales of spatial analysis, ranging from multiple countries to global. While the fifth level may seem ecologically far removed from the majority of established PAs, it has particular relevance for migratory species and related resources, such as wetlands that are important for migratory waterbirds and are supposedly covered by international conventions and agreements (e.g., CBD, Ramsar, AEWA; see United Nations 1992, Matthews 1993, Lenten 2001). Similarly, international conventions and agreements (or lack thereof) can have a strong influence at the level of a single tenure unit, as in the case of the management of species that are listed in Appendix I of the Convention in Trade and Endangered Species (CITES).

At each different scale and level, different temporal dynamics occur. The temporal scales that are relevant to the ecology of PAs range from short term processes such as predation and competition that occur on a daily scale, through seasonal processes such as breeding or wintering seasons for birds, to long-term processes such as atmospheric oscillations, ocean acidification, and climate change that take place at decadal and centennial scales. Similarly, the temporal scales within the social realm also vary from short-term initiatives to establish PAs to long-standing national assets such as Yellowstone National Park in the USA. Social history and pre-PA politics may also affect the resilience and social-ecological context of a PA. Both ecological and social processes act synergistically to produce outcomes and thus neither can be considered in isolation (Hughes et al. 2005).

FIGURE 2 HERE

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

The boundaries of a PA can span multiple nested institutional and ecological levels and scales. Each PA interacts directly with its immediate context (i.e., defined by the scale and level above), which becomes the main source of both inputs and outputs (e.g., information, finances) for any given PA. Since the number and diversity of people involved in SESs at different levels affects both ecological and social processes and the temporal periods over which they occur (Westley et al. 2002), it is crucial to take these interactions into consideration. Practitioners often speak of getting the different levels of governance aligned: for example, a decision made internationally, at say Ramsar Convention level, may or may not promote wetland sustainability, depending on whether the national government/s that are involved take action and are supported by local communities. In other cases, local communities may seek support at the international level for initiatives that lacked support from their own national governments. Alternatively, some of the policies funded and promoted by international donors and organizations may contribute to systematic disenfranchisement of local communities despite a supporting rhetoric of social justice (Blaikie 2006). Positive synergies among scale-dependent institutions therefore usually depend upon brokering organizations that facilitate (even make possible) the interactions between the various levels, including in the reverse direction so that the national governments and Ramsar convention receive the necessary confirmation or other feedback to assist future policy formulation. Worldviews, values, attitudes, and power are key factors that shape PA design and governance and shape the behaviors and practices of social groups operating at different spatial levels that directly or indirectly affect PAs.

Resources and biophysical processes exist over a range of different scales, and vary in their grain (or frequency) and their extent (or duration) in space and in time. The scale of socioeconomic processes depends heavily on the scale of economic and political organization and the level of international interest in a particular PA (Figure 3). It ranges from individuals to networks of organizations and includes the effective scales of social institutions (rules, laws, policies, and norms) that govern the extent of resource-related rights and management responsibilities (Cumming et al. 2006). For example, in creating an urban PA, a country's constitution may provide for national-level tenure rights that must then be applied within the local context of municipal policies and by-laws. Scale-dependent system components and interactions may influence PAs in different ways depending on how their relative magnitude and frequency change across different scales. It is often unclear where resilience, or a lack of resilience, resides - both within and between scales. The interactions between different spatial and temporal scales of both pattern and process, and their potential effects on resilience, are recurring themes in the ecological literature (e.g., Peterson et al. 1998, Cumming et al. 2006).

360

346

347

348

349

350

351

352

353

354

355

356

357

358

359

FIGURE 3 HERE

362

363

364

365

366

367

368

361

Dealing with the many scale dependencies of PAs is conceptually challenging. However, as Cash et al. (2006) point out, ignorance of cross-scale interactions tends to reveal itself in a wide range of management problems. Therefore, a useful starting point is to be explicit about the spatial and temporal elements of the problem and their key scales (Figure 4). PAs in contemporary conservation efforts are developed as networks (Vimal et al. 2012). They are planned and increasingly managed as part of local, regional and international conservation systems. For

example, in the French national park and biosphere reserve design approach, new PAs are designed as a set of zones that range from strictly protected areas (core area of national park or nature reserve) to integrated zones in surrounding areas where integrated management of natural resources is implemented with local stakeholders and landowners (Batisse 1997, Mathevet et al. 2010).

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

369

370

371

372

373

Social-ecological interactions occur most intensively within and between entities that operate at similar scales (Allen and Starr 1982, Levin 1992, 1999) (Figure 2). For example, in South Africa, provincial administrations such as Cape Nature or Ezemyelo-KZN Parks manage provincial parks, while national parks are regulated nationally by SANParks. At the same time, actors and processes at scales and levels above and below the focal scale influence patternprocess interactions via flows between nested elements. Matter includes the exchange of physical materials across scales and levels, such as water, carbon, and nitrogen. Organisms, including people as well as mobile animals and plant propagules, link scales and levels via their movements (e.g., labour, migration, transhumance). Information flows include the exchange of ideas, perceptions and skills across scales. These local to global flows and the ways in which they are mediated and managed can play an important role in the function and performance of the PA (Mathevet et al. 2010, Thompson et al. 2011) and may consolidate ecological and social interdependence in biodiversity policy that goes beyond park boundaries, such as the health of the tourism sector in and around PAs (Hall 2010, Biggs 2011). Rules link institutions and regulations across scales, for example with global treaties affecting regulations within PAs. In addition, flows of information, perceptions, and money across scales are central to the functioning of the nature-based tourism sector in many of the world's PAs (e.g., Biggs 2011).

392

The presence of different interlinked subsystems across different scales (Figure 4) suggests the presence of multiple action arenas where decisions related to PAs are made and a strong need to somehow align multiple subsystems to coordinate responses to common threats (e.g., climate change or an escalation of poaching activity). This observation aligns neatly with Holling's ideas about panarchies (Holling and Gunderson 2002), which suggest that some degree of synchrony in system cycles is a necessary pre-condition for effective interventions (Westley et al. 2002).

399

400

401

402

403

404

405

406

407

408

409

410

393

394

395

396

397

398

The adoption of a multi-scale, social-ecological perspective on the resilience of PAs (Figures 2-4) provides a useful way of organizing and thinking through their long-term sustainability. Over the last decade, conservation organizations have increasingly recognized that the protection of ecosystems requires that key ecosystem functions and processes be maintained at multiple scales (Poiani et al. 2000). Several of the world's largest conservation NGOs have developed stratified, ecoregional-based plans and approaches to formally structure the process of developing and maintaining PA networks (e.g., TNC 2003, Loucks et al. 2004). Poiani et al. (2000), for instance, developed a hierarchical classification for habitats (ranging from small patches through the matrix to entire regions) and associated species (ranging from small patch species through to regional and long-distance migratory species; Figure 4) as part of TNC's 'Conservation by Design' initiative.

411

412

FIGURE 4 HERE

413

414

415

We propose that a similar leap forward must be taken by recognizing that multi-scale socioeconomic (and further, social-ecological) 'functional landscapes' exist, and that they are integral to understanding and managing PAs for long-term sustainability. For example, the Man and the Biosphere Programme (UNESCO) integrates social and ecological goals and aims to ensure the sustainable use of natural resources, while also emphasizing the interdependencies of cultural and natural landscapes (Batisse 1971, IUCN 1979, Batisse 1997, German MAB National Committee 2005). Such areas are structured and organized at a range of social and ecological scales, depending on the particular set of negotiated goals and objectives. The new concept of ecological solidarity, a core feature of the 2006 law reforming national park policy in France, similarly stresses the need to reconnect people to their PAs. Ecological solidarity is both social and ecological; it is based on social recognition of the spatial interdependence among natural organisms, including people, and their physical environment. This sets the scene for a vision of nature conservation and management of PAs. Ecological solidarity offers a pragmatic compromise between ecocentric and anthropocentric ethics. It suggests that biodiversity conservation at different spatial and temporal scales needs to be collectively explored by local communities and stakeholders to give social meaning to the establishment of PAs, to the expansion of ecological networks, and to the integrated management of cultural landscapes (Mathevet 2012).

432

433

434

435

436

437

438

431

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

As an example, conservation of a 'local-scale species', such as an endemic butterfly, typically requires fewer resources and a much finer scale of management than that which is required to conserve species that use their landscape at a regional scale (e.g., migratory songbirds). In the same way, meeting local stakeholder needs and demands within and around a PA requires different and much finer-scale action than governmental resource policies, the international tourist market, or the international trade in animal products. These ideas can be summarized by

uniting the ecological approach of Poiani et al. (2000) with a socioeconomic perspective (Figure 5).

441

439

440

FIGURE 5 HERE

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

442

Figure 5 provides a way of conceptualizing and comparing the different scales and levels at which social and ecological systems are organized. It does not, however, provide a dynamic temporal perspective for understanding interactions between scales. One of the key components of incorporating scale and scaling in a framework for the analysis of PA resilience is that of understanding feedbacks, both within and between scales. Formally, a cross-scale feedback occurs if (1) A influences B and B influences A; and (2) A and B are system elements (whether human or not, but excluding interactions) that exist at different scales. For instance, global demand drives the prices of many commodities but production is often limited to a smaller subset of locations. If local conditions in the production location influence global prices, a local to global interaction occurs. If global prices also influence local actions, a cross-scale feedback occurs. Such feedbacks may be extremely difficult to manage given the inherent complexity of social-ecological systems (Berkes et al. 2006). For example, the Asian demand for rhinoceros horn is driven by cultural beliefs. Coupled with limits on local production (i.e., a small number of slowly-reproducing rhinos), it has created spiraling commodity price increases and a massive conservation problem for African PAs (Biggs et al. 2013). Our future ability to manage individual resources, or PAs as a whole, will depend on our ability to devise a system that can detect potentially harmful feedbacks and respond to them in a timely manner (Hughes et al. 2005, Biggs et al. 2013).

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

The different social-ecological system elements that determine the resilience of an individual PA may be connected in different ways and to varying degrees of strength (Figures 2-4). One of the challenges in analyzing PA resilience is to determine which influences are the strongest within the system and which are sufficiently weak that they can safely be ignored or disregarded during analysis (keeping in mind that sometimes, weak influences and dormant social networks can be important in times of crisis). Closed feedback loops (A influences B influences C influences A) are also of particular importance because they can produce surprising dynamics, such as dampening or exacerbation of local variability. In practice these feedbacks (and especially those that reinforce one another) are critical for system performance and must be considered in the design of environmental policies as they are directly responsible for the stability of a socialecological system in a given state. Conversely, if system change is desired, they must in some direct or indirect manner (e.g., through modifying other inter-linkages which feed into it but can be influenced) be overcome. For example, in the Eastern Cape case study discussed below and in Maciejewski & Kerley (in press), managers' perceptions of what tourists want to see provide a powerful driver for the ecological management of private PAs (PPAs). This influence is crossscale in the sense that tourists come from a far wider extent than the PA. By their actions, managers in turn influence the likelihood that tourists will visit the PA, setting up a cross-scale feedback that can result in harmful ecological effects (e.g., habitat alteration by excessive numbers of elephants, and resulting species loss) within PA boundaries. Breaking this feedback requires that managers be willing to accept data indicating that tourist numbers would be unaffected by lower stocking rates, and willing to take the risk of reducing population levels of charismatic megafauna such as elephants.

While many studies have implied or discussed the importance of scaling principles and cross-scale dynamics for PAs, few have explicitly analyzed them. Some exceptions include Jones et al. (2013) and Mills et al. (2010) (both on MPAs) and Guerrero et al. (2013). Guerrero et al. (2013) identified eight ways in which scale mismatches between actors and resources involved in the spatial planning process manifest themselves. These include (1) ecosystem or ecological processes that extend beyond governance boundaries; (2) the absence of resolution-appropriate data for decision making; (3) a lack of implementation capacity; (4) threats to ecological diversity that operate at diverse spatial and temporal scales; (5) mismatches between funding and the long-term nature of ecological processes; (6) rates of implementation that do not reflect the rate of change of the ecological system; (7) lack of appropriate indicators for monitoring activities; and (8) the occurrence of ecological change at scales smaller or larger than the scale of implementation of management actions or monitoring.

Among the most important questions in this context are (1) whether, and how, PAs may contribute to desirable regional resilience (e.g., Plumptre et al. 2007, Slotow and Hunter. 2009, Cantú-Salazar and Gaston. 2010, Laurance 2013, Sjöstedt 2013); and conversely, (2) how regional resilience may influence the resilience of individual PAs (González et al. 2008, Jones et al. 2013).

UNDERSTANDING THE RESILIENCE OF PROTECTED AREAS

In the preceding sections we have argued that (1) PAs are multi-scale and multi-level socialecological systems; and (2) an explicit recognition of scale and cross-scale interactions must be incorporated in analyses of PA resilience if we are to advance our understanding of their dynamics, manage them better, and ultimately foster their resilience. The third logical step in this line of argument is to consider empirical evidence that indicates whether, and how, cross-scale feedbacks may in practice influence the resilience of specific PAs, and how PAs may in turn influence regional resilience. The starting point is to define different scales and levels; this is followed by a more detailed consideration of system dynamics. We illustrate these steps below for three real-world examples, noting that space constraints and the goals of this article do not permit the next step, which would be a full resilience analysis of each case. As the subsequent discussion shows, the nature of the interaction between regional and local resilience may be quite strongly dependent on context-specific factors.

518

519

520

521

522

523

524

525

526

527

528

508

509

510

511

512

513

514

515

516

517

Case Study 1: Social-ecological dynamics of Private Protected Areas in the Eastern Cape Private Protected Areas (PPAs) constitute a high proportion of conservation land in South Africa. Exact figures are hard to obtain, but according to the PAs Act 57 of 2003, approximately 7% of the country's land is in statutory national parks and 17% in some form of private conservation area (Cousins et al. 2008). In southern Africa, ecotourism generates roughly the same revenue as farming, forestry and fisheries combined (Scholes and Biggs 2004). Growth in the ecotourism industry has had substantial impacts in the Eastern Cape, where large areas of marginal pastoral lands have given way to PPAs. Private PAs may fall within any of the IUCN categories. Some believe they are better represented under categories IV – VI, although many private PAs fit the management objectives of categories I – III (Dudley 2008).

529

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

At the **sub-tenure level and patch extent**, within PAs, former agricultural fields with interspersed natural areas have been converted into more economically viable game farms. This involves restoring the vegetation and reintroducing wildlife into the area. At the **PA level**, system dynamics and related ecological management decisions are heavily driven by economic processes. Private PAs aim to build populations of charismatic species at stocking levels that ensure tourist satisfaction. It has been estimated that during the establishment of PPAs, the introduction of species to the Eastern Cape cost between \$97,500 and \$1.8 million (Sims-Castley et al. 2005). Stocked animals are often extralimital species, such as giraffe, which did not historically occur in the Eastern Cape. These non-indigenous introductions have several negative effects including hybridization, degradation of habitat, and low survival rates and competitive exclusion of indigenous species (Chapin et al. 2000, Castley et al. 2001). Stocking charismatic species, such as the African elephant (Loxodonta africana), above ecological carrying capacity to meet social demands and ensure tourist satisfaction may also have negative ecological impacts. Numerous studies have documented significant impacts of elephants on biodiversity (e.g., Cumming et al. 1997, Blignaut et al. 2008, Kerley et al. 2008). At regional, national, and international levels and extents, the main driving forces are socialecological processes, represented by two conflicting trends. On one hand the land conversion

At regional, national, and international levels and extents, the main driving forces are social-ecological processes, represented by two conflicting trends. On one hand the land conversion trend increases ecotourism in the Eastern Cape, potentially leading to an increase in income and job opportunities, and resulting in social uplift and poverty alleviation in the rural communities surrounding the PPAs. On the other hand, the ecological carrying capacity of the PPAs places a threshold on the types and numbers of species that can be introduced. The habitat fragmentation and land degradation that can result from overstocking large herbivores may reduce the number

of national and international tourists visiting the area. In addition, as a consequence of South Africa's history, many areas of the Eastern Cape are contested (Cundill et al. 2005); reserve creation may engender social resentment and create political opposition to conservation, particularly if it entails the loss of jobs formerly provided by agriculture (Brooks et al. 2011).

557

558

559

560

561

562

563

564

565

566

567

568

553

554

555

556

Case Study 2: Man and the Biosphere (MAB) case: Regime shifts on the Causse Méjan The Causse Méjan is a limestone plateau (1000m average altitude) in the Cévennes Mountains of France. It is home to the largest steppe-like grassland in France (Fonderflick et al. 2013), and is part of the core area of the Cévennes National Park (372000 ha) and the Cévennes Biosphere Reserve. Both PAs were created to maintain a rural way of life (including sheep and cattle farming, and cheese production) as well as to support the conservation of indigenous grassland and several endangered species (e.g., vultures, Przewalski's horse). The Causse Méjan is an IUCN category VI PA. Farmers are the managers of open meadows and steppes, which cover 37% of the core area; the rest is forest (O'Rourke 1999, Etienne and Le Page 2002). The plateau is ecologically vulnerable to bush encroachment and invasion by pine, boxwood and juniper trees (Etienne 2001).

569

570

571

572

573

574

575

At the **patch scale**, the main ecological driving force is the pine seed rain intensity. There is a threshold of grazing pressure above which pine encroachment is impossible. Below this threshold, pine tree regeneration can be controlled by mechanical or manual removal of pine seedlings. The transformation of grassland to woodland represents a local ecological regime shift, but may not be a regime shift at the extent of an individual farm because the main driving forces at the **sub-tenure level** are economic. Here, social-ecological regime shifts are provoked

577

578

579

580

581

582

583

584

585

586

587

588

589

592

593

594

595

596

597

598

by changes in the percentage of a farm covered by pine forest but the threshold will differ according to the area of grassland per stock required by the farming system (Kinzig et al. 2006). The farmer will select which farming system to practice for cultural and economic reasons that are largely derived from higher levels, such as national prices for livestock (Etienne and Le Page 2002). Both vegetation patches and farms occur within the broader extent of the biosphere reserve. At the biosphere reserve (PA and PA network levels) and the regional extent, the main driving forces are both social and ecological, potentially producing two conflicting kinds of system change. On one hand, a regime shift may occur between cheese and timber production, depending on the unstable interactions between Roquefort cheese, Fedou cheese (a local cheese that like Roquefort, is produced from sheep's milk) and lamb meat producers, and timber producers (O'Rourke 1999, Kinzig et al. 2006). On the other hand, the pine-grassland dynamic may result in ecological regime shifts and the loss of open grassland biodiversity (Kinzig et al.

590 2006). Finally, **national level** institutions, policies and **international** commodity demands will

591 influence economic tradeoffs in this system.

Interestingly, two recent changes have provoked a new regime shift and management paradigm.

First, in June 2011, the Causse Méjan and some neighboring sites of the Cévennes, were declared

as world cultural heritage sites by UNESCO for their agro-pastoral Mediterranean landscape.

Second in the summer of 2012, and for the first time in the 21st Century, wolf attacks were

registered (n=36) and 22 ewes killed. The Causse Méjan social-ecological system is now at a

threshold. Will it switch to a bushy and abandoned farmland landscape, supporting the

establishment of a permanent wolf pack, or will it remain an agropastoral landscape, giving priority to sustainable sheep grazing practices and the conservation of open grassland? This dilemma questions the goals and practices of the national park, especially in its core area, as well as the adaptive capacity of farmers to integrate predators into their grazing system.

603

604

605

606

607

608

609

610

611

612

613

614

616

617

618

619

620

599

600

601

602

Case study 3: The Greater Limpopo Transfrontier Conservation Area

The Greater Limpopo Transfrontier Conservation Area (GLTFCA, c. 90,000 km²) includes adjacent parts of Mozambique, South Africa and Zimbabwe. The core PA comprises three national parks (IUCN Category II): Limpopo in Mozambique, Kruger in South Africa, and Gonarezhou in Zimbabwe. Other PAs, mostly IUCN Category VI, are included in each country, as are areas of communal and private land. State PAs cover 53% and communal lands 34% of the area, respectively (Cumming et al. 2013). The core GLTFCA, created by treaty between the three countries, is embedded within a provisional transfrontier conservation area that may serve as a buffer that increases the resilience of the GLTFCA. Within South Africa, the Kruger to Canyons Biosphere initiative (Coetzee et al. 2012) is extending the area under protection (IUCN Category VI).

615

Historically, Khoisan people occupied the area for millennia before Bantu agro-pastoralists arrived some 2,000 years ago. Livestock appeared in areas adjacent to and within the GLTFCA between 600 and 1200 AD and various species of antelope, and hippo and elephant were hunted and ivory was traded at the coast (Plug 2000). The period 1200 - 1800 was characterised by shifting tribal control of the region and Nguni invasions.

621

The entry of smallpox and measles into the region in the 1830s and the rinderpest pandemic in 1895 took their toll on both humans and ungulates respectively. The major transitions and drivers and associated social-ecological changes in the GLTFCA landscape are summarised in Fig. 6. At the patch scale and sub-tenure unit level, within the GLTFCA, changes have occurred in ecological habitat connectivity, disturbance regimes, water availability, and herbivore species composition and abundance as well as in the settlement patterns of people and their farming practices. At the protected area scale, numerous changes have occurred in boundaries and associated tenure rights (see detailed explanations in Cumming et al (2007), and Anderson and Cumming (2013)). The key feature of the changes that have occurred since 1830 is that they have largely been driven by political dictates at international and national levels. Initial change was driven by European colonisation of the three countries, and then by national policies of racial segregation resulting in the development of dual agricultural systems in South Africa and Zimbabwe. This resulted on the one hand in the development of large commercial farms on privately owned land and on the other, in increased densities of traditional small scale agropastoral farms in communal lands. Superimposed on this matrix were the formation of state PAs and the resulting displacement of people.

638

639

640

641

642

643

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

The continuing top-down influences of international and national policies and legislation on resource management in the GLTFCA continue, with significant impacts on the management of animal diseases (e.g. Foot and Mouth Disease) and the conservation and management of three species of charismatic mega-herbivores (elephant and black and white rhinoceros) (Biggs et al. 2013).

644

645 FIGURE 6 HERE 646 647 Key Elements within Case Studies 648 Despite the different locations and scales of each of the three case studies, they share 649 considerable commonality in their key drivers (Figure 7). 650 651 FIGURE 7 HERE 652 653 Interestingly, our case studies suggest that ecological processes are often most directly relevant to PAs at intermediate to finer scales. In the Eastern Cape case study, carrying capacity and 654 habitat fragmentation both occur at the patch and PA scales. Similarly, pine seed rain intensity, 655 656 grazing pressure, bush encroachment, grassland to woodland transformation, predator-prey 657 dynamics and species home-ranges are finer-scale elements in the Causse Méjan. In the 658 GLTFCA, ecological habitat connectivity, disturbance regimes and herbivore species composition and abundance are also patch- and protected-area scale processes and patterns. 659 660 661 At broader scales, the dominant processes that shape and alter PAs are primarily sociopolitical 662 and economic. In our case-studies, the top-down drivers were elements such as tourism demands (Eastern Cape case study), international policies and commodity demands (Causse Méjan), and 663 664 colonization and international and national policy changes (GLTFCA). 665 666 Sociopolitical and economic processes may of course impact ecosystems via impacts on the 667 abiotic environment, as in the case of anthropogenic climate change, which is driven by human

socioeconomic demands for such things as energy, transport, and manufactured goods. The main exception to this general pattern arises when migratory species are particularly important elements of a PA; this is not the case in any of our examples, but it is not uncommon. We could also envisage that regional ecological influences become relatively more important for smaller PAs that are more dependent on colonization from nearby natural areas that are not necessarily within the boundaries of the PA (Bengtsson et al. 2003).

674

675

676

677

678

668

669

670

671

672

673

Dynamic interactions within case studies

If we consider a more dynamic representation of cross-scale interactions, the different variables summarized in Figure 7 interact to drive change in PAs. In Figure 8 we graph the spatial scale of our case-study variables against a notional speed at which these processes typically operate.

679

FIGURE 8 HERE

681

682

683

684

680

As these graphical depictions show, temporal scales do not arrange as readily along a hierarchy as spatial scales, creating opportunities for spatial-temporal scale mismatches (Cumming et al. 2006).

685

686

687

688

689

690

Since system changes are usually driven by feedbacks, particularly cross-scale feedbacks (Walker et al. 2006), it follows that slower feedbacks and feedbacks from slower processes will take longer to drive change than feedbacks from faster processes. Our examples show that topdown, broad-scale processes like tourism demand can change over short time periods, while processes like habitat fragmentation manifest at a smaller spatial scale, but can take much longer

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

to manifest and drive change. If, as our PA examples show, ecological processes such as succession and trophic cascades generally occur at smaller, slower scales (i.e., more gradually and at smaller extents, noting that biophysical perturbations are not 'ecological processes') and socio-economic drivers occur at broader, faster ones, an emerging hypothesis is that because of differential selection, PA social-ecological systems gradually become better adapted to cope with changes that result from sociopolitical drivers than with feedbacks from ecological processes. As a result of inertia and cross-scale gradients, top-down sociopolitical processes may drive the system to develop along a trajectory that renders it less resilient to large shocks that may eventually manifest from cross-scale ecological feedbacks. For example, timber demand during and after the second world war led to forest fire management policies in the USA that were designed to save timber; resulting management approaches eventually led to the hugely destructive 1988 fire in Yellowstone National Park.

Case Study Insights

Figures 7 and 8 provide strong support for two general points that we have emphasized throughout the paper. First, PAs function as social-ecological systems, and hence understanding their social and economic components is as fundamental as understanding their ecology if we are to analyse and manage their resilience. Second, cross-scale processes are highly relevant to the resilience of PAs and should be considered as integral to any analysis, rather than treated as subordinate to analyses of system dynamics at a single scale.

710

711

712

713

In addition to current cross-scale influences, the history and current objectives of each case study PA play an important role in their current resilience. In the Causse Méjan, with its long history of human use and livelihood support, the PA is politically uncontested and is seen as a way of

715

716

717

718

719

720

721

722

723

724

725

726

727

maintaining its unique regional identity. In southern Africa, with its colonial history, PAs are sometimes seen as a form of neo-colonial land grab. This is particularly true in South Africa, where the memory of apartheid is still recent. About 40% of National and Provincial PAs in the Eastern Cape are under some form of land claim from historically dispossessed local communities, and the political acceptability of PPAs is unclear. The potential for land redistribution from conservation to agriculture, whether legally or through illegal occupation (as has occurred in Zimbabwe), therefore represents a very real possibility. PAs in South Africa must maintain their sociopolitical resilience by remaining accessible and continuing to cater for and support all strata of society, even if this reduces their overall potential economic viability. Similarly, although the GLTFCA was created after the end of apartheid, Kruger Park has a contested history and the greater PA was also created in a relatively top-down manner by intergovernmental agreements. Its continued viability as a conservation area thus depends heavily on maintaining its political acceptability.

728 It is obvious that PAs and their effectiveness in conserving biodiversity will be influenced by 729 regional changes, particularly in relation to politics, governance, and conflict. Uncertainty over 730 land tenure can definitively undermine conservation efforts. For example, ongoing conflict in the 731 Democratic Republic of the Congo, and the displacement of local communities from their land, 732 is endangering key populations of chimpanzees and gorillas; and poor governance and land 733 appropriations in Zimbabwe have undoubtedly contributed to declines in rhinoceros populations 734 in Zimbabwean PAs. One of the starkest conservation challenges in regions with poor 735 governance remains that of working out how to protect PAs against the winds of political 736 change.

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

Our case studies also suggest that PAs contribute to regional social-ecological dynamics and hence to regional social-ecological resilience. For example, in the Eastern Cape, if PAs maintain patches of indigenous vegetation that would otherwise be converted to agriculture, they may be able to cumulatively reduce local fragmentation and maintain a range of ecosystem services and natural processes (as has been shown in similar systems: e.g., seed dispersal by cavity-nesting birds that depend on dead wood in old, large trees; Joseph et al. (2011)) that contribute positively to human wellbeing (Cumming and Spiesman 2006). Similarly, in the Causse Méjan, the PA contributes to building and maintaining a regional identity that includes an awareness of the reliance of the community on ecosystems.

DISCUSSION

We have argued that if we are to understand and enhance the long-term resilience of PAs, we must adopt an inter- or trans-disciplinary perspective that incorporates (at a minimum) elements of ecology and social science. Similarly, our analysis shows that questions of scale and recognition of cross-scale influences are of fundamental importance for PAs. Our case studies illustrate the interlinked nature of PAs as social-ecological systems. Intriguingly, it is particularly at broader scales that social, political, and economic considerations become paramount. While this may be due in part to ways of thinking or management practices that are still rooted in the internationally validated and powerful wilderness discourse, it also reflects the broad-scale nature of socioeconomic processes and ongoing globalization.

Our case study analysis does not explicitly consider an additional element of scale-related problems and multi-scale interactions, that of emergent 'higher-level' system properties arising from the interactions of elements at a single scale. Many PAs belong to socioeconomic networks. These may be formal, as in the case of National and Provincial Parks, which are generally the responsibility of a governmental management agency; or informal, through exchanges of information and resources (e.g., Goss and Cumming 2013). PAs are also members of an ecological network that facilitates the propagation and movements of animals and plants. Membership in a network may increase the resilience of an individual PA (e.g., by providing additional options for problem solving) or decrease it if acts to serve the interests of local, regional and global elites (e.g., if membership in a network demands the imposition of locally inappropriate management practices). Clearly, network membership and its relevance for PA resilience will change with scale and should thus form part of any scaling analysis of PA resilience.

772

773

774

775

776

777

778

779

780

781

759

760

761

762

763

764

765

766

767

768

769

770

771

Although they remain propositions rather than established generalities, our cross-scale extension of Ostrom's SES framework suggests some general theoretical principles for the resilience of PAs. These propositions can serve as the basis for more specific hypotheses that future studies about social-ecological resilience of PAs can test. First, there is a relationship between the scales and levels at which different system elements exist and the frequency and/or magnitude of their interactions. This is a general principle that is derived from hierarchy theory and has been further reinforced by ecological research (Allen and Starr 1982, Levin 1992, 2005). Fine-scale processes may be irrelevant for understanding system dynamics at larger scales of analysis, or conversely may occur at speeds such that larger-scale dynamics are largely irrelevant for their outcomes. For 782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

example, the movements of individual atoms are inconsequential to understanding an animal's movement path; and continental drift has had a profound influence on global species composition but is largely irrelevant for understanding PAs at the time scales that are of interest to managers. It may also be easier to generalize about larger-scale pattern-process dynamics because a considerable amount of fine-scale variation is averaged out at broader scales (Levin 1992). Social-ecological feedbacks should therefore be most pronounced when they occur between a given functional scale of the ecosystem and the most closely aligned socioeconomic scale, and/or the scales immediately above or below the focal scale (see Figure 4). For analyses of PA resilience, this means that recognizing and making explicit the ways in which system scales and levels align and interact with one another should clarify the most important perturbations against which resilience and adaptive capacity must be built, and help in making decisions about management tradeoffs. For example, in the GLTFCA, threats to the area's protected status from higher-level political processes may suggest enhancing social acceptability and community engagement through providing greater access to parts of the PA or the resources that it contains (e.g., permitting mopane worm harvesting (Makhado et al. 2009, Gondo et al. 2010)), whereas threats from pathogens introduced by or transmitted to livestock in neighbouring areas may require greater segregation and reduced access (Rodwell et al. 2001, Caron et al. 2003). Second, the kinds of interactions and feedback loops in which PAs participate may have

differing consequences for system resilience, particularly in relation to the spatial and temporal scales of different actors and interactions. Although interactions between closely aligned ecological scales and socioeconomic levels (e.g., the extent of grassland that is necessary for

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

game viewing, the scale at which the manager can implement controlled burns, and the monthly gate revenue of the PA) may dominate the 'usual' dynamics of the PA, very broad-scale or very slow variables – acting either directly or indirectly - can have important implications for overall system resilience, regime shifts, and management (Carpenter and Gunderson 2001, Lundy and Montgomery 2010). For example, a gradual trend towards regional deforestation may affect rainfall and temperature patterns within a PA, potentially leading to irreversible changes in vegetation composition and long-term impacts on ecosystem service provision to surrounding human communities. A closely related phenomenon is that of the shifting baseline, where change that is slow by human standards may mean that degraded ecosystem states (e.g., reduced size of fishes in marine PAs, or lower levels of forage in a rangeland) become regarded as normal. Slow variables in particular can lead to surprises and push PAs into traps (i.e., states in which feedbacks maintain an undesirable system state, such as a low-diversity thicket in a savanna system) that can result in a loss of resilience and eventual collapse (Carpenter and Turner 2000).

Third, we would expect to find decay in the strengths of drivers (and related feedback effects) with both distance and time. For example, the numbers of tourists visiting a PA decline with increasing distance from airports and major cities (de Vos et al, in prep., this issue). Remote PAs thus experience lower human impacts and are managed differently from those that are more accessible. Similarly, while path dependencies may be important in understanding the current locations of PAs, their influence also diminishes with time. For example, many southern African PAs were originally set aside for hunting (rather than exploited for farming) because of the presence of sleeping sickness, malaria, and tick-borne diseases. Tsetse flies have been eradicated in some areas and their distributions, and those of malaria vectors and ticks, are likely to change

828

829

830

831

832

833

834

835

836

837

844

845

846

847

848

849

850

further as the global climate is altered by people (Rogers and Randolph 1993, 2000, Cumming and Van Vuuren 2006), making all three kinds of disease increasingly less relevant to the location of PAs. For PA resilience, the principle of time- and distance-based declines in driver and feedback strengths suggests that PA resilience will correlate with both ecological and socioeconomic connectivity, but in different ways for different drivers, depending on whether resilience is enhanced or reduced by the distance effect. Remoteness may result in fewer visitors and lower economic resilience, for example, but may also reduce the potential impacts of such factors as poaching, pesticide use on neighbouring farmland, and water extraction outside the PA.

Fourth, the resilience of a complex system should correlate to its size; larger and older PAs, and 838 839 those established areas that involve more people, should be more resilient (although not 840 inevitably so). Note that we use 'older' here to refer to PAs that have had natural habitat cover 841 for a longer period of time, and in contrast to areas that are reclaimed or restored from farmland 842 or other land uses; some newly proclaimed PAs may have 'old' ecosystems and 'young' social 843 systems. Larger, older PAs (1) will be more resilient to natural perturbations, such as fires or

or de-gazetted; (5) may have a history that invests them with greater cultural meaning (e.g., more people remember childhood holidays there, and it may have achieved 'iconic' status, like Yellowstone National Park or Kruger National Park); (6) will have larger sunk costs, in the form of infrastructure and investment in the park; (7) are more likely to contain multiple IUCN categories, thereby achieving multiple goals that different stakeholders might have; and (8) are less likely to experience the level of social change that is needed to transform their management or for them to be de-proclaimed. It is possible, of course, that 'revolt' processes occur that lead to change in larger PAs, and/or that their size makes them a more obvious target for land redistribution initiatives, but on average we would expect them to be more resilient.

861

862

863

864

865

866

867

868

869

870

871

872

873

851

852

853

854

855

856

857

858

859

860

Fifth, given the many different ways in which power relations work in different societies, the relative importance of top-down and bottom-up influences is likely to be asymmetrical and dependent on the context in which the PA exists. As we have shown in the three case studies, understanding context-dependent factors is essential to the proper functioning of a PA. Therefore, there are no governance panaceas for building PA resilience that can be applied with equal success to all situations (Ostrom and Cox 2010). For example, in a nation with a weak government, it may be very difficult to buffer PAs from higher-level influences (e.g., development pressures, resource acquisition by the rich and powerful, or regional conflicts) or to implement policies and laws at scales relevant for effective PA management. Normative issues, value systems, and attitudes will influence PA resilience. Incorporating stakeholders in building local resilience, even where regional resilience is low, should be a major focus of conservation efforts. Current thinking suggests that the growing role of NGOs, international agencies, scientific groups and private operators should be explored in the context of the development of

polycentric governance of PAs where community-based management, Integrated and Conservation Development Projects and adaptive co-management approaches are promoted and implemented. It is not clear yet, however, whether such consensus-based approaches will be sufficient to maintain PAs in the face of demographic and globalization processes.

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

874

875

876

877

Our framework and proposed principles have implications for PA management and planning, although the uptake and application of some of these insights might be challenging. Our case studies show that PA managers and planners cannot afford to ignore either ecological or social dynamics, or (more importantly) their interactions at scales and levels below and above that of the PA. Analyses of the key drivers of change will assist with identifying the relevant scales of processes that are likely to influence PA management and planning. Such analyses must be undertaken with a clear idea of the PA's social-ecological role, goals, and objectives. When new PAs are planned, emphasis on larger, multi-objective and multi-IUCN category PAs may lead to improved long-term viability of the area. Cross-scale institutional linkages have the potential to serve as a link between top-down and bottom-up influences. However, while incorporating these elements into management and planning would be desirable, national and international legal and political systems may not readily allow for adaptively managing PAs as interacting cross-scale SESs (Garmestani and Allen in press). Similarly, institutional and cultural constraints may further exacerbate the challenges of changing embedded management approaches. In particular, many PAs have a top-down, command-and-control history and approach to management (e.g., Andersson and Cumming 2013, Goss and Cumming 2013). Challenging these legal, political, institutional and cultural constraints is paramount for making PAs more resilient into the future.

896

We have argued throughout this paper that understanding PAs as social-ecological systems is integral to developing the approaches, and the science, that will be required to maintain PAs as functional and effective conservation tools into the next century. While awareness of the multifaceted nature of PAs has been gradually building in conservation biology for many years, our understanding of their dynamics is still weak in some areas, particularly in relation to quantifying and managing the ability of PAs to withstand shocks arising from socioeconomic and governance-related variance at higher and lower scales. Concepts from social-ecological systems research that explicitly address cross-scale feedback loops and resilience appear to offer a range of useful conclusions in this context, and we look forward to further growth in this important area of research.

ACKNOWLEDGEMENTS

We are grateful to the Resilience Alliance for enabling a meeting of the RA Protected Areas Working Group in South Africa, and to Karen Kotschy for her useful comments on an earlier draft of this manuscript. This research was supported by the South African National Research Foundation, a James S. McDonnell Foundation grant to GC, and the SCALES project (funded by the European Commission as a Large-scale Integrating Project within FP 7 under grant 226852). The Nebraska Cooperative Fish and Wildlife Research Unit is jointly supported by a cooperative agreement among the U.S. Geological Survey, the Nebraska Game and Parks Commission, the University of Nebraska, the U.S. Fish and Wildlife Service and the U.S. Wildlife Management Institute.

LITERATURE CITED

920 Adger, W. N. 2006. Vulnerability. Global Environmental Change 16:268-281. 921 Adger, W. N., K. Brown, and E. L. Tompkins. 2005. The political economy of cross-scale 922 networks in resource co-management. Ecology and Society 10. 923 Agrawal, A. 2005. Environmentality: technologies of government and the making of subjects. 924 Duke University Press Durham, NC. 925 Allen, T. F. H. and T. B. Starr. 1982. Hierarchy: Perspectives for ecological complexity. The 926 University of Chicago Press, Chicago. Anderies, J. M., M. A. Janssen, and E. Ostrom. 2004. A framework to analyze the robustness of 927 928 social-ecological systems from an institutional perspective. Ecology and Society 9. 929 Andersson, J. A. and D. H. M. Cumming, 2013. Defining the edge: boundary formation and 930 TFCAs in Southern Africa. In: Anderson J. A, de Garine-Wichatitsky, M, Cumming, D. 931 H. M., Dzingirai, V., and Giller, K. E. (eds.) Transfrontier Conservation Areas: People 932 Living on the Edge. Routledge. 933 Armitage, D., C. Béné, A. T. Charles, D. Johnson, and E. H. Allison. 2012. The interplay of 934 well-being and resilience in applying a social-ecological perspective. Ecology and 935 Society 17:15. http://dx.doi.org/10.5751/ES-04940-170415. 936 Ban, N. C., M. Mills, J. Tam, C. Hicks, S. Klain, N. Stoeckl, M. C. Bottrill, J. Levine, R. L. 937 Pressey, T. Satterfield, and K. M. A. Chan. 2013. Towards a social-ecological approach 938 for conservation planning: embedding social considerations. Frontiers in Ecology and the 939 Environment 11:194-202. 940 Batisse, M. 1971. Man and the Biosphere: an international research programme. Biological 941 Conservation **4**:173-179.

) 42	Batisse, M. 1997. Biosphere reserves: a challenge for biodiversity conservation and regional
943	development. Environment 39.
944	Bengtsson, J., P. Angelstam, T. Elmqvist, U. Emanuelsson, C. Folke, M. Ihse, F. Moberg, and
945	M. Nystrom. 2003. Reserves, resilience and dynamic landscapes. Ambio 32:389-396.
946	Berkes, F., J. Colding, and C. Folke, editors. 2003. Navigating Social-Ecological Systems:
947	Building Resilience for Complexity and Change. Cambridge University Press,
948	Cambridge.
949	Berkes, F., T. P. Hughes, R. S. Steneck, J. A. Wilson, D. R. Bellwood, B. Crona, C. Folke, L. H.
950	Gunderson, H. M. Leslie, J. Norberg, M. Nystrom, P. Olsson, H. Osterblom, M. Scheffer,
951	and B. Worm. 2006. Ecology - Globalization, roving bandits, and marine resources.
952	Science 311 :1557-1558.
953	Berkes, F. and H. Ross. 2013. Community Resilience: Toward an Integrated Approach. Society
954	& Natural Resources 26 :5-20.
955	Biggs, D. 2011. Understanding resilience in a vulnerable industry - the case of reef tourism on
956	Australia's Great Barrier Reef. Ecology & Society 16:30. Available online:
957	http://www.ecologyandsociety.org/vol16/iss31/art30/.
958	Biggs, D., F. Courchamp, R. Martin, and H. P. Possingham. 2013. Legal trade of Africa's rhino
959	horns. Science 339 :1038 – 1039.
960	Blaikie, P. 2006. Is small really beautiful? Community-based natural resource management in
961	Malawi and Botswana. World Development 34:1942-1957.
962	Blanco, E. 2011. A social-ecological approach to voluntary environmental initiatives: the case of
963	nature-based tourism. Policy Sciences 44:35-52.

964	Blignaut, J., M. De Wit, and J. Barnes. 2008. The economic value of elephants.in R. J. Scholes
965	and K. G. Mennell, editors. Elephant Management: A Scientific Assessment of South
966	Africa. Witwatersrand University Press, Johannesburg.
967	Brandon, K., K. H. Redford, and S. E. Sanderson. 1998. Parks in peril: people, politics, and
968	protected areas. Island Press.
969	Brooks, S., M. Spierenburg, L. Van Brakel, A. Kolk, and K. B. Lukhozi. 2011. Creating A
970	Commodified Wilderness: Tourism, Private Game Farming, And 'Third
971	Nature' Landscapes In Kwazulu Natal. Tijdschrift Voor Economische En Sociale
972	Geografie 102 :260-274.
973	Cantú-Salazar, L. and K. J. Gaston. 2010. Very large protected areas and their contribution to
974	terrestrial biological conservation Bioscience, 60:808-818.
975	Caron, A., P. C. Cross, and J. T. du Toit. 2003. Ecological implications of bovine tuberculosis in
976	African buffalo herds. Ecological Applications 13:1338-1345.
977	Carpenter, S., B. Walker, J. M. Anderies, and N. Abel. 2001. From Metaphor to Measurement:
978	Resilience of What to What? Ecosystems 4:765-781.
979	Carpenter, S. R. and L. H. Gunderson. 2001. Coping with collapse: Ecological and social
980	dynamics in ecosystem management. Bioscience 51 :451-457.
981	Carpenter, S. R. and M. G. Turner. 2000. Hares and tortoises: Interactions of fast and slow
982	variables in ecosystems. Ecosystems 3 :495-497.
983	Cash, D. W., W. N. Adger, F. Berkes, P. Garden, L. Lebel, P. Olsson, L. Pritchard, and O.
984	Young. 2006. Scale and cross-scale dynamics: governance and information in a
985	multilevel world. Ecology and Society 11:8.

986 Castley, J. G., A. F. Boshoff, and G. I. H. Kerley. 2001. Compromising South Africa's natural 987 biodiversity—inappropriate herbivore introductions. South African Journal of Science 988 **96**:365-378. 989 Chapin, F. S., E. S. Zavaleta, V. T. Eviner, R. L. Naylor, P. M. Vitousek, H. L. Reynolds, D. U. 990 Hooper, S. Lavorel, O. E. Sala, S. E. Hobbie, M. C. Mack, and S. Diaz. 2000. 991 Consequences of changing biodiversity. Nature **405**:234-242. 992 Chapin, I., F.S, A.L. Lovecraft, E. S. Zavaleta, J. Nelson, M. D. Robards, G. P. Kofinas, S. F. 993 Trainor, G. D. Peterson, H. P. Huntington, and R. L. Naylor. 2006. Policy strategies to 994 address sustainability of Alaskan boreal forests in response to a directionally changing 995 climate. Proceedings of the National Academy of Sciences of the United States of 996 America 103:16637–16643. 997 Clement, F. 2010. Analysing decentralised natural resource governance: proposition for a 998 "politicised" institutional analysis and development framework. Policy Sciences 43:129-999 156. 1000 Clement, F. and J. M. Amezaga. 2013. Conceptualising context in institutional reforms of land 1001 and natural resource management: the case of Vietnam. International Journal of the Commons 7. 1002 1003 Coetzee, M., H. C. Biggs, and S. Malan. 2012. Sharing the benefits of biodiversity: a regional 1004 action plan to nurture and sustain the contribution of biodiversity and ecosystem services 1005 to livelihoods and resilient economic development within the Kruger to Canyons 1006 Biosphere. Available at: http://www.kruger2canyons.org/consolidated technical report 1007 vfin.pdf. SANParks, Nelspruit.

1008	Cousins, J. A., J. P. Sadler, and J. Evans. 2008. Exploring the Role of Private Wildlife Ranching
1009	as a Conservation Tool in South Africa: Stakeholder Perspectives. Ecology and Society
1010	13:43. http://www.ecologyandsociety.org/vol13/iss42/art43/ .
1011	Cumming, D., H. Biggs, M. Kock, N. Shongwe, and S. and Osofsky. 2007. The AHEAD
1012	(Animal Health for the Environment And Development) - Great Limpopo Transfrontier
1013	Conservation Area (GLTFCA) Programme: Key questions and conceptual framework
1014	revisited. http://wcs-ahead.org/workinggrps_limpopo.html .
1015	Cumming, D. H. M., V. Dzingirai, and M. d. G. Wichatitsky. 2013. Land and natural resource-
1016	based livelihood opportunities in TFCAs.in d. GW. Anderson J. A, M, Cumming, D. H
1017	M., Dzingirai, V., and Giller, K. E., editor. Transfrontier Conservation Areas: People
1018	Living on the Edge. Routledge.
1019	Cumming, D. H. M., M. B. Fenton, I. L. Rautenbach, R. D. Taylor, G. S. Cumming, M. S.
1020	Cumming, J. M. Dunlop, A. G. Ford, M. D. Hovorka, D. S. Johnston, M. Kalcounis, Z.
1021	Mahlangu, and C. V. R. Portfors. 1997. Elephants, woodlands and biodiversity in
1022	southern Africa. South African Journal of Science 93:231-236.
1023	Cumming, G. S. 2011. Spatial Resilience in Social-Ecological Systems. Springer.
1024	Cumming, G. S. and J. Collier. 2005. Change and identity in complex systems. Ecology and
1025	Society 10:29 [online] URL: http://www.ecologyandsociety.org/vol10/iss21/art29/ .
1026	Cumming, G. S., D. H. M. Cumming, and C. L. Redman. 2006. Scale Mismatches in social-
1027	ecological systems: causes, consequences, and solutions. Ecology and Society 11:14.
1028	Cumming, G. S. and B. J. Spiesman. 2006. Regional problems need integrated solutions: Pest
1029	management and conservation biology in agroecosystems. Biological Conservation
1030	131 :533-543.

1031	Cumming, G. S. and D. P. Van Vuuren. 2006. Will climate change affect ectoparasite species
1032	ranges? Global Ecology and Biogeography 15:486-497.
1033	Cundill, G., C. Fabricius, and M. Neus. 2005. Foghorns to the future: using knowledge and
1034	transdisciplinarity to navigate complex systems. Ecology and Society 10:8.
1035	Dudley, N. 2008. Guidelines for Applying Protected Area Management Categories. IUCN, x +
1036	86pp.
1037	Etienne, M. 2001. Pine trees-invaders or forerunners in Mediterranean-type ecosystems: a
1038	controversial point of view. Journal of Mediterranean Ecology 2:221-232.
1039	Etienne, M. and C. Le Page. 2002. Modelling contrasted management behaviours of stakeholders
1040	facing a pine encroachment process: an agent-based simulation approach. Pages 208-213
1041	in Proceedings International Environmental Modelling and Software Society Conference,
1042	Lugano, Switzerland.
1043	Fischer, J., G. D. Peterson, T. A. Gardner, L. J. Gordon, I. Fazey, T. Elmqvist, A. Felton, C.
1044	Folke, and S. Dovers. 2009. Integrating resilience thinking and optimisation for
1045	conservation. Trends in Ecology & Evolution 24:549-554.
1046	Folke, C., C. S. Holling, and C. Perrings. 1996. Biological diversity, ecosystems, and the human
1047	scale. Ecological Applications 6 :1018-1024.
1048	Fonderflick, J., A. Besnard, and J. L. Martin. 2013. Species traits and the response of
1049	open habitat species to forest edge in landscape mosaics. Oikos 122:42-51.
1050	Garmestani, A. S. and C. R. Allen. in press. Social-ecological resilience and law. Uiversity of
1051	Columbia Press.
1052	German MAB National Committee. 2005. Full of Life: UNESCO Biosphere Reserves, Model
1053	Regions for Sustainable Development. Springer.

1054	Gondo, T., P. Frost, W. Kozanayi, J. Stack, and M. Mushongahande. 2010. Linking knowledge
1055	and practice: Assessing options for sustainable use of mopane worms (Imbrasia belina) in
1056	southern Zimbabwe. Journal of Sustainable Development in Africa 12:127-145.
1057	González, J. A., C. Montes, J. Rodríguez, and W. Tapia. 2008. Rethinking the Galapagos Islands
1058	as a complex social-ecological system: implications for conservation and management.
1059	Ecology and Society 13:13. [online] URL:
1060	http://www.ecologyandsociety.org/vol13/iss12/art13/.
1061	Goss, J. and G. S. Cumming. 2013. Networks of wildlife translocations in developing countries:
1062	an emerging conservation issue? Frontiers in Ecology and Environment 11:243-250.
1063	Guerrero, A. M., R. McALLISTER, J. Corcoran, and K. A. Wilson. 2013. Scale Mismatches,
1064	Conservation Planning, and the Value of Social Network Analyses. Conservation
1065	Biology 27 :35-44.
1066	Gunderson, L. and C. Holling, editors. 2002. Panarchy: understanding transformations in human
1067	and natural systems. Island Press, Washington, DC.
1068	Gutierrez, N. L., R. Hilborn, and O. Defeo. 2011. Leadership, social capital and incentives
1069	promote successful fisheries. Nature 470 :386-389.
1070	Haider, L. J., A. E. Quinlan, and G. D. Peterson. 2012. Interacting Traps: Resilience Assessment
1071	of a Pasture Management System in Northern Afghanistan. Planning Theory & Practice
1072	13 :299-333.
1073	Hall, C. M. 2010. Crisis events in tourism: subjects of crisis in tourism. Current Issues in
1074	Tourism 13 :401 - 417.
1075	Holling, C. S. 1973. Resilience and stability of ecological systems. Annual Review of Ecology
1076	and Systematics 4:1-23.

1077	Holling, C. S. 2001. Understanding the complexity of economic, ecological, and social systems.
1078	Ecosystems 4 :390-405.
1079	Holling, C. S. and L. H. Gunderson. 2002. Resilience and adaptive cycles. Pages 25-62 in L. H.
1080	Gunderson and C. S. Holling, editors. Panarchy: understanding transformations in human
1081	and natural systems. Island Press, Washington, DC.
1082	Holt, R. F., R. P. Bio, P. A. G. Utzig, and R. F. P. Pearce. 2012. Assessment and Decision-
1083	making for Climate Change: An Overview of Theory and Approaches. West Kootenay
1084	Climate Vulnerability and Resilience Project Report #2.
1085	URL: http://www.westkootenayresilience.org/Report2_Approaches_Final.pdf .
1086	Hughes, T. P., D. R. Bellwood, C. Folke, R. S. Steneck, and a. J. Wilson. 2005. New paradigms
1087	for supporting the resilience of marine ecosystems. Trends in Ecology & Evolution
1088	20 :380-386.
1089	IUCN. 1979. The Biosphere Reserve and Its Relationship to other Protected Areas. International
1090	Union for Conservation of Nature and Natural Resources, Gland, Switzerland.
1091	Janzen, D. H. 1983. No park is an island, increase in interference from outside as park size
1092	decreases. Oikos 41 :402–410.
1093	Jax, K. 2010. Ecosystem Functioning. Cambridge University Press, Cambridge, U.K.
1094	Jentoft, S. 2007. In the power of power: The understated aspect of fisheries and coastal
1095	management. Human Organization 66 :426-437.
1096	Jones, P. J. S., W. Qiu, and E. M. D. Santo. 2013. Governing marine protected areas: Social-
1097	ecological resilience through institutional diversity. Marine Policy 41:5-13, ISSN 0308-
1098	0597X, http://dx.doi.org/0310.1016/j.marpol.2012.0312.0026.

1099	Joseph, G., G. S. Cumming, D. H. M. Cumming, Z. Mahlangu, and C. Seymour. 2011. Large
1100	termitaria act as refugia for tall trees, deadwood and cavity-using birds in a miombo
1101	woodland. Landscape Ecology 26 :439-448.
1102	Kates, R. W., W. C. Clark, R. Corell, J. M. Hall, C. C. Jaeger, I. Lowe, J. J. McCarthy, H. J.
1103	Schellnhuber, B. Bolin, N. M. Dickson, S. Faucheux, G. C. Gallopin, A. Grubler, B.
1104	Huntley, J. Jager, N. S. Jodha, R. E. Kasperson, A. Mabogunje, P. Matson, and H.
1105	Mooney. 2001. Sustainability Science. Science 292.
1106	Kay, J. J. and M. Boyle. 2008. Self-organizing, holarchic, open systems (SOHOs).in D. Waltner-
1107	Toews, J. J. Kay, and NM. E. Lister, editors. The ecosystem approach: complexity,
1108	uncertainty, and managing for sustainability. Columbia University Press, New York.
1109	Kenward, R., R. Clarke, K. Hodder, and S. Walls. 2001. Density and linkage estimators of home
1110	range: nearest-neighbor clustering defines multinuclear cores. Ecology 82:1905-1920.
1111	Kerley, G. I. H., M. Landman, L. Kruger, and N. Owen-Smith. 2008. The effects of elephants on
1112	ecosystems and biodiversity.in R. J. Scholes and K. G. Mennell, editors. Elephant
1113	Management. A Scientific Assessment for South Africa. Witwatersrand University Press,
1114	Johannesburg.
1115	Kinzig, A. P., P. A. Ryan, M. Etienne, H. E. Allison, T. Elmqvist, and B. H. Walker. 2006.
1116	Resilience and regime shifts: assessing cascading effects. Ecology and Society 11.
1117	Laurance, W. F. 2013. Does research help to safeguard protected areas? Trends in ecology &
1118	evolution. In press.
1119	Lee, T. M. and W. Jetz. 2008. Future battlegrounds for conservation under global change.
1120	Proceedings of the Royal Society B-Biological Sciences 275:1261-1270.

1121	Lenten, B. 2001. A flying start for the agreement on the conservation of African Eurasian
1122	Migratory Waterbirds (AEWA). Journal of International Wildlife Law & Policy 4:159-
1123	164.
1124	Levin, S. A. 1992. The Problem of Pattern and Scale in Ecology. Ecology 73 :1943-1967.
1125	Levin, S. A. 1999. Fragile dominion: complexity and the commons. Perseus Books, Reading,
1126	MA.
1127	Levin, S. A. 2005. Self-organization and the emergence of complexity in ecological systems.
1128	Bioscience 55 :1075-1079.
1129	Loucks, C., J. Springer, S. Palminteri, J. Morrison, and H. Strand. 2004. From the Vision to the
1130	Ground: A guide to implementing ecoregion conservation in priority areas. Worldwide
1131	Fund for Nature, Washington, D.C.
1132	Lundy, M. G. and W. I. Montgomery. 2010. A multi-scale analysis of the habitat associations of
1133	European otter and American mink and the implications for farm scale conservation
1134	schemes. Biodiversity and Conservation 19:3849-3859.
1135	Maciejewski, K. and G. I. H. Kerley. in press. Elevated elephant density does not improve
1136	ecotourism opportunities, suggesting convergence in social and ecological objectives.
1137	Ecological Applications.
1138	Makhado, R. A., G. P. Von Maltitz, M. J. Potgieter, and D. C. Wessels. 2009. Contribution of
1139	woodland products to rural livelihoods in the northeast of Limpopo Province, South
1140	Africa. South African Geographical Journal 91:46-53.
1141	Mathevet, R. 2012. La solidarité écologique, Ce lien qui nous oblige. Actes Sud, Paris.
1142	Mathevet, R. and A. Mauchamp. 2005. Evidence-based conservation: dealing with social issues.
1143	Trends in Ecology and Evolution 20 :422-423.

1144	Mathevet, R., J. D. Thompson, O. Delanoe, M. Cheylan, C. Gil-Fourrier, and M. Bonnin. 2010.
1145	La solidarité écologique: un nouveau concept pour la gestion intégrée des parcs nationaux
1146	et des territoires. Natures Sciences Sociétés 18:424-433.
1147	Matthews, G. V. T. 1993. The Ramsar Convention on Wetlands: its History and Development.
1148	The Ramsar Convention, Gland, Switzerland.
1149	McGinnis, M. D. 2011. An Introduction to IAD and the Language of the Ostrom Workshop: A
1150	Simple Guide to a Complex Framework. Policy Studies Journal 39:169-183.
1151	Mills, M., R. Weeks, R. L. Pressey, S. Foale, and N. C. Ban. 2010. A mismatch of scales:
1152	challenges in planning for implementation of marine protected areas in the Coral
1153	Triangle. Conservation Letters 3 :291-303.
1154	Mitchell, R. C. 2011. Social-Ecological Inventories: Building Resilience to Environmental
1155	Change within Biosphere reserves. Climate Adaptation Workshop Report, Brock
1155 1156	Change within Biosphere reserves. Climate Adaptation Workshop Report, Brock University, Canada.
1156	University, Canada.
1156 1157	University, Canada. URL: http://www.resalliance.org/files/raprojects/11/BESRU_2011_SEI_Workshop_Repo
1156 1157 1158	University, Canada. URL: http://www.resalliance.org/files/raprojects/11/BESRU_2011_SEI_Workshop_Repo rt.pdf.
1156115711581159	University, Canada. URL: http://www.resalliance.org/files/raprojects/11/BESRU_2011_SEI_Workshop_Repo rt.pdf. O'Rourke, E. 1999. The Causse Méjan: changing relationships between agriculture, environment
1156 1157 1158 1159 1160	University, Canada. URL: htt.pdf . O'Rourke, E. 1999. The Causse Méjan: changing relationships between agriculture, environment and society within a French National Park. Landscape Research 24:141-165.
1156 1157 1158 1159 1160 1161	URL: http://www.resalliance.org/files/raprojects/11/BESRU 2011 SEI Workshop Report.pdf . O'Rourke, E. 1999. The Causse Méjan: changing relationships between agriculture, environment and society within a French National Park. Landscape Research 24:141-165. Ostrom, E. 1990. Governing the Commons. Cambridge University Press, New York.
1156 1157 1158 1159 1160 1161 1162	University, Canada. URL: http://www.resalliance.org/files/raprojects/11/BESRU 2011 SEI Workshop Report.pdf . O'Rourke, E. 1999. The Causse Méjan: changing relationships between agriculture, environment and society within a French National Park. Landscape Research 24:141-165. Ostrom, E. 1990. Governing the Commons. Cambridge University Press, New York. Ostrom, E. 2007. A diagnostic approach for going beyond panaceas. Proceedings of the National

1166	Ostrom, E. 2011. Reflections on "Some Unsettled Problems of Irrigation". American Economic
1167	Review 101 :49-63.
1168	Ostrom, E. and M. Cox. 2010. Moving beyond panaceas: a multi-tiered diagnostic approach for
1169	social-ecological analysis. Environmental Conservation 37:451-463.
1170	Ostrom, E., R. Gardner, and J. Walker. 1994. Rules, Games and Common-Pool Resources. Ann
1171	Arbor: University of Michigan Press.
1172	Peet, R. and M. Watts. 1993. Development Theory and Environment in an Age of Market
1173	Triumphalism. Economic Geography 69:227-253.
1174	Peterson, G., C. R. Allen, and C. S. Holling. 1998. Ecological resilience, biodiversity, and scale.
1175	Ecosystems 1:6-18.
1176	Pickett, S. T. A., C. Jones, and J. Kolasa. 2007. Ecological understanding: the nature of theory
1177	and the theory of nature. Academic Press, New York.
1178	Plug, I. 2000. Overview of iron age fauna from the Limpopo Valley. South Africa
1179	Archaeological Society Goodwin Series 8:117-126.
1180	Plumptre, A. J., D. Kujirakwinja, A. Treves, I. Owiunji, and H. Rainer. 2007. Transboundary
1181	conservation in the greater Virunga landscape: Its importance for landscape species.
1182	Biological Conservation 134:279-287.
1183	Poiani, K. A., B. D. Richter, M. G. Anderson, and H. E. Richter. 2000. Biodiversity conservation
1184	at multiple scales: functional sites, landscapes, and networks. Bioscience 50:133-146.
1185	Poteete, A., M. Janssen, and E. Ostrom. 2009. Multiple methods in practice: Collective action
1186	and the commons. Princeton University Press.
1187	Poteete, A. R., M. Janssen, and E. Ostrom. 2010. Working together: collective action, the
1188	commons, and multiple methods in practice. Princeton University Press.

1189	Resilience Alliance. 2007a. Assessing and Managing Resilience in Social-Ecological Systems: A
1190	Practitioners Workbook.
1191	Resilience Alliance. 2007b. Assessing Resilience in Social-Ecological Systems: A Scientist's
1192	Workbook. Available on-line at http://www.resalliance.org/3871.php . The Resilience
1193	Alliance.
1194	Resilience Alliance. 2010. Assessing resilience in social-ecological systems: Workbook for
1195	practitioners. Version 2.0. Resilience Alliance, Online:
1196	http://www.resalliance.org/3871.php.
1197	Robbins, P. 2004. Political ecology: a critical introduction. Blackwell Publishers, London, UK.
1198	Rodwell, T. C., N. P. Kriek, R. G. Bengis, I. J. Whyte, P. C. Viljoen, V. de Vos, and W. M.
1199	Boyce. 2001. Prevalence of bovine tuberculosis in African buffalo at Kruger National
1200	Park. Journal of Wildlife Diseases 37 :258-264.
1201	Rogers, D. J. and S. E. Randolph. 1993. Distribution of Tsetse and Ticks in Africa - Past, Present
1202	and Future. Parasitology Today 9:266-271.
1203	Rogers, D. J. and S. E. Randolph. 2000. The global spread of malaria in a future, warmer world.
1204	Science 289 :1763-1766.
1205	Schlager, E. 2007. A comparison of frameworks, theories, and models of policy processes.
1206	Westview Press, Boulder, Colorado.
1207	Scholes, R. J. and R. Biggs. 2004. Ecosystem services in southern Africa: a regional perspective.
1208	Council for Scientific and Industrial Research, Pretoria.
1209	Simon, H. A. 1991. Bounded Rationality and Organizational Learning. Organization Science
1210	2 :125-134.

1211	Sims-Castley, R., G. I. H. Kerley, B. Geach, and J. Langholz. 2005. Socio-economic significance
1212	of ecotourism-based private game reserves in South Africa's Eastern Cape Province.
1213	Parks 15 :6-18.
1214	Sjöstedt, B. 2013. The Role of Multilateral Environmental Agreements in Armed
1215	Conflict: 'Green-keeping' in Virunga Park. Applying the UNESCO World Heritage
1216	Convention in the Armed Conflict of the Democratic Republic of the Congo. Nordic
1217	Journal of International Law 82:129-153.
1218	Slotow, R. and L. T. B. Hunter. 2009. Reintroduction decisions taken at the incorrect social scale
1219	devalue their conservation contribution: the African lion in South Africa Pages 43-71
1220	The reintroduction of top-order predators
1221	Strickland-Munro, J. K., H. E. Allison, and S. A. Moore. 2010. Using resilience concepts to
1222	investigate the impacts of protected area tourism on communities. Annals of Tourism
1223	Research 37 :499-519.
1224	Thompson, J., R. Mathevet, O. Delanoë, M. Cheylan, C. Gil-Fourrier, and M. Bonnin. 2011.
1225	Ecological solidarity as a conceptual tool for rethinking ecological and social
1226	interdependence in conservation policy for protected areas and their surrounding
1227	landscape. Comptes Rendus de l'Académie des Sciences, serie Biologies 334:412-419.
1228	TNC. 2003. The Five-S Framework for Site Conservation: A Practitioner's Handbook for Site
1229	Conservation Planning and Measuring Conservation Success. The Nature Conservancy,
1230	Washington, D.C.
1231	Turner, B. L. I., R. E. Kasperson, P. A. Matson, J. J. McCarthy, R. W. Corell, L. Christensen, N.
1232	Eckley, J. X. Kasperson, A. Luers, M. L. Martello, C. Polsky, A. Pulsipher, and A.

1233	Schiller. 2003. A Framework for Vulnerability Analysis in Sustainability Science.
1234	Proceedings of the National Academy of Science 100:8074-8079.
1235	United Nations. 1992. Convention on Biological Diversity. Page 30 in U. Nations, editor.
1236	Vimal, R., R. Mathevet, and J. D. Thompson. 2012. The Changing Landscape of Ecological
1237	Networks. Journal for Nature Conservation 20:49-55.
1238	Walker, B., L. Gunderson, A. Kinzig, C. Folke, S. Carpenter, and L. Schultz. 2006. A handful of
1239	heuristics and some propositions for understanding resilience in social-ecological
1240	systems. Ecology and Society 11:13.
1241	Walker, B. H., N. Abel, J. M. Anderies, and P. Ryan. 2009. Resilience, adaptability, and
1242	transformability in the Goulburn-Broken Catchment, Australia. Ecology and Society
1243	14:12 [online] URL: http://www.ecologyandsociety.org/vol14/iss11/art12/ .
1244	Waltner-Toews, D., J. J. Kay, and NM. E. Lister. 2008. The Ecosystem Approach: Uncertainty
1245	and Managing for Sustainability. Columbia University Press, New York.
1246	Westley, F., S. R. Carpenter, W. A. Brock, C. S. Holling, and L. H. Gunderson. 2002. Why
1247	systems of people and nature are not just social and ecological systems. Pages 103-119 in
1248	L. H. Gunderson and C. S. Holling, editors. Panarchy: understanding transformations in
1249	human and natural systems. Island Press, Washington, DC.
1250	
1251	
1252	

1253 **Tables**

1254

Table 1: IUCN protected area categories. 1255



IOUN Category	Description
Ia Strict Nature Reserve	Category Ia are strictly protected areas set aside to protect biodiversity and also
	possibly geological/geomorphical features, where human visitation, use and impacts
	are strictly controlled and limited to ensure protection of the conservation values.
	Such protected areas can serve as indispensable reference areas for scientific research
	and monitoring.
Ib Wilderness Area	Category Ib protected areas are usually large unmodified or slightly modified areas,
	retaining their natural character and influence, without permanent or significant
	human habitation, which are protected and managed so as to preserve their natural
	condition.
II National Park	Category II protected areas are large natural or near natural areas set aside to protect
	large-scale ecological processes, along with the complement of species and
	ecosystems characteristic of the area, which also provide a foundation for
	environmentally and culturally compatible, spiritual, scientific, educational,
	recreational, and visitor opportunities.

value: and where safeguarding the integrity of this interaction is vital to protecting and A protected area where the interaction of people and nature over time has produced an cave, or even a living feature such as an ancient grove. They are generally quite small associated cultural values and traditional natural resource management systems. They are generally large, with most of the area in a natural condition, where a proportion is which can be a landform, sea mount, submarine cavern, geological feature, such as area of distinct character with significant ecological, biological, cultural and scenic regular, active interventions to address the requirements of particular species or to Category III protected areas are set aside to protect a specific natural monument, management reflects this priority. Many Category IV protected areas will need Category IV protected areas aim to protect particular species or habitats and Category VI protected areas conserve ecosystems and habitats together with sustaining the area and its associated nature conservation and other values. maintain habitats, but this is not a requirement of the category protected areas and often have high visitor value. VI Protected area with sustainable use of IV Habitat/Species Management Area III Natural Monument or Feature V Protected Landscape/ Seascape natural resources

under sustainable natural resource management and where low-level non-industrial

use of natural resources compatible with nature conservation is seen as one of the

main aims of the area.

(Source: http://www.iucn.org/about/work/programmes/gpap home/gpap quality/gpap pacategories/)

Comment citer ce document :
Cumming, G. S., Allen, C., Ban, N., Biggs, D., Biggs, H. C., Cumming, D. H., de Vos, A.,
Epstein, G., Etienne, M., Maciejewski, K., Mathevet, R., Moore, C., Nenadovic, M., Schoon, M. (2014).
Understanding protected area resilience: a multi-scale socioal-ecological approach.
Ecological Applications. 25 (2), 299-319. DOI: 10.1890/13-2113.1

Figure captions

Figure 1: A summary depiction of Ostrom's SES framework. Different components of

social-ecological systems (characterized as resource units, resource systems, governance

systems, and actors) interact to produce outcomes. Each component is composed of

numerous different elements. Although the framework indicates a role for cross-scale

dynamics, this aspect of it has not been well developed in most applications. We note

also that interactions --> outcomes includes interactions among (i) the ecological

components of the system (e.g., predator-prey dynamics), (ii) the social components of

the system (e.g., rulemaking), and (iii) the social and ecological components of the

system (e.g., harvesting).

Figure 2: A multi-scale perspective of PAs as social-ecological systems, showing the

relationships between the sizes, response times, and persistence times of different system

elements. Note that individual elements in this figure are nested within each other. At

each scale, Ostrom's SES framework captures some of the complexity of interactions

between and across different subsystems.

Figure 3: Protected areas in the Western Cape, South Africa, showing the proximate,

national, and international institutional context of each different area in geographic space.

These different institutional levels interact with ecological and social processes at

different geographic scales, as described in the text.

Figure 4: The depiction by Poiani et al. (2002) of the components of an ecologically functional landscape. Different species have different habitat requirements and if a full range of ecological function is to be retained, habitat conservation must be undertaken in a nested manner, with wide-ranging, regional species having access to high quality patches at local scales. Note that despite its emphasis on functional landscapes, this figure does not directly include people and the scales at which they modify landscapes.

Figure 5: Summary of social-ecological patterns and processes at different scales.

Pattern-process interactions across and between these different scales must be reconciled if effective, sustainable conservation is to occur. In addition, different actors and processes operating at the same scale may interact in important ways. This figure extends the depiction of Poiani et al. (2002) of the ecological components of a functional landscape.


Figure 6: Timeline showing changes in and tenure/land use and wildlife and livestock populations in the GLTFCA between about 1830 and 2010. The 1890 decline in wildlife and livestock was due to the rinderpest pandemic. The early period was characterised by increasing ecological and social fragmentation, followed by TFCA formation and moves to reconnect landscape elements for conservation.

Figure 7: Overview showing examples of issues identified as particularly important in each of the three case studies at different spatial scales in ecological, socio-political, and economic categories respectively. The case studies are indicated on the left of the

diagram (EC, Eastern Cape; CM, Causse Méjan; GLTFCA, Greater Limpopo Transfrontier Conservation Area). Note that (1) this list is not intended to be exhaustive; and (2) many of the issues that are indicated for individual case studies are also relevant to other case studies in the same compartment. For example, tourism & community upliftment are important in all three areas.

Figure 8: Diagrams presenting a dynamic perspective for each case study system: (a) Eastern Cape; (b) Causse Méjan; (c) Greater Limpopo Transfrontier Conservation Area. As indicated in the legend, the different colours for each box represent different kinds of system element (social, economic, and ecological) and arrows indicate interactions and feedbacks within and between scales. These elements are plotted on the notional spatial (on the x-axis) and temporal (on the y-axis) scales at which they exist. The lengths of the boxes are not drawn to scale.

Cross-Scale Social-Ecological Interactions



Cross-Scale Social-Ecological Interactions





Fig 3






Fig 5




Fig 6







Fig	7 Ecological	Scale	Socio-Political	Economic
S			Tourist demand, e.g. for viewing charismatic species	
S		International International institutional	Commodity demand, tourism	
GLTFCA	Migratory species – e.g., songbirds, fruit bats	contexts	Overriding political desire to form peace parks; environmental flow-based legislation in SA & Mocambique; central govt. pressure to promote wildlife economy (SA); disease control regulations to protect western hygiene/production ethic	International Tourist income, marketing expenses
EC	Landscape ecology and ecological carrying capacity	Regional	Tourism & community upliftment	Revenue; costs of developing access infrastructure
Σ	Grassland area & patchiness	Proximate and national institutional	Labeled products	Production system
GLTFCA	Concerns over endangered species	contexts	Catchment management agency imperative; in SA, biosphere institutional arrangements	National tourists (esp. SA and Mocambique, latter still esp. to Kruger Park in SA)
EC	Appropriate species introductions and species abundance	Protected	Tourist satisfaction	Income; investments in facilities and infrastructure
S	% woodland	Area Single-tenure unit	Food production systems	Farming system
GLTFCA	Overall abundance of large herbivores		Location of boundaries, particularly fences	Tourism revenues, infrastructure development, marketing
EC	Converting agricultural fields to natural indigenous state	Patch Sub-tenure unit	Introduction and stocking of charismatic species	Tourist satisfaction
S	Pine seed rain	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Grazing & tree cutting practices	Need for particular resources for stock production
GLTFCA	Conversion from stock farming to wildlife		Water for dignity programme	Water to support basic economic needs. Shared benefits from parks and from mainstreaming biodiversity-friendly practices (e.g., Working for Water)





Size

Fig 8(c) Greater Limpopo Transfrontier Conservation Area



Size