



**HAL**  
open science

# ON NONLOCAL QUASILINEAR EQUATIONS AND THEIR LOCAL LIMITS

Emmanuel Chasseigne, Espen R. Jakobsen

► **To cite this version:**

Emmanuel Chasseigne, Espen R. Jakobsen. ON NONLOCAL QUASILINEAR EQUATIONS AND THEIR LOCAL LIMITS. 2015. hal-01134618v1

**HAL Id: hal-01134618**

**<https://hal.science/hal-01134618v1>**

Preprint submitted on 24 Mar 2015 (v1), last revised 2 Dec 2016 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON NONLOCAL QUASILINEAR EQUATIONS AND THEIR LOCAL LIMITS

EMMANUEL CHASSEIGNE & ESPEN R. JAKOBSEN

ABSTRACT. We introduce a new class of quasilinear nonlocal operators and study equations involving these operators. The operators are degenerate elliptic and may have arbitrary growth in the gradient. Included are new nonlocal versions of  $p$ -Laplace,  $\infty$ -Laplace, mean curvature of graph, and even strongly degenerate operators. Our main results are non-trivial comparison, uniqueness, and existence results for viscosity solutions of linear and fully-linear equations involving these operators. We also identify the conditions under which the nonlocal operators converge to local quasilinear operators, and show that the solutions of the corresponding nonlocal equations converge to the solutions of the local limit equations. Finally, we give a (formal) stochastic representation formula for the solutions and provide many examples.

Loosely speaking our results imply that for “any” quasi-linear 2nd order local operator, “any” well-posed local equation, and “any” non-local Lévy type operator, there is a corresponding Lévy type quasilinear operator and a well-posed non-local equation. Moreover, the solution of any such local equation can be approximated by the solutions of a multitude of different non-local equations.

## 1. INTRODUCTION

In this paper we introduce a new class of gradient dependent Lévy type diffusion operators

$$L[u, Du] = (L_1 + L_2)[u, Du]$$

where

$$L_1[u, Du](x) = \int_{\mathbb{R}^P} u(x + j_1(Du, z)) - u(x) - j_1(Du, z) \cdot Du(x) \, d\mu_1(z), \quad (1.1)$$

$$L_2[u, Du](x) = \int_{\mathbb{R}^P} u(x + j_2(Du, z)) - u(x) \, d\mu_2(z), \quad (1.2)$$

and  $\mu_1, \mu_2$  are non-negative Lévy measures. In these operators the strength and direction of the non-local diffusion depend on the gradient, and hence as we explain below, they are natural generalizations of local (non-divergence form) quasi-linear operators of the form

$$L_0(Du, D^2u) = \frac{1}{2} \text{tr}(\sigma(Du)\sigma(Du)^T D^2u) + b(Du)Du.$$

The operators are allowed to degenerate ( $j_1 = 0$  or  $j_2 = 0$  in some set) and have arbitrary growth in the gradient, so  $\infty$ -Laplace,  $p$ -Laplace, and strongly degenerate operators are included.

---

*Date:* March 24, 2015.

*2010 Mathematics Subject Classification.* 35R09 35D40, 35J60, 47G20.

*Key words and phrases.* Nonlocal elliptic equation, quasilinear nonlocal operators, viscosity solutions, Lévy processes.

E.C. is partially supported by Spanish Project MTM2011-25287, and E.R.J. is partially supported by the NFR project Discrete Models in Mathematical Analysis (DIMMA).

We want to study equations involving the operator  $L$ , and to simplify and focus on the new issues, we study here the following special problem:

$$F(u, Du, L[u, Du]) = f(x) \quad \text{in} \quad \mathbb{R}^N, \quad (1.3)$$

where we assume  $F$  to be (degenerate) elliptic and strictly increasing in  $u$  ( $D_u F > 0$ ). But for this equation, we make an effort to push for very general results. First we obtain comparison, uniqueness, stability, and existence results for bounded solutions of (1.3), results which are highly non-trivial due to the implicit nature of our operators and our weak integrability assumptions. Then we identify the limit problems where non-local operators converge to local ones,

$$L_\varepsilon[u, Du] \rightarrow L_0(Du, D^2u) \quad \text{as} \quad \varepsilon \rightarrow 0,$$

and prove that the solutions  $u_\varepsilon$  of the corresponding non-local equations

$$F(u_\varepsilon, Du_\varepsilon, L_\varepsilon[u_\varepsilon, Du_\varepsilon]) = f(x) \quad \text{in} \quad \mathbb{R}^N, \quad (1.4)$$

converges locally uniformly to the solution of the local equation

$$F(u, Du, L_0(Du, D^2u)) = f(x) \quad \text{in} \quad \mathbb{R}^N. \quad (1.5)$$

We refer to Section 2 for the precise assumptions and results. Here we just remark that (i) the weak solution concept we use is bounded viscosity solutions, (ii) generators  $L$  of every pure jump Lévy processes are included as linear special cases, and (iii) a typical special case of (1.3) satisfying our assumptions is the quasi-linear equation

$$-L[u, Du](x) + u(x) = f(x) \quad \text{in} \quad \mathbb{R}^N, \quad (1.6)$$

with bounded uniformly continuous  $f$ .

Let us illustrate our results on  $\infty$ -Laplace type operators. In the local case (see e.g. [23]) this is a diffusion operator with “diffusion” (Brownian motion, generator  $(-\Delta)$ ) only in the gradient direction:

$$\Delta_\infty u(x) = \text{tr}[Du(x)Du(x)^T D^2u(x)] = (Du(x) \cdot D)^2 u(x). \quad (1.7)$$

Natural non-local generalizations are operators with e.g.  $\alpha$ -stable diffusion ( $\alpha \in (0, 2)$ ) along the gradient direction. The generator of the symmetric  $\alpha$ -stable process is the fractional Laplacian [3],

$$-(-\Delta)^{\alpha/2} u(x) = \int_{\mathbb{R}^N} u(x+z) - u(x) - (z \cdot Du(x)) \mathbb{1}_{|z|<1} \frac{c_\alpha dz}{|z|^{N+\alpha}},$$

and hence the corresponding non-local version of the  $\infty$ -Laplace operator would take the form

$$\mathcal{L}_{\Delta_\infty}^{\alpha/2}[u](x) = \int_{\mathbb{R}^1} u(x + Du(x)z) - u(x) - Du(x) \cdot Du(x)z \mathbb{1}_{|z|<1} \frac{c_\alpha dz}{|z|^{1+\alpha}}. \quad (1.8)$$

This operator is in the form  $L$  with  $j_1 = Du \cdot z = j_2$ ,  $\mu_1 = \mathbb{1}_{|z|<1}\mu$ , and  $\mu_2 = \mathbb{1}_{|z|\geq 1}\mu$ , where  $d\mu = \frac{c_\alpha dz}{|z|^{1+\alpha}}$ . By our results,  $L = \mathcal{L}_{\Delta_\infty}^{\alpha/2}$  gives rise to well-posed equations (1.3), and since

$$\mathcal{L}_{\Delta_\infty}^{\alpha/2}[u](x) \rightarrow \Delta_\infty u(x) \quad \text{as} \quad \alpha \rightarrow 2^-$$

for smooth bounded  $u$ , it also follows that (possibly non-smooth viscosity) solutions of (1.4) with  $L_\varepsilon = \mathcal{L}_{\Delta_\infty}^{1-\varepsilon}$  will converge as  $\varepsilon \rightarrow 0$  to the solution of (1.5) with  $L_0 = \Delta_\infty$ .

*A similar construction can be carried out for “any” local (non-divergence form) quasi-linear operator and “any” Lévy diffusion, thereby producing a corresponding quasi-linear Lévy diffusion.* Under our assumptions this new operator is well-posed, and can approximate the original local operator. This will be explained in Remark 2.7. In Section 3 we present a (formal) stochastic interpretation

of our equations and give many more examples. Included are several non-local versions of the  $\infty$ -Laplace, the  $p$ -Laplace, and the mean curvature of graph operators; versions that are modulations of singular integral operators and others based on bounded non-local operators. It is interesting to note that the limit operator  $L_0$  will include also a drift term ( $b \neq 0$ ) whenever the measures  $\mu_{2,\varepsilon}$  in the  $L_2$ -term has a non-zero mean value near  $z = 0$ , see assumption  $(\mathbf{M}_\varepsilon)$  in section 2.2. The reason is that in  $L_2$  this mean is not compensated by a first-order gradient term as in  $L_1$ .

The litterature on non-local quasi-linear equations seems to be rather recent. In the calculus of variations, such equations can be obtained as Euler-Lagrange equations by minimizing fractional Sobolev norms ( $W^{p,\frac{\alpha}{2}}$ -norms) [22, 15, 21] or truncated versions of such norms [2]. In the first three papers, (variational) fractional  $p$  and  $\infty$ -Laplace operators are introduced. In [14], a different “variational” type of non-local operators is studied by non-variational viscosity solution techniques. Non-variational non-local  $\infty$ -Laplace type operators are introduced in [9, 8], and shown in [9] to be connected to a sequence of Tug of War games. But none of these operators have an implicit form as our operators do. Our operators are not variational, and among existing work they resemble most closely the operators of [9, 8], especially [9]. However, whereas the operators in [9, 8] have bounded dependence on the gradient but are discontinuous where it is zero, our operators are continuous but may have arbitrary growth in the gradient. The operators in [9, 8] correspond to normalized  $\infty$ -Laplacians, which in the local case take the form (see e.g. [24, 23])

$$\frac{1}{|Du(x)|^2} \Delta_\infty u(x) = \left( \frac{Du(x)}{|Du(x)|} \cdot D \right)^2 u(x),$$

while our version (1.8) corresponds to an unnormalized one (i.e. to  $\Delta_\infty u$ ).

In this paper we work with viscosity solutions. This weak solution concept is not distributional and does not involve integration. It is very well adapted to the implicit and degenerate form of our equations. The solution concept was introduced by Crandall and Lions in the early 1980s to get uniqueness of solutions of first order Hamilton-Jacobi equations. Later it has been extended to wide range of problems, including many non-local ones. The standard reference for local problems is [13]. For non-local problems, we only refer to [7, 18] for the basic well-posedness theory for problems posed in the whole space. But we mention that there is a large literature on regularity and properties of solutions, asymptotic problems, boundary conditions, approximations and numerics, relation to stochastic processes, applications etc.. The problems we consider here represent a natural class of non-local quasi-linear equations where the viscosity solution techniques still apply and give comparison and uniqueness.

In fact we have optimized the assumptions to allow for very general dependence on the gradients in  $L$  and  $F$  at the cost of no dependence on the variable  $x$ ! We have also made an effort to optimize the assumption on  $j_i$  and  $\mu_i$ . In both cases our assumptions are much more general than in [7, 18]. In the doubling of variables argument of the comparison proof, these differences to [7, 18] are e.g. reflected in a different choice of test function and two of the limits being taken in the reverse order. Existence, on the other hand, does not follow from clever modifications of commonly used arguments. Because of the implicit form of the equations, with the gradient dependence in  $j_1$  and  $j_2$ , compactness arguments do not work and it seems not possible to adapt Perron’s method either. Instead we propose a new non-trivial argument based on a so-called *Sirtaki method* inspired by [5]. The argument involves several regularization and approximation arguments, a Schauder fixed point argument, and several limit problems. In each limit problem, we show that a limit exists and solves the relevant equation by the half relaxed limit method combined with strong comparison results for the limit equations.

At the cost of longer and more technical arguments, one can of course relax the assumptions of this paper. For example we could consider less degenerate equations without the assumption that  $D_u F > 0$ , like uniformly elliptic or even  $p$  and  $\infty$ -Laplace equations. We could also have considered equations on domains with boundary conditions. In section 6, we give the extension to the parabolic case (Cauchy problems). Finally, we mention that in the upcoming paper [1], we will study the local limits of non-local equations under assumptions that are optimized w.r.t. the  $x$ -dependence. In this case we also give explicit convergence rates.

**Outline.** We present the main results in Section 2 and give several examples and a stochastic interpretation in Section 3. Then, precise definitions of viscosity solutions appear in Section 4 and the proofs of the comparison, existence and concentration results are given in Section 5. In Section 6 we extend our results to parabolic problems, and in the appendix at the end of the paper, we give the proofs of some technical results we need.

**Notation.** The notation  $UC(\mathbb{R}^N)$  denotes the set of uniformly continuous functions defined on  $\mathbb{R}^N$  and  $BUC(\mathbb{R}^N)$  is the space of bounded, uniformly continuous functions;  $usc$  [resp.  $lsc$ ] stands for upper semicontinuous [resp. lower semicontinuous]; the spaces  $C^1/C^2$  are the spaces of functions having continuous first-order / second-order derivatives;  $C^{0,\alpha}, C^{1,\alpha}$  stand for the usual Hölder spaces;  $C_b$  denotes the space of continuous, bounded functions;  $\limsup^*$  and  $\liminf_*$  are the half-relaxed limits (more precise definitions in the text where they are used); we denote by  $\mathbb{1}_A$  the indicator function of the set  $A$ ; a modulus of continuity is a subadditive function  $\omega : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that  $\lim_{s \rightarrow 0^+} \omega(s) = 0$ ; the notation  $a \wedge b$  stands for the min of  $a$  and  $b$ ,  $a \vee b$  is for the max and  $s^+ = \max(s, 0)$ . Note that in this paper  $x \in \mathbb{R}^N$  for  $N \geq 1$  while  $z \in \mathbb{R}^P$  for  $P \geq 1$ ; finally,  $\mathbb{R}^{P \times Q}$  denotes the space of matrices with  $P$  rows and  $Q$  columns.

## 2. THE MAIN RESULTS

The results of this section essentially implies that for “any” quasi-linear 2nd order local operator  $L_0$ , “any” well-posed local equation (1.5), and “any” non-local Lévy type operator, there is a corresponding Lévy type quasilinear operator  $L$  and a *well-posed* non-local equation (1.3). Moreover, the solution of any such local equation can be *approximated locally uniformly* by the solutions of a multitude of different non-local equations.

**2.1. Comparison, uniqueness, and existence.** Let us first list the assumptions under which we construct a general existence and uniqueness theory for (1.3):

(M)  $\mu_1$  and  $\mu_2$  are non-negative Radon measures on  $\mathbb{R}^P \setminus \{0\}$  satisfying

$$\int_{|z|>0} |z|^2 d\mu_1(z) + \int_{|z|>0} d\mu_2(z) < \infty.$$

(J1)  $j_1(p, z)$  and  $j_2(p, z)$  are Borel measurable functions from  $\mathbb{R}^N \times \mathbb{R}^P$  into  $\mathbb{R}^N$ , continuous in  $p$  for a.a.  $z \in \mathbb{R}^P$ , and for any  $r > 0$  there is a  $C_{j,r} > 0$  such that for all  $|p| < r$ ,

$$\int_{|z|>0} |j_1(p, z)|^2 d\mu_1(z) \leq C_{j,r}.$$

(J2) For any  $r > 0$ , there is a modulus of continuity  $\omega_{j,r}$  such that for all  $|p|, |q| < r$ ,

$$\int_{|z|>0} |j_1(p, z) - j_1(q, z)|^2 d\mu_1(z) \leq \omega_{j,r}(p - q).$$

**(J3)** There exists  $\delta_0 > 0$  such that for any  $r > 0$  and  $\varepsilon > 0$  there exists  $\eta > 0$  such that

$$\sup_{|p| < r} \int_A |j_1(p, z)|^2 d\mu_1(z) < \varepsilon$$

for every Borel set  $A \subset \{0 < |z| < \delta_0\}$  such that  $\int_A |z|^2 \mu_1(dz) < \eta$ .

**(F1)**  $F$  is continuous, and for any  $u, p$ , and  $\ell \leq \ell'$ ,  $F(u, p, \ell) \geq F(u, p, \ell')$ .

**(F2)** For any  $M > 0$ , there exist  $\gamma_M > 0$  such that for all  $p \in \mathbb{R}^N$ ,  $|u|, |v| \leq M$ , and  $l \in \mathbb{R}$ ,

$$F(u, p, l) - F(v, p, l) \geq \gamma_M(u - v)^+.$$

**(F3)** For any  $M, r > 0$ , there exists a modulus of continuity  $\omega_{M,r}$  such that for any  $|u| \leq M$  and  $|p|, |q|, |\ell|, |\ell'| \leq r$ ,

$$\left| F(u, p, \ell) - F(u, q, \ell') \right| \leq \omega_{M,r}(|p - q| + |\ell' - \ell|).$$

**(F4)**  $f \in UC(\mathbb{R}^N)$ .

**(F5)**  $f \in BUC(\mathbb{R}^N)$  and all quantities in **(F2)**, **(F3)** are independent of  $M$ .

We give now the precise results and refer to Section 5 for the proofs.

**Theorem 2.1.** (Comparison results)

(a) [Quasi-linear case] Assume **(M)**, **(J1)**–**(J2)**, and **(F4)**. If  $u : \mathbb{R}^N \rightarrow \mathbb{R}$  is a bounded usc subsolution of (1.6) and  $v : \mathbb{R}^N \rightarrow \mathbb{R}$  is a bounded lsc supersolution of (1.6), then  $u \leq v$  in  $\mathbb{R}^N$ .

(b) [Fully non-linear case] Assume **(M)**, **(J1)**–**(J2)**, **(F1)**–**(F4)** hold. If  $u : \mathbb{R}^N \rightarrow \mathbb{R}$  is a bounded usc viscosity subsolution of (1.3) and  $v : \mathbb{R}^N \rightarrow \mathbb{R}$  is a bounded lsc viscosity supersolution of (1.3), then  $u \leq v$  in  $\mathbb{R}^N$ .

We have the following immediate consequences of this comparison result.

**Corollary 2.2.** Under the assumptions of Theorem 2.1:

(a) [Uniqueness] There is a most one solution  $u \in C_b(\mathbb{R}^N)$  of (1.3) (respectively of (1.6)).

(b) [Uniform continuity] Any solution  $u \in C_b(\mathbb{R}^N)$  of (1.3) (respectively of (1.6)) belongs to  $BUC(\mathbb{R}^N)$  and

$$\gamma_M \omega_u(h) \leq \omega_f(h) \quad (\text{respectively } \omega_u(h) \leq \omega_f(h)),$$

where  $M = \|u\|_\infty$  and  $\omega_\phi(r) = \sup_{x \in \mathbb{R}^N, |y| < r} |\phi(x+y) - \phi(x)|$  denotes the modulus of continuity of  $\phi(x)$ .

(c) [ $L^\infty$ -bound] If also **(F5)** holds with  $\gamma_M = \gamma$  (independent of  $M$ ), then any solution  $u \in C_b(\mathbb{R}^N)$  of (1.3) (respectively of (1.6)) satisfies

$$\gamma \|u\|_\infty \leq \|f\|_\infty \quad (\text{respectively } \|u\|_\infty \leq \|f\|_\infty).$$

*Proof.* (a) is immediate from Theorem 2.1, while (c) follows since  $\pm \frac{1}{\gamma} \|f\|_\infty$  are super and subsolutions of (1.3). To prove (b), note that  $v_\pm(x) = u(x+h) \pm \frac{1}{\gamma_M} \omega_f(|h|)$  is a super and subsolution of (1.3). By Theorem 2.1,  $v_-(x) \leq u(x) \leq v_+(x)$ , and hence  $|u(x) - u(x+h)| \leq \frac{1}{\gamma_M} \omega_f(|h|)$ .  $\square$

**Theorem 2.3** (Existence). Under the assumptions of Theorem 2.1, **(J3)**, and **(F5)**, there exists a unique bounded viscosity solution  $u \in BUC(\mathbb{R}^N)$  of (1.3) (respectively of (1.6)).

Let us now briefly comment on the assumptions.

*Remark 2.4.* (i)  $\mu_1$  and  $\mu_2$  are Lévy measures [3] by **(M)**. Conversely, any Lévy measure  $\mu$  can be written as  $\mu_1 + \mu_2$  for  $\mu_1$  and  $\mu_2$  satisfying **(M)**:

$$\mu = \mu \mathbb{1}_{|z| < 1} + \mu \mathbb{1}_{|z| \geq 1} =: \mu_1 + \mu_2.$$

(ii) Assumptions on  $j$  are optimized w.r.t. the dependence in  $p$  at the cost of no dependence on  $x$ ! A typical example is

$$j_i(p, z) = j(p)z,$$

where  $z \in \mathbb{R}^P$  and  $j : \mathbb{R}^N \rightarrow \mathbb{R}^{N \times P}$  only needs to be continuous. **(J1)** and **(J3)** follow from the stronger assumption  $|j_i(p, z)| \leq c|z|$  for  $z$  near 0. **(J3)** implies that  $\left\{ \frac{|j_i(p, z)|^2}{|z|^2} \right\}_{|p| < r}$  is  $|z|^2 \mu_1(dz)$  equi-integrable on  $\{0 < |z| < \delta\}$  for any  $\delta \leq \delta_0$ , cf. Appendix A. We need it to construct solutions under our general assumptions but not for comparison.

(iii) By **(M)**, **(J1)**, and a Taylor expansion,  $L[\phi, D\phi](x)$  is well-defined for any  $\phi \in C^2(\mathbb{R}^N) \cap C_b(\mathbb{R}^N)$ .

(iv) **(F2)** implies degenerate ellipticity and strict monotonicity in  $u$ , while **(F3)** allows for very general  $p$ -dependence at the cost no  $x$ -dependence. Compare **(F3)** to e.g. assumption (3.14) in [13].

(v) The assumptions on integrability and  $p$ -dependence of  $j$  and the  $(p, l)$ -dependence of  $F$  of this paper are much more general than e.g. in [18, 7].

**2.2. Local limits.** We also study the convergence of solutions of the nonlocal equation (1.4) to the local equation (1.5), including separate results for the quasi-linear case where (1.4) and (1.5) take the simpler forms

$$-L_\varepsilon[u_\varepsilon, Du_\varepsilon](x) + u_\varepsilon(x) = f(x) \quad \text{in } \mathbb{R}^N, \quad (2.1)$$

$$-L_0(Du, D^2u) + u(x) = f(x) \quad \text{in } \mathbb{R}^N, \quad (2.2)$$

where the local operator  $L_0$  is precisely defined in Definition 2.5 below. Concerning (2.1), we use the decomposition  $L_\varepsilon = L_{1,\varepsilon} + L_{2,\varepsilon}$  with

$$L_{i,\varepsilon}[\phi, D\phi](x) = \int_{\mathbb{R}^N} \phi(x + j_i(D\phi(x), z)) - \phi(x) - \delta_{i,1} j_i(D\phi(x), z) \cdot D\phi(x) \, d\mu_{i,\varepsilon}, \quad i = 1, 2,$$

where  $\delta_{i,1} = 1$  if  $i = 1$  and 0 otherwise. In order to prove the convergence result as  $\varepsilon \rightarrow 0$  we need the following additional assumptions:

**(M $_\varepsilon$ )**  $\mu_\varepsilon = (\mu_{1,\varepsilon}, \mu_{2,\varepsilon})$  satisfies **(M)** for every  $\varepsilon > 0$ , and there exists  $A_1, A_2 \in \mathbb{R}^{P \times N}$  and  $a \in \mathbb{R}^P$  such that for every  $Y \in \mathbb{R}^{P \times P}$ ,  $q \in \mathbb{R}^P$ , and  $\delta > 0$ , as  $\varepsilon \rightarrow 0$

$$\begin{aligned} \int_{|z| < \delta} z^T Y z \, d\mu_{1,\varepsilon} &\rightarrow \text{tr}[A_1^T Y A_1], \\ \int_{|z| < \delta} (z^T Y z + q \cdot z) \, d\mu_{2,\varepsilon} &\rightarrow \text{tr}[A_2^T Y A_2] + a \cdot q, \\ \int_{|z| > \delta} (d\mu_{1,\varepsilon} + d\mu_{2,\varepsilon}) &\rightarrow 0. \end{aligned}$$

**(J4)** For  $i = 1, 2$ , the function  $(p, z) \mapsto j_i(p, z)$  is continuous,  $z$ -differentiable at  $z = 0$  locally uniformly in  $p$ ,  $j_i(p, 0) = 0$ , and the function  $\sigma_i : \mathbb{R}^N \rightarrow \mathbb{R}^{N \times P}$ , defined by

$$\sigma_i(p) := D_z j_i(p, 0), \quad \text{is continuous.}$$

**Definition 2.5.** For any vector  $p \in \mathbb{R}^N$  and matrix  $X \in \mathbb{R}^{N \times N}$ , we define:

$$L_0(p, X) := \frac{1}{2} \text{tr}[\tilde{\sigma}_1(p) \tilde{\sigma}_1(p)^T X] + \frac{1}{2} \text{tr}[\tilde{\sigma}_2(p) \tilde{\sigma}_2(p)^T X] + b(p) \cdot p,$$

where  $\tilde{\sigma}_i(p) := \sigma_i(p)A_i$ ,  $i = 1, 2$ ,  $b(p) := a^T \sigma_2(p)$ , for  $A_1, A_2, a, \sigma_1, \sigma_2$  given by  $(\mathbf{M}_\varepsilon)$  and  $(\mathbf{J4})$ .

The limit result is the following:

**Theorem 2.6.** (Local limits) *Let  $L_0$  be given by Definition 2.5.*

(a) [Quasi-linear case] *Assume  $(\mathbf{M}_\varepsilon)$ ,  $(\mathbf{J1})$ – $(\mathbf{J4})$  and  $(\mathbf{F5})$ . Then any sequence of solutions  $u_\varepsilon$  of (2.1) converges locally uniformly as  $\varepsilon \rightarrow 0$  to the solution  $u$  of (2.2).*

(b) [Fully nonlinear case] *Assume  $(\mathbf{M}_\varepsilon)$ ,  $(\mathbf{J1})$ – $(\mathbf{J4})$ , and  $(\mathbf{F1})$ – $(\mathbf{F5})$ . Then any sequence of solutions  $u_\varepsilon$  of (1.4) converges locally uniformly as  $\varepsilon \rightarrow 0$  to the solution  $u$  of (1.5).*

*Remark 2.7.* (i)  $(\mathbf{M}_\varepsilon)$  is a concentration assumption implying e.g.  $z^T Y z \mu_{1,\varepsilon}(dz) \rightarrow \text{tr}[A_1^T Y A_1] \delta_0$  in measure. This is a convergence result for measures in  $\mathbb{R}^P$  and not in  $\mathbb{R}^N$ . Note that  $a$  plays a role only for the  $L_2$ -part of  $L$ . Illustrative examples are the following singular and truncated  $(2 - \varepsilon)$ -stable like Lévy measures:

$$\begin{aligned} \mu_{1,\varepsilon}(dz) &= \varepsilon \frac{g(z)}{|z|^{N+2-\varepsilon}} \mathbb{1}_{|z|<1} dz & \text{where} & \quad \lim_{z \rightarrow 0} g(z) = g(0) \neq 0, \\ \mu_{2,\varepsilon}(dz) &= \varepsilon \frac{g(z)}{|z|^{N+2-\varepsilon}} \mathbb{1}_{\varepsilon < |z| < 1} dz & \text{where} & \quad g \text{ is } C^1 \text{ at } z = 0 \text{ and } g(0) \neq 0. \end{aligned}$$

Both satisfy  $(\mathbf{M}_\varepsilon)$ :  $\mu_{1,\varepsilon}$  with  $A_1 = g(0)I$ ,  $A_2 = 0$ ,  $a = 0$  and  $\mu_{2,\varepsilon}$  with  $A_1 = 0$ ,  $A_2 = g(0)I$ ,  $a = Dg(0)$ .

(ii) By  $(\mathbf{J4})$  and Definition 2.5,

$$L_0(p, X) = \text{tr}[A_1 Y_1 A_1] + \text{tr}[A_2 Y_2 A_2] + a \cdot q$$

for  $Y_i = \sigma_i(p)^T X \sigma_i(p) \in \mathbb{R}^{P \times P}$ ,  $i = 1, 2$ , and  $q = \sigma_2(p)p \in \mathbb{R}^P$ .

(ii) If  $(\mathbf{F5})$  and  $(\mathbf{J4})$  (and  $(\mathbf{F1})$ – $(\mathbf{F3})$ ) hold, there exists a unique viscosity solution of (2.2) (and of (1.5)) satisfying the strong comparison principle, cf. Theorem 5.1 in [13] and Lemma 5.14 below.

(iii) We may specify (“any”)  $\sigma(p)$  first, and then for every Lévy measure  $(\mu_{1,\varepsilon}, \mu_{2,\varepsilon})$  satisfying the concentration assumption  $(\mathbf{M}_\varepsilon)$ , we get a non-local approximation  $L_\varepsilon$  of the local operator  $L_0$ . Moreover, the corresponding equations, (1.4) and (1.3), are well-posed with solutions that converge to one another under very general assumptions.

### 3. STOCHASTIC INTERPRETATION AND EXAMPLES

**3.1. Stochastic interpretation.** Formally equation (1.3) is always the Dynamic Programming Equation of an implicitly defined stochastic control problem or game. E.g. the solution  $u$  of (1.6) satisfies formally

$$u(x) = \mathbb{E}^x \left( \int_0^\infty e^{-t} f(X_t) dt \right) \quad (3.1)$$

where  $X_t$  is a pure jump Lévy-Ito process satisfying

$$X_t = x + \int_0^t \int_{|z|>0} j_1(Du(X_{s-}), z) \tilde{N}_1(dz, ds) + \int_0^t \int_{|z|>0} j_2(Du(X_{s-}), z) N_2(dz, ds), \quad (3.2)$$

where  $\tilde{N}_1$  is a compensated Poisson random measure,  $N_2$  is a finite intensity Poisson random measure, and  $\mathbb{E}^x$  is the expectation w.r.t. the law of  $X$  (which starts at  $x$ ). By a Lévy-Ito process we mean a *Lévy type stochastic integral* defined in Chapter 4.3.3 in [3], and we refer to e.g. [3, 12] for definitions



of the other probabilistic terms mentioned above. Formally, the generator  $A$  of  $X_t$  is given by the formula

$$Au(x) = L[u, Du](x)$$

for  $u$  in the domain of  $A$  (equation (6.36) in [3]). Moreover,  $X_t$  generates a semigroup  $T_t$  defined by  $T_t\phi(y) = \mathbb{E}^y\phi(X_t^y)$  with the convention that  $X_0^y = y$  almost surely, and  $u$  is then the 1-resolvent  $R_1$  (chapter 3 in [3]) of this the semi-group applied to  $f$ , *i.e.*  $R_1f(x) = 1u(x)$ . By the resolvent identity,

$$u - Au = (I - A)R_1f(x) = f(x) \quad \text{in} \quad \mathbb{R}^N,$$

*i.e.*  $u$  satisfies equation (1.6) at least formally. To make this discussion rigorous, we need the assumptions of section 2 and some additional ones including smoothness of  $u$ . Following chapter 6.7 in [3], it suffices to assume in addition that Assumptions 6.6.1 and 6.7.1 of [3] hold. We do not state them here, we only remark that they are satisfied if *e.g.*

$$j_1(p, z) = j(p)z \text{ with } j \in W_{\text{loc}}^{1, \infty}, \quad 0 \leq \mu(dz) \leq \frac{C dz}{|z|^{N+\alpha}} \text{ with } \alpha \in (0, 2), \quad u \in C_0^2 \text{ and } f \in C_0.$$

Note that then  $Du$  is bounded and Lipschitz. In this case it follows from Theorem 6.7.4 of [3] that  $T_t$  is a Feller semi-group with generator  $A$  as above and that  $u$  is in the domain of  $A$ . By Theorem 3.2.9 of [3] the resolvent  $R_1$  exists and satisfies the resolvent identity above for any  $f \in C_0(\mathbb{R}^N)$ .

We have the following result:

**Proposition 3.1.** *If  $u$  and  $X_t$  satisfy the assumptions mentioned above and (3.1) and (3.2) hold (in the strong sense), then  $u$  is a classical solution of (1.6).*

**3.2. Examples.** We introduce now some classes of quasi-linear non-local operators with special focus on operators of  $p$ -Laplacian,  $\infty$ -Laplacian, and mean curvature of graph type. Recall the definitions of the local and fractional  $\infty$ -Laplacian in (1.7) and (1.8). To define other non-local operators we need the following Lemma.

**Lemma 3.2.** *Let  $p \geq 1$ ,  $r_p = -1 + \sqrt{p-1}$  and  $I$  be  $N \times N$  identity matrix.*

$$(a) \quad I + (p-2) \frac{\xi \otimes \xi}{|\xi|^2} = a_p(\xi) a_p^T(\xi) \quad \text{where} \quad a_p(\xi) := I + r_p \frac{\xi \otimes \xi}{|\xi|^2}.$$

$$(b) \quad I - \frac{\xi \otimes \xi}{1 + |\xi|^2} = \tilde{a}(\xi) \tilde{a}^T(\xi) \quad \text{where} \quad \tilde{a}(\xi) := I - \frac{\xi \otimes \xi}{|\xi|^2} \left( 1 - \frac{1}{\sqrt{1 + |\xi|^2}} \right).$$

$$(c) \quad \text{The functions } \tilde{a}(\xi) \text{ and } |\xi|^{\frac{p-2}{2}} a_p(\xi), \quad p \geq 2, \text{ are continuous in } \mathbb{R}^N.$$

The proof is straightforward, using that  $r_p^2 + r_p = p - 2$ . In view of the lemma,

$$\begin{aligned} \Delta_p u(x) &= \operatorname{div} \left( |Du(x)|^{p-2} Du(x) \right) = |Du(x)|^{p-2} \left( \Delta u(x) + (p-2) \frac{\Delta_\infty u(x)}{|Du(x)|^2} \right) \\ &= \operatorname{tr} [ |Du(x)|^{p-2} D^2 u(x) ] + (p-2) \operatorname{tr} [ |Du(x)|^{p-4} Du(x) Du(x)^T D^2 u(x) ] \\ &= \operatorname{tr} [ \sigma_p(Du(x)) \sigma_p^T(Du(x)) D^2 u(x) ] \quad \text{where} \quad \sigma_p(\xi) = |\xi|^{\frac{p-2}{2}} a_p(\xi), \end{aligned}$$

$$\begin{aligned} H[u](x) &= \operatorname{div} \left( \frac{Du(x)}{\sqrt{1 + |Du(x)|^2}} \right) = \frac{1}{\sqrt{1 + |Du(x)|^2}} \left( \Delta u(x) - \frac{\Delta_\infty u(x)}{1 + |Du(x)|^2} \right) \\ &= \operatorname{tr} \left( \tilde{\sigma}(Du(x)) \tilde{\sigma}^T(Du(x)) D^2 u(x) \right) \quad \text{where} \quad \tilde{\sigma}(\xi) = \frac{\tilde{a}(\xi)}{(1 + |\xi|^2)^{\frac{1}{4}}}, \end{aligned}$$

where  $\tilde{\sigma}$  and  $\sigma_p$  are continuous for  $p \geq 2$ .

*First type of examples:* Quasilinear versions of every generator of pure jump Lévy processes [3]. E.g. non-local fractional Laplace type operators,

$$\begin{aligned} \mathcal{L}_{\Delta_\infty}^{\alpha/2}[u](x) & \text{ already defined in (1.8),} \\ \mathcal{L}_{\Delta_p}^{\alpha/2}[u](x) &= \int_{\mathbb{R}^N} u(x + \sigma_p(Du(x))z) - u(x) - \sigma_p(Du(x))z \cdot Du(x) \mathbb{1}_{|z|<1} \frac{c_\alpha dz}{|z|^{N+\alpha}}, \\ \tilde{\mathcal{L}}_{\Delta_p}^{\alpha/2}[u](x) &= \int_{\mathbb{R}^N} u(x + |Du(x)|^{\frac{p-2}{2}}z) - u(x) - |Du(x)|^{\frac{p-2}{2}}z \cdot Du(x) \mathbb{1}_{|z|<1} \frac{c_\alpha dz}{|z|^{N+\alpha}} \\ & \quad + (p-2) \int_{\mathbb{R}^1} u\left(x + |Du(x)|^{\frac{p-4}{2}}Du(x)z\right) - u(x) - |Du(x)|^{\frac{p-4}{2}}Du(x)z \cdot Du(x) \mathbb{1}_{|z|<1} \frac{c_\alpha dz}{|z|^{1+\alpha}}, \\ \mathcal{L}_H^{\alpha/2}[u](x) &= \int_{\mathbb{R}^N} u(x + \tilde{\sigma}(Du(x))z) - u(x) - \tilde{\sigma}(Du(x))z \cdot Du(x) \mathbb{1}_{|z|<1} \frac{c_\alpha dz}{|z|^{N+\alpha}} \end{aligned}$$

where  $p \geq 2$  and  $c_\alpha = O(2 - \alpha)$  is the constant of  $\Delta^{\alpha/2}$ . The fractional Laplacian is the generator of the symmetric  $\alpha$ -stable process, and the above non-local versions can be seen “generators” of gradient dependent modulations of this process. To be more precise,  $\mathcal{L}_{\Delta_\infty}^{\alpha/2}$  is a nonlocal version of the infinite Laplacian operator; both  $\mathcal{L}_{\Delta_p}^{\alpha/2}$  and  $\tilde{\mathcal{L}}_{\Delta_p}^{\alpha/2}$  are nonlocal versions of the  $p$ -Laplace operator, depending on how we write it; finally  $\mathcal{L}_H^{\alpha/2}$  is a nonlocal version of the curvature operator  $H$ . Note that these operators are of the form  $L = L_1 + L_2$  where both  $L_1 \neq 0$  ( $|z| < 1$ ) and  $L_2 \neq 0$  ( $|z| > 1$ ).

*Second type of examples:* Quasilinear versions of the generators of some Lévy-Ito jump-processes defined by stochastic differential equations (SDEs) driven by pure jump Lévy processes [3, 12]. An example is the operator from the CGMY model for the price of a European option in Finance [12],

$$\mathcal{L}u(x) = \int_{\mathbb{R}^1} u(x+z) - u(x) - Du(x)(e^z - 1) \frac{C e^{-Mz^+ - Gz^-} dz}{|z|^{1+Y}},$$

for  $C, G, M > 0$ ,  $Y \in (0, 2)$ , and the following new non-local infinity Laplacian (compare to  $\mathcal{L}_{\Delta_\infty}^{\alpha/2}$ ):

$$\mathcal{J}_{\Delta_\infty}^{\alpha/2}[u](x) = \int_{\mathbb{R}^1} u(x + Du(x)z) - u(x) - Du(x) \cdot Du(x)(e^z - 1) \frac{c_\alpha e^{-Mz^+ - Gz^-} dz}{|z|^{1+\alpha}}.$$

In this case  $L_1 \neq 0$  and  $L_2 = 0$ , and  $L = \mathcal{J}_{\Delta_\infty}^{\alpha/2}$  is a gradient dependent modulation of  $\mathcal{L}$ . Here  $\mathcal{L}$  is not the generator of a Lévy process, but the exponential of a Lévy process [12] (after a transformation). The driving (Lévy) process here is a tempered  $\alpha$ -stable process [12]. Other quasilinear versions ( $p$ -Laplace etc.) can be easily be constructed as above.

*Remark 3.3.* Since we do not allow for  $x$ -dependence in  $j_1$  and  $j_2$  at the level of the PDE (1.3), we can only consider generators of very special SDEs. In the example above the coefficients in the SDE will depend on  $X_t$ , but after a change of variables this dependence is lost in the corresponding PDE.

*Third type of examples:* Versions of the above non-local operators with truncated and hence non-singular measures. Simply replace  $d\mu(z)$  in the definition of  $L$  by  $\mathbb{1}_{|z|>r} d\mu(z)$ , e.g.

$$\mathcal{L}_{\Delta_\infty}^{\alpha/2,r}[u](x) = \int_{\mathbb{R}^1} \left( u(x + Du(x)z) - u(x) - Du(x) \cdot Du(x)(e^z - 1) \right) \mathbb{1}_{|z|>r} \frac{c_\alpha g(z) dz}{|z|^{1+\alpha}},$$

where  $g(0) \neq 0$  and  $g$  is  $C^1$  at  $z = 0$ . Note that here  $L = \mathcal{L}_{\Delta_\infty}^{\alpha/2,r}$  with  $L_1 = 0$  and  $L_2 \neq 0$ , and  $(\mathbf{M}_\varepsilon)$  holds with  $A_2 = g(0)I$  and  $a = Dg(0)$ .

### 3.3. Remarks.

(a) [Continuity in  $\alpha, p$ ] All the operators above will be continuous in  $(\alpha, p) \in (0, 2) \times [2, \infty)$ . For example for any bounded  $C^2$  function  $\phi$  and sequence  $(\alpha', p') \rightarrow (\alpha, p) \in (0, 2) \times [2, \infty)$ ,

$$\mathcal{L}_{\Delta_{p'}}^{\alpha'/2} \phi \longrightarrow \mathcal{L}_{\Delta_p}^{\alpha/2} \phi \quad \text{in } \mathbb{R}^N.$$

(b) [The limit  $\alpha \rightarrow 2$ ] If  $u$  is smooth and bounded, then by easy computations,

$$\mathcal{L}_{\Delta_\infty}^{\alpha/2}[u], \mathcal{J}_{\Delta_\infty}^{\alpha/2}[u] \longrightarrow \Delta_\infty u, \quad \mathcal{L}_{\Delta_p}^{\alpha/2}[u], \tilde{\mathcal{L}}_{\Delta_p}^{\alpha/2}[u] \longrightarrow \Delta_p u, \quad \mathcal{L}_H^{\alpha/2}[u] \longrightarrow H[u],$$

$$\text{and} \quad \mathcal{L}_{\Delta_\infty}^{\alpha/2, 2-\alpha}[u] \longrightarrow g(0)\Delta_\infty u + Dg(0)Du$$

point-wise as  $\alpha \rightarrow 2$ . Hence *all* of these operators converge to their local counterparts including the truncated ones. These latter operators also give rise to a drift term (when  $\mu$  is non-symmetric!). Note that in these examples assumption  $(\mathbf{M}_\varepsilon)$  hold with  $\alpha = 2 - \varepsilon$ ,  $A = I$  or  $A = g(0)I$ , and  $a = 0$  or  $a = Dg(0)$ .

(c) [Growth assumptions] Our assumptions allow for extreme growth in the gradient and non-local terms. Our results cover the equation

$$u - F(L[u, Du](x)) = f(x)$$

for any continuous nondecreasing function  $F$  and any good operator  $L$  as above, e.g.

$$u - \left( e^{\mathcal{L}_{\Delta_p}^{\alpha/2}[u]} - 1 \right) = f(x) \quad \text{for any } p \geq 2.$$

## 4. VISCOSITY SOLUTIONS

In this section, we introduce the good notion of weak solution for equation (1.3). We prove that we have two equivalent definitions and that the solution concept is stable with respect to pointwise limits of uniformly bounded solutions.

We start by splitting  $L_1$  in (1.1) into two parts:  $L_1 = L_\delta + L^\delta$  for  $\delta > 0$ , where

$$L_\delta[\phi, D\phi](x) := \int_{|z| < \delta} \phi(x + j_1(D\phi(x), z)) - \phi(x) - j_1(D\phi(x), z) \cdot D\phi(x) \, d\mu_1(z), \quad (4.1)$$

$$L^\delta[u, p](x) := \int_{|z| \geq \delta} u(x + j_1(p, z)) - u(x) - j_1(p, z) \cdot p \, d\mu_1(z) \quad (p \in \mathbb{R}^N).$$

In view of  $(\mathbf{M})$ ,  $L_\delta$  is well-defined for any  $C^2$  function  $\phi$  and  $L^\delta$  for any bounded function  $u$ . Likewise, the operator  $L_2[u, p]$  is also well-defined for any  $p \in \mathbb{R}^N$  and bounded measurable function  $u$ . Now we can introduce the concept of solutions that we will use in this paper.

### Definition 4.1.

(a) A bounded usc function  $u$  is a viscosity subsolution of (1.3) if for any  $\delta > 0$ , any  $C^2$  function  $\phi$ , and any global maximum point  $x$  of  $u - \phi$ ,

$$F\left(u(x), D\phi(x), L_\delta[\phi, D\phi](x) + L^\delta[u, D\phi](x) + L_2[u, D\phi](x)\right) \leq f(x). \quad (4.2)$$

(b) A bounded lsc function  $u$  is a viscosity supersolution of (1.3) if for any  $\delta > 0$ , any  $C^2$  function  $\phi$ , and any global minimum point  $x$  of  $u - \phi$ ,

$$F\left(u(x), D\phi(x), L_\delta[\phi, D\phi](x) + L^\delta[u, D\phi](x) + L_2[u, D\phi](x)\right) \geq f(x). \quad (4.3)$$

(c) A viscosity solution is a bounded continuous function  $u$  which is both a subsolution and a supersolution.

Another possible definition is the following:

**Definition 4.2.**

(a) A bounded usc function  $u$  is a viscosity subsolution of (1.3) if for any bounded  $C^2$  function  $\phi$ , and any global maximum point  $x$  of  $u - \phi$ ,

$$F\left(u(x), D\phi(x), L[\phi, D\phi](x)\right) \leq f(x). \quad (4.4)$$

(b) A bounded lsc function  $u$  is a viscosity subsolution of (1.3) if for any bounded  $C^2$  function  $\phi$ , and any global minimum point  $x$  of  $u - \phi$ ,

$$F\left(u(x), D\phi(x), L[\phi, D\phi](x)\right) \geq f(x). \quad (4.5)$$

(c) A viscosity solution is a bounded continuous function  $u$  which is both a subsolution and a supersolution.

*Remark 4.3.* We may assume without loss of generality that the extrema of  $u - \phi$  are strict and that  $\phi = u$  at the extremal point. The latter comes from shifting the test function by a constant. To make an extremum (say a maximum) point  $x$  strict, we replace  $\phi$  by  $\phi + \delta\psi$  where  $\delta > 0$  and

$$\psi \in C^2(\mathbb{R}^N) \cap W^{2,\infty}(\mathbb{R}^N), \quad \psi = 0 \text{ and } D\psi = 0 \text{ at } x, \quad \text{and} \quad \psi > 0 \text{ elsewhere,}$$

and send  $\delta \rightarrow 0$  in the final step of the proof. As opposed to the local case, the  $\delta$ -terms will now be visible throughout the computations and vanish only in the final step.

**Lemma 4.4.** *If (M), (J1), (F1), and (F4) hold, then Definitions 4.1 and 4.2 are equivalent.*

*Proof.* The proof is pretty standard [26, 4, 18, 7]. Since  $(u - \phi)$  has a max in  $x$ ,  $L^\delta[u, D\phi](x) \leq L^\delta[\phi, D\phi](x)$  and  $L_2[u, D\phi](x) \leq L_2[\phi, D\phi](x)$ , and hence by (F1), (F4), and since  $L = L_\delta + L^\delta + L_2$ , inequality (4.4) follows from (4.2). Conversely, we may assume the max is strict (see Remark 4.3). Then there exists a smooth and uniformly bounded function  $\phi_\varepsilon$  such that  $u \leq \phi_\varepsilon \leq \phi$  and  $\phi_\varepsilon \rightarrow u$  a.e. as  $\varepsilon \rightarrow 0$ . It immediately follows that also  $u - \phi_\varepsilon$  and  $\phi_\varepsilon - \phi$  have maximum points at  $x$ . Hence, since  $D\phi_\varepsilon(x) = D\phi(x)$  and by the definition of  $L$  (monotonicity and  $L = L_\delta + L^\delta + L_2$ ),

$$L[\phi_\varepsilon, D\phi_\varepsilon](x) \leq L_\delta[\phi, D\phi](x) + L^\delta[\phi_\varepsilon, D\phi](x) + L_2[\phi_\varepsilon, D\phi](x).$$

Hence, by inequality (4.4) with  $\phi_\varepsilon$  replacing  $\phi$ , inequality (4.2) with  $\phi_\varepsilon$  replacing  $u$  follows. Now we conclude by sending  $\varepsilon \rightarrow 0$ , using (M), (J1), (F1), (F4), and the dominated convergence theorem.  $\square$

Next, we show that this solution concept is stable with respect to local uniform limits, to so-called half-relaxed limits, and more generally to very general perturbations of the equation. Consider

$$F_\varepsilon(u_\varepsilon, Du_\varepsilon, L_\varepsilon[u_\varepsilon, Du_\varepsilon]) = f_\varepsilon(x) \quad \text{in} \quad \mathbb{R}^N, \quad (4.6)$$

where  $(f_\varepsilon, F_\varepsilon, L_\varepsilon := L_{1,\varepsilon} + L_{2,\varepsilon})$  satisfy (F4), (F1), (M), and (J1) for each fixed  $\varepsilon > 0$ , and where

$$\begin{aligned} L_{1,\varepsilon}[u, Du](x) &= \int_{\mathbb{R}^N} u(x + j_{1,\varepsilon}(Du, z)) - u(x) - j_{1,\varepsilon}(Du, z) \cdot Du(x) \, d\mu_{1,\varepsilon}(z), \\ L_{2,\varepsilon}[u, Du](x) &= \int_{\mathbb{R}^N} u(x + j_{2,\varepsilon}(Du, z)) - u(x) \, d\mu_{2,\varepsilon}(z). \end{aligned}$$

Then we define the “half-relaxed limits”:

$$\bar{u}(x) := \limsup_{y \rightarrow x, \varepsilon \rightarrow 0} u_\varepsilon(y) \quad \text{and} \quad \underline{u}(x) := \liminf_{y \rightarrow x, \varepsilon \rightarrow 0} u_\varepsilon(y). \quad (4.7)$$

$$\bar{f}(x) := \limsup_{y \rightarrow x, \varepsilon \rightarrow 0} f_\varepsilon(y) \quad \text{and} \quad \underline{f}(x) := \liminf_{y \rightarrow x, \varepsilon \rightarrow 0} f_\varepsilon(y). \quad (4.8)$$

$$\bar{F}(u, p, l) := \limsup_{\substack{(v, q, m) \rightarrow (u, p, l) \\ \varepsilon \rightarrow 0}} F_\varepsilon(v, q, m) \quad \text{and} \quad \underline{F}(u, p, l) := \liminf_{\substack{(v, q, m) \rightarrow (u, p, l) \\ \varepsilon \rightarrow 0}} F_\varepsilon(v, q, m). \quad (4.9)$$

**Lemma 4.5** (Stability 1). *Assume  $\{f_\varepsilon, F_\varepsilon, L_\varepsilon\}_\varepsilon$  satisfy **(M)**, **(J1)**, **(F1)**, **(F4)** for any  $\varepsilon > 0$ ,*

$$\liminf_{\varepsilon \rightarrow 0} L_\varepsilon[\phi, D\phi](x_\varepsilon) \leq L[\phi, D\phi](x) \quad (\text{resp. } \limsup_{\varepsilon \rightarrow 0} L_\varepsilon[\phi, D\phi](x_\varepsilon) \geq L[\phi, D\phi](x)),$$

for all bounded  $\phi \in C^2$  and all sequences  $x_\varepsilon \rightarrow x$ , and that  $\{u_\varepsilon\}_{\varepsilon > 0}$  is a sequence of uniformly bounded subsolutions (resp. supersolutions) of (4.6).

Then  $\bar{u}(x)$  is a subsolution (resp.  $\underline{u}(x)$  supersolution) of (1.3) with  $(f, F)$  replaced by  $(\bar{f}, \underline{F})$  (resp.  $(\underline{f}, \bar{F})$ ).

We also have the following stability result.

**Lemma 4.6** (Stability 2). *Assume **(M)**, **(J1)**, **(F1)**, **(F4)** hold, and  $\{u_a\}_{a \in A}$ , for a set  $A$ , is a family of uniformly bounded subsolutions (resp. supersolutions) of (1.3).*

(a) *If for any  $n$ ,  $u_{a_n}$  is continuous and  $u_{a_n} \rightarrow u$  locally uniformly as  $n \rightarrow \infty$ , then  $u$  is a continuous bounded subsolution (resp. supersolution) of (1.3).*

(b)  *$u := \sup_{a \in A} u_a$  is a subsolution (resp.  $v = \inf_{a \in A} u_a$  is a supersolution) of (1.3).*

The proofs follow after the next remark.

*Remark 4.7.* (i) Similar type of results can be found in [7], but without variation in  $L$ .

(ii) Compare Lemma 4.6-(a) to the no stability w.r.t. local uniform convergence result of [9]. In [8] there is stability, but the non-local operators are more different from ours than in [9].

*Proof of Lemma 4.5.* The proof is quite standard, see e.g. [7] (Theorem 2) for a similar proof. We only do the subsolution case since the supersolution case is similar. Assume  $\phi$  is  $C^2$  and bounded and  $\bar{u} - \phi$  has a global maximum at  $x$ , we will show that inequality (4.4) holds and we are done.

Modifying the test function if necessary (as in Remark 4.3, assuming also  $\psi(y) = 1$  for  $|y| > 1$ ), we may assume the maximum is unique, strict, and can not be attained at infinity. In fact, we may assume that

$$(\bar{u} - \phi)(x) > \sup_{|y-x| > r} (\bar{u} - \phi)(y) \quad \text{for any } r > 0. \quad (4.10)$$

Then we take a subsequence such that  $\bar{u}(x) = \lim_\varepsilon u_\varepsilon(x_\varepsilon)$ , and note that by (4.10) and classical arguments [10, Lemma V.1.6], we may find a sequence  $\{y_\varepsilon\}_\varepsilon$  such that

$$u_\varepsilon - \phi \quad \text{has a global maximum at } y_\varepsilon, \quad y_\varepsilon \rightarrow x, \quad \text{and} \quad u_\varepsilon(y_\varepsilon) \rightarrow \bar{u}(x).$$

Since  $u_\varepsilon$  is a subsolution of (4.6),

$$F_\varepsilon(u_\varepsilon(y_\varepsilon), D\phi(y_\varepsilon), l_\varepsilon) \leq f_\varepsilon(y_\varepsilon) \quad \text{where} \quad l_\varepsilon = L_\varepsilon[\phi, D\phi](y_\varepsilon).$$

By the construction of  $y_\varepsilon$  and the assumption of the Lemma,

$$\liminf_\varepsilon l_\varepsilon \leq L[\phi, D\phi](x). \quad (4.11)$$

Hence if we take a further subsequence in  $\varepsilon$  such that  $l_\varepsilon \rightarrow \liminf_\varepsilon l_\varepsilon$ , then by the definition of  $(\bar{f}, \underline{F})$  and continuity, **(F1)** and **(F4)**,

$$\underline{F}(\bar{u}(x), D\phi(x), \liminf_\varepsilon l_\varepsilon) \leq \bar{f}(x).$$

and inequality (4.4) then follows from (4.11) and monotonicity **(F1)**.  $\square$

*Proof of Lemma 4.6.* (a) Let  $\phi$  be  $C^2$  and bounded and  $x_\varepsilon \rightarrow x$ . By assumptions **(M)** and **(J1)**, and the dominated convergence theorem,

$$\lim_{x_\varepsilon \rightarrow x} L[\phi, D\phi](x_\varepsilon) = L[\phi, D\phi](x).$$

Hence by Lemma 4.5,  $\bar{u}(x) = \limsup_{y \rightarrow x, n \rightarrow \infty} u_{a_n}(y)$  is a (bounded) subsolution of (1.3). Since  $u_{a_n}$  is continuous and  $u_{a_n} \rightarrow u$  locally uniformly, it follows that  $\bar{u} = u$  and  $u$  is continuous.

(b) The proof is similar to the proof of Lemma 4.5 and we only do the subsolution case. Assume  $u - \phi$  has a strict global max at  $x$ . By the definition of the supremum, there is a sequence  $u_{a_k}(x_k) \rightarrow u(x)$  as  $k \rightarrow \infty$ . As in the previous proof we may find a sequence  $\{y_k\}_k$  such that

$$u_{a_k} - \phi \text{ has a global maximum at } y_k, \quad y_k \rightarrow x, \quad \text{and} \quad u_{a_k}(y_k) \rightarrow u(x).$$

Since  $u_{a_k}$  is a subsolution of (1.3),

$$F(u_{a_k}(y_k), D\phi(y_k), l_k) \leq f(y_k) \quad \text{where} \quad l_k = L[\phi, D\phi](y_k).$$

By the construction of  $y_k$ , assumptions **(M)** and **(J1)**, and the dominated convergence theorem,

$$\lim_k l_k = L[\phi, D\phi](x),$$

and then by the continuity, **(F1)** and **(F4)**, inequality (4.4) holds.  $\square$

## 5. PROOFS OF THE MAIN RESULTS

### 5.1. Proof of Theorem 2.1 (comparison).

*Proof of Theorem 2.1(a).* We proceed by contradiction, assuming that  $M := \sup(u - v) > 0$ .

Let  $\varepsilon, R > 0$  and define

$$\Phi_{\varepsilon, R}(x, y) := u(x) - v(y) - \phi(x, y), \tag{5.1}$$

where

$$\phi(x, y) = \frac{1}{\varepsilon^2} \varphi(x - y) + \psi\left(\frac{x}{R}\right) + \psi\left(\frac{y}{R}\right), \tag{5.2}$$

and  $\varphi, \psi$  are smooth bounded radially symmetric and radially non-decreasing functions such that

$$\varphi(x) = \begin{cases} |x|^2 & \text{for } |x| < 1 \\ 2 & \text{for } |x| > 4 \end{cases} \quad \text{and} \quad \psi(x) = \begin{cases} 0 & \text{for } |x| < \frac{1}{2} \\ 2(\|u\|_\infty + \|v\|_\infty) + 1 & \text{for } |x| > 1 \end{cases}$$

By penalization (the  $\psi$ -terms) the supremum of  $\Phi_{\varepsilon, R}$  is attained at a point  $(\bar{x}, \bar{y})$ , and since  $M > 0$  this supremum is positive when  $R$  is big enough (see **1**) below):

$$M_{\varepsilon, R} := \max \Phi_{\varepsilon, R} = \Phi(\bar{x}, \bar{y}) > 0.$$

For the sake of simplicity we drop the reference to  $\varepsilon, R$  for the maximum point. By the inequality  $\Phi(\bar{x}, \bar{x}) + \Phi(\bar{y}, \bar{y}) \leq 2\Phi(\bar{x}, \bar{y})$ , it follows that  $\frac{2}{\varepsilon^2} \varphi(\bar{x} - \bar{y}) \leq u(\bar{x}) - u(\bar{y}) + v(\bar{x}) - v(\bar{y})$ , and hence

$$\varphi(\bar{x} - \bar{y}) \leq (\|u\|_\infty + \|v\|_\infty) \varepsilon^2. \tag{5.3}$$

By taking  $\varepsilon > 0$  small enough, we can always assume that

$$\varphi(\bar{x} - \bar{y}) = |\bar{x} - \bar{y}|^2 \quad \text{and} \quad (D\varphi)(\bar{x} - \bar{y}) = 2(\bar{x} - \bar{y}).$$

In particular,  $|\bar{x} - \bar{y}| \leq (\|u\|_\infty + \|v\|_\infty)\varepsilon$  and this estimate is independent of  $R$ .

From the maximum of  $\Phi_{\varepsilon,R}$  it follows that  $u(x) - \phi(x, \bar{y})$  has a global maximum point at  $\bar{x}$  and  $v(y) - (-\phi)(\bar{x}, y)$  has a global minimum point at  $\bar{y}$ . Subtracting the corresponding viscosity inequalities for  $u$  and  $v$  (cf. Definition 4.1) gives for any  $\delta > 0$  that

$$\begin{aligned} 0 &\geq - \left( L_\delta[\phi(\cdot, \bar{y}), D_x\phi](\bar{x}) - L_\delta[(-\phi)(\bar{x}, \cdot), D_y(-\phi)](\bar{y}) \right) \\ &\quad - \left( L^\delta[u, D_x\phi](\bar{x}) - L^\delta[v, D_y(-\phi)](\bar{y}) \right) \\ &\quad - \left( L_2[u, D_x\phi](\bar{x}) - L_2[v, D_y(-\phi)](\bar{y}) \right) \\ &\quad - (f(\bar{x}) - f(\bar{y})) + (u(\bar{x}) - v(\bar{y})) \\ &\geq -I_\delta - I^\delta - I_2 - \omega_f(\bar{x} - \bar{y}) + (u(\bar{x}) - v(\bar{y})). \end{aligned} \tag{5.4}$$

The strategy is now to estimate  $I_\delta$ ,  $I^\delta$ , and  $I_2$ , and prove that when sending first  $\delta \rightarrow 0$ , then  $R \rightarrow \infty$ , and finally  $\varepsilon \rightarrow 0$ ,

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{R \rightarrow \infty} \limsup_{\delta \rightarrow 0} (I_\delta + I^\delta + I_2) \leq 0.$$

We will also show that

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{R \rightarrow \infty} \limsup_{\delta \rightarrow 0} (u(\bar{x}) - v(\bar{y})) \geq M, \tag{5.5}$$

and hence by the viscosity inequality (5.4) we get the contradiction that concludes the proof:

$$0 \geq M.$$

We proceed in 4 steps:

**1)** We show that (5.5) holds. First note that  $u(\bar{x}) - v(\bar{y}) = M_{\varepsilon,R} + \phi(\bar{x}, \bar{y})$  do not depend on  $\delta$ . Then by the maximum point property, it follows that

$$M_{\varepsilon,R} \xrightarrow{\varepsilon} := \sup \left( u(x) - v(y) - \frac{1}{\varepsilon^2} \varphi(x - y) \right) \quad \text{and} \quad \psi\left(\frac{\bar{x}}{R}\right) + \psi\left(\frac{\bar{y}}{R}\right) \rightarrow 0$$

as  $R \rightarrow \infty$  (see Lemma 2.3 in [19]). Observe now that  $M \leq M_\varepsilon \leq M_{\varepsilon'}$  for  $\varepsilon \leq \varepsilon'$ , and hence by monotone convergence,  $M_\varepsilon \searrow \tilde{M}$  for some  $\tilde{M} \geq M$ . Since  $\tilde{M} = \limsup_{\varepsilon \rightarrow 0} \limsup_{R \rightarrow \infty} \limsup_{\delta \rightarrow 0} (u(\bar{x}) - v(\bar{y}))$ , we are done.

**2)** To estimate the  $I_\delta$ -term, we Taylor expand to find that

$$\begin{aligned} &\int_{|z| < \delta} \phi(\bar{x} + j_1(D_x\phi(\bar{x}, \bar{y}), z), \bar{y}) - \phi(\bar{x}, \bar{y}) - j_1(D_x\phi(\bar{x}, \bar{y}), z) \cdot D_x\phi(\bar{x}, \bar{y}) \, d\mu_1(z) \\ &\leq \|D^2\phi\|_\infty \int_{|z| < \delta} |j_1(D_x\phi(\bar{x}, \bar{y}), z)|^2 \, d\mu_1(z) = o_\delta(1) \end{aligned}$$

for fixed  $\varepsilon, R > 0$ . Here the  $o_\delta(1)$  comes from assumption **(J1)** and dominated convergence as  $\delta \rightarrow 0$ . After a similar estimate for  $L_\delta[-\phi, D_y(-\phi)]$ , we conclude that  $I_\delta \rightarrow 0$  as  $\delta \rightarrow 0$  and  $\varepsilon, R > 0$  are fixed.

**3)** We estimate  $I^\delta$ . Using the notation  $j_{\bar{x}}(z) := j_1(D_x\phi(\bar{x}, \bar{y}), z)$  and  $j_{\bar{y}}(z) := j_1(D_y(-\phi)(\bar{x}, \bar{y}), z)$ , and the maximum point property of  $\Phi_{\varepsilon, R}$ ,

$$\Phi_{\varepsilon, R}(\bar{x} + j_{\bar{x}}, \bar{y} + j_{\bar{y}}) \leq \Phi_{\varepsilon, R}(\bar{x}, \bar{y}),$$

we see that

$$\begin{aligned} I^\delta &= \int_{\delta \leq |z|} \left( \left( u(\bar{x} + j_{\bar{x}}(z)) - u(\bar{x}) \right) - \left( v(\bar{y} + j_{\bar{y}}(z)) - v(\bar{y}) \right) \right. \\ &\quad \left. + j_{\bar{x}}(z) \cdot D_x\phi(\bar{x}, \bar{y}) + j_{\bar{y}}(z) \cdot D_y\phi(\bar{x}, \bar{y}) \right) d\mu_1(z) \\ &\leq \int_{\delta \leq |z|} \left( \phi(\bar{x} + j_{\bar{x}}(z), \bar{y} + j_{\bar{y}}(z)) - \phi(\bar{x}, \bar{y}) - D_x\phi(\bar{x}, \bar{y}) \cdot j_{\bar{x}}(z) - D_y\phi(\bar{x}, \bar{y}) \cdot j_{\bar{y}}(z) \right) d\mu_1(z). \end{aligned}$$

Since  $D^2\varphi$  is bounded and  $|D^2\psi(\frac{x}{R})| \leq \frac{1}{R^2}\|D^2\psi\|_\infty < \infty$ , a short computation using Taylor expansions shows that

$$I^\delta \leq \int_{\delta \leq |z|} \left( \frac{1}{2\varepsilon^2}\|D^2\varphi\|_\infty |j_{\bar{x}}(z) - j_{\bar{y}}(z)|^2 + \frac{1}{2R^2}\|D^2\psi\|_\infty (|j_{\bar{x}}(z)|^2 + |j_{\bar{y}}(z)|^2) \right) d\mu_1(z).$$

To proceed we compute the gradients,

$$D_x\phi(\bar{x}, \bar{y}) = p_\varepsilon + \frac{1}{R}D\psi\left(\frac{\bar{x}}{R}\right), \quad D_y(-\phi)(\bar{x}, \bar{y}) = p_\varepsilon - \frac{1}{R}D\psi\left(\frac{\bar{y}}{R}\right), \quad p_\varepsilon = \frac{2(\bar{x} - \bar{y})}{\varepsilon^2},$$

and note that for fixed  $\varepsilon > 0$ , they are uniformly bounded for  $R > 1$  by estimate (5.3). Hence, there is  $r_\varepsilon > 0$  such that  $|D\phi(\bar{x}, \bar{y})| \leq r_\varepsilon$  for all  $\delta > 0$  and  $R > 1$ , and then by assumptions **(J1)** and **(J2)**,

$$\begin{aligned} I^\delta &\leq O\left(\frac{1}{\varepsilon^2}\right)\omega_{j, r_\varepsilon}(D_x\phi(\bar{x}, \bar{y}) - D_y(-\phi)(\bar{x}, \bar{y})) + O\left(\frac{1}{R^2}\right)C_{j, r_\varepsilon} \\ &\leq O\left(\frac{1}{\varepsilon^2}\right)\omega_{j, r_\varepsilon}\left(O\left(\frac{1}{R}\right)\right) + O\left(\frac{1}{R^2}\right)C_{j, r_\varepsilon}. \end{aligned}$$

We first send  $\delta \rightarrow 0$  since nothing depends on  $\delta$  on the right-hand side, and then we send  $R \rightarrow \infty$  and find that

$$\limsup_{R \rightarrow \infty} \limsup_{\delta \rightarrow 0} I^\delta \leq 0.$$

**4)** Finally, we estimate  $I_2$ . First note that by the maximum point property, the positivity of  $\phi$ , the calculations of gradients in (c), and estimate (5.3),

$$\begin{aligned} I_2 &\leq \int_{|z| > 0} \phi\left(\bar{x} + j_2\left(p_\varepsilon + \frac{1}{R}D\psi\left(\frac{\bar{x}}{R}\right), z\right), \bar{y} + j_2\left(p_\varepsilon - \frac{1}{R}D\psi\left(\frac{\bar{y}}{R}\right), z\right)\right) - \phi(\bar{x}, \bar{y}) \, d\mu_2(z) \\ &\leq \frac{1}{\varepsilon^2} \int_{|z| > 0} \sup_{x, y \in \mathbb{R}^N} \left\{ \varphi\left(x + j_2\left(p_\varepsilon + \frac{1}{R}D\psi\left(\frac{x}{R}\right), z\right) - \left(y + j_2\left(p_\varepsilon - \frac{1}{R}D\psi\left(\frac{y}{R}\right), z\right)\right) \right. \right. \\ &\quad \left. \left. - \varphi(x - y) \right\} d\mu_2(z) \\ &\quad + \int_{|z| > 0} \psi\left(\frac{\bar{x} + j_2\left(p_\varepsilon + \frac{1}{R}D\psi\left(\frac{\bar{x}}{R}\right), z\right)}{R}\right) + \psi\left(\frac{\bar{y} + j_2\left(p_\varepsilon - \frac{1}{R}D\psi\left(\frac{\bar{y}}{R}\right), z\right)}{R}\right) \, d\mu_2(z) \\ &:= J_1 + J_2. \end{aligned}$$

Now we send  $R \rightarrow \infty$  in  $J_1$ . Then by compactness,  $p_\varepsilon$  will up to a subsequence converge to a limit that we also call  $p_\varepsilon$ . By the boundedness of  $D\psi$  and  $p$ -continuity of  $j_2(p, z)$  for a.e.  $z$  in **(J1)**,

$$\lim_{R \rightarrow \infty} \sup_{x \in \mathbb{R}^N} \left| j_2\left(p_\varepsilon \pm \frac{1}{R}D\psi\left(\frac{x}{R}\right), z\right) - j_2(p_\varepsilon, z) \right| = 0 \quad \text{for a.e. } z.$$



Hence since  $\varphi$  is a Lipschitz continuous function,

$$\begin{aligned} & \sup_{x,y \in \mathbb{R}^N} \left| \varphi \left( x - y - j_2 \left( p_\varepsilon + \frac{1}{R} D\psi \left( \frac{x}{R} \right), z \right) + j_2 \left( p_\varepsilon - \frac{1}{R} D\psi \left( \frac{y}{R} \right), z \right) \right) - \varphi(x - y) \right| \\ & \leq \|D\varphi\|_\infty \sup_{x,y \in \mathbb{R}^N} \left| j_2 \left( p_\varepsilon + \frac{1}{R} D\psi \left( \frac{x}{R} \right), z \right) - j_2 \left( p_\varepsilon - \frac{1}{R} D\psi \left( \frac{y}{R} \right), z \right) \right| \rightarrow 0 \end{aligned}$$

as  $R \rightarrow \infty$  for a.e.  $z$ . Hence, the  $J_1$ -integrand is a uniformly bounded function converging to 0 as  $R \rightarrow \infty$  for a.e.  $z$ . Hence by the dominated convergence theorem (for fixed  $\varepsilon$ ),

$$\limsup_{R \rightarrow \infty} J_1 \leq 0.$$

Now we send  $R \rightarrow \infty$  in  $J_2$ . Here we use the fact that

$$\psi \left( \frac{\bar{x}}{R} \right) + \psi \left( \frac{\bar{y}}{R} \right) \rightarrow 0 \quad \text{as} \quad R \rightarrow \infty,$$

which is a simple consequence of the maximum point property (see Lemma 2.3 in [19]). Since  $j_2$  is locally bounded for a.e. fixed  $z$  by **(J1)** and  $\psi$  is continuous,

$$\frac{j_2 \left( p_\varepsilon + \frac{1}{R} D\psi \left( \frac{\bar{y}}{R} \right), z \right)}{R} \rightarrow 0 \quad \text{for a.a. } z$$

as  $R \rightarrow \infty$ , and hence

$$\psi \left( \frac{\bar{x} + j_2 \left( p_\varepsilon + \frac{1}{R} D\psi \left( \frac{\bar{x}}{R} \right), z \right)}{R} \right) \rightarrow \psi(0) = 0 \quad \text{for a.a. } z$$

as  $R \rightarrow \infty$ . Since  $\psi$  is bounded, we can use the dominated convergence theorem to conclude that

$$\limsup_{R \rightarrow \infty} J_2 = 0.$$

Since  $I_2$  is independent of  $\delta$ , we can now conclude that

$$\limsup_{R \rightarrow \infty} \limsup_{\delta \rightarrow 0} I_2 \leq 0,$$

and the proof is complete.  $\square$

*Proof of Theorem 2.1-(b).* Part of the proof is similar to the previous proof. We start by assuming that  $M := \sup\{u(x) - v(x)\} > 0$  and consider the maximum  $M_{\varepsilon,R}$  of

$$\Phi_{\varepsilon,R}(x, y) := u(x) - v(y) - \phi(x, y),$$

where  $\phi$  was defined in the proof of Theorem 2.1-(a). Since  $M_{\varepsilon,R} \rightarrow M_\varepsilon \geq M$  as  $R \rightarrow \infty$ , we may assume that  $M_{\varepsilon,R} \geq M/2$  and  $u(\bar{x}) > v(\bar{y})$ . Since  $u - \phi(\cdot, \bar{y})$  has a global max in  $\bar{x}$  and  $v - (-\phi)(\bar{x}, \cdot)$  has a global min in  $\bar{y}$ , we subtract the corresponding viscosity inequalities and find that

$$\begin{aligned} & F \left( u(\bar{x}), p_\varepsilon + O\left(\frac{1}{R}\right), \underbrace{L_\delta[\phi(\cdot, \bar{y}), D_x \phi](\bar{x}) + L^\delta[u, D_x \phi](\bar{x}) + L_2[u, D_x \phi](\bar{x})}_{I_x} \right) \\ & - F \left( v(\bar{y}), p_\varepsilon + O\left(\frac{1}{R}\right), \underbrace{L_\delta[-\phi(\bar{x}, \cdot), D_y(-\phi)](\bar{y}) + L^\delta[v, D_y(-\phi)](\bar{y}) + L_2[v, D_y(-\phi)](\bar{y})}_{I_y} \right) \quad (5.6) \\ & \leq f(\bar{x}) - f(\bar{y}), \end{aligned}$$

where  $p_\varepsilon = 2(\bar{x} - \bar{y})/\varepsilon$ .

We now estimate the different terms. By the estimates in the proof of Theorem 2.1-(a),

$$\begin{aligned} & \left| L_\delta[\phi(\cdot, \bar{y}), D_x \phi](\bar{x}) - L_\delta[\phi(\bar{x}, \cdot), D_y(-\phi)](\bar{y}) \right| = \frac{1}{\varepsilon^2} o_\delta(1), \\ & \left| L^\delta[u, D_x \phi](\bar{x}) - L^\delta[v, D_y(-\phi)](\bar{y}) \right| + \left| L_2[u, D_x \phi](\bar{x}) - L_2[v, D_y(-\phi)](\bar{y}) \right| = \left(1 + \frac{1}{\varepsilon^2}\right) o_R(1), \end{aligned}$$

and hence

$$|I_x - I_y| = \left(1 + \frac{1}{\varepsilon^2}\right) (o_R(1) + o_\delta(1)).$$

By the order we will take the limits, we may and will always assume that terms on the right hand sides are bounded (by 1 for example). moreover, by the estimates in the proof of Theorem 2.1-(a),

$$\begin{aligned} |D\phi(\bar{x}, \bar{y})| & \leq C \left( \frac{1}{\varepsilon} + \frac{1}{R} \right), \\ |D^2\phi(\bar{x}, \bar{y})| & \leq C \left( \frac{1}{\varepsilon^2} + \frac{1}{R^2} \right), \\ |L_\delta[\phi(\cdot, \bar{y}), D_x \phi](\bar{x})| & \leq \|D^2\phi\|_\infty \int_{|z| < \delta} |j_1(D_x \phi(\bar{x}, \bar{y}), z)|^2 d\mu_1(z), \\ |L_\delta[-\phi(\bar{x}, \cdot), D_y(-\phi)](\bar{y})| & \leq \|D^2\phi\|_\infty \int_{|z| < \delta} |j_1(D_y(-\phi)(\bar{x}, \bar{y}), z)|^2 d\mu_1(z), \\ |L_2[u, D_x \phi](\bar{x})| + |L_2[v, D_y(-\phi)](\bar{y})| & \leq 2(\|u\|_\infty \vee \|v\|_\infty) \mu_2(\mathbb{R}^N), \end{aligned}$$

and by the maximum point property,

$$\begin{aligned} L^\delta[u, D_x \phi](\bar{x}) & \leq L^\delta[\phi(\cdot, \bar{y}), D_x \phi(\bar{x}, \bar{y})](\bar{x}) \\ & \leq \|D_x^2\phi\|_\infty \int_{|z| > 0} |j(D_x \phi(\bar{x}, \bar{y}), z)|^2 d\mu_1(z) - \|D_y^2\phi\|_\infty \int_{|z| > 0} |j(D_y(-\phi)(\bar{x}, \bar{y}), z)|^2 d\mu_1(z) \\ & \leq L^\delta[v, D_y(-\phi)](\bar{y}). \end{aligned}$$

If  $\varepsilon > 0$  is fixed, then by **(M)** and **(J1)**, these terms are uniformly bounded for  $R > 1$  and  $\delta > 0$ . These and the previous bounds then implies that there is a  $C_\varepsilon > 0$  such that

$$-C_\varepsilon \leq L^\delta[v, D_y(-\phi)](\bar{y}) \leq L^\delta[u, D_x \phi](\bar{x}) + |L^\delta[v, D_y(-\phi)](\bar{y}) - L^\delta[u, D_x \phi](\bar{x})| \leq C_\varepsilon,$$

and similarly we can show that  $|L^\delta[u, D_x \phi](\bar{x})| \leq C_\varepsilon$ . Hence there is  $r_\varepsilon > 0$  such that

$$|D\phi(\bar{x}, \bar{y})| + |I_x| + |I_y| \leq r_\varepsilon \quad \text{for all } R > 1, \delta > 0.$$

By (5.6) and the previous estimates, **(F2)**, **(F3)**, **(F4)**, we see

$$\begin{aligned} \gamma \frac{M}{2} & \leq F(u(\bar{x}), p_\varepsilon - O(\frac{1}{R}), I_y) - F(u(\bar{x}), p_\varepsilon + O(\frac{1}{R}), I_x) + f(\bar{x}) - f(\bar{y}) \\ & \leq \omega_{\|u\|_\infty \vee \|v\|_\infty, r_\varepsilon} \left( O(\frac{1}{R}) + \left(1 + \frac{1}{\varepsilon^2}\right) (o_R(1) + o_\delta(1)) \right) + o_\varepsilon(1). \end{aligned}$$

Sending first  $\delta \rightarrow 0$ , then  $R \rightarrow \infty$ , and finally  $\varepsilon \rightarrow 0$ , we get again  $M \leq 0$ . This is a contradiction and the result follows.  $\square$

**5.2. Proof of Theorem 2.3 (existence).** A major challenge we face when we want to prove existence, is the implicit nature of equation (1.3) with a gradient dependence inside the  $j$  functions. It seems non-trivial to use Perron's method for such equations, and since fixed point iterations require convergence of the full sequence to get the equation in the limit, compactness argument (yielding subsequences) can not work. We have been able to overcome the problem by a nontrivial approximation procedure, which is inspired by the "Sirtaki method" of [5], along with a fixed point argument using Schauder's fixed point theorem. We start by proving existence for an approximate problem in a bounded set, and then pass to the limit using the method of half-relaxed limits and strong comparison of the limit equation.

We begin with the linear case (1.6). The simple adaptations for the general case are given at the end of this section. Consider now the following approximate problem: find  $u \in C^2(\overline{B}_R)$  such that

$$\begin{cases} u - T_M[L_k^R[u, Du]] - \varepsilon \Delta u = f(x), & x \in B_R(0), \\ u = 0, & x \in \partial B_R(0), \end{cases} \quad (5.7)$$

where  $L_k^R[v, Dv] = L_{1,k}^R[v, Dv] + L_{2,k}^R[v, Dv]$  for

$$\begin{aligned} L_{1,k}^R[v, Dv](x) &= \int_{\mathbb{R}^P} v(P_R(x + j_1(Dv(x), z))) - v(x) - Dv(x) \cdot j_1(Dv(x), z) \mathbb{1}_{|z| < 1} \, d\mu_{1,k}(z), \\ L_{2,k}^R[v, Dv](x) &= \int_{\mathbb{R}^P} v(P_R(x + j_2(Dv(x), z))) - v(x) \, d\mu_{2,k}(z), \end{aligned}$$

$T_M$  is a truncation and  $P_R$  the orthogonal projection onto  $\overline{B}_R$ ,

$$T_M[f] := \min(\max(f, -M), M), \quad P_R(x) := \begin{cases} x & \text{if } |x| \leq R, \\ \frac{R}{|x|}x & \text{if } |x| > R, \end{cases}$$

and the measures

$$\mu_{1,k} := \rho_k * (\mu_1 \cdot \mathbb{1}_{1/k < |z| < k}) \quad \text{and} \quad \mu_{2,k} := \rho_k * \mu_2,$$

for a mollifier  $\rho_k(z) = k^P \rho(kz)$ ,  $0 \leq \rho \in C^\infty(\mathbb{R}^P)$  is symmetric with support in  $B_1$  and  $\int \rho = 1$ .

*Remark 5.1.* (i) The truncated mollified measures  $\mu_{1,k}$  and  $\mu_{2,k}$  are absolutely continuous with respect to the Lebesgue measure with bounded densities, see Lemma 5.3 below.

(ii) From the definition it follows that

$$|P_R(x) - P_R(y)| \leq |x - y| \quad \text{for all } x, y \in \mathbb{R}^N.$$

The projection allows us to look for solutions that are defined only in  $\overline{B}_R$  and not in all of  $\mathbb{R}^N$  as in equation (1.3). The new nonlocal term is of Neuman-type, corresponding to jump processes that are projected back to the boundary of the domain immediately upon leaving it (cf.[5, 6]).

To prove existence we first strengthen the assumptions on  $j$  and  $f$ , later we do the general case.

**(J1')**  $j_1(p, z)$  and  $j_2(p, z)$  are Borel measurable, locally bounded, continuous in  $p$  for a.e.  $z$ , and for every  $r > 0$  there is  $C_r$  such that for all  $|p| \leq r$  and  $|z| < 1$ ,

$$|j_1(p, z)| \leq C_r |z|$$

**(J2')** for any  $K > 0$ , there exists  $C = C(K)$  such that for any  $|p|, |q|, |z| \leq K$ ,

$$|j_i(p, z) - j_i(q, z)| \leq C |p - q|.$$

(F5')  $f : \mathbb{R}^N \rightarrow \mathbb{R}$  is bounded and Lipschitz continuous.

*Remark 5.2.* For any  $x \in B_R$  and  $u \in C^2(\overline{B_R})$ , all terms in (5.7) are well-defined and the equation holds in the classical sense (since  $\mu_{i,k}$  are bounded and  $P_R$  is continuous, the integral terms are well-defined because of (J1')). Note that (J1') and (M) implies (J1), while  $\mu_1$  bounded and compactly supported and (J2') implies (J2).

We state the properties of  $\mu_{i,k}$  that we will need later. The proof is given in Appendix B.

**Lemma 5.3.** *Assume (M) holds and  $\delta > 0$ .*

(a) *The measures  $\mu_{1,k}$  and  $\mu_{2,k}$  have densities  $\bar{\mu}_{1,k}$  and  $\bar{\mu}_{2,k}$  with respect to the Lebesgue measure on  $\mathbb{R}^N$  such that*

$$\|\bar{\mu}_{1,k}\|_\infty, \|\bar{\mu}_{2,k}\|_\infty < \infty \quad \text{and} \quad \int_{|z|<\delta} |z|^2 \bar{\mu}_{1,k}(z) dz \leq 4 \int_{|z|<\delta} |z|^2 \mu_1(dz).$$

*From now on, let  $(\psi_k)$  be a sequence of functions and  $C > 0$  constants independent of  $p, z$  that differ from line to line.*

(b) *If  $\|\psi_k\|_\infty \leq C$  for all  $k$ , and  $\sup_{\delta \leq |z| < K} |\psi_k(z) - \psi(z)| \xrightarrow{k \rightarrow \infty} 0$  for any  $K > \delta$ , then*

$$\int_{|z| \geq \delta} \psi_k(z) d\mu_{1,k}(z) \xrightarrow{k \rightarrow \infty} \int_{|z| \geq \delta} \psi(z) \mu_1(dz).$$

(c) *If  $|\psi_k(z)| \leq C|z|^2$  for all  $k, z$ , and  $\sup_{|z| < \delta} |\psi_k(z) - \psi(z)| \xrightarrow{k \rightarrow \infty} 0$ , then*

$$\int_{0 < |z| < \delta} \psi_k(z) d\mu_{1,k}(z) \xrightarrow{k \rightarrow \infty} \int_{0 < |z| < \delta} \psi(z) \mu_1(dz).$$

(d) *If  $\|\psi_k\|_\infty \leq C$  for all  $k$ , and  $\sup_{|z| < K} |\psi_k(z) - \psi(z)| \xrightarrow{k \rightarrow \infty} 0$  for any  $K > 0$ , then*

$$\int_{|z| > 0} \psi_k(z) d\mu_{2,k}(z) \xrightarrow{k \rightarrow \infty} \int_{|z| > 0} \psi(z) \mu_2(dz).$$

We also need the following results.

**Lemma 5.4.** *Assume (M), (J1'), (J2') and let  $v \in C^{1,\theta}(\overline{B_R})$  for some  $\theta \in (0, 1)$ . Then*

(a) *the function  $x \mapsto L_k^R[v, Dv](x)$  belongs to  $C^{0,\theta}(\overline{B_R})$ ;*

(b) *if  $v_n \rightarrow v$  in  $C^{1,\theta}(\overline{B_R})$ , then  $L_k^R[v_n, Dv_n] \rightarrow L_k^R[v, Dv]$  in  $C^{0,\theta}(\overline{B_R})$ .*

*Proof.* (a) We only do the proof for the  $L_1$ -term since the  $L_2$ -case is similar but easier. Below  $x, y \in \overline{B_R}$ ,  $\frac{1}{k} < |z| < k$  (the support of  $\mu_{1,k}$ ), and  $C_*$  will denote all constants (that may vary from line to line) depending only on  $R, k, \varepsilon, v, j_i, \mu_i, N$ . Since  $v$  is  $C^{1,\theta}$ ,

$$|v(x)| + |Dv(x)| + \frac{|Dv(x) - Dv(y)|}{|x - y|^\theta} \leq \|v\|_{C^{1,\theta}}.$$

Then,  $|v(x) - v(y)| \leq C_*|x - y|$  and by Lipschitz continuity of  $P_R$  and assumption (J2'),

$$\begin{aligned} \left| v\left(P_R(x + j_1(Dv(x), z))\right) - v\left(P_R(y + j_1(Dv(y), z))\right) \right| &\leq C_* (|x - y| + |Dv(x) - Dv(y)|), \\ |Dv(x) \cdot j_1(Dv(x), z) - Dv(y) \cdot j_1(Dv(y), z)| &\leq C_* |Dv(x) - Dv(y)|. \end{aligned}$$

Thus, all these quantities are controled by  $C_*|x - y|^\theta$ . Since the measure  $\mu_{1,k}$  is bounded and supported in  $\frac{1}{k} < |z| < k$ , it then follows that

$$|L_{1,k}^R[v, Dv](x) - L_{1,k}^R[v, Dv](y)| \leq C_*|x - y|^\theta \mu_{1,k}(\mathbb{R}^N),$$

and the proof of (a) is complete.

(b) By assumption

$$|v_n(x) - v(y)| + \frac{|v_n(x) - v(y)|}{|x - y|} + \frac{|Dv_n(x) - Dv(y)|}{|x - y|^\theta} \leq \|v_n - v\|_{C^{1,\theta}} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

By similar computations as above, we end up with

$$|L_k^R[v_n, Dv_n](x) - L_k^R[v, Dv](y)| \leq C_* \|v_n - v\|_{C^{1,\theta} \mu_{1,k}(\mathbb{R}^N)} |x - y|^\theta,$$

for a  $C_*$  independent of  $x, y, n$ . It follows that  $L_k^R[v_n, Dv_n] \rightarrow L_k^R[v, Dv]$  in  $C^{0,\theta}(\overline{B}_R)$  as  $n \rightarrow \infty$ .  $\square$

We can now prove an existence result for the approximate problem (5.7).

**Proposition 5.5.** *Assume (M), (J1'), (J2'), (F5'), and let  $\varepsilon, R, M > 0, k \in \mathbb{N}$ . Then there exists a classical solution  $u \in C^2(\overline{B}_R)$  of (5.7).*

*Proof.* The proof is based on Schauder's fixed point theorem (cf. e.g. [17, Corollary 11.2]).

1) Let  $X := C^{1,\theta_0}(\overline{B}_R)$  for a fixed  $\theta_0 \in (0, 1)$ , and define

$$\mathcal{C} = \{w \in X : \|w\|_\infty \leq M + \|f\|_\infty\}.$$

Note that  $\mathcal{C}$  is a convex and closed subset of  $X$ . On  $\mathcal{C}$  we now define a map  $\mathcal{T} = \mathcal{T}_{\varepsilon, R, M, k}$  in the following way: for every  $v \in \mathcal{C}$ ,  $u = \mathcal{T}(v)$  is the classical solution of the Dirichlet problem

$$\begin{cases} u - \varepsilon \Delta u = T_M [L_k^R[v, Dv]] + f(x) & \text{in } B_R, \\ u = 0 & \text{on } \partial B_R. \end{cases} \quad (5.8)$$

When  $v \in X$ ,  $L_k^R[v, Dv] \in C^{0,\theta_0}$  by Lemma 5.4, and then by the definition of  $T_M$  and (F5'),

$$w := T_M [L_k^R[v, Dv]] + f(x) \in C^{0,\theta_0}(\overline{B}_R).$$

Since  $B_R$  is a smooth domain, classical results ([17, Corollary 6.9]) then tell us that there exists a unique classical solution  $u \in C^{2,\theta_0}(\overline{B}_R) (\subset X)$  of (5.8). Moreover, by the maximum principle and the definition of  $T_M$ ,  $\|u\|_\infty \leq M + \|f\|_\infty$ . We conclude that  $\mathcal{T}$  is a well-defined map from  $\mathcal{C}$  into  $\mathcal{C}$ .

2) We show that  $\mathcal{T} : \mathcal{C} \rightarrow \mathcal{C}$  is continuous with respect to norm of  $X = C^{1,\theta_0}(\overline{B}_R)$ . Take a sequence  $\{v_n\} \subset \mathcal{C}$  such that  $v_n \rightarrow v$  in  $X$ . By subtracting the equations for  $u_n$  and  $u_p$ , we see that  $w := u_n - u_p$  is a classical solution of

$$\begin{cases} w - \Delta w = T_M [L_k^R[v_n, Dv_n]] - T_M [L_k^R[v_p, Dv_p]] =: g_{n,p}, & \text{in } B_R, \\ w = 0, & \text{in } \partial B_R. \end{cases}$$

By the maximum principle, we then find that

$$\|w\|_{C^0(\overline{B}_R)} \leq \|g_{n,p}\|_{C^0(\overline{B}_R)},$$

and by standard  $C^{2,\theta_0}$ -theory (e.g. [17, Thm 6.6]),

$$\|u_n - u_p\|_{C^{2,\theta_0}(\overline{B}_R)} = \|w\|_{C^{2,\theta_0}(\overline{B}_R)} \leq C \left( \|w\|_{C^0(\overline{B}_R)} + \|g_{n,p}\|_{C^{0,\theta_0}(\overline{B}_R)} \right).$$

Hence, since  $T_M[L_k^R[v_n, Dv_n]] \rightarrow T_M[L_k^R[v, Dv]]$  in  $C^{0, \theta_0}(\overline{B_R})$  by Lemma 5.4 (b), it follows that  $(u_n)$  is a Cauchy sequence in  $C^{2, \theta_0}(\overline{B_R})$  and hence also in  $X$ . By completeness, the limit  $u$  exists and belongs to  $C^{2, \theta_0}(\overline{B_R}) \cap \mathcal{C}$  since  $\mathcal{C}$  is closed in  $X$ .

By the  $C^{2, \theta_0}$  convergence of  $u_n$ , the  $C^{1, \theta_0}$  convergence of  $v_n$ , and Lemma 5.4, we can pass to the limit in the equation to see that  $u = \mathcal{T}(v)$ . It follows that  $\mathcal{T}(v_n) \rightarrow \mathcal{T}(v)$  in  $X$ , and we conclude that  $\mathcal{T}$  is continuous in  $\mathcal{C}$ .

**3)** We now show that  $\mathcal{T}(\mathcal{C})$  is relatively compact. Take any sequence  $\{u_n\} \subset \mathcal{T}(\mathcal{C})$ . Then there exists a sequence  $\{v_n\} \subset \mathcal{C}$  such that  $u_n = \mathcal{T}(v_n)$ . It follows that

$$u_n - \varepsilon \Delta u_n = g_n,$$

where  $|g_n| \leq M + \|f\|_\infty$ . By the maximum principle and  $W^{2,p}$ -theory (e.g. [17, Thm 9.11]), we have the following two a priori estimates for any  $1 < p < \infty$ ,

$$\begin{aligned} \|u_n\|_{L^\infty(B_R)} &\leq M + \|f\|_\infty, \\ \|u_n\|_{W^{2,p}(B_R)} &\leq C \left( \|u_n\|_{L^p(B_R)} + \|g_n\|_{L^p(B_R)} \right) \leq 2C|B_R|^{1/p} (M + \|f\|_\infty). \end{aligned}$$

By compact embeddings of Sobolev spaces [17, Thm 7.26], we can extract a subsequence  $u_{\varphi(n)}$  converging in  $C^{1, \theta}(\overline{B_R})$  for any  $\theta \in (0, 1)$ , in particular for  $\theta = \theta_0$ . This provides a subsequence which converges in  $X$ , and proves the claim.

**4)** By Schauder's fixed point theorem, there exists a function  $u \in \mathcal{C}$  such that  $u = \mathcal{T}(u)$ , which means that we have a  $C^2(\overline{B_R})$ -solution of (5.7) and the proof is complete.  $\square$

We proceed to prove existence under the restrictive assumptions **(J1')** and **(J2')**. We will need the following result.

**Lemma 5.6.** *Assume **(M)**, **(J1')**,  $f, g \in C^0(\overline{B_R})$ . Let  $u$  and  $v$  be  $C^2(\overline{B_R})$  solutions of*

$$u - T_M[L_k^R[u, Du]] - \varepsilon \Delta u \leq f(x) \quad \text{and} \quad v - T_M[L_k^R[v, Dv]] - \varepsilon \Delta v \geq g(x) \quad \text{in } B_R.$$

*If  $u \leq v$  on  $\partial B_R$  and  $f \leq g$  in  $B_R$ , then  $u \leq v$  in  $\overline{B_R}$ .*

*Remark 5.7.* The result is a comparison result for smooth sub and supersolutions of (5.7). It implies uniqueness of classical solutions of (5.7).

*Proof.* Let  $w := u - v$  and  $\bar{m} := \max_{\overline{B_R}} w$ . If this max is attained at  $x_0 \in \partial B_R$ , since  $u \leq v$  there, we have  $\bar{m} \leq 0$ . Otherwise, there is an interior point  $x_0 \in B_R$  such that  $m = w(x_0)$ . By assumption,

$$w(x_0) - \left\{ T_M[L_k^R[u, Du]](x_0) - T_M[L_k^R[v, Dv]](x_0) \right\} - \varepsilon \Delta w(x_0) \leq (f - g)(x_0).$$

Since  $x_0$  is a maximum point, and  $u, v$  are smooth,  $Dw(x_0) = (Du - Dv)(x_0) = 0$ ,  $\Delta w(x_0) \geq 0$ , and  $(u - v)(x_0) \geq (u - v)(y)$  for all  $y \in \overline{B_R}$ . By the latter inequality,

$$u\left(P_R(x_0 + j(Du(x_0, z)))\right) - u(x_0) \leq v\left(P_R(x_0 + j(Dv(x_0, z)))\right) - v(x_0),$$

and hence

$$L_k^R[u, Du](x_0) - L_k^R[v, Dv](x_0) \leq 0.$$

Since  $T_M$  is a non-decreasing function and  $f \leq g$ , we can conclude that

$$\bar{m} = w(x_0) \leq 0,$$

and the proof is complete because in either case, we get  $\bar{m} \leq 0$ .  $\square$

**Corollary 5.8.** *If  $u_{R,M,k,\varepsilon}$  is the solution of (5.7), then  $\|u_{R,M,k,\varepsilon}\|_{L^\infty(B_R)} \leq \|f\|_{L^\infty(\mathbb{R}^N)}$ .*

*Proof.* Follows from Lemma 5.6 with  $-\|f\|_\infty/u/\|f\|_\infty$  as subsolution/solution/supersolution.  $\square$

**Proposition 5.9.** *Assume (M), (J1'), (J2'), (J2), and (F5'). Then there exists a unique viscosity solution  $u \in C_b(\mathbb{R}^N)$  of (1.6).*

*Proof.* Let  $R > 0$ ,  $k = M = 1/\varepsilon = n \in \mathbb{N}$ , and  $u_{R,n}$  be the corresponding solution of (5.7) given by Proposition 5.5. Using the ‘‘half relaxed limit’’ method, we first we send  $R \rightarrow \infty$  and then  $n \rightarrow \infty$  and show that we can obtain from  $u_{R,n}$  a function  $u$  which is the viscosity solution of (1.6).

1) *Claim:* For every  $n \in \mathbb{N}$ , the functions

$$\bar{u}_n(x) := \limsup_{y \rightarrow x, R \rightarrow \infty} u_{R,n}(y) \quad \text{and} \quad \underline{u}_n(x) := \liminf_{y \rightarrow x, R \rightarrow \infty} u_{R,n}(y)$$

are bounded viscosity sub- and supersolutions respectively of

$$u - \frac{1}{n} \Delta u - T_n \left[ L_n[u, Du] \right] = f \quad \text{in} \quad \mathbb{R}^N. \quad (5.9)$$

where  $L_k$  is defined as  $L$  in (1.1), but with the measure  $\mu_{i,k}$  replacing  $\mu_i$  for  $i = 1, 2$ . Moreover,

$$-\|f\|_\infty \leq \underline{u}_n(x) \leq \bar{u}_n(x) \leq \|f\|_\infty \quad \text{in} \quad \mathbb{R}^N.$$

*Proof of Claim:* First note that  $\bar{u}_n$  and  $\underline{u}_n$  are defined for every  $x \in \mathbb{R}^N$  since  $R \rightarrow \infty$ , they are semicontinuous and  $\underline{u}_n \leq \bar{u}_n$  by definition and bounded by  $\|f\|_\infty$  by Corollary 5.8. We show that  $\bar{u}$  is a subsolution of (5.9) according to Definition 4.4. Take any bounded test function  $\phi$  and any point  $x \in \mathbb{R}^N$  such that  $\bar{u} - \phi$  has a global maximum at  $x$ . We may as usual assume the maximum is strict. Then there exists a sequence  $y_R \rightarrow x$  of maximum points of  $u_{R,n} - \phi$  in  $\bar{B}_R$ , such that  $u_{R,n}(y_R) \rightarrow \bar{u}(x)$ . Take  $R$  big enough such that  $R > |x|$  and  $y_R \in B_R$  (since  $y_R \rightarrow x$ ,  $y_R$  cannot be located on  $\partial B_R$  for  $R$  big enough). Since  $y_R$  is a maximum point and  $u_{R,n}$  is smooth,  $Du_{R,n}(y_R) = D\phi(y_R)$ ,  $\Delta u_{R,n}(y_R) \leq \Delta\phi(y_R)$ , and (cf. the proof of Lemma 5.6)

$$L_n^R[u_{R,n}, Du_{R,n}](y_R) \leq L_n^R[\phi, D\phi](y_R).$$

Since  $u_{R,n}$  satisfies equation (5.7) at the point  $y_R$ , it then follows that

$$u_{R,n}(y_R) - \frac{1}{n} \Delta\phi(y_R) - T_n \left[ L_n^R[\phi, D\phi](y_R) \right] \leq f(y_R). \quad (5.10)$$

By the boundedness of  $y_R$ , the regularity of  $\phi$ , (J1'), and the definition of  $P_R$ ,

$$\phi \left( P_R(y_R + j_i(D\phi(y_R), z)) \right) \rightarrow \phi(x + j_i(D\phi(x), z)) \quad \text{as} \quad R \rightarrow \infty, \quad \text{for} \quad a.e. \quad z \quad \text{and} \quad i = 1, 2.$$

Hence, since this term is uniformly bounded,  $\mu_{i,k}$  is bounded, and (M) holds, we can use the dominated convergence theorem to conclude that

$$L_n^R[\phi, D\phi](y_R) \rightarrow L_n[\phi, D\phi](x) \quad \text{as} \quad R \rightarrow \infty.$$

By the regularity of  $\phi$  and the continuity of  $T_n$ , we can then pass to the limit as  $R \rightarrow \infty$  in (5.10) and find that

$$\bar{u}(x) - \frac{1}{n} \Delta\phi(x) - T_n \left[ L_n[\phi, D\phi](x) \right] \leq f(x).$$

We conclude that  $\bar{u}$  is a viscosity subsolution of (5.9), and in a similar way we can show that  $\underline{u}$  is viscosity supersolution of (5.9). The claim is proved.

2) We now pass to the limit as  $n \rightarrow \infty$ . We proceed as before, defining

$$\bar{u}(x) := \limsup_{y \rightarrow x, n \rightarrow \infty} \bar{u}_n(y) \quad \text{and} \quad \underline{u}(x) := \liminf_{y \rightarrow x, n \rightarrow \infty} \underline{u}_n(y),$$

where  $\bar{u}_n$  and  $\underline{u}_n$  are the uniformly bounded sub and supersolutions of (5.9) given by part 1). It immediately follows that  $-\|f\|_\infty \leq \underline{u} \leq \bar{u} \leq \|f\|_\infty$ .

We prove that  $\bar{u}$  is viscosity subsolution of (1.6). Again we take any smooth bounded test function  $\phi$  and global strict maximum point  $x$  of  $\bar{u} - \phi$  in  $\mathbb{R}^N$ . We have a sequence  $y_n \rightarrow x$  of maximum points of  $u_n - \phi$  such that  $u_n(y_n) \rightarrow \bar{u}(x)$ . Fix  $\delta > 0$  and choose  $n$  big enough so that  $1/n < \delta$ . We split  $L_{1,n}$  into  $L_{1,n} = L_{\delta,n} + L_n^\delta$ , where  $L_\delta$  and  $L^\delta$  are defined in the beginning of section 4. Since  $\bar{u}$  is a subsolution of (5.9), it follows that

$$u_n(y_n) - \frac{1}{n} \Delta \phi(y_n) - T_n [L_n[\phi, D\phi](y_n)] \leq f(y_n). \quad (5.11)$$

We pass to the limit in the different terms of this inequality. First, let

$$\psi_n(z) = \phi(y_n + j_1(D\phi(y_n), z)) - \phi(y_n) - D\phi(y_n) \cdot j_1(D\phi(y_n), z) \mathbf{1}_{|z| < 1}.$$

Since  $D^2\phi$  is continuous,  $y_n \rightarrow x$  and  $D\phi(y_n) \rightarrow D\phi(x)$  are bounded, a Taylor expansion and **(J1')** reveals that there is a  $C$  independent of  $n$  and  $z$  such that

$$|\psi_n(z)| \leq \sup_{t \in (0,1)} \frac{1}{2} \left| D^2\phi \left( y_n + t j_1(D\phi(y_n), z) \right) \right| |j_1(D\phi(y_n), z)|^2 \leq C|z|^2 \quad \text{for } |z| < 1.$$

By regularity of  $\phi$  and **(J2')**, the  $\psi_n$  converge uniformly for  $|z| \leq \delta$  to

$$\psi(z) := \phi(x + j_1(D\phi(x), z)) - \phi(x) - D\phi(x) \cdot j_1(D\phi(x), z) \mathbf{1}_{|z| < 1},$$

and hence we use Lemma 5.3 (c) to conclude that

$$L_{\delta,n}[\phi, D\phi](y_n) \rightarrow L_\delta[\phi, D\phi](x).$$

Simpler but similar arguments, this time using Lemma 5.3 (b) and (d), show that

$$L_n^\delta[\phi, D\phi](y_n) \rightarrow L^\delta[\phi, D\phi](x) \quad \text{and} \quad L_{2,n}[\phi, D\phi](y_n) \rightarrow L_2[\phi, D\phi](x).$$

We conclude that  $L_n[\phi, D\phi](y_n) \rightarrow L[\phi, D\phi](x)$ , and since this limit is bounded, for  $n$  big enough

$$T_n[L_n[\phi, D\phi](y_n)] = L_n[\phi, D\phi](y_n) \rightarrow L[\phi, D\phi](x) \quad \text{as } n \rightarrow \infty.$$

In view of the regularity of  $\phi$  and  $f$ , we can then pass to the limit as  $n \rightarrow \infty$  in (5.11) to find that

$$\bar{u}(x) - L[\phi, D\phi](x) \leq f(x).$$

Hence  $\bar{u}$  is a viscosity subsolution of (1.6). Similar arguments show that  $\underline{u}$  is viscosity supersolution.

**3)** By the comparison result for semicontinuous viscosity solutions, Theorem 2.1 (a), it follows that  $\bar{u} \leq \underline{u}$ . Note that this result requires **(J2)**, see also Remark 5.2. Since the opposite inequality holds by part 2),  $\bar{u} = \underline{u} =: u$ , and this function is continuous, bounded by  $\|f\|_\infty$ , and a viscosity solution of (1.6). The proof is complete.  $\square$

Now we show how to remove assumptions **(J1')**, **(J2')**, and **(F5')** and obtain an existence result in the general case. We do it in two steps, starting by removing **(J2')** and **(F5')** by a regularization argument (mollification), and then we remove **(J1')** by a truncation argument. We will need the following lemma, whose proof is given in Appendix C.

**Lemma 5.10.** *Assume **(M)**, **(J1')**, and **(J2)** and define  $j_{1,k}$  and  $j_{2,k}$  by*

$$j_{i,k}(p, z) := (\rho_k * j_i(\cdot, z))(p) = \int_{\mathbb{R}^N} \rho_k(q-p) j_i(q, z) dq \quad \text{for } i = 1, 2,$$

for a mollifier  $\rho_k(p) = k^N \rho(kp)$ ,  $0 \leq \rho \in C^\infty(\mathbb{R}^N)$  is symmetric with support in  $B_1$  and  $\int \rho = 1$ . Then the following holds:



(a) For any  $k \in \mathbb{N}$ ,  $j_{1,k}$  and  $j_{2,k}$  are locally bounded, and for any  $|p| \leq r$ ,  $|z| < 1$ , and  $k \in \mathbb{N}$ ,

$$|j_{1,k}(p, z)| \leq C_{r+1/k}|z|, \quad \text{where } C_r \text{ is the constant from (J1')}.$$

(b) For every  $k \in \mathbb{N}$  and  $r > 0$ , there is  $C_{k,r}$  such that for all  $|p|, |q|, |z| \leq r$ ,

$$|j_{i,k}(p, z) - j_{i,k}(q, z)| \leq C_{k,r}|p - q| \quad \text{for } i = 1, 2.$$

(c) For any  $r > 0$ ,  $|p|, |q| < r$  and  $k \in \mathbb{N}$ ,

$$\int_{|z|>0} |j_{1,k}(p, z) - j_{1,k}(q, z)|^2 d\mu_1(z) \leq \omega_{j,r+\frac{1}{k}}(p - q) \quad \text{where } \omega_{j,r} \text{ is the modulus from (J2)}.$$

(d) There exists  $\delta_0 > 0$  such that for any  $r > 0$  and  $\varepsilon > 0$  there exists  $\eta > 0$  such that

$$\sup_{|p|<r, k \in \mathbb{N}} \int_A |j_{1,k}(p, z)|^2 d\mu_1(z) < \varepsilon$$

for every Borel set  $A \subset \{0 < |z| < \delta_0\}$  such that  $\int_A |z|^2 \mu_1(dz) < \eta$ .

(e) Let  $L_{1,k}$  and  $L_{2,k}$  be defined as in (1.1) and (1.2) with  $j_1$  and  $j_2$  replaced by  $j_{1,k}$  and  $j_{2,k}$ . For all  $\phi \in C_b \cap C^2$  and  $x_k \rightarrow x$ ,

$$L_{i,k}[\phi, D\phi(x_k)](x_k) \rightarrow L_i[\phi, D\phi(x)](x) \quad \text{for } i = 1, 2.$$

Here is the general existence result for equation (1.6).

**Proposition 5.11.** *Assume (M), (J1)–(J3) and (F1)–(F5). Then there exists a viscosity solution of (1.6).*

*Proof.* We proceed in two steps.

1) Assume in addition (J1') holds. We approximate  $j_i$  and  $f$  by

$$j_{i,k} \text{ as defined in Lemma 5.10} \quad \text{and} \quad f_k = \rho_k * f,$$

where  $k \in \mathbb{N}$  and  $\rho_k$  is the mollifier defined in Lemma 5.10. By Lemma 5.10,  $j_{i,k}$  satisfy (J1)–(J3) and (J1')–(J2'), while assumption (F5) and properties of mollifiers imply that  $f_k$  is bounded and Lipschitz continuous in  $\mathbb{R}^N$  (so (F5') holds) and converges uniformly to  $f$  in  $\mathbb{R}^N$  as  $k \rightarrow \infty$ .

Let  $L_k$  be defined as  $L$  with  $j_{i,k}$  replacing  $j_i$ , and consider the problem

$$u - L_k[u, Du] = f_k \quad \text{in } \mathbb{R}^N.$$

By the above discussion, this problem satisfies all the assumptions of Proposition 5.9 for every  $k \in \mathbb{N}$ , and hence there exist solutions  $u_k$  of this problem for every  $k \in \mathbb{N}$ . By the comparison principle (Theorem 2.1 (a)),

$$\|u_k\|_\infty \leq \|f_k\|_\infty \leq \|f\|_\infty.$$

As in the proof of Proposition 5.9, we then define the half-relaxed limits

$$\bar{u}(x) = \limsup_{y \rightarrow x, k \rightarrow \infty} u_k(y) \quad \text{and} \quad \underline{u}(x) = \liminf_{y \rightarrow x, k \rightarrow \infty} u_k(y),$$

prove that they are viscosity sub and supersolutions of (1.6), and conclude by the comparison result Theorem 2.1 (a) that  $\bar{u} = \underline{u} = u$  is a viscosity solution of (1.6). We just sketch the argument for  $\bar{u}$  being a subsolution. Take a smooth test-function  $\phi$  such that  $\bar{u} - \phi$  has a strict maximum at  $x \in \mathbb{R}^N$ . Then there exists a sequence  $y_k \rightarrow x$  such that  $u_k - \phi$  has a maximum at  $x_k$ . Using the

subsolution property for  $u_k$ , the fact that  $L_k[\phi, D\phi](x_k) \rightarrow L[\phi, D\phi](x)$  (Lemma 5.10 (e)) and that  $f_k(x_k) \rightarrow f(x)$ , we arrive at

$$\bar{u}(x) - L[\phi, D\phi](x) \leq f(x),$$

hence  $\bar{u}$  is a subsolution of (1.6). This completes the existence proof under the assumptions **(M)**, **(J1')**, **(J2)**, **(J3)**, and **(F5)**.

**2)** We remove the **(J1')** condition through a truncation procedure for the  $j_1$ -term:

$$j_1^M(p, z) := |z| T_M\left(\frac{j_1(p, z)}{|z|}\right),$$

where  $T_M(x) = x$  if  $|x| \leq M$  and  $T_M(x) = M\frac{x}{|x|}$  if  $|x| > M$ . In this case is easy to see that  $j_1^M$  satisfies **(J1)**–**(J3)** and in addition **(J1')** with  $C_r = M$ . Furthermore, for almost any  $z$ ,

$$j_1^M(p, z) \rightarrow j_1(p, z) \quad \text{locally uniformly in } p.$$

Hence by part 1), for any  $M > 0$  there exists a solution  $u_M$  of

$$u - L^M[u, Du] = f \quad \text{in } \mathbb{R}^N,$$

where  $L^M$  is defined as  $L$  (cf. (1.1)) but with  $(j_1^M, j_2)$  replacing  $(j_1, j_2)$ . As in part 1), we define the half-relaxed limits

$$\bar{u}(x) = \limsup_{y \rightarrow x, M \rightarrow \infty} u_M(y) \quad \text{and} \quad \underline{u}(x) = \liminf_{y \rightarrow x, M \rightarrow \infty} u_M(y),$$

prove that they are viscosity sub and supersolutions of (1.6), and conclude by the comparison result Theorem 2.1 (a) that  $\bar{u} = \underline{u} = u$  is a viscosity solution of (1.6). In view of previous arguments, the only thing we need to check is that

$$L_1^M[\phi, D\phi](x_M) \rightarrow L_1[\phi, D\phi](x) \quad \text{when } x_M \rightarrow x.$$

To do so, we use note that the integrand converges pointwise a.e. by the regularity of  $\phi$  and the local uniform convergence of  $j_1^M$  in the  $p$ -variable. Since  $|j_1^M(p, z)| \leq |j_1(p, z)|$ ,  $j_1^M$  satisfies the equi-integrability condition **(J3)** uniformly in  $M$ . Hence we may use Vitali's convergence theorem (cf. Appendix A and Remark 2.4-(b)) to pass the limit  $M \rightarrow \infty$  inside the integral over  $|z| < \delta$ . Passage to the limit in the integral over  $|z| \geq \delta$  is done using the dominated convergence theorem since  $\phi$  is bounded and  $\mu_1$  is not singular on this domain. The proof is complete.  $\square$

*Proof of Theorem 2.3.* Uniqueness and uniform continuity follow from Corollary 2.2. Existence of solutions of (1.6) follows from Proposition 5.11. Existence for (1.3) follows in a similar way as for (1.6) where all difficulties are already present. We only give a very brief sketch of the proof. We start by an approximate problem (5.7). In the general case it is to find  $u \in C^2(\bar{B}_R)$  such that

$$\begin{cases} \gamma u + T_M \left[ -\gamma u + F(u, Du, L_k^R[u, Du]) \right] - \varepsilon \Delta u = f(x), & x \in B_R(0), \\ u = 0, & x \in \partial B_R(0), \end{cases}$$

where  $L_k^R$  is defined below (5.7). Under the same assumptions as the linear case in addition to the assumption that  $F$  is also locally Lipschitz, we obtain existence of solutions of this problem following step by step the fixed point argument of the proof of Proposition 5.5. To get an existence result for equation (1.3), we follow the approximation and limit procedures given in the proofs of Proposition 5.9 and 5.11. The only slight difference is that we also need an approximation argument for  $F$  (e.g. by mollification again), and when we pass to the limit, we use this time the strong comparison result Theorem 2.1-(b) for the limit equation. It is straight forward to check that this will work out and produce a bounded viscosity solution of (1.3).  $\square$

**5.3. Proof of Theorem 2.6 (local limits).** Theorem 2.6 is a consequence of the half-relaxed limit method and comparison. Define

$$\bar{u}(x) := \limsup_{\varepsilon \rightarrow 0, y \rightarrow x} u_\varepsilon(y) \quad \text{and} \quad \underline{u}(x) := \liminf_{\varepsilon \rightarrow 0, y \rightarrow x} u_\varepsilon(y).$$

In the quasi-linear case we have the following result.

**Lemma 5.12.** *Assume the assumptions of Theorem 2.6-(a) hold. Then  $\bar{u}$  is a viscosity subsolution of (2.2) and  $\underline{u}$  is a viscosity supersolution of (2.2).*

*Proof.* Let  $\phi$  be a smooth bounded function such that  $\bar{u} - \phi$  has a global maximum at  $x$ . We may assume the maximum is strict and that there is a sequence  $\{y_\varepsilon\}_\varepsilon$  of global maximum points of  $u_\varepsilon - \phi$  such that  $y_\varepsilon \rightarrow x$  and  $u_\varepsilon(y_\varepsilon) \rightarrow \bar{u}(x)$  as  $\varepsilon \rightarrow 0$ . We let  $L_{1,\varepsilon} = L_{\varepsilon,\delta} + L_\varepsilon^\delta$  as in (4.1), and to see the localization effect, we also decompose  $L_{2,\varepsilon} = \tilde{L}_{\varepsilon,\delta} + \tilde{L}_\varepsilon^\delta$ . By the maximum point property  $\tilde{L}_{\varepsilon,\delta}[u_\varepsilon, D\phi](y_\varepsilon) \leq \tilde{L}_{\varepsilon,\delta}[\phi, D\phi](y_\varepsilon)$ , and then since  $u_\varepsilon$  is a subsolution of (2.1),

$$-L_{\varepsilon,\delta}[\phi, D\phi](y_\varepsilon) - L_\varepsilon^\delta[u_\varepsilon, D\phi](y_\varepsilon) - \tilde{L}_{\varepsilon,\delta}[\phi, D\phi](y_\varepsilon) - \tilde{L}_\varepsilon^\delta[u_\varepsilon, Du_\varepsilon](y_\varepsilon) + u_\varepsilon(y_\varepsilon) \leq f(y_\varepsilon). \quad (5.12)$$

Then by a Taylor expansion and **(J4)**, for  $i = 1, 2$ ,

$$j_i(D\phi(y_\varepsilon), z) = \sigma_i(D\phi(y_\varepsilon))z + o(z) \quad \text{as} \quad z \rightarrow 0,$$

where  $o(z)$  is independent of  $\varepsilon > 0$ . Another Taylor expansion and hypotheses **(M $_\varepsilon$ )** applied to  $Y = \sigma_1(D\phi(x))^T D^2\phi(x) \sigma_1(D\phi(x)) \in \mathbb{R}^{P \times P}$  then gives that

$$\begin{aligned} L_{\varepsilon,\delta}[\phi, D\phi](y_\varepsilon) &= \frac{1}{2} \int_{|z| < \delta} z^T \left[ \sigma_1(D\phi(y_\varepsilon))^T D^2\phi(y_\varepsilon) \sigma_1(D\phi(y_\varepsilon)) \right] z \, d\mu_{1,\varepsilon}(z) + o_\delta(1) \\ &= \frac{1}{2} \text{tr} \left[ A_1^T \left[ \sigma_1(D\phi(x))^T D^2\phi(x) \sigma_1(D\phi(x)) \right] A_1 \right] + o_\varepsilon(1) + o_\delta(1) \\ &= \frac{1}{2} \text{tr} \left[ \tilde{\sigma}_1(D\phi(x)) \sigma_1(D\phi(x))^T D^2\phi(x) \right] + o_\varepsilon(1) + o_\delta(1), \end{aligned}$$

where for the last line, we use that  $\tilde{\sigma}_1(x, p) = \sigma_1(x, p)A_1$  is a  $N \times N$  matrix as is  $D^2\phi(x)$ , so that we can use the property  $\text{tr}(M_1M_2) = \text{tr}(M_2M_1)$ . Here the  $o_\delta(1)$ -term is independent of  $\varepsilon$  since the measure  $|z|^2 \mu_{1,\varepsilon}(dz)$  has a uniformly bounded mass. Similarly, by **(J4)** and **(M $_\varepsilon$ )**,

$$\begin{aligned} \tilde{L}_{\varepsilon,\delta}[\phi, D\phi](x) &= \frac{1}{2} \int_{|z| < \delta} z^T \left[ \sigma_2(D\phi(x))^T D^2\phi(x) \sigma_2(D\phi(x)) \right] z \, d\mu_{2,\varepsilon}(z) \\ &\quad + \int_{|z| < \delta} D\phi(x) \sigma_2(D\phi(x)) z \, d\mu_{2,\varepsilon}(z) + o_\varepsilon(1) + o_\delta(1) \\ &= \frac{1}{2} \text{tr} \left[ A_2^T \left[ \sigma_2(D\phi(x))^T D^2\phi(x) \sigma_2(D\phi(x)) \right] A_2 \right] \\ &\quad + \sigma_2(D\phi(x)) D\phi(x) \cdot a + o_\varepsilon(1) + o_\delta(1) \\ &= \frac{1}{2} \text{tr} \left[ \tilde{\sigma}_2(D\phi(x)) \tilde{\sigma}_2(D\phi(x)) D^2\phi(x) \right] + b(D\phi(x)) \cdot D\phi(x) + o_\varepsilon(1) + o_\delta(1) \end{aligned}$$

By **(J1)** and continuity,  $D\phi(y_\varepsilon)$  and  $j(D\phi(y_\varepsilon), z)$  are uniformly bounded in  $\varepsilon > 0$  for  $|z| < 1$ , and by **(F5)** and Corollary 2.2-(c),  $u_\varepsilon$  is also uniformly bounded:

$$\|u_\varepsilon\|_\infty \leq \|f\|_\infty.$$

By  $(\mathbf{M}_\varepsilon)$  it then follows that

$$L_\varepsilon^\delta[u_\varepsilon, D\phi](y_\varepsilon) + \tilde{L}_\varepsilon^\delta[u_\varepsilon, D\phi](y_\varepsilon) = O\left(\int_{|z|\geq\delta} d\mu_\varepsilon(z)\right) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

and  $\delta > 0$  is fixed.

Sending first  $\varepsilon \rightarrow 0$  and then  $\delta \rightarrow 0$  in (5.12) then leads to

$$-L_0(D\phi(x), D^2\phi(x)) + \bar{u}(x) \leq f(x),$$

and hence  $\bar{u}$  is a viscosity subsolution of (2.2). In a similar way we can show that  $\underline{u}$  is a viscosity supersolution of (2.2). We omit the proof.  $\square$

Then it is enough to invoke the comparison principle for the limit (local) equation to conclude that Theorem 2.6-(a) holds:

*Proof of Theorem 2.6-(a).* By Lemma 5.12  $\bar{u}$  is a subsolution and  $\underline{u}$  is a supersolution of (2.2), hence  $\bar{u} \leq \underline{u}$  by the comparison principle for (2.2), see [13] or Lemma 5.14 below. Since  $\underline{u} \leq \bar{u}$  by definition,  $\bar{u} = \underline{u} =: u$ , and hence  $u_\varepsilon \rightarrow u$  point-wise,  $u$  is continuous and a viscosity solution of (2.2). Now we show that the convergence is locally uniform. Fix any  $R > 0$  and take  $x_\varepsilon \in B_R(0)$  such that

$$\max_{|x|\leq R} (u_\varepsilon(x) - u(x)) = u_\varepsilon(x_\varepsilon) - u(x_\varepsilon)$$

for any  $\varepsilon > 0$ . Since  $|x_\varepsilon| \leq R$ , there exists a convergent subsequence  $x_\varepsilon \rightarrow \bar{x}$  for some  $|\bar{x}| \leq R$ . By the continuity of  $u$ , the definition of  $\bar{u}$ , and the fact that  $u = \bar{u}$ , it follows that

$$\limsup_{\varepsilon \rightarrow 0} \max_{|x|\leq R} (u_\varepsilon(x) - u(x)) = \limsup_{\varepsilon \rightarrow 0} (u_\varepsilon(x_\varepsilon) - u(x_\varepsilon)) \leq \bar{u}(\bar{x}) - u(\bar{x}) = 0.$$

A similar argument shows that  $\liminf_{\varepsilon \rightarrow 0} \max_{|x|\leq R} (u_\varepsilon(x) - u(x)) \geq 0$ . Combined with similar arguments for  $-(u_\varepsilon - u)$ , this shows that

$$\lim_{\varepsilon \rightarrow 0} \max_{|x|\leq R} |u_\varepsilon(x) - u(x)| = 0 \quad \text{for all } R > 0,$$

and we are done.  $\square$

In the fully nonlinear case, the strategy is the same and we begin with a half-relaxed limit result.

**Lemma 5.13.** *Assume the assumptions of Theorem 2.6-(b) hold. Then  $\bar{u}$  is a viscosity subsolution of (1.5) and  $\underline{u}$  is a viscosity supersolution of (1.5).*

*Proof.* Take a test-function  $\phi$  such that  $\bar{u} - \phi$  has a maximum at  $\bar{x}$  that we can assume to be strict. Hence  $u_\varepsilon - \phi$  also has a maximum at some  $x_\varepsilon$ , and  $x_\varepsilon \rightarrow \bar{x}$  and  $u_\varepsilon(x_\varepsilon) \rightarrow \bar{u}(\bar{x})$  as  $\varepsilon \rightarrow 0$ . Hence  $\tilde{L}_{\varepsilon,\delta}[u_\varepsilon, D\phi](x_\varepsilon) \leq \tilde{L}_{\varepsilon,\delta}[\phi, D\phi](x_\varepsilon)$ , and since  $u_\varepsilon$  is a subsolution and  $(\mathbf{F1})$  holds,

$$F(u_\varepsilon(x_\varepsilon), D\phi(x_\varepsilon), L_{\varepsilon,\delta}[\phi, D\phi](x_\varepsilon) + L_\varepsilon^\delta[u_\varepsilon, D\phi](x_\varepsilon) + \tilde{L}_{\varepsilon,\delta}[\phi, D\phi](x_\varepsilon) + \tilde{L}_\varepsilon^\delta[u_\varepsilon, D\phi](x_\varepsilon)) \leq f(x_\varepsilon).$$

As in the proof of Lemma 5.12,

$$L_\varepsilon\text{-terms} \rightarrow L_0(D\phi(\bar{x}), D^2\phi(\bar{x})) \quad \text{as } \varepsilon \rightarrow 0.$$

By the continuity of  $F$ , we then send  $\varepsilon \rightarrow 0$  to find that

$$F(\bar{u}(\bar{x}), D\phi(\bar{x}), L_0(D\phi(\bar{x}), D^2\phi(\bar{x}))) \leq f(\bar{x}),$$

which means that  $\bar{u}$  is a subsolution of (1.5). In a similar way we can show that  $\underline{u}$  is a supersolution of (1.5).  $\square$

Then we prove a comparison result for the limit equation.

**Lemma 5.14.** *Under the assumptions of Theorem 2.6-(b), if  $u$  is a bounded usc subsolution of (1.5) and  $v$  is a bounded lsc supersolution of (1.5), then  $u \leq v$  in  $\mathbb{R}^N$ .*

To prove this result, first note that by **(F1)** and **(F2)**, the non-linearity in (1.5),

$$H(x, u, p, X) := F(u, p, L_0(p, X)) - f(x), \quad (5.13)$$

is strictly increasing in  $u$  and nonincreasing in  $X$ . Moreover, it is straightforward to check that we also have the following result.

**Lemma 5.15.** *Assume **(J4)** and **(F3)** hold,  $H$  is defined in (5.13), and  $M, r, \hat{r}, R, \varepsilon > 0$ . Let  $x, y \in \mathbb{R}^N$ ,  $|u| \leq M$ ,  $p = 2(x - y)/\varepsilon^2 + o_R(1)$ ,  $|p| < r$ ,  $X, Y \in \mathcal{S}_N$ , and  $-\hat{r} \leq X \leq Y \leq \hat{r}$  such that*

$$\left(-\frac{8}{\varepsilon^2} + o_R(1)\right) \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \leq \begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix} \leq \frac{1}{\varepsilon^2} \begin{pmatrix} I & -I \\ -I & I \end{pmatrix} + o_R(1) \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix}.$$

*Then there are moduli of continuity  $\omega_M, \omega_{M, r, \hat{r}}$  such that*

$$H\left(y, u, p - O\left(\frac{1}{R}\right), Y\right) - H\left(x, u, p + O\left(\frac{1}{R}\right), X\right) \leq \omega_M(x - y) + \frac{1}{\varepsilon^2} \omega_{M, r, \hat{r}}(o_R(1)) + o_R(1).$$

*Proof of Lemma 5.14.* In view of our assumptions and Lemma 5.15, the proof is standard. It can be obtained by following the line of reasoning of the proof of Theorem 5.1 in [13]. We omit the details.  $\square$

*Proof of Lemma 2.6-(b).* In view of Lemma 5.13 and 5.14, we can conclude the proof exactly as for Theorem 2.6-(a) above.  $\square$

## 6. PARABOLIC PROBLEMS

In this section we extend the results to the case of quasilinear and fully nonlinear parabolic equations,

$$u_t - L[u, Du] = f(x, t) \quad \text{in} \quad Q_T := \mathbb{R}^N \times (0, T), \quad (6.1)$$

$$u_t + F(u, Du, L[u, Du]) = f(x, t) \quad \text{in} \quad Q_T, \quad (6.2)$$

with initial data

$$u(x, 0) = u_0(x) \quad \text{in} \quad \mathbb{R}^N.$$

These extensions are straightforward since the time variable does not play an important role here. We keep the same notation, definitions, and most of the assumptions as in the previous sections. However, we need the following parabolic versions of **(F4)** and **(F5)**:

**(F6)**  $u_0 \in UC(\mathbb{R}^N)$  and  $f \in UC(Q_T)$ .

**(F6')**  $u_0$  and  $f$  satisfy **(F6)**, are bounded, and the quantities in **(F2)**, **(F3)** are independent of  $M$ .

In the parabolic case, we could also relax assumption **(F2)** in the classical way. We have the following parabolic version of the existence and comparison results.

**Theorem 6.1.** (Comparison results)

(a) [Quasi-linear case] Assume **(M)**, **(J1)**–**(J2)** and **(F6)**. If  $u$  is a bounded usc subsolution of (6.1),  $v$  a bounded lsc supersolution of (6.1), and  $u(x, 0) \leq v(x, 0)$  in  $\mathbb{R}^N$ , then  $u \leq v$  in  $\overline{Q}_T$ .

(b) [Fully non-linear case] Assume **(M)**, **(J1)**–**(J2)**, **(F1)**–**(F3)**, and **(F6)**. If  $u$  is a bounded usc viscosity subsolution of (6.2),  $v$  a bounded lsc viscosity supersolution of (6.2), and  $u(x, 0) \leq v(x, 0)$  in  $\mathbb{R}^N$ , then  $u \leq v$  in  $\overline{Q}_T$ .

*Sketch of proof.* We need to double the variables in time as well as in space and consider

$$\Phi_{\varepsilon, \beta, R, c}(x, y, s, t) := u(x, t) - v(y, s) - \phi(x, t, y, s)$$

where

$$\phi(x, t, y, s) = \frac{1}{\varepsilon^2} \varphi(x - y) + \frac{|t - s|^2}{\beta^2} + \psi\left(\frac{x}{R}\right) + \psi\left(\frac{y}{R}\right) + \frac{c}{T - t},$$

and  $\varphi$  and  $\psi$  are defined in the proof of Theorem 2.1.

We assume by contradiction that  $\sup_{\mathbb{R}^N \times [0, T]}(u - v) > 0$ . It follows that  $\sup_{\mathbb{R}^N \times [0, T]} \Phi_{\varepsilon, \beta, R, c} > 0$  for  $\varepsilon, \beta, c$  small enough and  $R$  big enough. By definition,  $\Phi_{\varepsilon, \beta, R, c}$  will attain its supremum at some point  $(\bar{x}, \bar{t}, \bar{y}, \bar{s})$ , and since  $u(x, 0) \leq v(x, 0)$ , we may assume that both  $\bar{t}, \bar{s} > 0$ . Because of the term  $c/(T - t)$ , we may also assume that  $\bar{t}, \bar{s} < T$ . Hence we may use the viscosity inequalities for  $u$  and  $v$ . After we have subtracted these inequalities, we send  $\beta \rightarrow 0$  to obtain  $\bar{s} = \bar{t} \in (0, T)$ . Since  $0 \leq \phi_t - \phi_s$ , we can drop this term and we are essentially back to inequality (5.6). At this stage we proceed as in the proof of Theorem 2.1, and show that

$$\lim_{c \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \lim_{R \rightarrow \infty} \lim_{\beta \rightarrow 0} \Phi_{\varepsilon, \beta, R, c} \leq 0.$$

This is a contradiction to our original assumption and the proof is complete.  $\square$

**Theorem 6.2** (Existence). *Under the assumptions of Theorem 6.1, **(F6')**, and **(J3)**, there exists a bounded viscosity solution of (6.2).*

*Sketch of proof.* We first assume **(J1')**, **(J2')** and that  $(x, t) \mapsto f(x, t)$ ,  $x \mapsto u_0(x)$  are bounded and Lipschitz continuous. The general result will follow from passage to the limit as in Proposition 5.11. Then we consider the parabolic version of the approximation we used in Section 5.2:

$$\begin{cases} u_t + T_M \left[ F(u, Du, L_k^R[u, Du]) \right] - \varepsilon \Delta u = f(x, t), & (x, t) \in B_R(0) \times (0, T), \\ u(x, t) = 0, & (x, t) \in \partial B_R(0) \times (0, T), \\ u(x, 0) = u_0(x), & x \in B_R(0). \end{cases}$$

Assuming this problem has a solution  $u_{M, R, k, \varepsilon}$ , we pass to the limit as  $R \rightarrow \infty$ ,  $k \rightarrow \infty$ ,  $M \rightarrow \infty$ ,  $\varepsilon \rightarrow 0$  as in the elliptic case. Using half-relaxed limits and comparison for the limit equation, we show that the sequence of solutions has a limit which is a solution of (6.2). Existence is then proved.

To prove that there exists a solution of the approximate problem we use Schauder's fixed point theorem and the argument given in Proposition 5.5 with some small modifications:

1) We use Schauder's fixed point theorem in the Banach space

$$X := H_{1+\theta_0}([0, T] \times \overline{B}_R)$$

for some  $\theta_0 \in (0, 1)$ . The space  $H_{1+\theta_0} (= C^{1+\theta_0, \frac{1+\theta_0}{2}})$  is a standard parabolic Hölder space where  $u \in H_{1+\theta_0}$  e.g. implies that  $Du \in H_{\theta_0}$ . See page 46 in [20] or Section 1.2.3 in [27] for the definition.

**2)** The time-dependent version of Lemma 5.4 remains valid: if  $v \in X$ , then  $(x, t) \mapsto L_k^R[v, Dv](x, t)$  belongs to  $H_{\theta_0}(\overline{B}_R \times [0, T])$ , and if  $v_n \rightarrow v$  in  $X$ , then  $L_k^R[v_n, Dv_n] \rightarrow L_k^R[v, Dv]$  in  $H_{\theta_0}(\overline{B}_R \times [0, T])$ .

**3)** The  $C^{2, \theta_0}$  regularity result that we use in Lemma 5.5 (step 2) is replaced by the parabolic  $H_{2+\theta_0}$  version in Theorem 4.28 in [20].

**4)** In Lemma 5.5 (step 3), instead of the  $W^{2,p}$ -theory we use the parabolic  $W_p^{2,1}$ -theory of Theorem 7.17 in [20]. We also use the compact embedding of  $W_p^{2,1}$  into  $X$  for  $p$  big enough, see Theorem 1.4.1 in [27].  $\square$

For the local limit result, we introduce the local parabolic equations

$$u_t - L_0(Du, D^2u) = f(x, t) \quad \text{in} \quad Q_T, \quad (6.3)$$

$$u_t + F(u, Du, L_0(Du, D^2u)) = f(x, t) \quad \text{in} \quad Q_T, \quad (6.4)$$

with an initial data  $u(x, 0) = u_0(x)$ . The result is the following:

**Theorem 6.3.** (Localization)

(a) [Quasi-linear case] *Under the assumptions of Theorem 6.1-(a),  $(\mathbf{F6}')$ ,  $(\mathbf{M}_\varepsilon)$ , and  $(\mathbf{J4})$ , any sequence of solutions  $u_\varepsilon$  of (2.1) converge locally uniformly in  $\overline{Q}_T$  as  $\varepsilon \rightarrow 0$  to the solution  $u$  of (6.3).*

(b) [Fully-nonlinear case] *Under the assumptions of Theorem 6.1-(b),  $(\mathbf{F6}')$ ,  $(\mathbf{M}_\varepsilon)$ ,  $(\mathbf{J4})$ , any sequence of solutions  $u_\varepsilon$  of (1.4) converge locally uniformly in  $\overline{Q}_T$  as  $\varepsilon \rightarrow 0$  to the solution  $u$  of (6.4).*

*Sketch of Proof.* We use uniform boundedness of  $u_\varepsilon$  and the half-relaxed limits

$$\overline{u}(x) := \limsup_{\varepsilon \rightarrow 0, y \rightarrow x, s \rightarrow t} u_\varepsilon(y, s) \quad \text{and} \quad \underline{u}(x) := \liminf_{\varepsilon \rightarrow 0, y \rightarrow x, s \rightarrow t} u_\varepsilon(y, s),$$

and prove that  $\overline{u}$  and  $\underline{u}$  are respectively sub and supersolutions of the local limit problem. Local uniform convergence is then obtained after proving that the limit problem satisfies the comparison principle. The proofs the comparison principles for (6.3) and (6.4) are similar to the proofs in stationary case with standard modification such as doubling also the time variables.  $\square$

#### ACKNOWLEDGEMENT

We would like to thank Luca Rossi for pointing out a critical error in the existence proof of an earlier version of this paper.

#### APPENDIX A. EQUI-INTEGRABILITY AND CONVERGENCE

We give the definition equi-integrability (also called uniform integrability) and a result that we have used many times in this paper, a generalization of the dominated convergence theorem due to Vitali. Our presentation follow [25] page 133.

**Definition A.1.** *Let  $(\Omega, \mathcal{E}, \mu)$  be a positive measure space. A family  $(f_i)_{i \in I} \subset L^1(\Omega, \mu)$  is equi-integrable if for any  $\varepsilon > 0$  there exists  $\eta > 0$  such that*

$$\sup_{i \in I} \int_A |f_i(z)| d\mu(z) < \varepsilon$$

*for every  $A \in \mathcal{E}$  such that  $\mu(A) < \eta$ .*

The Vitali convergence theorem is the following result:

**Proposition A.2.** *Assume  $(\Omega, \mathcal{E}, \mu)$  is a positive finite measure space and  $(f_n)_{n \in \mathbb{N}} \subset L^1(\Omega, \mu)$  an equi-integrable family such that  $f_n \rightarrow f$   $\mu$ -a.e.. Then  $f_n \rightarrow f$  in  $L^1(\Omega, \mu)$ .*

Note that the dominated convergence theorem is a consequence of this result (on finite measure spaces!) since domination by a fixed, integrable function implies equi-integrability.

### APPENDIX B. PROOF OF LEMMA 5.3

(a) Note that  $\mu_1 \mathbb{1}_{\frac{1}{k} < |z| < k}$  has a finite mass by **(M)**. An application of Fubini's theorem then shows that the convolution  $\mu_{1,k} = (\mu_1 \mathbb{1}_{\frac{1}{k} < |z| < k}) * \rho_k$  is a measure which has density  $\bar{\mu}_{1,k}$  with respect to the Lebesgue measure given by

$$\bar{\mu}_{1,k}(x) = \int_{\mathbb{R}^N} \rho_k(x-z) \mathbb{1}_{\frac{1}{k} < |z| < k} \mu_1(dz).$$

This function is bounded for each  $k$ :

$$\|\bar{\mu}_{1,k}\|_\infty \leq \|\rho_k\|_\infty \mu_1\left(\left\{\frac{1}{k} < |z| < k\right\}\right) < \infty,$$

A similar argument shows the existence and boundedness of  $\bar{\mu}_{2,k}$ .

Let  $g(z) := |z|^2 \mathbb{1}_{|z| < \delta}$ , then by Fubini's theorem and symmetry of  $\rho_k$ ,

$$\int_{|z| < \delta} |z|^2 \bar{\mu}_{1,k}(z) dz = \int_{\mathbb{R}^N} g(z) \int_{\mathbb{R}^N} \rho_k(z-y) \mathbb{1}_{\frac{1}{k} < |y| < k} \mu_1(dy) dz = \int_{\mathbb{R}^N} (g * \rho_k)(y) \mathbb{1}_{\frac{1}{k} < |y| < k} \mu_1(dy).$$

Note that  $\rho_k * g$  is continuous with support in  $\{|z| < \delta + 1/k\}$ . By Hölder's inequality,

$$|\rho_k * g(z)| \leq \max_{y \in z + \overline{B}_{1/k}} |g(y)| \cdot 1 \leq (|z| + 1/k)^2 \leq 4|z|^2 \quad \text{for } |z| > 1/k,$$

and hence

$$\int_{|z| < \delta} |z|^2 \bar{\mu}_{1,k}(z) dz \leq 4 \int_{|z| < \delta} |z|^2 \mu_1(dz).$$

The proof of (a) is complete.

(b) Let  $\varepsilon > 0$  be given, and split the integral in two using a  $K > \delta$  to separate the domains:

$$\begin{aligned} I &:= \int_{|z| \geq \delta} \psi_k(z) d\mu_{1,k}(z) = \int_{\delta \leq |z| \leq K} \psi_k(z) d\mu_{1,k}(z) + \int_{|z| \geq K} \psi_k(z) d\mu_{1,k}(z) = I_1 + I_2, \\ J &:= \int_{|z| \geq \delta} \psi(z) d\mu_1(z) = \int_{\delta \leq |z| \leq K} \psi(z) d\mu_1(z) + \int_{|z| \geq K} \psi(z) d\mu_1(z) = J_1 + J_2. \end{aligned}$$

We will show that if we take  $K$  big enough, then  $|I_2| + |J_2| < \varepsilon$  for all  $k$ , and then if  $k$  is big enough,  $|I_1 - J_1| < \varepsilon$ . The conclusion is that  $|I - J| \leq 2\varepsilon$  and the proof is complete.

Consider first  $I_2$  and  $J_2$ . By the definition of  $\mu_{1,k}$  and Fubini's theorem,  $\mu_{1,k}(\{|z| > K\}) \leq \mu_1(\{|z| > K - 1/k\})$ , and then by the dominated convergence theorem and  $\mu_1(\{|z| > 1\}) < \infty$ ,

$$0 \leq \mu_{1,k}(\{|z| > K\}) \leq \mu_1(\{|z| > K - 1/k\}) \rightarrow 0 \quad \text{as } K \rightarrow \infty.$$

Note that this convergence is uniform in  $k$ . Hence since  $\psi_k$  and  $\psi$  are uniformly bounded in  $k$ , it follows that  $I_2, J_2 \rightarrow 0$  as  $K \rightarrow \infty$  uniformly in  $k$ .



We complete the proof by showing that  $|I_1 - J_1| \rightarrow 0$  as  $k \rightarrow \infty$  for any fixed  $K$ . Note that

$$|I_1 - J_1| \leq \int_{\delta \leq |z| \leq K} |\psi_k - \psi| \mu_{1,k}(dz) + \left| \int_{\delta \leq |z| \leq K} \psi \mu_{1,k}(dz) - \int_{\delta \leq |z| \leq K} \psi \mu_1(dz) \right|.$$

Consider the first term on the right hand side. Since  $\sup_{\delta \leq |z| \leq K} |\psi_k - \psi| \rightarrow 0$  by assumption, and  $\mu_{1,k}(\{\delta \leq |z| \leq K\}) \leq \mu_1(\{|z| \geq \delta - \frac{1}{k}\})$  as in the  $|z| > K$  case, for  $k > \frac{1}{\delta}$  we get

$$\int_{\delta \leq |z| \leq K} |\psi_k - \psi| \mu_{1,k}(dz) \leq \sup_{\delta \leq |z| \leq K} |\psi_k - \psi| \mu_1(\{|z| \geq \delta - \frac{1}{k}\}) \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

For the second term, let  $g(z) = \psi(z) \mathbb{1}_{\delta \leq |z| \leq K}$  and use Fubini's theorem to see that

$$\int_{\delta \leq |z| \leq K} \psi \mu_{1,k}(dz) = \int_{\mathbb{R}^N} (\rho_k * g)(z) \mathbb{1}_{\frac{1}{k} < |z| < k} \mu_1(dz).$$

In the last integral, the support of the convolution is  $\{\delta - \frac{1}{k} \leq |z| \leq K + \frac{1}{k}\}$  so we need  $k > \frac{1}{\delta}$ . Since  $\rho_k * g \rightarrow g$  and  $\mathbb{1}_{\frac{1}{k} < |z| < k} \rightarrow 1$  pointwise (almost everywhere) both functions are uniformly bounded, we can pass to the limit using dominated convergence and find that

$$\left| \int_{\delta \leq |z| \leq K} \psi \mu_{1,k}(dz) - \int_{\delta \leq |z| \leq K} \psi \mu_1(dz) \right| \rightarrow 0 \quad \text{as } k \rightarrow \infty. \quad (\text{B.1})$$

The proof of (b) is complete.

(c) Let  $\varepsilon > 0$  be given, and split the integral in two using a  $0 < r < \min(\delta, \delta_0)$  to separate the domains:

$$\begin{aligned} I &:= \int_{0 < |z| \leq \delta} \psi_k(z) d\mu_{1,k}(z) = \int_{|z| < r} \psi_k(z) d\mu_{1,k}(z) + \int_{r < |z| \leq \delta} \psi_k(z) d\mu_{1,k}(z) = I_1 + I_2, \\ J &:= \int_{0 < |z| \leq \delta} \psi(z) d\mu_1(z) = \int_{|z| < r} \psi(z) d\mu_1(z) + \int_{r < |z| \leq \delta} \psi(z) d\mu_1(z) = J_1 + J_2. \end{aligned}$$

We will show that if we take  $r$  small enough, then  $|I_2| + |J_2| < \varepsilon$  for all  $k$ , and then if  $k$  is big enough,  $|I_1 - J_1| < \varepsilon$ . The conclusion is that  $|I - J| \leq 2\varepsilon$  and the proof is complete.

The estimate  $|I_2| + |J_2| < \varepsilon$  for  $r$  small, follows by the assumptions on  $\psi$ , part (a), and dominated convergence and (M). For example,

$$\int_{0 < |z| < r} |\psi_k(z)| \mu_{1,k}(dz) \leq C \int_{0 < |z| < r} |z|^2 \mu_{1,k}(dz) \leq 4C \int_{0 < |z| < r} |z|^2 \mu_1(dz) \rightarrow 0 \quad \text{as } r \rightarrow 0.$$

Consider now  $|I_1 - J_1|$ . We first introduce the functions

$$\tilde{\psi}_k(z) := \frac{\psi_k(z)}{|z|^2} \quad \text{and} \quad \tilde{\psi}(z) := \frac{\psi(z)}{|z|^2},$$

and measures

$$\tilde{\mu}_{1,k}(dz) := |z|^2 \mu_{1,k}(dz) \quad \text{and} \quad \tilde{\mu}_1(dz) := |z|^2 \mu_1(dz).$$

By the assumptions and (M),  $\tilde{\psi}_k$  and  $\tilde{\psi}$  are uniformly bounded,  $\tilde{\psi}_k \rightarrow \tilde{\psi}$  uniformly on  $r < |z| < \delta$ , and  $\tilde{\mu}_{1,k}$  and  $\tilde{\mu}_1$  are bounded measures on  $0 < |z| < \delta$ . It follows that

$$|I_1 - J_1| \leq \int_{r < |z| < \delta} |\tilde{\psi}_k - \tilde{\psi}| \tilde{\mu}_{1,k}(dz) + \left| \int_{r < |z| < \delta} \tilde{\psi} \tilde{\mu}_{1,k}(dz) - \int_{r < |z| < \delta} \tilde{\psi} \tilde{\mu}_1(dz) \right|.$$

The first term converges by uniform convergence of  $\tilde{\psi}_k$  and uniform boundedness of  $\mu_{1,k}(\{r < |z| < \delta\})$ . For the second term, we note that (see part (a))

$$\int_{r < |z| < \delta} \tilde{\psi} \tilde{\mu}_{1,k}(dz) = \int_{r < |z| < \delta} \psi \mu_{1,k}(dz) = \int_{\mathbb{R}^N} (\rho_k * (\psi(\cdot) \mathbb{1}_{r < |\cdot| < \delta}))(z) \mathbb{1}_{\frac{1}{k} < |z| < k} \mu_1(dz).$$

The integrand is uniformly bounded (by  $\|\psi(\cdot) \mathbb{1}_{r < |\cdot| < \delta}\|_\infty$ ) and converges pointwise to  $\psi(z) \mathbb{1}_{r < |z| < \delta}$  for a.a.  $z$ , so by **(M)** and the dominated convergence theorem, we can conclude that  $|I_1 - J_1| \rightarrow 0$  as  $k \rightarrow \infty$ . The proof of (c) is complete.

(d) This proof is similar to the proof of (b), we omit it.

### APPENDIX C. THE PROOF OF LEMMA 5.10

(a) Let  $K \subset \mathbb{R}^N \times \mathbb{R}^P$  be any bounded set, then by the definition of  $\rho_k$ , Hölder's inequality, and  $\int \rho_k(p) dp = 1$ ,  $\|j_{i,k}\|_{L^\infty(K)} \leq \|j_i\|_{L^\infty(K_{1/k})}$  where

$$K_{1/k} = \{(q, z) \in \mathbb{R}^N \times \mathbb{R}^P : \exists p \in \mathbb{R}^N \text{ such that } (p, z) \in K \text{ and } |q - p| < 1/k\}.$$

By Hölder's inequality and **(J1')**, we also find that for  $|p| \leq r$  and  $|z| < 1$ ,

$$|j_{1,k}(p, z)| \leq \max_{|q-p| < 1/k} |j(q, z)| \int \rho_k(p) dp \leq C_{r+1/k} |z|,$$

and the proof of (a) is complete.

(b) By similar arguments, for  $|p|, |q|, |z| < r$ ,

$$|j_{i,k}(p, z) - j_{i,k}(q, z)| \leq \int_{|s| < r+1/k} |j(s, z)| |\rho_k(p-s) - \rho_k(q-s)| ds \leq \sup_{\substack{|s| \leq r+1/k \\ |z| \leq r}} |j(s, z)| \|D\rho_k\|_{L^1} |p-q|,$$

and the proof of (b) is complete by **(J1')** and the standard estimate  $\|D\rho_k\|_{L^1} \leq k\|D\rho\|_{L^1}$ .

(c) By the definition of  $j_{1,k}$ , properties of mollifiers and Jensen's inequality, Fubini and **(J2)**,

$$\begin{aligned} \int_{|z| > 0} |j_{1,k}(p, z) - j_{1,k}(q, z)|^2 d\mu_1(z) &\leq \int_{|z| > 0} \int_{y \in \mathbb{R}^N} \rho_k(y) |j_1(p-y, z) - j_1(q-y, z)|^2 dy d\mu_1(z) \\ &\leq \int_{y \in \mathbb{R}^N} \rho_k(y) \left( \int_{|z| > 0} |j_1(p-y, z) - j_1(q-y, z)|^2 d\mu_1(z) \right) dy \\ &\leq \int_{y \in \mathbb{R}^N} \rho_k(y) \omega_{j, r+\frac{1}{k}}(p-q) dy \leq \omega_{j, r+\frac{1}{k}}(p-q). \end{aligned}$$

(d) Let  $A \subset \{0 < |z| < \delta_0\}$  be a Borel set. Then as in part (c), we use properties of mollifiers and Jensen's inequality, Fubini and **(J2)**, to see that

$$\int_A |j_{1,k}(p, z)|^2 d\mu(z) \leq \int_A \int_{|q| < \frac{1}{k}} \rho_k(q) |j_1(p-q, z)|^2 dq d\mu_1(z) \leq \max_{|q| < \frac{1}{k}} \int_A |j_1(p-q, z)|^2 d\mu_1(z).$$

Now (d) follows from **(J3)** applied with  $r+1$ , which is bigger than  $r+1/k$  for any  $k \geq 1$ .

(e) First fix a  $z$  such that  $j_i(p, z)$  is continuous in  $p$ , cf. **(J1)**. Hence  $j_{i,k}(\cdot, z)$  converges locally uniformly to  $j_i(\cdot, z)$  as  $k \rightarrow \infty$  (see for instance Appendix C, Theorem 6 in [16]). Moreover,  $j_i(\cdot, z)$  is locally uniformly continuous in  $p$ , say with a modulus  $\omega_r$  for  $|p|, |q| \leq r$ . Then

$$|j_{i,k}(p, z) - j_{i,k}(q, z)| \leq \int_{y \in \mathbb{R}^N} |j(p-y, z) - j(q-y, z)| \rho_k(y) dy \leq \omega_r(|p-q|) \cdot 1,$$

and  $j_{i,k}(\cdot, z)$  is equicontinuous in  $p$ . Combining these two results, it follows that for every  $p_k \rightarrow p$ ,

$$j_{i,k}(p_k, z) \rightarrow j_i(p, z) \quad \text{as } k \rightarrow \infty \quad \text{for a.e. } z. \quad (\text{C.1})$$

Then we let  $p_k = D\phi(x_k)$  and  $p = D\phi(x)$ , and consider  $L_{2,k}[\phi, p_k](x_k) = \int \phi(x_k + j_{2,k}(p_k, z)) - \phi(x_k) \mu_2(dz)$ . By (C.1) and continuity of  $\phi$  and  $D\phi$ ,

$$\phi(x_k + j_{2,k}(p_k, z)) - \phi(x_k) \rightarrow \phi(x + j_2(p, z)) - \phi(x) \quad \text{for a.e. } z,$$

and since the integrand is uniformly bounded by the  $\mu_2$ -integrable function  $2\|\phi\|_\infty$  (cf. **(M)**), the dominated convergence theorem implies that

$$L_{2,k}[\phi, p_k](x_k) \rightarrow L_2[\phi, p](x) \quad \text{as } k \rightarrow \infty.$$

Note that  $L_{1,k} = L_{1,\delta,k} + L_{1,k}^\delta$ . For the  $L_{1,k}^\delta$ -term the proof is more or less the same as for the  $L_{2,k}$ -term (see above). It only remains to consider the  $L_{1,\delta}$ -term. As above, we see that the integrand converges to  $\phi(x + j_1(D\phi(x), z)) - \phi(x) - j_1(D\phi(x), z) \cdot D\phi(x)$  for a.e.  $z$ . An application of Taylor's theorem and **(J1')**, show that the integrand uniformly bounded by the  $\mu_1$ -integrable function  $\|D^2\phi\|_{L^\infty(B_{R_2})} C_{R_1} |z|^2$  where  $R_1 = \max(\delta, \max_k |D\phi(x_k)|)$  and

$$R_2 = \max_{k \in \mathbb{N}} |x_k| + \max_{|s|, |z| \leq R_1} |j(s, z)|.$$

Hence we conclude by the dominated convergence theorem that

$$L_{1,\delta,k}[\phi, D\phi(x_k)](x_k) \rightarrow L_{1,\delta}[\phi, D\phi(x)](x) \quad \text{as } k \rightarrow \infty.$$

The proof of (e) is complete.

## REFERENCES

- [1] N. Alibaud, E. Chasseigne, E. Jakobsen: *work in preparation*.
- [2] F. Andreu-Vaillio, J.M. Mazón, J.D. Rossi, J.J. Toledo: *Nonlocal Diffusion Problems*, American Mathematical Society. Mathematical Surveys and Monographs (2010)
- [3] D. Applebaum. *Lévy Processes and Stochastic Calculus*. Cambridge University Press, 2009.
- [4] G. Barles, R. Buckdahn, and E. Pardoux. *Backward stochastic differential equations and integral-partial differential equations*. Stochastics Stochastics Rep. 60 (1997), no. 1-2, 57–83.
- [5] G. Barles, E. Chasseigne, C. Georgelin, E.R. Jakobsen, *On Neumann-type problems for nonlocal equations set in a half space*. Trans. Amer. Math. Soc. 366 (2014), 4873–4917.
- [6] G. Barles, C. Georgelin, and E. R. Jakobsen. *On Neumann and oblique derivatives boundary conditions for nonlocal elliptic equations*. J. Differential Equations. 256(4): 1368–1394, 2014.
- [7] G. Barles, C. Imbert *Second order elliptic integro-differential Equations: viscosity solutions's theory revisited.*, Ann. Inst. H. Poincaré Anal. non linéaire 25(2008), 567-585.
- [8] C. Bjorland, L. Caffarelli, A. Figalli. *Non-Local Gradient Dependent Operators*, Adv. Math., v.230, 2012, p. 46.
- [9] C. Bjorland, L. Caffarelli, and A. Figalli. *Nonlocal tug-of-war and the infinity fractional Laplacian*. Comm. Pure Appl. Math. 65 (2012), no. 3, 337–380.
- [10] I. Capuzz-Dolcetta and M. Bardi. *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman equations*. Birkhäuser, 1997.
- [11] A. Chambolle, E. Lindgren, R. Monneau *Hölder infinity Laplacian*. ESAIM: Control, Optimisation and Calculus of Variations 18.3 (2012)
- [12] R. Cont and P. Tankov. *Financial modelling with jump processes*. Chapman & Hall/CRC, 2004.
- [13] .G Crandall, H.Ishii and P.L Lions: *User's guide to viscosity solutions of second order Partial differential equations*. Bull. Amer. Soc. 27 (1992), pp 1-67.
- [14] H. Ishii and G. Nakamura. *A class of integral equations and approximation of p-Laplace equations*. Calc. Var. Partial Differential Equations 37 (2010).
- [15] A. Di Castro, T. Kuusi, and G. Palatucci. *Local behavior of fractional p-minimizers*. Preprint, 2013.
- [16] L. C. Evans. *Partial Differential Equations*. Graduate studies in Mathematics, Vol. 19, AMS, 2010.
- [17] D. Gilbarg and N. Trudinger. *Elliptic partial differential equations of second order*. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001.

- [18] E. R. Jakobsen and K. H. Karlsen. *A Maximum principle for semicontinuous functions applicable to integro-partial differential equations* Nonlinear Differential Equations and Applications 13, 2006.
- [19] E. R. Jakobsen and K. H. Karlsen. *Continuous dependence estimates for viscosity solutions of fully nonlinear degenerate elliptic equations*. Electron. J. Diff. Eqns. 2002(39): 1-10, 2002.
- [20] G.M. Lieberman, *Second order parabolic differential equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [21] E. Lindgren. *Hölder estimates for viscosity solutions of equations of fractional  $p$ -Laplace type*. Preprint, <http://arxiv.org/abs/1405.6612>.
- [22] E. Lindgren and P. Lindqvist. *Fractional eigenvalues*. Calc. Var. Partial Differential Equations 49 (2014), no. 1-2, 795-826.
- [23] P. Lindqvist. *Notes on the Infinity-Laplace Equation*. Lecture Note. Preprint, <http://arxiv.org/abs/1411.1278>.
- [24] Y. Peres, O. Schramm, S. Sheffield, and D. B. Wilson. Tug-of-war and the infinity Laplacian *J. Amer. Math. Soc.* 22 (2009), 167-210
- [25] W. Rudin. *Real and Complex Analysis*, 3rd edition. cGraw-Hill, 1987.
- [26] A. Sayah. *Equations d'Hamilton-Jacobi du premier ordre avec termes integro-différentiels. I. Unicité des solutions de viscosité*. Comm. Partial Differential Equations 16 (1991), no. 6-7, 1057-1074.
- [27] Z. Wu, J. Yin, and C. Wang. *Elliptic and Parabolic Equations* World Scientific Publishing, 2006.

EMMANUEL CHASSEIGNE  
 LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE (UMR CNRS 7350)  
 FÉDÉRATION DENIS POISSON (FR CNRS 2964)  
 UNIVERSITÉ F. RABELAIS - TOURS  
 PARC DE GRANDMONT  
 37200 TOURS, FRANCE

*E-mail address:* [emmanuel.chasseigne@univ-tours.fr](mailto:emmanuel.chasseigne@univ-tours.fr)

*URL:* <http://www.lmpt.univ-tours.fr/~manu>

ESPEN R. JAKOBSEN  
 DEPARTMENT OF MATHEMATICAL SCIENCES  
 NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY  
 7491 TRONDHEIM, NORWAY

*E-mail address:* [erj@math.ntnu.no](mailto:erj@math.ntnu.no)

*URL:* <http://www.math.ntnu.no/~erj>