

Magmatism on rift flanks: insights from Ambient-Noise Phase-velocity in Afar region

Félicie Korostelev, Cornelis Weemstra, Sylvie Leroy, Lapo Boschi, Derek Keir, Yong Ren, Irene Molinari, Abdulhakim Ahmed, Graham W. Stuart, Frédérique Rolandone, et al.

▶ To cite this version:

Félicie Korostelev, Cornelis Weemstra, Sylvie Leroy, Lapo Boschi, Derek Keir, et al.. Magmatism on rift flanks: insights from Ambient-Noise Phase-velocity in Afar region. Geophysical Research Letters, 2015, 42 (7), pp.2179-2188. 10.1002/2015GL063259 . hal-01136108

HAL Id: hal-01136108 https://hal.science/hal-01136108

Submitted on 26 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Magmatism on rift flanks: insights from Ambient-Noise Phase-velocity in Afar region

Félicie Korostelev,^{1,2} Cornelis Weemstra,³ Sylvie Leroy,^{1,2} Lapo Boschi^{1,2}, Derek Keir⁴, Yong Ren⁵, Irene Molinari⁶, Abdulhakim Ahmed^{1,2,7}, Graham W. Stuart⁵, Frédérique Rolandone^{1,2}, Khaled Khanbari⁸, James O. S. Hammond⁹, J. M. Kendall¹⁰, Cécile Doubre¹¹, Ismail Al Ganad¹², Berhe Goitom¹³, and Atalay Ayele¹⁴

Corresponding author: Félicie Korostelev, (felicie.korostelev@gmail.com)

¹Sorbonne Universités, UPMC Univ Paris

DRAFT

February 27, 2015, 9:20am

During the breakup of continents in magmatic settings, the extension of 3 the rift valley is commonly assumed to initially occur by border faulting and 4 progressively migrate in space and time towards the spreading axis. Mag-5 matic processes near the rift flanks are commonly ignored. We present phase-6 velocity maps of the crust and uppermost mantle of the conjugate margins 7 of the southern Red Sea (Afar and Yemen) using ambient noise tomography 8 to constrain crustal modification during breakup. Our images show that the q low seismic velocities characterize not only the upper crust beneath the ax-10 ial volcanic systems, but also both upper and lower crust beneath the rift 11 flanks where ongoing volcanism and hydrothermal activity occur at the sur-12 face. Magmatic modification of the crust beneath rift flanks likely occurs for 13 a protracted period of time during the breakup process, and may persist through 14 to early seafloor spreading. 15

06, UMR 7193, Institut des Sciences de la

DRAFT

February 27, 2015, 9:20am

1. Introduction

During the breakup of continents, stretching and thinning of the plate commonly causes 16 decompression melting and volcanism. In the resultant magnatically active rift valleys 17 it is widely thought that extension is initially accommodated mainly by border faulting, 18 and progressively localizes to relatively narrow axial volcanic segments as the rift valley 19 widens [e.g. *Ebinger and Casey*, 2001]. However, it is becoming increasingly more recog-20 nized that magma intrusion and volcanism can occur on the rift flanks at an early stage 21 of rifting [e.g. Maccaferri et al., 2014]. These rift flank magmatic systems accommodate 22 extension through diking [Rooney et al., 2014], and thermally and compositionally mod-23 ify the lithosphere [Daniels et al., 2014]. Despite the importance of magmatic processes 24 during continental extension, we have few constraints on their spatial and temporal vari-25 ability. In order to address this issue we use ambient seismic noise tomography to image the Rayleigh wave phase-velocity structure of the crust in a region of late stage breakup 27 at the conjugate margins of the southern Red Sea in Afar and Yemen. 28

Geochronological constraints in Ethiopia suggest rifting began 29-31 Ma on the western Afar margin [e.g. *Ayalew et al.*, 2006; *Wolfenden et al.*, 2005, Fig.1], approximately coeval with \sim 35 Ma faulting along large portions of the Gulf of Aden to the east [*Leroy et al.*, 2010]. Rifting was associated with the development of large offset border faults that currently define \sim 2000-3000 m of relief between the submarine Red Sea and subaerial Afar depression with the uplifted Ethiopian and Yemeni plateaus [*Wolfenden et al.*, 2004]. Extension is thought to have occurred above warm mantle with a potential temperature

Terre Paris (iSTeP), F-75005 Paris, France.

DRAFT

of ~ 1450 degrees [Rooney et al., 2012], associated with voluminous flood basalts on the Ethiopian and Yemeni plateaus synchronous with the onset of extension [Wolfenden et al., 37 2004], and associated with ongoing magmatism [Ferguson et al., 2013]. At \sim 21-23 Ma, 38 magmatism occurred through dike intrusions along most of the eastern margin of the 39 Red Sea [Bosworth et al., 2005]. Magmatism on the rift flanks is ongoing, with the 40 Quaternary to Recent volcanic centers of Sana'a, Dhamar and Marib located in Yemen 41 [Manetti et al., 1991; Korostelev et al., 2014]. In addition, thermal hotsprings are present 42 along the conjugate southern Afar margin [Keir et al., 2009]. Magma intrusion and 43 volcanism is also common within the rift valley. Since ~ 10 Ma in Afar, extension via 44 diking progressively localized to the rift axis [e.g. Wolfenden et al., 2005; Rooney et al., 45 2011], with the current locus of strain being \sim 70-km-long, \sim 20-km-wide axial volcanic 46 segments such as the Dabbahu-Manda-Hararo segment in central-west Afar [e.g. Hayward 47 and Ebinger, 1996]. Here, episodic intrusion of dikes fed from crustal magma chambers at both the segments centers and tips accommodates the majority of extension [e.g. Keir 49 et al., 2009; Grandin et al., 2010, 2011]. 50

⁵¹ Current opening across the kinematically complex southern Red Sea rift is constrained ⁵² with relatively high-density GPS [e.g. ArRajehi et al., 2010; McClusky et al., 2010] and ⁵³ InSAR measurements[e.g. Pagli et al., 2014]. These data show that south of ~16°N, the ⁵⁴ rift bifurcates into two branches: the main Red Sea and the subaerial Red Sea rift in Afar ⁵⁵ (Danakil Depression). Partitioning of extension between rift branches varies along-strike.

²CNRS, UMR 7193, Institut des Sciences

DRAFT

North of $\sim 16^{\circ}$ N, all the extension is accommodated in the main Red Sea rift, spreading 56 at ~ 15 mm/yr. Moving south of 16°N, the extension is progressively accommodated in 57 the Afar Depression reaching $\sim 20 \text{ mm/yr}$ at 13°N [McClusky et al., 2010; Vigny et al., 58 2006]. The crust beneath Afar varies from 25 km thick beneath most of Afar, to 15 km 59 thick beneath the Danakil depression (Afdera-Erta'Ale segment) in the north [Makris and 60 Ginzburg, 1987; Bastow and Keir, 2011, Fig.1]. The crustal thickness is ~ 25 km thick 61 beneath the Danakil block and increases to 40-45 km beneath the Ethiopian and Yemeni 62 Plateaus [Hammond et al., 2011; Ahmed et al., 2013, Fig.1]. 63

2. Data

Our dataset is based on continuous recordings from 89 seismic stations. Only a limited 64 number of high-quality permanent seismic stations span the Afar-southern Red Sea mar-65 gins and so temporary experiments using portable broad-band equipment are our major 66 source of information on the structure of the area. A seismic deployment was conducted 67 between March 2009 and March 2010 as part of the YOCMAL (Young Conjugate Margins 68 Laboratory) project, with 23 stations covering western Yemen during one year [Korostelev 69 et al., 2014; Corbeau et al., 2014, Fig.2]. We also use data from 41 stations in the Afar 70 Consortium network (UK and US, from March 2007 to November 2009) [e.g. Keir et al., 71 2011, five stations of the Horn of Africa network in Yemen and Ethiopia [from June 1999 72 to December 2002, e.g. Sicilia et al., 2008] and six temporary stations recording from 73 May 2011 to September 2012 in Eritrea [e.g. Hammond et al., 2013, Fig.2]. One station 74

de la Terre Paris (iSTeP), F-75005 Paris,

DRAFT

⁷⁵ from the Djibouti temporary network was added to our dataset, together with permanent
⁷⁶ seismic stations in Djibouti, Yemen and Ethiopia.

The ambient-noise cross-correlation technique relies on having simultaneous recordings 77 of the noise field at two seismic stations so that the Green's function between them can 78 be estimated [Shapiro and Campillo, 2004; Wapenaar and Fokkema, 2006; Halliday and 79 *Curtis*, 2008]. Because the different deployments of portable instruments occurred at 80 different times, we are not able to estimate Green's functions for all receiver pairs. We 81 partly compensate for this, however, by utilizing permanent stations from the IRIS and 82 GEOSCOPE networks, providing data over a period during which several of the mentioned 83 portable arrays were active. 84

3. Method

The ambient noise technique to study Earth structure is free of limitations imposed by the distribution of natural earthquakes. Extracting travel times from a multitude of station-station correlations therefore allows for relatively high-resolution tomographic inversions [e.g. *Shapiro et al.*, 2005]. We follow the approach of *Ekström et al.* [2009], discussed in further detail in section 3.2 of *Boschi et al.* [2013], to estimate phase velocity from the ambient signal recorded at two stations.

The background seismic noise is to a large extent generated by the coupling of oceans with the solid Earth [e.g. *Longuet-Higgins*, 1950; *Hillers et al.*, 2012]. Because this area is almost surrounded by seas or oceans (Red Sea, Gulf of Aden, Indian Ocean), it is particularly suitable for ambient noise surface wave retrieval.

France.

DRAFT

February 27, 2015, 9:20am

DRAFT

Х-6

To maximize data quality, we (i) only used the pairs of stations that recorded simultaneously for at least 6 months, and (ii) compared measured and predicted Green's function for all station pairs, and discarded pairs that clearly showed a bad fit (see figure in supplementary material). The duration of cross-correlated signal varies by 6-36 months depending on the station pair. These long durations guarantee that all seasons and hence

¹⁰⁰ all possible azimuths of noise propagation are sampled [e.g. *Stehly et al.*, 2006]. Data ¹⁰¹ processing was limited to whitening, as reasonable dispersion curves could be obtained ¹⁰² without any filtering and/or "one-bit" amplitude compression.

4. Resolution

95

96

97

98

99

4.1. Station-to-station paths

To assess the resolving power of our inversion, we first show in Figure 2 the station-tostation paths corresponding to ambient Rayleigh-wave observations at each period. The solid gray line delimitates the area with good coverage, and therefore the zone of best resolution.

4.2. Random tests

¹⁰⁷ We perform two random resolution tests [e.g. *Verbeke et al.*, 2012] to assess the reliability ¹⁰⁸ of the tomographic inversion: one with structures smaller than 100 km (Fig.3.a) and a ¹⁰⁹ second one with structures larger than 100 km (Fig.3.b). The input synthetic random

DRAFT

³Department of Geoscience and

velocity model consists of alternating random structures of opposite sign with a maximum
velocity variation of 1.5% relative to the reference velocity.

¹¹² Synthetic phase velocities were computed between the same station pairs as in the ¹¹³ observed database. Figures 3.a and b show the input velocity models and the retrieved ¹¹⁴ velocity models from these tests for periods of 9, 15.5 and 20.5 s. These synthetic tests ¹¹⁵ indicate that our inversion can resolve most of the Afar-southern Red Sea margins region, ¹¹⁶ with some degradation of the recovered solutions near the edges of the illuminated area. ¹¹⁷ The tomography algorithm is that utilized e.g. by *Verbeke et al.* [2012]. Pixel-size is ¹¹⁸ $0.1^{\circ}x0.1^{\circ}$.

Our synthetic tests (Fig.3) serve to validate both pixel size and select the values of regularization parameters that allow us to represent heterogeneities of scale-length such as in figure 3.a (left). The same parameterization and regularization is applied to real data in the following. Notice that resolution changes across the region of study, so that a unique resolution limit cannot be specified.

5. Results

¹²⁴ We compute Rayleigh-wave phase-velocity maps for periods between 9 s and 25.5 s, ¹²⁵ and present examples at 9, 15.5 and 20.5 s (Fig.4; see supplementary material for other ¹²⁶ periods). According to e.g. *Lebedev and Van Der Hilst* [2008] and *Fry et al.* [2010], 9-s ¹²⁷ Rayleigh waves are most sensitive to depths < 20 km (upper and mid crust), while 15.5-¹²⁸ s are most sensitive to 10 - 40 km depth (primarily the lower crust). 20.5-s Rayleigh ¹²⁹ waves can sample down to 70 km, at the top of the upper mantle. The locus of major

Engineering, Delft University of Technology,

D R A F T February 27, 2015, 9:20am D R A F T

X - 8

velocity anomalies is fairly constant from 9 s to 15.5 s (Fig. 4). We image positive velocity 130 perturbations beneath the border faults of the eastern flank of the Red Sea in Yemen, 131 beneath the Danakil Horst, and in central western Afar in the region between the rift 132 margin and the axial volcanic segments (Sullu Adu area, Fig.1). We also see positive 133 velocity perturbations beneath the western Afar margin north of 12°N. The main slow 134 anomalies are located beneath Dabbahu Manda-Hararo axial volcanic system, beneath 135 Durrie off-axis volcano and the southern axis extension to Kurub volcano (Fig.1). We 136 also find slow anomalies associated with the volcanic systems 150 to 200 km east of the 137 rift margin in Yemen, and beneath the western Afar margin south of 12°N (Fig.4). 138

The magnitude of several of the distinct velocity perturbations varies subtly with period. For example, beneath the eastern rift margin (Tihama Plain, Yemen, Fig.1), the positive anomaly increases in magnitude from 5% at 9 s to 7% at 20.5 s (Fig. 4). The slow anomaly beneath the western Afar margin flank south of 12N is mostly more than -3% at 9 s, whereas at 15.5 and 20.5 s a larger proportion of the anomaly is -4 to -6% (Fig.4).

The slow anomalies beneath Yemen and beneath the axial volcanic segment of Dabbahu-Manda-Hararo in Afar correlate well with the locus of surface volcanism (Fig.4). In addition, figure 5 shows the surface distribution of known thermal springs in the region [*Keir et al.*, 2009]. The slow anomaly beneath the western Afar margin is beneath the locus of thermal springs on the western Afar margin, whereas north of 12°N, thermal springs are absent and the crust is faster than average. The spatial extent of slow anomalies imaged

Stevinweg 1, 2628 CN Delft, The

DRAFT

¹⁵⁰ using ambient noise also correlates well with the spatial extent of high Vp/Vs ratios in ¹⁵¹ the crust constrained using P-S receiver functions [*Hammond et al.*, 2011].

6. Discussion

Seismic wave velocity is known to be affected by the temperature and chemical com-152 position of the medium of propagation (crustal rocks), as well as by the concentration of 153 fluids, such as partial melt that might be present within crustal rocks [e.g. Christensen 154 and Mooney, 1995; Karato et al., 2003. We image slow velocities beneath axial regions of 155 localized magma intrusion, consistent with the hypothesis that major surface-wave slow 156 anomalies are associated with magmatism (Fig.4). In addition, the lowest velocities in 157 our images are beneath zones of active volcanism and geothermal activity near the flanks 158 of the southern Red Sea conjugate margins (Fig. 4 and 5). The magnitude of the anoma-159 lies and spatial association with regions where either partial melt or fluids released from 160 cooling magmatic systems are present suggests the crust beneath the flanks of the rift is 161 currently being modified by magmatic processes. Ongoing magmatism occurs at the rift 162 flanks in spite of the majority of strain having shifted to the rift axis since the onset of 163 rifting at 11 Ma [Wolfenden et al., 2004]. Beneath the western Afar margin, where slow 164 anomalies are associated with geothermal systems rather than known volcanoes, geologi-165 cal studies suggest early border faulting at 30 Ma was associated with spatially localized 166 volcanism in the marginal graben systems [Ayalew et al., 2006]. Our velocity maps suggest 167 that the magmatic systems beneath the rift flanks that were active during the onset of 168 rifting, remain magnetically active throughout the breakup process either through con-169

Netherlands.

DRAFT

February 27, 2015, 9:20am

tinued minor accumulation of partial melt in reservoirs, dike intrusion, and / or ongoing conductive cooling leading to release of fluids such as water [*Keir et al.*, 2009; *Holtzman* et al., 2010].

The low-velocity anomalies in our phase-velocity maps under the rift axis are observed 173 with higher amplitude in the upper crust (Fig.4, period = 9 seconds), whereas the low-174 velocity anomalies located beneath the rift flanks are observed both in the upper and 175 lower crust, but with higher amplitude in the lower crust (Fig.4, period = 15.5 seconds). 176 This is consistent with the proposed plumbing systems of axial and flank volcanic systems 177 of the nearby Main Ethiopian Rift, where petrological constraints on flank volcanism are 178 good. The volcanic products observed on the flanks and at the axis of the MER are not 179 identical: they consist mainly of trachytes for the flanks, and mainly of rhyolites and 180 basalts for the axis [Peccerillo et al., 2007]. Petrological models indicate that the origin of 181 the off-axis trachytes is probably high-pressure fractional crystallization of asthenosphere-182 derived basalts, with this fractionation occurring at the base of the crust [Peccerillo et al., 183 2007]. Rooney et al. [2005, 2007] suggest that these off-axis volcanic products result of 184 moderate-degree partial melting at 50-90 km depth and undergo fractional crystalliza-185 tion in complex plumbing systems spanning depths throughout the crust [Rooney et al., 186 2011]. The volcanic rocks at the axis are asthenospheric basalts produced by rift-related 187 decompressional melting, rather than other potential sources such as melting in the crust. 188 The axial basalts undergo fractional crystallization mostly in the upper crust [Peccerillo 189

DRAFT

⁴National Oceanography Centre

¹⁹⁰ et al., 2007]. Thus, the axial magmatic chamber is shallow (in the upper crust), and the ¹⁹¹ melt ascension from the asthenosphere is probably rapid. At the flanks, however, there ¹⁹² is a complex plumbing system with stacked-reservoirs both in the upper and lower crust ¹⁹³ [Rooney et al., 2011]. The geothermal systems of the flanks are probably fed or heated ¹⁹⁴ by such a complex plumbing system (Fig.5). Our surface-wave velocity maps are consis-¹⁹⁵ tent with this model and therefore suggest similar magmatic plumbing systems for the ¹⁹⁶ southernmost Red Sea.

According to *Medynski et al.* [2015], the magma supply has decreased in the Dabbahu-Manda-Hararo axis reservoir since 15 kyr. An off-axis reservoir, located 15 km to the west of the Dabbahu-Manda-Hararo rift beneath Durrie volcano has been actively fed since 15 kyr, and is currently imaged using magneto-telluric techniques (Fig.1) [*Desissa et al.*, 2013]. It is consistent with our phase-velocity maps, where the maximum amplitude for the northern Dabbahu-Manda-Hararo seems to be slightly to the west of the rift, beneath Durrie volcano (Fig.4).

In the past, geodynamic models of breakup ignored the presence and impact of maintained magmatism at rift flanks on the thermal and subsidence history of the rift during late stage breakup and early seafloor spreading. At the southern Red Sea, where seafloor spreading is young, our new crustal-velocity maps coupled with surface expression of volcanism (Fig.4) show clear evidence for ongoing magmatism beneath the rift flanks in Afar and Yemen (Sana'a, Dhamar and Marib volcanic fields, Fig.1) [*Korostelev et al.*, 2014; *Corbeau et al.*, 2014]. Similarly, there is evidence for ongoing dike intrusion further north

Southampton, University of Southampton,

DRAFT

²¹¹ along the eastern Red Sea flank from InSAR (interferometric synthetic aperture radar)
²¹² and seismicity studies at Harrat Lunayyir volcanic system in Saudi Arabia [e.g. *Pallister*²¹³ et al., 2010; Ebinger et al., 2010]. There, localized subsidence, horizontal opening and
²¹⁴ earthquakes in April to May 2009 are best modeled by intrusion of a dike and induced
²¹⁵ normal faulting. These studies, combined with the evidence presented by our new surface²¹⁶ wave velocity maps, demonstrate that rift flank magmatism during late stage breakup may
²¹⁷ be more common than previously assumed.

7. Conclusions

Our study provides new high-resolution phase-velocity maps of the crust and uppermost 218 mantle of the conjugate margins of the southern Red Sea (Afar and Yemen) using ambient 219 noise tomography to constrain crustal evolution during breakup. Low-velocity anomalies 220 are imaged in the crust beneath the axial volcanic systems, but also in the upper and 221 lower crust beneath rift flanks where hydrothermal activity and ongoing volcanism are 222 observed at the surface. Our results show that the crust beneath the southern Red sea 223 rift flanks is currently being modified by magmatic processes, and that this activity is 224 continuous from the onset of rifting. We therefore demonstrate that rift flank magmatism 225 after breakup may be more common than it was previously thought in context of margins 226 with excess magmatism. 227

Acknowledgments. This project was funded by the ANR-07-BLAN-0135 YOCMAL, CNRS-INSU-PICS Yemen, GSMRB Yemen and is in the framework of the Actions Southampton, SO14 3ZH, U.K.

DRAFT

X - 14

Marges program. Seismic equipment from SEIS-UK is funded by NERC under agreement 230 R8/H10/64. We thank David Hawthorn, Alex Brisbourne and Victoria Lane for their 231 efforts during the deployment and servicing of network, the French Embassy in Yemen, 232 local governers and the people of the Yemen governorates for their help during the field 233 work. The authors thank Raphael Pik, Nicolas Bellahsen and Martin Stab for discus-234 sions about the Afar region. DK is supported by NERC grant NE/L013932/1. CW is 235 supported by the Netherlands Research Centre for Integrated Solid Earth Science (ISES). 236 Djibouti instruments belong to the French national pool of portable seismic instruments 237 Sismob-RESIF and IPGS/EOST (University of Strasbourg). 238

References

- ²³⁹ Ahmed, A., C. Tiberi, S. Leroy, G. W. Stuart, D. Keir, J. Sholan, K. Khanbari, I. Al-
- Ganad, and C. Basuyau (2013), Crustal structure of the rifted volcanic margins and
 uplifted plateau of Western Yemen from receiver function analysis, *Geophysical Journal International*, 193(3), 1673–1690.
- ²⁴³ ArRajehi, A., S. McClusky, R. Reilinger, M. Daoud, A. Alchalbi, S. Ergintav, F. Gomez,
- J. Sholan, F. Bou-Rabee, G. Ogubazghi, et al. (2010), Geodetic constraints on present-
- day motion of the Arabian Plate: Implications for Red sea and Gulf of Aden rifting,
- ²⁴⁶ *Tectonics*, 29(3), TC3011.

⁵School of Earth and Environment,

DRAFT

February 27, 2015, 9:20am

- Ayalew, D., C. Ebinger, E. Bourdon, E. Wolfenden, G. Yirgu, and N. Grassineau (2006),
- ²⁴⁸ Temporal compositional variation of syn-rift rhyolites along the western margin of the
- ²⁴⁹ southern Red Sea and northern Main Ethiopian Rift, *Geological Society, London, Special*
- $_{250}$ Publications, 259(1), 121-130.
- ²⁵¹ Bastow, I. D., and D. Keir (2011), The protracted development of the continent-ocean ²⁵² transition in Afar, *Nature Geoscience*, 4(4), 248–250.
- ²⁵³ Boschi, L., C. Weemstra, J. Verbeke, G. Ekström, A. Zunino, and D. Giardini (2013), On
- measuring surface wave phase velocity from station-station cross-correlation of ambient
 signal, *Geophysical Journal International*, 192(1), 346–358.
- Bosworth, W., P. Huchon, and K. McClay (2005), The Red Sea and Gulf of Aden Basins,
 Journal of African Earth Sciences, 43(1-3), 334–378.
- ²⁵⁸ Christensen, N. I., and W. D. Mooney (1995), Seismic velocity structure and composition
 of the continental crust: A global view, *Journal of Geophysical Research: Solid Earth* (1978–2012), 100(B6), 9761–9788.
- ²⁶¹ Corbeau, J., F. Rolandone, S. Leroy, A. Al-Lazki, D. Keir, G. Stuart, and A. Stork (2014),
- ²⁶² Uppermost mantle velocity from Pn tomography in the Gulf of Aden, *Geosphere*.
- ²⁶³ Daniels, K., I. Bastow, D. Keir, R. Sparks, and T. Menand (2014), Thermal models of
- ²⁶⁴ dyke intrusion during development of continent-ocean transition, *Earth and Planetary* ²⁶⁵ Science Letters, 385, 145–153.
- ²⁶⁶ Desissa, M., N. Johnson, K. Whaler, S. Hautot, S. Fisseha, and G. Dawes (2013), A
- ²⁶⁷ mantle magma reservoir beneath an incipient mid-ocean ridge in Afar, Ethiopia, *Nature*

University of Leeds, Leeds, U.K.

DRAFT

February 27, 2015, 9:20am

- $_{268}$ geoscience, 6(10), 861–865.
- Drake, C. L., and R. Girdler (1964), A geophysical study of the Red Sea, *Geophysical Journal International*, 8(5), 473–495.
- ²⁷¹ Ebinger, C., and M. Casey (2001), Continental breakup in magmatic provinces: An ²⁷² Ethiopian example, *Geology*, 29(6), 527–530.
- ²⁷³ Ebinger, C., D. Keir, A. Ayele, E. Calais, T. Wright, M. Belachew, J. Hammond, E. Camp-
- ²⁷⁴ bell, and W. Buck (2008), Capturing magma intrusion and faulting processes during
 ²⁷⁵ continental rupture: seismicity of the Dabbahu (Afar) rift, *Geophysical Journal Inter-*²⁷⁶ national, 174(3), 1138–1152.
- 277 Ebinger, C., A. Ayele, D. Keir, J. Rowland, G. Yirgu, T. Wright, M. Belachew, and
- I. Hamling (2010), Length and timescales of rift faulting and magma intrusion: the Afar
 rifting cycle from 2005 to present, Annual Review of Earth and Planetary Sciences, 38,
 439–466.
- 281 Egloff, F., R. Rihm, J. Makris, Y. Izzeldin, M. Bobsien, K. Meier, P. Junge, T. Noman,
- and W. Warsi (1991), Contrasting structural styles of the eastern and western margins of
- the southern Red Sea: the 1988 SONNE experiment, Tectonophysics, 198(2), 329-353.
- ²⁸⁴ Ekström, G., G. A. Abers, and S. C. Webb (2009), Determination of surface-wave phase
- velocities across USArray from noise and Aki's spectral formulation, *Geophysical Re-*
- ²⁸⁶ search Letters, 36(18), L18,301.

DRAFT

⁶Istituto Nazionale di Geofisica e

Ferguson, D. J., J. Maclennan, I. Bastow, D. Pyle, S. Jones, D. Keir, J. Blundy, T. Plank,

and G. Yirgu (2013), Melting during late-stage rifting in Afar is hot and deep, Nature,

Vulcanologia, via di Vigna Murata 605,

00143 Roma, Italy.

⁷Seismological and Volcanological

Observatory Center, Dhamar, Yemen.

⁸Sana'a University, Yemen Remote

Sensing and GIS Center, Sana'a, Yemen.

⁹Imperial College London, London, SW7,

U.K.

¹⁰University of Bristol, Bristol, BS8, U.K.

¹¹Institut de Physique du Globe de

Strasbourg; UMR 7516, Université de

Strasbourg/EOST, CNRS, 5 rue René

Descartes, F-67084 Strasbourg Cedex.

¹²Yemen Geological Survey and mineral

Resources Board, Sana'a, Yemen.

¹³School of Earth Sciences, University of

Bristol, Bristol, UK.

¹⁴Institute of Geophysics, Space Science and Astronomy, Addis Ababa University.

February 27, 2015, 9:20am

X - 18

- $_{289}$ 499(7456), 70–73.
- Fry, B., F. Deschamps, E. Kissling, L. Stehly, and D. Giardini (2010), Layered azimuthal
 anisotropy of Rayleigh wave phase velocities in the European Alpine lithosphere inferred
 from ambient noise, *Earth and Planetary Science Letters*, 297(1), 95–102.
- Grandin, R., A. Socquet, E. Jacques, N. Mazzoni, J.-B. de Chabalier, and G. King (2010),
 Sequence of rifting in Afar, Manda-Hararo rift, Ethiopia, 2005–2009: Time-space evolution and interactions between dikes from interferometric synthetic aperture radar and
 static stress change modeling, *Journal of Geophysical Research: Solid Earth (1978–2012)*, 115(B10).
- ²⁹⁸ Grandin, R., E. Jacques, A. Nercessian, A. Ayele, C. Doubre, A. Socquet, D. Keir, M. Kas-
- sim, A. Lemarchand, and G. King (2011), Seismicity during lateral dike propagation:
 Insights from new data in the recent Manda Hararo–Dabbahu rifting episode (Afar,
 Ethiopia), *Geochemistry, Geophysics, Geosystems, 12*(4).
- Halliday, D., and A. Curtis (2008), Seismic interferometry, surface waves and source distribution, *Geophysical Journal International*, 175(3), 1067–1087.
- Hammond, J., J.-M. Kendall, G. Stuart, D. Keir, C. Ebinger, A. Ayele, and M. Belachew
 (2011), The nature of the crust beneath the Afar triple junction: evidence from receiver
 functions, *Geochemistry, Geophysics, Geosystems, 12*(12).
- ³⁰⁷ Hammond, J., J.-M. Kendall, G. Stuart, C. Ebinger, I. Bastow, D. Keir, A. Ayele,
- M. Belachew, B. Goitom, G. Ogubazghi, and T. Wright (2013), Mantle upwelling and
- initiation of rift segmentation beneath the Afar Depression, Geology, 41(6), 635–638.

DRAFT

- Hayward, N., and C. Ebinger (1996), Variations in the along-axis segmentation of the Afar Rift system, *Tectonics*, 15(2), 244–257.
- Hillers, G., N. Graham, M. Campillo, S. Kedar, M. Landès, and N. Shapiro (2012),
 Global oceanic microseism sources as seen by seismic arrays and predicted by wave
 action models, *Geochemistry, Geophysics, Geosystems*, 13(1).
- ³¹⁵ Holtzman, B. K., J. Kendall, et al. (2010), Organized melt, seismic anisotropy, and plate ³¹⁶ boundary lubrication, *Geochemistry*, *Geophysics*, *Geosystems*, 11(12).
- ³¹⁷ Karato, S. i., et al. (2003), *The dynamic structure of the deep earth: an interdisciplinary* ³¹⁸ *approach*, Princeton University Press.
- ³¹⁹ Keir, D., I. Bastow, K. Whaler, E. Daly, D. Cornwell, and S. Hautot (2009), Lower crustal
- earthquakes near the Ethiopian rift induced by magmatic processes, *Geochem. Geophys.* Geosyst, 10(10).
- Keir, D., C. Pagli, I. Bastow, and A. Ayele (2011), The magma-assisted removal of Arabia in Afar: Evidence from dike injection in the Ethiopian rift captured using InSAR and seismicity, *Tectonics*, 30(2), TC2008.
- ³²⁵ Korostelev, F., C. Basuyau, S. Leroy, C. Tiberi, A. Ahmed, G. W. Stuart, D. Keir,
- F. Rolandone, I. Ganad, K. Khanbari, et al. (2014), Crustal and upper mantle structure
- ³²⁷ beneath south-western margin of the arabian peninsula from teleseismic tomography,
- Geochemistry, Geophysics, Geosystems, 15(7), 2850–2864.
- Laughton, A., and C. Tramontini (1969), Recent studies of the crustal structure in the Gulf of Aden, *Tectonophysics*, 8(4), 359–375.

DRAFT

X - 20

- Lebedev, S., and R. D. Van Der Hilst (2008), Global upper-mantle tomography with the automated multimode inversion of surface and S-wave forms, *Geophysical Journal*
- $_{333}$ International, 173(2), 505-518.
- ³³⁴ Leroy, S., F. Lucazeau, E. d'Acremont, L. Watremez, J. Autin, S. Rouzo, N. Bellahsen,
- ³³⁵ C. Tiberi, C. Ebinger, M. Beslier, et al. (2010), Contrasted styles of rifting in the ³³⁶ eastern Gulf of Aden: A combined wide-angle, multichannel seismic, and heat flow ³³⁷ survey, *Geochemistry Geophysics Geosystems*, 11(7), Q07,004.
- Longuet-Higgins, M. S. (1950), A theory of the origin of microseisms, *Philosophical Trans- actions of the Royal Society of London. Series A, Mathematical and Physical Sciences*,
 243(857), 1–35.
- Maccaferri, F., E. Rivalta, D. Keir, and V. Acocella (2014), Off-rift volcanism in rift zones
 determined by crustal unloading, *Nature Geoscience*.
- Makris, J., and A. Ginzburg (1987), The Afar Depression: transition between continental rifting and sea-floor spreading, *Tectonophysics*, 141(1), 199–214.
- Manetti, P., G. Capaldi, S. Chiesa, L. Civetta, S. Conticelli, M. Gasparon, L. Volpe,
 and G. Orsi (1991), Magmatism of the eastern Red Sea margin in the northern part of
 Yemen from Oligocene to present, *Tectonophysics*, 198(2), 181–202.
- McClusky, S., R. Reilinger, G. Ogubazghi, A. Amleson, B. Healeb, P. Vernant, J. Sholan,
- S. Fisseha, L. Asfaw, R. Bendick, et al. (2010), Kinematics of the southern Red Sea–Afar
 Triple Junction and implications for plate dynamics, *Geophys. Res. Lett*, 37, L05,301.
- ³⁵¹ Medynski, S., R. Pik, P. Burnard, C. Vye-Brown, L. France, I. Schimmelpfennig,
- K. Whaler, N. Johnson, L. Benedetti, D. Ayelew, et al. (2015), Stability of rift axis

DRAFT

February 27, 2015, 9:20am

- magma reservoirs: Spatial and temporal evolution of magma supply in the dabbahu rift
 segment (afar, ethiopia) over the past 30 kyr, *Earth and Planetary Science Letters*, 409,
 278–289.
- ³⁵⁶ Pagli, C., H. Wang, T. J. Wright, E. Calais, and E. Lewi (2014), Current plate boundary
- deformation of the Afar rift from a 3-D velocity field inversion of InSAR and GPS, Journal of Geophysical Research: Solid Earth, 119(11), 8562–8575.
- ³⁵⁹ Pallister, J., W. McCausland, S. Jónsson, Z. Lu, H. Zahran, S. El Hadidy, A. Aburukbah,
- I. Stewart, P. Lundgren, R. White, et al. (2010), Broad accommodation of rift-related
 extension recorded by dyke intrusion in Saudi Arabia, *Nature Geoscience*, 3(10), 705–
 712.
- Peccerillo, A., C. Donati, A. Santo, A. Orlando, G. Yirgu, and D. Ayalew (2007), Pet rogenesis of silicic peralkaline rocks in the Ethiopian rift: geochemical evidence and
 volcanological implications, *Journal of African Earth Sciences*, 48(2), 161–173.
- ³⁶⁶ Prodehl, C., and J. Mechie (1991), Crustal thinning in relationship to the evolution of the
- Afro-Arabian rift system: a review of seismic-refraction data, *Tectonophysics*, 198(2),
 311–327.
- Reed, C. A., S. Almadani, S. S. Gao, A. A. Elsheikh, S. Cherie, M. G. Abdelsalam, A. K.
- Thurmond, and K. H. Liu (2014), Receiver function constraints on crustal seismic veloc-
- ities and partial melting beneath the Red Sea rift and adjacent regions, Afar Depression,
 Journal of Geophysical Research: Solid Earth, 119(3), 2138–2152.
- ³⁷³ Rooney, T., T. Furman, I. Bastow, D. Ayalew, and G. Yirgu (2007), Lithospheric mod-
- ification during crustal extension in the Main Ethiopian Rift, J. Geophys. Res, 112,

B10,201. 375

X - 22

- Rooney, T., I. Bastow, and D. Keir (2011), Insights into extensional processes during 376
- magma assisted rifting: Evidence from aligned scoria cones, Journal of Volcanology and 377 Geothermal Research, 201(1-4), 83-96. 378
- Rooney, T., C. Herzberg, and I. Bastow (2012), Elevated mantle temperature beneath 379 East Africa, Geology, 40(1), 27–30. 380
- Rooney, T. O., T. Furman, G. Yirgu, and D. Avalew (2005), Structure of the Ethiopian 381
- lithosphere: Xenolith evidence in the Main Ethiopian Rift, Geochimica et Cosmochimica 382 Acta, 69(15), 3889–3910. 383
- Rooney, T. O., I. D. Bastow, D. Keir, F. Mazzarini, E. Movsesian, E. B. Grosfils, J. R. 384
- Zimbelman, M. S. Ramsey, D. Ayalew, and G. Yirgu (2014), The protracted develop-385 ment of focused magmatic intrusion during continental rifting, Tectonics. 386
- Ruegg, J. (1975), Main results about the crustal and upper mantle structure of the Dji-387 bouti region (TFAI), Afar Depression of Ethiopia, 1, 120–134. 388
- Shapiro, N. M., and M. Campillo (2004), Emergence of broadband Rayleigh waves from 389 correlations of the ambient seismic noise, *Geophysical Research Letters*, 31(7). 390
- Shapiro, N. M., M. Campillo, L. Stehly, and M. H. Ritzwoller (2005), High-resolution 391 surface-wave tomography from ambient seismic noise, Science, 307(5715), 1615–1618. 392
- Sicilia, D., J. Montagner, M. Cara, E. Stutzmann, E. Debayle, J. Lépine, J. Lévêque, 393
- E. Beucler, A. Sebai, G. Roult, et al. (2008), Upper mantle structure of shear-waves 394
- velocities and stratification of anisotropy in the Afar Hotspot region, Tectonophysics, 395 462(1-4), 164-177.

DRAFT

396

February 27, 2015, 9:20am

- ³⁹⁷ Stab, M., N. Bellahsen, R. Pik, S. Leroy, and D. Ayalew (2014), Mode of rifting in ³⁹⁸ magmatic-rich setting: Tectono-magmatic evolution of the central afar rift system, in
- ³⁹⁹ EGU General Assembly Conference Abstracts, vol. 16, p. 14484.
- ⁴⁰⁰ Stehly, L., M. Campillo, and N. Shapiro (2006), A study of the seismic noise from its long-
- range correlation properties, Journal of Geophysical Research: Solid Earth (1978–2012),
 111 (B10).
- Tramontini, C., and D. Davies (1969), A seismic refraction survey in the Red Sea, Geo-*physical Journal International*, 17(2), 225–241.
- ⁴⁰⁵ Verbeke, J., L. Boschi, L. Stehly, E. Kissling, and A. Michelini (2012), High-resolution
- A06 Rayleigh-wave velocity maps of central Europe from a dense ambient-noise data set,
- 407 Geophysical Journal International, 188(3), 1173–1187.
- ⁴⁰⁸ Vigny, C., P. Huchon, J.-C. Ruegg, K. Khanbari, and L. M. Asfaw (2006), Confirmation of
- Arabia plate slow motion by new GPS data in Yemen, Journal of Geophysical Research:
 Solid Earth (1978–2012), 111 (B2).
- ⁴¹¹ Wapenaar, K., and J. Fokkema (2006), Green's function representations for seismic inter-⁴¹² ferometry, *Geophysics*, 71(4), SI33–SI46.
- Wolfenden, E., C. Ebinger, G. Yirgu, A. Deino, and D. Ayalew (2004), Evolution of the
 northern Main Ethiopian rift: birth of a triple junction, *Earth and Planetary Science Letters*, 224 (1), 213–228.
- Wolfenden, E., C. Ebinger, G. Yirgu, P. Renne, and S. Kelley (2005), Evolution of a
 volcanic rifted margin: Southern Red Sea, Ethiopia, *Geological Society of America*Bulletin, 117(7-8), 846–864.

DRAFT

February 27, 2015, 9:20am



Figure 1. Structure of the Afar and southern Red Sea region. The crustal thicknesses are displayed by colored dots and based on the Moho depths, which are obtained from *Egloff et al.* [1991]; *Tramontini and Davies* [1969]; *Drake and Girdler* [1964]; *Prodehl and Mechie* [1991]; *Laughton and Tramontini* [1969]; *Ruegg* [1975]; *Hammond et al.* [2011]; *Ahmed et al.* [2013]; *Reed et al.* [2014]. Structures are modified from *Ebinger et al.* [2008] and *Stab et al.* [2014]. D R A F T Bathymetry is not represented. DMH: Dabbahu-Manda-Hararo volcano-tectonic segment; AEA: Afdera-Erta'Ale volcano-tectonic segment; G. Tadj.: Gulf of Tadjura; DbV: Dabbahu volcano; DrV: Durrie volcano; KV: Kurub volcano; TGD: Tendaho-Goba'ad Discontinuity; SVF: Sana'a volcanic field; MVF: Marib volcanic field; DVF: Dhamar volcanic field.



Figure 2. Map of the station pairs used for the tomographic inversion. The red lines show the station-to-station paths. The solid gray line delimitates the best constrained area. The green triangles are the stations.

February 27, 2015, 9:20am



Figure 3. Result of two reconstruction synthetic tests with randomly distributed velocity anomalies of various size as input. a. Small-scale synthetic anomalies; b. large-scale synthetic anomalies. The left image displays the synthetic input, whereas the right image displays the output model.

DRAFT

February 27, 2015, 9:20am



Figure 4. Maps of phase velocity anomalies (% with respect to average) resulting from tomographic inversion of ambient noise dispersion data. The average velocity for each period is In RhA bfotTom right of each map (Fre/s)uaFlye 27 lid20145, lide 20 diminitates the best construinfed AaFeaT The green triangles are the stations.

Anomalies (%)



Figure 5. Distribution of thermal wells, fumaroles and thermal springs in the Afar triple junction region [*Keir et al.*, 2009]. No data was available in Yemen. The green triangles are the stations.

DRAFT

February 27, 2015, 9:20am