



HAL
open science

Full statistics of energy conservation in two times measurement protocols

Tristan Benoist, Vojkan Jaksic, Annalisa Panati, Yan Pautrat, Claude-Alain
Pillet

► **To cite this version:**

Tristan Benoist, Vojkan Jaksic, Annalisa Panati, Yan Pautrat, Claude-Alain Pillet. Full statistics of energy conservation in two times measurement protocols. *Physical Review E: Statistical, Nonlinear, and Soft Matter Physics*, 2015, 92 (3), pp.032115. 10.1103/PhysRevE.92.032115 . hal-01137957

HAL Id: hal-01137957

<https://hal.science/hal-01137957>

Submitted on 29 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Full statistics of energy conservation in two times measurement protocols

Tristan Benoist,^{1,2} Vojkan Jakšić,¹ Annalisa Panati,^{1,3} Yan Pautrat,⁴ and Claude-Alain Pillet³

¹*Department of Mathematics and Statistics, McGill University,
805 Sherbrooke Street West, Montreal, QC, H3A 2K6, Canada*

²*CNRS, Laboratoire de Physique Théorique, IRSAMC,
Université de Toulouse, UPS, F-31062 Toulouse, France*

³*Aix-Marseille Université, CNRS, CPT, UMR 7332, Case 907, 13288 Marseille, France
Université de Toulon, CNRS, CPT, UMR 7332, 83957 La Garde, France*

⁴*Laboratoire de Mathématiques, Université Paris-Sud, 91405 Orsay Cedex, France*

The first law of thermodynamics states that the average total energy current between different reservoirs vanishes at large times. In this note we examine this fact at the level of the full statistics of two times measurement protocols also known as the Full Counting Statistics. Under very general conditions, we establish a tight form of the first law asserting that the fluctuations of the total energy current computed from the energy variation distribution are exponentially suppressed in the large time limit. We illustrate this general result using two examples: the Anderson impurity model and a 2D spin lattice model.

PACS numbers: 02.50.Cw, 03.65.-w, 05.30.-d

Recent technical advances in the control of nanoscale systems have enabled the experimental study of out of equilibrium thermodynamics in the quantum regime [1–9]. These new experiments allow for the assessment of fluctuations in addition to the mean heat and particle currents, thus leading to a renewed theoretical investigation of the related quantum thermodynamic laws.

One of the basic questions in this context concerns the energy flow between two initially isolated large systems A and B . The purpose of this note is to study some consequences of energy conservation on the statistical properties of this flow.

By the first law, the average work performed by the interaction coupling the two systems is equal to the average heating of the combined system:

$$\Delta W_t = \Delta Q_t = \Delta Q_t^A + \Delta Q_t^B.$$

In the case of a sudden switching on/off of the interaction V , the average heating is given by

$$\Delta Q_t = -\langle V \rangle_t + \langle V \rangle_0 \quad (1)$$

where $\langle \cdot \rangle_t$ denotes the expectation with respect to a suitable system state at time t . Whenever V is bounded, (1) gives

$$\lim_{t \rightarrow \infty} \frac{\Delta Q_t}{t} = 0. \quad (2)$$

The individual energy currents $\Delta Q_t^{A/B}/t$ are also expected to reach steady values $J^{A/B}$. They satisfy $J^A = -J^B$, and are non-vanishing for systems out of equilibrium.

This note concerns the statistical character of the first law related to the thermodynamics of open quantum systems at the mesoscopic scale. Our main result is a refinement of relation (2). It states that the fluctuations of the total energy current are exponentially suppressed in the large time limit.

The nature of work in quantum physics is more subtle than in classical physics [10]. In the 1990's Lesovik and Levitov introduced the concept of the Full Counting Statistics (FCS) in the study of charge transport [11]. The use of the FCS in the definition of work in quantum physics appeared in the early 2000's in the works of J. Kurchan and H. Tasaki on the extension of the fluctuation relations to quantum systems [12, 13]. The emerging idea is that in quantum mechanics work should not be understood as an observable. Instead, the work performed during a given time period is identified with the energy variation ΔE observed in a repeated measurement protocol where the system energy is measured at the beginning and at the end of the period. The distribution of the measured energy variation, $\mathbb{P}_t(\Delta E)$, is the work FCS (we comment on terminology in footnote [14]). This change of perspective opened a whole new area of research [15, 16]. In particular, it allowed for the extension of the fluctuation relations to quantum systems [12, 13, 15–18].

The fluctuation relations are intimately related to the second law of thermodynamics and have been extensively studied [12, 13, 15–20]. Regarding the first law, the well known identity

$$\mathbb{E}_t(\Delta E) = \Delta Q_t$$

and (2) give

$$\lim_{t \rightarrow \infty} \mathbb{E}_t \left(\frac{\Delta E}{t} \right) = 0 \quad (3)$$

where \mathbb{E}_t denotes the expectation with respect to the FCS distribution \mathbb{P}_t [10, 18]. In this note we sharpen (3) by showing that, under very general conditions, the exponential moment

$$\mathbb{E}_t \left(e^{\alpha_m |\Delta E|} \right)$$

remains bounded as $t \rightarrow \infty$ where the constant $\alpha_m > 0$ is a measure of the regularity of the interaction V (see (5) below).

Until recently, the first law and energy conservation in the FCS setting have received little attention in the literature. In the case where A and B are thermal reservoirs, the FCS of the total energy current was previously studied theoretically in [21]. The works [22, 23] concern the FCS of energy transfer in the thermalization process of a finite level quantum system in contact with a thermal bath, a problem which is radically different from the one considered here. We also emphasize that here we are only interested in the FCS of the total energy, and not in the FCS of the individual energy variations $\Delta E^{A/B}$.

We start with a system described by a finite dimensional Hilbert space $\mathcal{H}^{(L)}$ where the superscript L refers to the size of the system. Taking $L \rightarrow \infty$ corresponds to the thermodynamic limit. The limiting objects will be denoted without the superscript. Let $H^{(L)} = H_A^{(L)} + H_B^{(L)}$ be the Hamiltonian of the joint but non-interacting system $A+B$. The evolution between the two measurements of $H^{(L)}$ is generated by $H_V^{(L)} = H^{(L)} + V^{(L)}$, where $V^{(L)}$ denotes the interaction coupling A and B . The initial state is described by the density matrix $\rho^{(L)}$.

Let $P_e^{(L)}$ denote the projection on the eigenspace associated to the eigenvalue e in the spectrum $\text{sp}(H^{(L)})$. The measurement of $H^{(L)}$ at initial time $t = 0$ gives e with probability $\text{tr}(P_e^{(L)} \rho^{(L)})$. After the measurement the system is in the projected state

$$P_e^{(L)} \rho^{(L)} P_e^{(L)} / \text{tr}(P_e^{(L)} \rho^{(L)}).$$

The second measurement of $H^{(L)}$ at a later time t gives e' with probability

$$\text{tr} \left(P_{e'}^{(L)} e^{-itH_V^{(L)}} P_e^{(L)} \rho^{(L)} P_e^{(L)} e^{itH_V^{(L)}} \right) / \text{tr}(P_e^{(L)} \rho^{(L)}).$$

It follows that the probability of observing the energy variation ΔE in this measurement protocol is

$$\mathbb{P}_t^{(L)}(\Delta E) = \sum_{e' - e = \Delta E} \text{tr} \left(P_{e'}^{(L)} e^{-itH_V^{(L)}} P_e^{(L)} \rho^{(L)} P_e^{(L)} e^{itH_V^{(L)}} \right).$$

The moment generating function of the Full Counting Statistics $\mathbb{P}_t^{(L)}$ is

$$\begin{aligned} \chi_t^{(L)}(\alpha) &= \int_{\mathbb{R}} e^{\alpha \Delta E} d\mathbb{P}_t^{(L)}(\Delta E) \\ &= \text{tr} \left(e^{\alpha H^{(L)}} e^{-itH_V^{(L)}} e^{-\alpha H^{(L)}} \tilde{\rho}^{(L)} e^{itH_V^{(L)}} \right) \end{aligned}$$

where

$$\tilde{\rho}^{(L)} = \sum_{e \in \text{sp}(H^{(L)})} P_e^{(L)} \rho^{(L)} P_e^{(L)}.$$

We assume that for α purely imaginary, the limit

$$\lim_{L \rightarrow \infty} \chi_t^{(L)}(\alpha) = \chi_t(\alpha) \quad (4)$$

exists and is a continuous function of α . This assumption is harmless and easy to verify in most concrete models of physical interest. By Levy's continuity theorem [24], (4) implies that the thermodynamic limit $\lim_{L \rightarrow \infty} \mathbb{P}_t^{(L)} = \mathbb{P}_t$ exists. The probability distribution \mathbb{P}_t is the FCS of the thermodynamic system.

Let

$$R^{(L)}(\alpha) = 2|\alpha| \max_{-1 \leq s \leq 1} \|e^{s \frac{\alpha}{2} H^{(L)}} V^{(L)} e^{-s \frac{\alpha}{2} H^{(L)}}\|$$

and

$$R(\alpha) = \sup_L R^{(L)}(\alpha).$$

Note that $R(\alpha)$ takes values in $[0, \infty]$ and is an even function. Moreover, $R(\alpha) \geq R(\alpha')$ if $\alpha \geq \alpha' \geq 0$. Our regularity condition is that there exists $\alpha_m > 0$ such that

$$R(\alpha_m) < \infty. \quad (5)$$

We emphasize that (5) is the only regularity assumption we require and that no further hypothesis on the dynamical behaviour of the system is needed. We also make no assumptions on the initial state of the system.

Our main result is the following strengthening of (3):

Theorem For all $t > 0$,

$$\mathbb{E}_t \left(e^{\alpha_m |\Delta E|} \right) \leq 2e^{R(\alpha_m)}. \quad (6)$$

An immediate consequence of this result and Chebyshev's inequality [24] is that for any $\epsilon > 0$,

$$\mathbb{P}_t \left(\frac{|\Delta E|}{t} \geq \epsilon \right) \leq 2e^{-t\epsilon\alpha_m + R(\alpha_m)}. \quad (7)$$

Note that if $R(\alpha) < \infty$ for all α , then

$$\mathbb{P}_t \left(\frac{|\Delta E|}{t} \geq \epsilon \right) \leq 2e^{R(C/\epsilon) - Ct} \quad (8)$$

for any $C > 0$.

The estimates (7) and (8) can be interpreted in terms of the large deviation theory [25] (see [26]). For example, (8) implies that the large deviation rate function of the random variable $|\Delta E|/t$ satisfies $I(s) = \infty$ for $s \neq 0$, and that the large deviations are completely suppressed in the large time limit.

The main novelty of our proof is the derivation of a time independent bound for $\chi_t^{(L)}$ inspired by the bounds proposed in [21]. The derivation is based on two well-known inequalities. The first is

$$\text{tr}(XY) \leq \|X\| \text{tr}(Y)$$

which holds for any two non-negative matrices X, Y . The second states that for any two self-adjoint matrices T, S ,

$$\|e^{T+S} e^{-T}\| \leq e^{\max_{0 \leq s \leq 1} \|e^{sT} S e^{-sT}\|}. \quad (9)$$

To prove this inequality, let $\Gamma(s) = e^{s(T+S)}e^{-sT}$. Then one has

$$\partial_s \Gamma(s) = \Gamma(s)e^{sT}Se^{-sT}, \quad \Gamma(0) = \mathbb{I}.$$

Using

$$\|\partial_s \Gamma(s)\| \leq \|\Gamma(s)\| \|e^{sT}Se^{-sT}\|$$

and Gronwall's inequality we obtain (9). The bound (9) is similar but unrelated to the bound (3.10) of [27].

The proof of (6) proceeds as follows. For α real we set

$$X = e^{-\frac{\alpha}{2}H_V^{(L)}} e^{\alpha H^{(L)}} e^{-\frac{\alpha}{2}H_V^{(L)}}$$

and

$$Y = e^{-itH_V^{(L)}} e^{\frac{\alpha}{2}H_V^{(L)}} e^{-\frac{\alpha}{2}H^{(L)}} \tilde{\rho}^{(L)} e^{-\frac{\alpha}{2}H^{(L)}} e^{\frac{\alpha}{2}H_V^{(L)}} e^{itH_V^{(L)}}$$

(note that $\tilde{\rho}^{(L)}$ and $H^{(L)}$ commute). Observe that

$$\chi_t^{(L)}(\alpha) = \text{tr}(XY)$$

and that X, Y are non-negative matrices. We then use the first inequality to derive the estimate

$$\chi_t^{(L)}(\alpha) \leq \|X\| \text{tr}(Y)$$

where

$$\|X\| = \|e^{-\frac{\alpha}{2}H_V^{(L)}} e^{\frac{\alpha}{2}H^{(L)}}\|^2$$

and

$$\text{tr}(Y) = \text{tr}\left(e^{\frac{\alpha}{2}H_V^{(L)}} e^{-\frac{\alpha}{2}H^{(L)}} \tilde{\rho}^{(L)} e^{-\frac{\alpha}{2}H^{(L)}} e^{\frac{\alpha}{2}H_V^{(L)}}\right).$$

The cyclicity of the trace gives

$$\text{tr}(Y) = \text{tr}\left(e^{-\frac{\alpha}{2}H^{(L)}} e^{\alpha H_V^{(L)}} e^{-\frac{\alpha}{2}H^{(L)}} \tilde{\rho}^{(L)}\right).$$

Applying the first inequality once again and using that $\text{tr}(\tilde{\rho}^{(L)}) = 1$, we derive

$$\text{tr}(Y) \leq \|e^{\frac{\alpha}{2}H_V^{(L)}} e^{-\frac{\alpha}{2}H^{(L)}}\|^2.$$

Hence

$$\chi_t^{(L)}(\alpha) \leq \|e^{-\frac{\alpha}{2}H_V^{(L)}} e^{\frac{\alpha}{2}H^{(L)}}\|^2 \|e^{\frac{\alpha}{2}H_V^{(L)}} e^{-\frac{\alpha}{2}H^{(L)}}\|^2.$$

Using the second inequality with

$$T = \mp \frac{\alpha}{2}H^{(L)}, \quad S = \mp \frac{\alpha}{2}V^{(L)},$$

we obtain

$$\chi_t^{(L)}(\alpha) \leq e^{R^{(L)}(\alpha)}.$$

The regularity assumption (5), the existence of the limit (4) for purely imaginary α 's, and Vitali's convergence

theorem (see Appendix B in [18]) give that for all complex α with real part $\text{Re}(\alpha)$ in $(-\alpha_m, \alpha_m)$, the limit $\lim_{L \rightarrow \infty} \chi_t^{(L)}(\alpha) = \chi_t(\alpha)$ exists. Moreover, for such α 's,

$$\chi_t(\alpha) = \int_{\mathbb{R}} e^{\alpha \Delta E} d\mathbb{P}_t(\Delta E)$$

and

$$|\chi_t(\alpha)| \leq e^{R(\text{Re}(\alpha))}.$$

It follows that

$$|\chi_t(\pm \alpha_m)| \leq e^{R(\alpha_m)}.$$

The last estimate gives

$$\mathbb{E}_t\left(e^{\alpha_m |\Delta E|}\right) \leq \chi_t(-\alpha_m) + \chi_t(\alpha_m) \leq 2e^{R(\alpha_m)}$$

and the theorem follows.

a. Spin-fermion models. Electronic transport through a 1D-lattice containing a single magnetic impurity is a typical problem involving bounded interactions. The Anderson model [28, 29] commonly used to study this question is a specific example of a general class of spin-fermion models to which our main theorem applies.

The study of the FCS of charge transport through the impurities in such models is an active field of research [30–34]. We emphasize, however, that here we are only concerned with the statistics of the total energy.

The impurity is described by a quantum dot supporting four different eigenstates: empty, occupied by a single electron with either spin up or spin down, or occupied by two electrons with opposite spins. The remaining parts of the lattice, regarded as fermionic (say left and right) reservoirs at different chemical potentials, are described in the tight binding approximation.

Here, the subsystem A is the left side of the lattice together with the impurity. The lattice right side is the subsystem B .

The operator $c_{l/r,\sigma}^*(x)$ ($c_{l/r,\sigma}(x)$) creates (annihilates) an electron with spin σ at the lattice site x of the left ($x < 0$)/right ($x > 0$) reservoir. Similarly, the operator d_σ^* (d_σ) creates (annihilates) an electron with spin σ in the dot. The anti-commutation relations $\{c_{l/r,\sigma}(x), c_{l/r,\sigma'}^*(x')\} = \delta_{x,x'} \delta_{\sigma,\sigma'}$ and $\{d_\sigma, d_{\sigma'}^*\} = \delta_{\sigma,\sigma'}$ hold while the c operators commute with the d operators. We use the shorthand $c_{l/r,\sigma}(\phi) = \sum_x \bar{\phi}(x) c_{l/r,\sigma}(x)$. The reservoir Hamiltonians are

$$H_l = \sum_{\substack{\sigma=\pm; x,x'<0 \\ |x-x'|=1}} c_{l,\sigma}^*(x) c_{l,\sigma}(x'),$$

with a similar expression for H_r . Let $h_{l/r}$ be the discrete Laplacian of the left/right part of the lattice. Since $h_{l/r}$ is a bounded operator,

$$e^{\alpha H_{l/r}} c_{l/r,\sigma}(\phi) e^{-\alpha H_{l/r}} = c_{l/r,\sigma}(e^{\alpha h_{l/r}} \phi)$$

for all real α . In particular, for all α ,

$$\|e^{\alpha H_{l/r}} c_{l/r,\sigma}(\phi) e^{-\alpha H_{l/r}}\| < \infty. \quad (10)$$

The total Hamiltonian is

$$H = H_S + H_l + H_r$$

where $H_S = \epsilon \sum_{\sigma} d_{\sigma}^* d_{\sigma} + U d_{+}^* d_{+} d_{-}^* d_{-}$ is the Hamiltonian of the dot. Regarding the subdivision in A/B subsystem, we have $H_A = H_l + H_S$ and $H_B = H_r$. The coupling of the conduction electrons with the dot is described by

$$V = \sum_{\sigma} (d_{\sigma}^* (c_{l,\sigma}(v_{l,\sigma}) + c_{r,\sigma}(v_{r,\sigma})) + \text{h.c.})$$

for some coupling functions $v_{l/r,\sigma}(x)$. In the context of the Anderson model, the superscript L refers to the confinement of the reservoirs to the finite part of the lattice defined by $|x| \leq L$. Such confinement is necessary to allow for a meaningful definition of the repeated measurement protocol leading to the FCS. The limit $L \rightarrow \infty$ restores the extended reservoirs. It follows from relation (10) that $R(\alpha)$ is finite for all α , and that our theorem holds for all $\alpha_m > 0$. Hence we have inequality (8):

$$\mathbb{P}_t \left(\frac{|\Delta E|}{t} \geq \epsilon \right) \leq 2e^{R(C/\epsilon) - Ct}$$

for any $\epsilon > 0$ and any $C > 0$.

We also note that one can consider instead the FCS of $H' = H_l + H_r$ by setting $V' = H_S + V$. Then $H'_A = H_l$ and $H'_B = H_r$. One then obtains the same result by replacing ΔE with $\Delta E'$. The energy of the impurity is irrelevant in the large time limit.

b. Spin systems. Another popular class of models involving bounded interactions are locally interacting spin systems. In [26] we prove that, under general conditions, our theorem applies to locally interacting spin systems in arbitrary dimension. Moreover, for 1D spin systems with finite range interactions, Araki's results [35] give that $R(\alpha) < \infty$ for all α , and hence that our theorem holds for all $\alpha_m > 0$. We restrict ourselves to the description of a simple example.

Consider a 2D square lattice of $\frac{1}{2}$ -spins. Let $\Lambda_L \subset \mathbb{Z}^2$ be the finite sub-lattice of size $2L \times 2L$. We denote by Λ_L^{\pm} its left/right half. Subsystems A and B are the spins in Λ_L^- and Λ_L^+ respectively (see Figure 1).

The system Hilbert space is $\mathcal{H}^{(L)} = \bigotimes_{x \in \Lambda_L} \mathbb{C}^2$. The Hamiltonian is that of an XY-spin model where the spins on Λ_L^- do not interact with that on Λ_L^+ [36]:

$$H^{(L)} = H^{(L,-)} + H^{(L,+)},$$

with

$$H^{(L,\pm)} = -\frac{J}{2} \sum_{\substack{x,y \text{ nearest} \\ \text{neighbors in } \Lambda_L^{\pm}}} \left(\sigma_x^{(1)} \sigma_y^{(1)} + \sigma_x^{(2)} \sigma_y^{(2)} \right),$$




FIG. 1. A partitioned finite spin system $A + B$. Solid lines represent the nearest neighbour coupling J and dashed lines the interaction $K_{x,y}$ between the 2 subsystems.

where J is a coupling constant. The interaction is

$$V^{(L)} = -\frac{1}{2} \sum_{x \in \Lambda_L^-, y \in \Lambda_L^+} K_{x,y} (\sigma_x^{(1)} \sigma_y^{(1)} + \sigma_x^{(2)} \sigma_y^{(2)}),$$

where

$$K_{x,y} = \frac{\epsilon}{1 + x_2^2}$$

if $x = (0, x_2) \in \Lambda_L^-$ and $y = (1, x_2) \in \Lambda_L^+$ and $K_{x,y} = 0$ otherwise. The boundary between the two halves of the lattice is between the lines $x_1 = 0$ and $x_1 = 1$. Note that the interaction intensity decreases as one moves away from $(0, 0)$. An assumption of this type is necessary if $V^{(L)}$ is to remain bounded in the thermodynamic limit $L \rightarrow \infty$.

For this model one can show that there exists $\alpha_m > 0$ such that (5) holds and that our theorem applies. Hence we have inequality (7):

$$\mathbb{P}_t \left(\frac{|\Delta E|}{t} \geq \epsilon \right) \leq 2e^{-t\epsilon\alpha_m + R(\alpha_m)}$$

for any $\epsilon > 0$.

c. Discussion. Under a general condition on the regularity of the interaction evolution in imaginary time, we have proven a sharp form of the first law of thermodynamics for the FCS of energy variation.

Our result holds for any initial state of the system. If one assumes that systems A and B are initially in thermal equilibrium at temperatures T_A and T_B , then the suppression of the fluctuations of the total energy current can be also proven by following the arguments of [21].

Under additional assumptions it is possible to deal with cases where several reservoirs drive the joint system towards a non-equilibrium steady state and to derive properties of the joint distribution of the energy variations in each part of the system. A more strict condition on $R(\alpha)$ allows for the generalization of a symmetry of the limiting cumulant generating function proposed in [21]. Combined with time reversal invariance this leads to Onsager's reciprocity relations. We investigate these topics in [26].

In the present note we have limited ourselves to bounded interactions. The case of unbounded interactions (an example is the spin-boson model) is more technical and requires a separate analysis based on an application of Ruelle’s quantum transfer operators [18]. Although the physical picture emerging from this analysis is of an independent interest, the final results are much less general than in the case of bounded interactions [37, 38].

ACKNOWLEDGMENTS

The research of T.B. was partly supported by ANR project RMTQIT (grant ANR-12-IS01-0001-01). T.B.

also wishes to thank Technische Universität München and Pr. M. M. Wolf for his hospitality during the last weeks of work on this note. The research of T.B. and Y.P. was partly supported by ANR contract ANR-14-CE25-0003-0. Y.P. also wishes to thank UMI-CRM for financial support, and McGill University for its hospitality. The research of T.B. and V.J. was partly supported by NSERC. The research of A.P. was partly supported by NSERC and ANR (grant 12-JS01-0008-01). The research of C.-A.P has been carried out in the framework of the Labex Archimède (ANR-11-LABX-0033) and of the A*MIDEX project (ANR-11-IDEX-0001-02), funded by the “Investissements d’Avenir” French Government programme managed by the French National Research Agency (ANR).

-
- [1] J. P. Pekola, *Nature Phys.* **11**, 118 (2015).
 [2] C. Bustamante, J. Liphardt, and F. Ritort, *Phys. Today* **58**, 43 (2005).
 [3] Y. Dubi and M. Di Ventra, *Rev. Mod. Phys.* **83**, 131 (2011).
 [4] S. Ciliberto, A. Imparato, A. Naert, and M. Tanase, *Phys. Rev. Lett.* **110**, 180601 (2013).
 [5] S. Jezouin, F. D. Parmentier, A. Anthore, U. Gennser, A. Cavanna, Y. Jin, and F. Pierre, *Science* **342**, 601 (2013).
 [6] A. Brut, A. Arakelyan, A. Petrosyan, S. Ciliberto, R. Dillenschneider, and E. Lutz, *Nature* **483**, 187 (2012).
 [7] S. Toyabe, T. Sagawa, M. Ueda, E. Muneyuki, and M. Sano, *Nature Phys.* **6**, 988 (2010).
 [8] J. V. Koski, T. Sagawa, O.-P. Saira, Y. Yoon, A. Kutvonen, P. Solinas, M. Mtnen, T. Ala-Nissila, and J. P. Pekola, *Nature Phys.* **9**, 644 (2013).
 [9] B. Kng, C. Rssler, M. Beck, M. Marthaler, D. S. Golubev, Y. Utsumi, T. Ihn, and K. Ensslin, *Phys. Rev. X* **2**, 011001 (2012).
 [10] P. Talkner, E. Lutz, and P. Hänggi, *Phys. Rev. E* **75**, 050102 (2007).
 [11] L. Levitov and G. Lesovik, *JETP Lett.* **58**, 230 (1993).
 [12] J. Kurchan, arXiv:cond-mat/0007360 (2000).
 [13] H. Tasaki, arXiv:cond-mat/0009244 (2000).
 [14] The use of term *Counting* in the above context is slightly misleading. In non-trivial cases, the energy variation (or work) ΔE is not a discrete quantity in the thermodynamic limit. Nevertheless the name *Full Counting Statistics* is usually used in the literature for the distribution emerging from the repeated measurement protocol we just described.
 [15] M. Esposito, U. Harbola, and S. Mukamel, *Rev. Mod. Phys.* **81**, 1665 (2009).
 [16] M. Campisi, P. Hänggi, and P. Talkner, *Rev. Mod. Phys.* **83**, 771 (2011).
 [17] C. Jarzynski and D. K. Wójcik, *Phys. Rev. Lett.* **92**, 230602 (2004).
 [18] V. Jakšić, Y. Ogata, Y. Pautrat, and C.-A. Pillet, “Quantum theory from small to large scales,” (Oxford University Press, Oxford, 2012) Chap. Entropic fluctuations in quantum statistical mechanics – an introduction.
 [19] G. Crooks, *J. Stat. Mech.*, P10023 (2008).
 [20] D. J. Evans and D. J. Searles, *Phys. Rev. E* **50**, 1645 (1994).
 [21] D. Andrieux, P. Gaspard, T. Monnai, and S. Tasaki, *New J. Phys.* **11**, 043014 (2009).
 [22] V. Jakšić, J. Panangaden, A. Panati, and C.-A. Pillet, arXiv:1409.8610 (to appear in *Lett. Math. Phys.*) (2014).
 [23] T. Benoist, M. Fraas, and V. Jakšić, (2015), in preparation.
 [24] P. Billingsley, *Convergence of Probability Measures* (Wiley, 1968).
 [25] A. Dembo and O. Zeitouni, *Large deviations techniques and applications*, Vol. 38 (Springer Science & Business Media, 2009).
 [26] T. Benoist, V. Jakšić, A. Panati, Y. Pautrat, and C.-A. Pillet, (2015), in preparation.
 [27] M. Lenci and L. Rey-Bellet, *J. Stat. Phys.* **119**, 715 (2005).
 [28] P. W. Anderson, *Phys. Rev.* **124**, 41 (1961).
 [29] A. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, 1993).
 [30] A. O. Gogolin and A. Komnik, *Phys. Rev. Lett.* **97**, 016602 (2006).
 [31] A. O. Gogolin and A. Komnik, *Phys. Rev. B* **73**, 195301 (2006).
 [32] T. L. Schmidt, A. O. Gogolin, and A. Komnik, *Phys. Rev. B* **75**, 235105 (2007).
 [33] T. L. Schmidt, A. Komnik, and A. O. Gogolin, *Phys. Rev. B* **76**, 241307 (2007).
 [34] R. Sakano, A. Oguri, T. Kato, and S. Tarucha, *Phys. Rev. B* **83**, 241301 (2011).
 [35] H. Araki, *Commun. Math. Phys.* **14**, 120 (1969).
 [36] The matrices $\sigma_x^{(j)}$ act non trivially only on the site x copy of \mathbb{C}^2 with the corresponding Pauli matrix: $\sigma_x^{(j)} = \bigotimes_{y \in \mathbb{Z}^2, x \neq y} \text{Id} \otimes \sigma^{(j)}$.

$$\sigma^{(1)} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma^{(2)} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma^{(3)} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

 [37] W. De Roeck, *Rev. Math. Phys.* **21**, 549 (2009).
 [38] V. Jakšić, A. Panati, Y. Pautrat, and C.-A. Pillet, “Non-equilibrium statistical mechanics of pauli-fierz systems,” (2015), in preparation.