



**HAL**  
open science

## Stereocontrolled lithiation/trapping of chiral 2-alkylideneaziridines: investigation into the role of the aziridine nitrogen stereodynamics

Rosmara Mansueto, Leonardo Degennaro, Jean-François Brière, Karen Griffin,  
Michael Shipman, Saverio Florio, Renzo Luisi

► **To cite this version:**

Rosmara Mansueto, Leonardo Degennaro, Jean-François Brière, Karen Griffin, Michael Shipman, et al.. Stereocontrolled lithiation/trapping of chiral 2-alkylideneaziridines: investigation into the role of the aziridine nitrogen stereodynamics. *Organic & Biomolecular Chemistry*, 2014, 12 (42), pp.8505-8511. 10.1039/c4ob01457j . hal-01141457

**HAL Id: hal-01141457**

**<https://hal.science/hal-01141457>**

Submitted on 13 Apr 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Cite this: *Org. Biomol. Chem.*, 2014, **12**, 8505

## Stereocontrolled lithiation/trapping of chiral 2-alkylideneaziridines: investigation into the role of the aziridine nitrogen stereodynamics†‡

Rosmara Mansueto,<sup>a,c</sup> Leonardo Degennaro,<sup>a</sup> Jean-François Brière,<sup>b</sup> Karen Griffin,<sup>c</sup> Michael Shipman,<sup>c</sup> Saverio Florio<sup>a</sup> and Renzo Luisi\*<sup>a</sup>

The origin of the stereoselectivity in the lithiation/trapping of 2-alkylideneaziridines bearing a chiral group as the nitrogen substituent was investigated. Optimal reaction conditions were discovered by *in situ* FT-IR monitoring. In addition, it has been found that the solvent and the alkene substitution pattern are important factors able to impart a switch in stereoselectivity. While lithiation of the alkylideneaziridine ring flanked by either a fully substituted or a *Z*-configured alkene pendant occurs stereoselectively in THF, in contrast unsubstituted 2-methyleneaziridine undergoes lithiation in toluene with the opposite sense of stereoselection. Lithiation experiments, on deuterium labelled 2-alkylideneaziridines, confirmed the configurational stability of the lithiated intermediates. A model based on complexation and proximity effects was proposed to rationalize the reactivity. This model assumes that slowly equilibrating *N*-invertomers undergo deprotonation (lithiation) at different rates and that the stereochemical outcome is established during the deprotonation step.

Received 10th July 2014,  
Accepted 26th August 2014  
DOI: 10.1039/c4ob01457j

www.rsc.org/obc

### Introduction

Aziridines are important building blocks in organic synthesis, and much research over the past few decades has been devoted to the development of effective synthetic methodologies for their preparation.<sup>1</sup> A special class of functionalized aziridines is represented by 2-alkylideneaziridines. These highly strained three-membered heterocycles possess a rich and varied chemistry.<sup>2</sup> For example, 2-methyleneaziridines participate in multi-component reactions,<sup>3</sup> Lewis acid promoted [3 + 2] and [4 + 3] cycloadditions,<sup>4</sup> oxidative rearrangements,<sup>5</sup> radical cascades,<sup>6</sup> and a growing number of transition metal catalysed processes.<sup>7</sup> An effective method for the preparation of these *N*-heterocycles relies on the intramolecular cyclization of readily available 2-bromoallylamines with sodium amide in liquid ammonia.<sup>8</sup> Whilst this method tolerates substantial variation in the substitution pattern of the

methyleneaziridine framework, it is not suitable for the synthesis of derivatives bearing substituents at C-3 (*i.e.*  $R^3 \neq H$ ). A potentially simple method to make such derivatives exploits the aziridinyl anion strategy<sup>9</sup> wherein C-3 of the 2-methyleneaziridine is functionalized through deprotonation, with a suitable base, followed by trapping with an electrophilic species (Scheme 1).<sup>10</sup> Using this method, Shipman and coworkers accessed functionalized aziridines bearing different substituents on the aziridine nitrogen and at the double bond.<sup>4a,c,11</sup> Using alkylideneaziridines bearing a chiral group as the *N*-substituent, the stereoselectivity of this lithiation/functionalization sequence was addressed, and it was found that the stereoselectivity of this process was dependent on the geometry (*i.e.* *E* or *Z*) and the pattern of substituents on the alkene double bond.<sup>12</sup> Here we report detailed optimization studies, including *in situ* FT-IR experiments that provide a better mechanistic understanding of the process, and reaction conditions that can deliver improved levels of stereocontrol.


<sup>a</sup>Department of Pharmacy – Drug Sciences University of Bari “A. Moro”, Via E. Orabona 4, 70125 Bari, Italy. E-mail: renzo.luisi@uniba.it

<sup>b</sup>Normandie Univ, COBRA, UMR 6014 et FR 3038; Univ Rouen; INSA Rouen; CNRS, IRCOF, 1 rue Tesnière, 76821 Mont Saint Aignan Cedex, France

<sup>c</sup>Department of Chemistry, University of Warwick, Gibbet Hill Road, Coventry, CV4 7AL, UK

† This work is dedicated to Prof. Richard Taylor on the occasion of his 65<sup>th</sup> birthday.

‡ Electronic supplementary information (ESI) available. See DOI: 10.1039/c4ob01457j




Scheme 1 Overview of strategies to 2-alkylideneaziridines.

## Results and discussion

To begin our investigation, the previously obtained results with chiral non-racemic aziridines **1a–c** were considered.<sup>12</sup> As reported in Scheme 2, upon electrophilic trapping of lithiated methyleneaziridines **1a–c-Li**, a new stereogenic centre is created, and a mixture of diastereomers results. Moreover, it was observed that high levels of stereoselectivity (dr >93 : 7) could be achieved only with a substituent installed on the double bond as in the case of (*S*)-**Z-1b** and (*S*)-**1c**. Curiously, with aziridine (*S*)-**E-1b** despite the presence of an alkene substituent, a lower level of stereoselectivity resulted. Similarly low levels of stereocontrol were seen in the case of unsubstituted aziridine (*S*)-**1a**. Such evidence prompted us to assume that a *cis*-relationship between the aziridine nitrogen and the substituent R<sup>1</sup> was a requirement for high stereoselectivity (Scheme 2). However, we felt that further experimentation with respect to the reaction parameters (*i.e.* ligand, temperature, solvent, time, *etc.*) was merited to determine whether the stereoselectivity of the lithiation of aziridines (*S*)-**1a** and (*S*)-**E-1b** could be improved.

First the effect of the ligand *N,N,N',N'*-tetramethylethylenediamine (TMEDA) was investigated, and the obtained results are reported in Table 1. Aziridine (*S*)-**1a** was lithiated in either the presence or absence of TMEDA and the resulting lithiated intermediates were trapped with electrophiles. The diastereomeric ratios were ascertained by <sup>1</sup>H-NMR analysis of the crude reaction mixtures, and compared with those obtained under the previously optimized conditions, that is, in the presence of the ligand. Aziridinyl lithium (*S*)-**1a-Li** proved to be chemically stable over a period of several hours even in the absence of TMEDA; indeed, it could be quenched with deuterium furnishing aziridine (*S*)-**1a-D** almost quantitatively (95% D) and without decomposition (Table 1, entries 1 and 2). The lithiated intermediate also reacted with different electrophiles to give products **2a–c** with good yields in either the presence or absence of TMEDA (Table 1, entries 3–8). Because the same diastereoselectivity was observed working in either the

**Table 1** Effect of the ligand on the stereoselectivity of the lithiation/trapping sequence of (*S*)-**1a**




Entry	TMEDA (equiv.)	E <sup>+</sup>	Product <sup>a</sup>	Conversion (%)	dr <sup>c,d</sup>
1	1.2	CD <sub>3</sub> OD	( <i>S</i> )- <b>1a-D</b> <sup>b</sup>	83	80/20
2	—	CD <sub>3</sub> OD	( <i>S</i> )- <b>1a-D</b> <sup>b</sup>	>98	80/20
3	1.2	BnBr	<b>2a</b> / <i>diast-2a</i>	90	70/30
4	—	BnBr	<b>2a</b> / <i>diast-2a</i>	>98	70/30
5	1.2	Me <sub>3</sub> SiCl	<b>2b</b> / <i>diast-2b</i>	90	80/20
6	—	Me <sub>3</sub> SiCl	<b>2b</b> / <i>diast-2b</i>	90	80/20
7 <sup>d</sup>	1.2	MeI	<b>2c</b> / <i>diast-2c</i>	87	70/30
8 <sup>d</sup>	—	MeI	<b>2c</b> / <i>diast-2c</i>	67	70/30

<sup>a</sup> Aziridines **2a–c** were obtained as a mixture of inseparable diastereomers. <sup>b</sup> Purification by distillation. <sup>c</sup> The dr values were determined by <sup>1</sup>H NMR analysis of the crude reaction mixture. <sup>d</sup> Assumed as a (*S,R*)/(*S,S*) diastereomeric ratio (**2**/*diast-2*).

presence or in the absence of TMEDA, it is reasonable to assume that the ligand is not playing an important role. Therefore, all the subsequent studies were performed without this additive.

The reaction temperature was the next parameter investigated. The lithiation reactions of (*S*)-**1a** were performed under the conditions reported in Table 2. Benzyl bromide and benzophenone were used as the electrophiles. The stereochemistry for the adducts **2a,d** and *diast-2a,d* was assigned by analogy to those of **2e** and **2f** whose stereochemistries have been deduced previously by X-ray diffraction (*vide infra*).<sup>13</sup> The lithiated intermediates were found to be chemically stable over a range of temperatures between −98 °C and −42 °C (Table 2). In the trapping reactions with benzyl bromide (entries 1–6), the stereoselectivity of the reactions was dependent on both the temperature *T*<sub>1</sub> for the generation of the lithiated aziridine, and temperature of quenching *T*<sub>2</sub>: the lower the temperature, the higher the diastereoselectivity. Moreover, a switch in diastereoselectivity occurs upon generation and/or trapping at higher temperatures (Table 2, entries 4 and 5). We ascribe this behaviour to the electrophile; benzyl bromide is known to react with lithiated species with both retention and inversion of configuration.<sup>14</sup> The possibility of a temperature promoted radical pathway (by SET) is also not ruled out. When a warm/cool protocol was applied, achieved by warming the solution to −42 °C for 30 min followed by trapping with BnBr at −78 °C, no inversion of the diastereoselectivity was observed (entry 6). Using benzophenone as the electrophile, which reacts predominantly with retention of configuration,<sup>15</sup> little change in stereoselectivity was observed over a range of lithiation/quenching temperatures (entries 7–10). We conclude that slightly improved stereoselectivity can be observed by running these reactions at a temperature as low as −98 °C.




**Scheme 2** Observed stereoselectivity in the lithiation/trapping sequence of alkylideneaziridines.

**Table 2** Effect of the temperature on the stereoselectivity of the lithiation/trapping sequence of (*S*)-**1a**

Entry	E <sup>+</sup>	T <sub>1</sub> <sup>a</sup> (°C)	T <sub>2</sub> <sup>b</sup> (°C)	Conversion (%)	dr 2/ diast-2 <sup>c</sup>
1	BnBr	-78	-78	>98	70 : 30
2	BnBr	-78	-98	73	88 : 12
3	BnBr	-98	-98	94	83 : 17
4	BnBr	-78	-42	93	30 : 70
5	BnBr	-42	-42	50	30 : 70
6	BnBr	-78	-42 → -78 <sup>d</sup>	50	75 : 25
7	Ph <sub>2</sub> CO	-78	-78	>98	80 : 20
8	Ph <sub>2</sub> CO	-78	-98	85	80 : 20
9	Ph <sub>2</sub> CO	-78	-42	85	87 : 13
10	Ph <sub>2</sub> CO	-98	-98	90	90 : 10


<sup>a</sup> T<sub>1</sub> is the lithiation temperature. <sup>b</sup> T<sub>2</sub> is the temperature of the trapping with the electrophile. All the reactions were kept for 30 min at T<sub>2</sub> before addition of the electrophile. <sup>c</sup> Diastereomeric ratio calculated by <sup>1</sup>H NMR of the crude reaction mixture. <sup>d</sup> The lithiated intermediate was stirred at -42 °C for 30 min and cooled to -78 °C before quenching.

To explore the influence of reaction time, we monitored the lithiation of aziridines (*S*)-**1a** and (*S*)-**1c** by *in situ* FT-IR.<sup>16,17</sup> This was done by monitoring the vibration of the exocyclic C=C double bond during the course of the reaction. In neutral aziridine (*S*)-**1a**, this signal was detected at 1780 cm<sup>-1</sup> in 0.2 M THF solution at -78 °C (Fig. 1). Upon addition of *s*-BuLi, a new band at 1710 cm<sup>-1</sup>, assigned to the lithiated intermediate (*S*)-**1a-Li**, appeared. The transformation was complete within 30 min with no further changes observed after 1 h. Upon addition of CD<sub>3</sub>OD as a deuterium source, aziridine (*S*)-**1a-D** was obtained in a 80/20 diastereomeric ratio (98% D) confirming that the detected signal belongs to the lithiated intermediate (Fig. 1). Similarly, in the lithiation of aziridine (*S*)-**1c** in THF, the signal for the stretching of the C=C double bond at 1810 cm<sup>-1</sup> quickly disappeared upon addition of *s*-BuLi, and a new signal, likely belonging to lithiated aziridine (*S*)-**1c-Li** appeared at 1760 cm<sup>-1</sup> (Fig. 1). From these experiments, we conclude that these lithiations are complete within 30 min at -78 °C in THF. Trapping of (*S*)-**1c-Li** with benzophenone furnished the expected adduct **2f** as the major diastereomer (dr = 95 : 5) in 80% yield. The lithiation of (*S*)-**1c** was also undertaken in toluene, and was complete within 30 min, as ascertained by *in situ* FT-IR monitoring. Moreover, quenching of (*S*)-**1c-Li** with benzophenone furnished a mixture of diastereomeric aziridines **2f** and *diast*-**2f** in a 70 : 30 ratio (Fig. 1). In all cases, the new signal was shifted to lower wavenumber by 50–60 cm<sup>-1</sup> but still in a range compatible with strained exocyclic double bonds.<sup>18</sup> These FT-IR experiments provided two important pieces of information. Firstly, that the lithiation does not require long reaction times,

**Fig. 1** *In situ* monitoring of the lithiation process by FT-IR.

being completed in just 30 min. Secondly, that the solvent seems to affect the stereoselectivity of the process. In particular, lower levels of stereoselectivity were obtained with the use of a less polar solvent such as toluene in the trapping of (*S*)-**1c-Li**.


To explore these observations, (*S*)-**1a**, (*S*)-**E-1b** and (*S*)-**1c** were lithiated in toluene and quenched with a range of electrophiles. Solvent-dependent stereoselectivity was observed in the lithiation/substitution reactions of (*S*)-**1a** (Scheme 3). Trapping of lithiated aziridine (*S*)-**1a** with CD<sub>3</sub>OD led to deuterated aziridine (*S*)-**1a-D** in a 80 : 20 diastereomeric ratio. <sup>1</sup>H NMR analysis of (*S*)-**1a-D** confirmed the opposite stereochemical preference in this substitution reaction with respect to that run in THF (see ESI†).<sup>13</sup> Similarly, trapping of (*S*)-**1a-Li**, generated in toluene, with benzophenone, Me<sub>3</sub>SiCl and 4,4'-dichlorobenzophenone furnished adducts **2b,d,g** and *diast*-**2b,d,g**. In each case, the major diastereomer observed in toluene had opposite stereochemistry with respect to that obtained in THF (*cf.* Table 1, entry 6 and Table 2, entry 7). These data suggest that this solvent-induced reversal in stereoselectivity is independent of the nature of the electrophile and arises in the lithiation step.<sup>19</sup>



**Scheme 3** Solvent-dependent stereoselectivity in the lithiation/trapping of methyleneaziridine (*S*)-**1a**.


The trapping of lithiated (*S*)-**E-1b** with diaryl ketones furnished, in toluene, a mixture of diastereomeric aziridines **2h-j** and *diast-2h-j* in moderate diastereomeric ratios but again with opposite stereoselectivity with respect to the THF conditions (Scheme 4).<sup>10</sup> Beside the lower levels of stereoselectivity observed in the reactions of (*S*)-**E-1b**, in comparison with (*S*)-**1a**, the solvent effect is remarkable with *diast-2h-j* being favored. The stereochemical course for the lithiation/trapping sequence of (*S*)-**1c** in toluene is also reported in Scheme 4. In this case, the same diastereomer **2f** was obtained in both THF and toluene. However, consistent with our other observations, the amount of *diast-2f* increased in toluene.

In order to rationalize the observed stereoselectivity, we turned our attention to the configurational stability of the lithiated intermediates. In fact, it is reasonable to assume that the presence of two diastereomeric adducts could be the result of a configurational instability of the lithiated intermediates. Based on some previous evidence on the regioselective lithia-



**Scheme 4** Solvent-dependent stereoselectivity in the lithiation/trapping of alkyldeneaziridines.

tion of *N*-alkyl-2,3-diphenylaziridines,<sup>20,21</sup> we decided to exploit the kinetic isotope effect (KIE) to assess the configurational stability of lithiated alkyldeneaziridines.<sup>22</sup> We reasoned that, in the presence of an appreciable KIE (that is different rates in proton and deuterium abstraction), the preferential removal of the residual proton would lead to lithiated intermediates having the opposite stereochemistry with respect to those generated from the fully protonated parent aziridines **1a-c** (that are **1a-c-Li** in Scheme 2). Upon reaction with the electrophile, two situations could be envisaged: (a) if the lithiated intermediates equilibrate, the diastereomeric ratios in the adducts should be similar to those observed in the lithiation/trapping of fully protonated parent aziridines; (b) conversely, in the case of a configurational stability of the lithiated intermediates, upon trapping with the electrophile, a different (if not opposite) diastereomeric ratio should be observed for the adducts. Moreover, in both cases an appreciable degree of deuteration is expected in the final products. Thus, simply evaluating the diastereomeric ratios resulting from the lithiation/trapping of deuterated aziridines, evidence on the configurational stability or instability of the lithiated intermediates would be provided. In order to test this hypothesis, we selected deuterated aziridines that furnished a different but appreciable degree of stereoselectivity in the lithiation/trapping sequence, namely aziridine (*S*)-**1a-D** and (*S*)-**Z-1b-D**. When deuterated aziridine (*S*)-**1a-D** (dr 80 : 20, 98% D) was subjected to lithiation with *s*-BuLi followed by trapping with benzophenone, adducts **2d-D** (75% D) and *diast-2d-D* (98% D) were formed in a 35/65 ratio (Scheme 5). Applying the same protocol to aziridine (*S*)-**Z-1b-D** (dr 98 : 2, 98% D), a 1 : 1 diastereomeric mixture of **2e** (0% D) and *diast-2e-D* (98% D) was obtained. In both cases, the diastereomeric ratios differ from that resulting from lithiation/trapping of parent aziridine (*S*)-**1a** (dr 80 : 20) and (*S*)-**Z-1b** (dr 93 : 7). In addition, in the reaction of (*S*)-**1a-D** opposite stereoselectivity was observed, with *diast-2d-D* being the major adduct, whilst in the reaction of (*S*)-**Z-1b-D** the adduct **2e** was completely non-deuterated. Such results indicate a significant KIE effect in these deproto-




**Scheme 5** Experiments proving the configurational stability of lithiated alkyldeneaziridines.

nations and configurational stability of the lithiated intermediates.

At this point, further questions arise: (a) how could the solvent induce the stereoselectivity-switch? (b) why was this stereoselectivity-switch not observed in the lithiation trapping of (*S*)-**1c** (Scheme 4)? A possible explanation for the above results could be given if we take into consideration the aziridine nitrogen stereodynamics, complexation and proximity effects. In fact, in the case of *N*-alkyl-2-arylaziridines, such reasoning allowed for rationalizing the solvent- and temperature-dependent regio and stereoselectivity of the lithiation/trapping sequences.<sup>16,17</sup> The model depicted in Scheme 6 could apply in the case of 2-alkylideneaziridines in which the aziridine nitrogen undergoes slow pyramidal inversion. As a consequence of this slow motion, the <sup>1</sup>H NMR spectra of aziridines (*S*)-**1a–c** showed very broad signals at room temperature (Fig. 2). Nevertheless, the <sup>1</sup>H NMR spectra acquired at low temperature (200 K) clearly showed the signals for the two slowly equilibrating invertomers (Fig. 2).

Analysis of the low-temperature (200 K) <sup>1</sup>H NMR spectra revealed an invertomer ratio of 62 : 38 for aziridines (*S*)-**1a** and (*S*)-*E*-**1b**, and a 71 : 29 ratio for aziridines (*S*)-*Z*-**1b** and (*S*)-**1c**. However, even though we were able to reveal the presence of



Scheme 6 Model based on the nitrogen stereodynamics.

two nitrogen invertomers, it was not possible to establish the relative configurations of the major invertomers. Nevertheless, with reference to Scheme 6, for the sake of argument,





Fig. 2 Stereodynamics of alkylideneaziridines by VT NMR under moderate and slow exchange regimes.

we assume that a preferential deprotonation *syn* to the nitrogen lone pair of invertomers **A** and **B** would lead to diastereomeric lithiated intermediates **3-Li** and *diast*-**3-Li**. In addition, on the basis of the experiments on deuterated aziridines (Scheme 5), lithiated intermediates **3-Li** and *diast*-**3-Li** should also be configurationally stable, and a retentive electrophilic substitution would furnish adducts **2** and *diast*-**2** respectively. Thus, a possible explanation for the observed stereoselectivity could rely on a difference between the rates of deprotonation ( $k_3$ ,  $k_4$  in Scheme 6) with respect to the rates of inversion ( $k_1$ ,  $k_2$  in Scheme 6). However, regardless of the equilibria involving the lithiating agent, which could have its own aggregation state,<sup>9b</sup> at least three different conditions are foreseen: (a) with  $k_3 \approx k_4 \gg k_1, k_2$  the ratio between **3-Li** and *diast*-**3-Li** would reflect that of the neutral invertomers; (b) with  $k_1, k_2 \gg k_3$  (or  $k_4$ ) >  $k_4$  (or  $k_3$ ) one prevalent lithiated intermediate should form; (c) with  $k_3$  (or  $k_4$ ) >  $k_4$  (or  $k_3$ )  $\approx k_1, k_2$  a mixture of **3-Li** and *diast*-**3-Li** with an unpredictable ratio is expected. The condition (b) could likely explain the high stereoselectivity observed in the lithiation/trapping sequences on aziridines (*S*)-**Z-1b** and (*S*)-**1c** in THF and on (*S*)-**1a** in toluene. On the other hand, condition (a) or (c) could apply to (*S*)-**1a**, (*S*)-**E-1b**, in THF where a lower level of stereoselectivity resulted.<sup>23</sup>

Moreover, the model proposed in Scheme 6 could explain the stereoselectivity observed in the lithiation/trapping of deuterated aziridines (*S*)-**1a-D** and (*S*)-**Z-1b-D**. In Scheme 7 the model is applied to the lithiation of (*S*)-**Z-1b-D** where two invertomers in a *ca.* 70 : 30 ratio are present. In this case, the KIE would make the rates  $k_3'$  and  $k_4'$  different enough to allow for a competitive removal of proton  $H_a$ ; thus the ratio of the lithiated intermediates will vary accordingly leading, upon trapping with benzophenone, to the observed 1 : 1 diastereomeric mixture of **2e** (0% D) and *diast*-**2e-D** (98% D).

This model seems to support the idea that nitrogen coordination and proximity effects could be responsible for the highly regio and stereoselective lithiation of 2-alkylideneaziridines.



Scheme 7 Nitrogen stereodynamics and KIE.

## Conclusions

This study intends to demonstrate the origin of the stereoselectivity in the lithiation/trapping sequence of 2-alkylideneaziridines. New conditions for conducting the lithiation reaction have been discovered by *in situ* FT-IR monitoring. In addition, it has been found that the solvent and the alkene substitution are important factors that lead to a switch in stereoselectivity. Lithiation of fully substituted or *Z*-configured 2-alkylideneaziridines, such as (*S*)-**1c** or (*S*)-**Z-1b**, occurred stereoselectively in THF. In striking contrast, unsubstituted 2-methyleneaziridines, such as (*S*)-**1a**, underwent lithiation with an opposite stereoselectivity in toluene. A model based on complexation and proximity effects has been proposed to rationalize this reactivity. This model assumes that slowly equilibrating N-invertomers undergo deprotonation with different rates and that the stereochemical outcome is established during the deprotonation step. Lithiation experiments on deuterium labeled 2-alkylideneaziridines revealed the configurational stability of the lithiated intermediates. Further stereochemical and spectroscopic studies are underway to support the proposed model, and will be reported in due course.

## Acknowledgements

We thank the National Project “FIRB-Futuro in Ricerca” (code: CINECA RBF083M5N), Regione Puglia: “Reti di Laboratori Pubblici di Ricerca” Project code 20, the University of Bari and the Consortium CINMPIS for supporting this research under the framework of the project PON Ricerca Industriale (code PON01\_00862). “Laboratorio SISTEMA” code: PONA300369 financed by the Italian MIUR; The support of the Engineering and Physical Sciences Research Council (EP/D035384/1) is also gratefully acknowledged.

## Notes and references

- (a) L. Degennaro, P. Trincherà and R. Luisi, *Chem. Rev.*, 2014, **114**, 7881–7929; (b) A. Padwa, in *Comprehensive Heterocyclic Chemistry III*, ed. A. R. Katritzky, C. A. Ramsden, E. F. V. Scriven and R. J. K. Taylor, Elsevier, 2008, vol. 1, pp. 1–104; (c) A. Padwa and S. S. Murphee, *Prog. Heterocycl. Chem.*, 2003, **15**, 75; (d) J. B. Sweeney, in *Science of Synthesis*, ed. E. Schaumann and D. Enders, Georg Thieme Verlag, Stuttgart, New York, 2008, vol. 40a, p. 643; (e) Aziridines in natural product synthesis, C. Botuha, F. Chemla, F. Ferreira and A. Perez-Luna, in *Heterocycles in Natural Product Synthesis*, ed. K. C. Majumdar and S. K. Chattopadhyay, Wiley-VCH, Weinheim, 2011, pp. 3–39; (f) P. Somfai and J. Ahman, *Targets Heterocycl. Syst.*, 1999, **3**, 341.
- For reviews concerning the reactivity of methyleneaziridines, see: (a) M. Shipman, *Synlett*, 2006, 3205–3217;

- (b) K. A. Tehrani and N. De Kimpe, *Curr. Org. Chem.*, 2009, **13**, 854–877.
- 3 (a) J. F. Hayes, M. Shipman and H. Twin, *Chem. Commun.*, 2001, 1784–1785; (b) J. F. Hayes, M. Shipman and H. Twin, *J. Org. Chem.*, 2002, **67**, 935–942; (c) J. F. Margathe, M. Shipman and S. C. Smith, *Org. Lett.*, 2005, **7**, 4987–4990; (d) C. Montagne, J. J. Shiers and M. Shipman, *Tetrahedron Lett.*, 2006, **47**, 9207–9209; (e) J. J. Shiers, G. J. Clarkson, M. Shipman and J. F. Hayes, *Chem. Commun.*, 2006, 649–651; (f) C. C. A. Cariou, G. J. Clarkson and M. Shipman, *J. Org. Chem.*, 2008, **73**, 9762–9764; (g) P. M. Mumford, G. J. Tarver and M. Shipman, *J. Org. Chem.*, 2009, **74**, 3573–3575.
- 4 (a) G. Prié, N. Prévost, H. Twin, S. A. Fernandes, J. F. Hayes and M. Shipman, *Angew. Chem., Int. Ed.*, 2004, **43**, 6517–6519; (b) J. Robertson, G. C. Feast, L. V. White, V. A. Steadman and T. D. W. Claridge, *Org. Biomol. Chem.*, 2010, **8**, 3060–3063; (c) K. Griffin, C. Montagne, C. T. Hoang, G. J. Clarkson and M. Shipman, *Org. Biomol. Chem.*, 2012, **10**, 1032–1039; (d) H. Takahashi, S. Yasui, S. Tsunoi and I. Shibata, *Eur. J. Org. Chem.*, 2013, 40–43; (e) H. Takahashi, S. Yasui, S. Tsunoi and I. Shibata, *Org. Lett.*, 2014, **16**, 1192–1195.
- 5 J. W. Rigoli, I. A. Guzei and J. M. Schomaker, *Org. Lett.*, 2014, **16**, 1696–1699.
- 6 (a) N. Prévost and M. Shipman, *Org. Lett.*, 2001, **3**, 2383–2385; (b) N. Prévost and M. Shipman, *Tetrahedron*, 2002, **58**, 7165–7175.
- 7 (a) H. Alper and N. Hamel, *Tetrahedron Lett.*, 1987, **28**, 3237–3240; (b) B. H. Oh, I. Nakamura and Y. Yamamoto, *Tetrahedron Lett.*, 2002, **43**, 9625–9628; (c) A. I. Siriwardana, K. K. A. D. S. Kathriarachchi, I. Nakamura, I. D. Gridnev and Y. Yamamoto, *J. Am. Chem. Soc.*, 2004, **126**, 13898–13899; (d) B. H. Oh, I. Nakamura and Y. Yamamoto, *J. Org. Chem.*, 2004, **69**, 2856–2858; (e) K. K. A. D. S. Kathriarachchi, A. I. Siriwardana, I. Nakamura and Y. Yamamoto, *Tetrahedron Lett.*, 2007, **48**, 2267–2270; (f) B. Pan, C. Wang, D. Wang, F. Wu and B. Wan, *Chem. Commun.*, 2013, **49**, 5073–5075.
- 8 J. J. Shiers, M. Shipman, J. F. Hayes and A. M. Z. Slawin, *J. Am. Chem. Soc.*, 2004, **126**, 6868–6869.
- 9 (a) S. Florio and R. Luisi, *Chem. Rev.*, 2010, **110**, 5128–5157; (b) *Lithium Compounds in Organic Synthesis – From Fundamentals to Applications*, ed. R. Luisi and V. Capriati, Wiley-VCH, Weinheim, 2014; (c) D. M. Hodgson and C. D. Bray, in *Aziridine and Epoxides in Organic Synthesis*, ed. A. K. Yudin, Wiley-VCH, Weinheim, 2006, ch. 5, pp. 145–184; (d) *The Chemistry of Organolithium Compounds*, ed. Z. Rappoport and I. Marek, Wiley-VCH, Weinheim, 2004.
- 10 For seminal studies see: H. Quast and C. A. Weise Vélez, *Angew. Chem., Int. Ed. Engl.*, 1974, **13**, 342.
- 11 J. F. Hayes, N. Prévost, I. Prokes, M. Shipman, A. M. Z. Slawin and H. Twin, *Chem. Commun.*, 2003, 1344.
- 12 C. Montagne, N. Prévost, J. J. Shiers, G. Prie, S. Rahman, J. Ince, J. F. Hayes and M. Shipman, *Tetrahedron*, 2006, **62**, 8447.
- 13 We assume that the reactions with electrophiles occur with retention of configuration at the lithiated centre. The stereochemistry for adducts **2a–d**, **2g** and (*S*)-**1a–D** is assigned by analogy to the stereochemistry of adducts **2e** and **2f** ascertained by X-ray analysis, see ref. 8 and 11.
- 14 R. E. Gawley, E. Low, Q. Zhang and R. Harris, *J. Am. Chem. Soc.*, 2000, **122**, 3344.
- 15 A. Basu and S. Thayumanavan, *Angew. Chem., Int. Ed.*, 2002, **41**, 717–738.
- 16 G. Barker, J. L. McGrath, A. Klapars, D. Stead, G. Zhou, K. R. Campos and P. O'Brien, *J. Org. Chem.*, 2011, **76**, 5936–5953.
- 17 (a) D. Stead, G. Carbone, P. O'Brien, K. R. Campos, I. Coldham and A. Sanderson, *J. Am. Chem. Soc.*, 2010, **132**, 7260–7261; (b) N. S. Sheikh, D. Leonori, G. Barker, J. D. Firth, K. R. Campos, A. J. H. M. Meijer, P. O'Brien and I. Coldham, *J. Am. Chem. Soc.*, 2012, **134**, 5300–5308.
- 18 For a related example concerning the lithiation of a strained heterocycle see: V. Capriati, S. Florio, R. Luisi, F. M. Perna and A. Spina, *J. Org. Chem.*, 2008, **73**, 9552–9564.
- 19 A solvent-dependent stereoselectivity has been observed in the lithiation of other *N*-alkylaziridines, see: (a) R. Luisi, V. Capriati, S. Florio and B. Musio, *Org. Lett.*, 2007, **9**, 1263–1266; (b) M. C. de Ceglie, B. Musio, A. Francesco, A. Moliterni, A. Altomare, S. Florio and R. Luisi, *Chem. – Eur. J.*, 2011, **17**, 286–296.
- 20 F. Affortunato, S. Florio, R. Luisi and B. Musio, *J. Org. Chem.*, 2008, **73**, 9214–9220.
- 21 V. Capriati, S. Florio, R. Luisi, L. Mazzanti and B. Musio, *J. Org. Chem.*, 2008, **73**, 3197–3204.
- 22 For a recent example on the KIE see: L. Řehová and U. Jahn, *Eur. J. Org. Chem.*, 2014, **21**, 4610–4623.
- 23 The invertomer ratio is not solvent dependent. The same invertomer ratio has been found in toluene-*d*<sub>8</sub> under slow exchange conditions.