



HAL
open science

ON THE NUMBER OF ISOLATED ZEROS OF PSEUDO-ABELIAN INTEGRALS: DEGENERACIES OF THE CUSPIDAL TYPE

Aymen Braghtha

► **To cite this version:**

Aymen Braghtha. ON THE NUMBER OF ISOLATED ZEROS OF PSEUDO-ABELIAN INTEGRALS: DEGENERACIES OF THE CUSPIDAL TYPE. 2015. hal-01145681

HAL Id: hal-01145681

<https://hal.science/hal-01145681>

Preprint submitted on 25 Apr 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE NUMBER OF ISOLATED ZEROS OF PSEUDO-ABELIAN INTEGRALS: DEGENERACIES OF THE CUSPIDAL TYPE

Aymen Braghtha

June 19, 2015

Abstract

We consider a multivalued function of the form $H_\varepsilon = P_\varepsilon^{\alpha_0} \prod_{i=1}^k P_i^{\alpha_i}$, $P_i \in \mathbb{R}[x, y]$, $\alpha_i \in \mathbb{R}_+^*$, which is a Darboux first integral of polynomial one-form $\omega = M_\varepsilon \frac{dH_\varepsilon}{H_\varepsilon} = 0$, $M_\varepsilon = P_\varepsilon \prod_{i=1}^k P_i$. We assume, for $\varepsilon = 0$, that the polycycle $\{H_0 = H = 0\}$ has only cuspidal singularity which we assume at the origin and other singularities are saddles.

We consider families of Darboux first integrals unfolding H_ε (and its cuspidal point) and pseudo-Abelian integrals associated to these unfolding. Under some conditions we show the existence of uniform local bound for the number of zeros of these pseudo-Abelian integrals.

Keywords. integrable systems, blowing-up, singular foliations, singularities, abelian functions

1 Formulation of main results

In this paper, we study a non generic case. Other non generic cases have been studied in [1,3,4,5]. Pseudo-Abelian integrals appear as the linear principal part of the displacement function in polynomial perturbation of Darboux integrable case.

More precisely consider Darboux integrable system ω given by

$$\omega = M d \log H, \tag{1}$$

where

$$M = \prod_{i=0}^k P_i, \quad H = \prod_{i=0}^k P_i^{\alpha_i}, \quad \alpha_i > 0, \quad P_i \in \mathbb{R}[x, y]. \tag{2}$$

Now we consider an unfolding ω_ε of Darboux integrable system ω , where ω_ε are one-forms with first integral

$$H_\varepsilon = P_\varepsilon^\alpha \prod_{i=1}^k P_i^{\alpha_i}, \quad \omega_\varepsilon = M_\varepsilon d \log H_\varepsilon, \quad M_\varepsilon = P_\varepsilon \prod_{i=1}^k P_i. \tag{3}$$

where the polynomial P_0 has a cuspidal singularity at $p_0 = (0, 0)$, i.e. $P_0(x, y) = y^2 - x^3 + \mathcal{O}((x, y)^4)$. For non zero ε , the polynomial $P_\varepsilon = y^2 - x^3 - \varepsilon x^2 + \mathcal{O}((x, y, \varepsilon)^4)$.

Choose a limit periodic set i.e. bounded component of $\mathbb{R}^2 \setminus \{\prod_{i=0}^k P_i = 0\}$ filled cycles $\gamma(h) \subset \{H = h\}$, $h \in (0, a)$. Denote by $D \subset H^{-1}(0)$ the polycycle which is in the boundary of this limit periodic set.

Consider the unfolding $\omega_\varepsilon = M_\varepsilon d \log H_\varepsilon$ of the form ω . The foliation ω_ε has a maximal nest of cycles $\gamma(\varepsilon, h) \subset \{H_\varepsilon = h\}$, $h \in (0, a(\varepsilon))$, filling a connected component of $\mathbb{R}^2 \setminus \{H_\varepsilon = 0\}$ whose boundary is a polycycle D_ε close to D . Assume moreover that the foliation $\omega_\varepsilon = 0$ has no singularities on $\text{Int}D_\varepsilon$.

Consider pseudo-Abelian integrals of the form

$$I(\varepsilon, h) := \int_{\gamma(\varepsilon, h)} \eta_2, \quad \eta_2 = \frac{\eta_1}{M_\varepsilon} \tag{4}$$

where η_1 is a polynomial one-form of degree at most n .

This integral appears as the linear term with respect to β of the displacement function of a polynomial perturbation

$$\omega_{\varepsilon,\beta} = \omega_\varepsilon + \beta\eta_1 = 0. \quad (5)$$

We assume the following genericity assumptions

1. The level curves $P_i = 0, i = 1, \dots, k$ are smooth and $P_i(0, 0) \neq 0$.
2. The level curves $P_\varepsilon = 0, P_i = 0, i = 1, \dots, k$, intersect transversally two by two.

Theorem 1. *Under the genericity assumptions there exists a bound for the number of isolated zeros of the pseudo-Abelian integrals $I(\varepsilon, h) = \int_{\gamma(\varepsilon, h)} \eta_2$ in $(0, a(\varepsilon))$. The bound is locally uniform with respect to all parameters in particular in ε .*

Let $\mathcal{F}_1 : \{\omega_\varepsilon = 0\}, \mathcal{F}_2 : \{d\varepsilon = 0\}$ are the foliations of dimension two in complex space of dimension three with coordinates (x, y, ε) .

Let \mathcal{F} be the foliation of dimension one on the complex space of dimension three with coordinates (x, y, ε) which is given by the intersection of leaves of \mathcal{F}_1 and \mathcal{F}_2 (i.e. given by the 2-form $\Omega = \omega_\varepsilon \wedge d\varepsilon$). This foliation has a cuspidal singularity at the origin (a cusp).

We want to study the analytical properties of the foliation \mathcal{F} in a neighborhood of the cusp. For this reason we make a global blowing-up of the cusp of the product space (x, y, ε) of phase and parameter spaces. We want our blow-up to separate the two branches of the cusp. This requirement leads to the quasi-homogeneous blowing-up of weight $(2, 3, 2)$.

Remark 1. *In term of first integrals, the foliation \mathcal{F} is given by two first integrals*

$$H(x, y, \varepsilon) = h, \quad \varepsilon = s.$$

2 Quasi-homogeneous blowing-up of \mathcal{F}

Recall the construction of the quasi-homogeneous blowing-up. We define the weighted projective space $\mathbb{C}\mathbb{P}_{2:3:2}^2$ as factor space of \mathbb{C}^3 by the \mathbb{C}^* action $(x, y, \varepsilon) \mapsto (t^2x, t^3y, t^2\varepsilon)$. The quasi-homogeneous blowing-up of \mathbb{C}^3 at the origin is defined as the incidence three dimensional manifold $W = \{(p, q) \in \mathbb{C}\mathbb{P}_{2:3:2}^2 \times \mathbb{C}^3 : \exists t \in \mathbb{C} : (q_1, q_2, q_3) = (t^2p_1, t^3p_2, t^2p_3)\}$, where $(q_1, q_2, q_3) \in \mathbb{C}$ and $[(p_1, p_2, p_3)] \in \mathbb{C}\mathbb{P}_{2:3:2}^2$.

The quasi-homogeneous blowing-up $\sigma : W \rightarrow \mathbb{C}^3$ is just the restriction to W of the projection $\mathbb{C}\mathbb{P}_{2:3:2}^2 \times \mathbb{C}^3$.

We will need explicit formula for the blow-up in the standard affine charts of W . The projective space $\mathbb{C}\mathbb{P}_{2:3:2}^2$ is covered by three affine charts: $U_1 = \{x \neq 0\}$ with coordinates (y_1, z_1) , $U_2 = \{y \neq 0\}$ with coordinates (x_2, z_2) and $U_3 = \{\varepsilon \neq 0\}$ with coordinates (x_3, y_3) .

The transition formula follow from the requirement that the points $(1, y_1, z_1), (x_2, 1, z_2)$ and $(x_3, y_3, 1)$ lie on the same orbit of the action:

$$F_2 : (y_1, z_1) \mapsto \left(x_2 = 1/y_1^{2/3}, z_2 = z_1/y_1\sqrt{y_1} \right)$$

$$F_3 : (y_1, z_1) \mapsto \left(x_3 = 1/z_1, y_3 = y_1/z_1\sqrt{z_1} \right).$$

These affine charts define affine charts on W , with coordinates $(y_1, z_1, t_1), (x_2, z_2, t_2)$ and (x_3, y_3, t_3) . The blow-up σ is written as

$$\sigma_1 : \quad x = t_1^2, \quad y = t_1^3 y_1, \quad \varepsilon = t_1^2 z_1 \quad (6)$$

$$\sigma_2 : \quad x = t_2^2 x_2, \quad y = t_2^3, \quad \varepsilon = t_2^2 z_2 \quad (7)$$

$$\sigma_3 : \quad x = t_3^2 x_3, \quad y = t_3^3 y_3, \quad \varepsilon = t_3^2. \quad (8)$$

We apply this blow-up σ to the one-dimensional foliation \mathcal{F} . Let $\sigma^{-1}\mathcal{F}$ the lifting of the foliation \mathcal{F} to the complement. This foliation has a cuspidal singularity at the origin. The pull-back foliation $\sigma^*\mathcal{F}$ will be called the strict transform of the foliation \mathcal{F} is defined by the pull-back $\sigma^*\Omega = \sigma^*(\omega_\varepsilon \wedge d\varepsilon)$ divided by a suitable power of the function defining the exceptional divisor. In this charts $U_j, j = 1, 2, 3$ we have

$$\sigma_1^*\Omega = x^2\Omega_1, \quad \sigma_2^*\Omega = y^3\Omega_2, \quad \sigma_3^*\Omega = \varepsilon^2\Omega_3,$$

where

$$\Omega_1 = (6y_1^2 - 6 - 4z_1)dx \wedge dz_1 + 4y_1z_1dy_1 \wedge dx + 2xy_1dy_1 \wedge dz_1, \quad (9)$$

$$\Omega_2 = (6 - 6x_2^3 - 4x_2^2z_2)dy \wedge dz_2 + (-6z_2x_2^2 - 4x_2z_2^2)dx_2 \wedge dy \quad (10)$$

$$+ (-3yx_2^2 - 2yx_2z_2)dx_2 \wedge dz_2, \quad (11)$$

$$\Omega_3 = (-6x_3^2 - 4x_3)dx_3 \wedge d\varepsilon + 4y_3dy_3 \wedge d\varepsilon. \quad (12)$$

Remark 2. In term of first integrals, the foliation $\sigma^*\mathcal{F}$ is given by two first integrals

$$\sigma^*H(x, y, \varepsilon) = h, \quad \sigma^*\varepsilon = s,$$

In particular in a neighborhood of the exceptional divisor the restrictions of the foliation $\sigma^*\mathcal{F}$ to the charts U_1 and U_3 are given respectively, by

$$\psi_1 = H(t_1^2, t_1^3y_1, t_1^2z_1) = x^3(y_1^2 - 1) = h, \quad \varphi_1 = xz_1 = s, \quad (13)$$

$$\psi_3 = H(t_3^2x_3, t_3^3y_3, t_3^2) = \varepsilon^3(y_3^2 - x_3^2 - x_3^3) = h, \quad \varphi_3 = \varepsilon = s, \quad (14)$$

where $\{x = 0\}$ and $\{\varepsilon = 0\}$ are local equations of the exceptional divisor respectively.

3 Singular locus of the foliation $\sigma^*\mathcal{F}$

In this section, we compute the singular locus of the pull-back $\sigma^*\Omega$ in a neighborhood of the exceptional divisor $\mathbb{CP}_{2:3:2}^2$. We check it in each chart separately.

In the chart U_1 , the zeros locus of the form Ω_1 in a neighborhood of the exceptional divisor $\{x = 0\}$ consists of germs of two curves $\{y_1 = \pm 1, z_1 = 0\}$ and a two singular points $p_1 = (0, 1, 0), p_2 = (0, -1, 0)$ generated by the quasi-homogeneous blowing-up.

In the chart U_3 , the zeros locus of the form Ω_3 in a neighborhood of the exceptional divisor $\{\varepsilon = 0\}$ consists of $p_3 = (0, 0, 0)$ (Morse point) and $p_4 = (-\frac{2}{3}, 0, 0)$ (center). The singularities of this foliation are the line of Morse points $x_3 = 0, y_3 = 0$, the lines of centers $x_3 = -\frac{2}{3}, y_3 = 0$ and the transform strict of $\{y^2 - x^3 - x^2\varepsilon = 0\}$.

Proposition 1. *The singularities of $\sigma^*\mathcal{F}$ are located at the points p_1, p_2, p_3 and p_4 . The points p_1, p_2 and p_3 are linearisable saddles and the point p_4 is a center.*

Proof. Since $\sigma : W \rightarrow \mathbb{C}^3$ is a biholomorphism outside the exceptional divisor $\mathbb{CP}_{2:3:2}^2$, all singularities of $\sigma^*\mathcal{F}$ on $\mathbb{C}^3 \setminus \{x = 0\}$ correspond to singularities of \mathcal{F} . Thus, it suffices to compute the singularities of $\sigma^*\mathcal{F}$ on the exceptional divisor $\{x = 0\}$. More precisely, we consider the foliation on neighborhood of $\mathbb{CP}_{(2:3:2)}^2$ (the exceptional divisor) generated by the blown-up one-form $\sigma^*\Omega$. Let ψ_1, ψ_3 are the functions given in (13) and (14).

(1) In the chart U_1 , near the divisor exceptional and for $|z_1| \leq \epsilon$ for ϵ sufficiently small, the foliation $\sigma^*\mathcal{F}$ is given by two first integrals

$$G_1 = \varphi_1^3\psi_1^{-1} = z_1^3(y_1^2 - (1 + z_1))^{-1}V^{-1} = s^3h^{-1}, \quad \varphi_1 = xz_1 = s.$$

where V is analytic function such that $V(0,0,0) \neq 0$. In particular on the exceptional divisor $\{x=0\}$ the foliation $\sigma^*\mathcal{F}$ is given by the levels $G_1 = s^3h^{-1} = t$.

Now we calculate the eigenvalues at p_1 and p_2 . The vector field V_1 generating the foliation $\sigma^*\mathcal{F}$ is given by

$$V_1(x, y_1, z_1) = \beta_1 x \frac{\partial}{\partial x} + \beta_2 y_1 \frac{\partial}{\partial y_1} + \beta_3 z_1 \frac{\partial}{\partial z_1},$$

where the vector $(\beta_1, \beta_2, \beta_3)$ satisfies the following equations

$$\langle (\beta_1, \beta_2, \beta_3), (3, 1, 0) \rangle = 0, \langle (\beta_1, \beta_2, \beta_3), (1, 0, 1) \rangle = 0$$

here \langle, \rangle be the usual scalar product on \mathbb{C}^3 . By simple computation, we obtain $\beta_1 = 1, \beta_2 = -3$ and $\beta_3 = -1$.

(2) In the chart U_3 , near the exceptional divisor $\{\varepsilon = 0\}$, the foliation $\sigma^*\mathcal{F}$ is given by

$$G_3 = \varphi_3^3 \psi_3^{-1} = (y_3^2 - x_3^2(1 + x_3))^{-1} = s^3h^{-1}, \quad \varphi_3 = \varepsilon = s.$$

In particular the restriction of this foliation to the exceptional divisor $\{\varepsilon = 0\}$, by Morse lemma we can put the function $1/G_3$ to the normal form $y_3^2 - z_3^2$ in a neighborhood of p_3 (we put the variable change $z_3 = \pm x_3(1 + x_3)^{1/2}$). On other hand the Hessian matrix of $1/G_3$ at the point p_4 has two positive eigenvalues. \square

4 The different scaled variations of $\delta(s, t)$

In this section, we compute the scaled variations with respect to different variables s and t of the integrals of the blown-up one form $\sigma_1^*\eta_2$ along the different relative cycles using the same technics of [5].

Proposition 2. *The computation of the different scaled variations of the cycle $\delta(s, t)$ us gives*

1. For $t \in [0, 2N]$, the cycle $\delta(s, t)$ satisfies a iterated scaled variations with respect to t of the form

$$\mathcal{V}ar_{(t,3)} \circ \mathcal{V}ar_{(t,-1)} \circ \mathcal{V}ar_{(t,-\alpha_1)} \circ \dots \circ \mathcal{V}ar_{(t,-\alpha_k)} \delta(s, t) = 0. \quad (15)$$

2. For $t \in [N, +\infty]$, the cycle $\delta(s, t)$ satisfies a iterated scaled variations with respect to $1/t$ of the form

$$\mathcal{V}ar_{(1/t,-3)} \circ \mathcal{V}ar_{(1/t,1)} \circ \mathcal{V}ar_{(1/t,1)} \circ \mathcal{V}ar_{(1/t,\alpha_1)} \circ \dots \circ \mathcal{V}ar_{(1/t,\alpha_k)} \delta(s, 1/t) = 0. \quad (16)$$

3. Near $s = 0$, we have

$$\mathcal{V}ar_{(s,1)} \circ \mathcal{V}ar_{(s,1)} \delta(s, t) = \mathcal{V}ar_{(s,1)}(\tilde{\delta}(s, t)) = 0, \quad (17)$$

where $\mathcal{V}ar_{(s,1)} \delta(s, t) = \tilde{\delta}(s, t)$ is a figure eight cycle.

Proof. As in [5], there exist a some local chart with coordinates (u, v, w) defined in a some neighborhood of each separatrix of polycycle such that the foliation $\sigma^*\mathcal{F}$ is defined by two first integrals. Precisely:

1. for $t \in [0, 2N]$, there exist a local chart $(V_{div}, (u, v, w))$ defined in neighborhood of the separatrix δ_{div} such the foliation $\sigma^*\mathcal{F}$ by two first integrals

$$F_1 = w^3(v-1)^{-1}(v+1)^{-1} = t, \quad F_2 = uw = s,$$

2. for $t \in [N, +\infty]$, there exists a local chart $(V_{div}^+, (u, v, w))$ defined in neighborhood of the separatrix δ_{div}^+ such that the foliation $\sigma_1^* \mathcal{F}$ is defined by two first integrals

$$F_1 = w^3(v+2)^{-1}v^{-1} = t, \quad F_2 = uw = s,$$

3. for $t \in [N, +\infty]$, there exists a local chart $(V_{div}^-, (u, v, w))$ defined in a neighborhood of the separatrix δ_{div}^- such that the foliation $\sigma_1^* \mathcal{F}$ is defined by two first integrals

$$F_1 = w^3(v-2)^{-1}v^{-1} = t, \quad F_2 = uw = s.$$

In second step we prove that each relative cycle can be chosen as a lift of a path contained in the separatrix associated to this relative cycle. Precisely:

1. on the chart $(V_{div}, (u, v, w))$, the linear projection $\pi(u, v, w) = v$ is every where transverse to the levels of the foliation $\sigma^* \mathcal{F}$ which corresponds simply to the graphs of the multivalued functions

$$v \mapsto (u, w) = \left(st^{-\frac{1}{3}}(v-1)^{-\frac{1}{3}}(v+1)^{-\frac{1}{3}}, t^{\frac{1}{3}}(v-1)^{\frac{1}{3}}(v+1)^{\frac{1}{3}} \right),$$

2. on the chart $(V_{div}^+, (u, v, w))$, the linear projection $\pi(u, v, w) = v$ is every where transverse to the levels of the foliation $\sigma^* \mathcal{F}$ which corresponds simply to the graphs of the multivalued functions

$$v \mapsto (u, w) = \left(st^{-\frac{1}{3}}v^{-\frac{1}{3}}(v+2)^{-\frac{1}{3}}, t^{\frac{1}{3}}v^{\frac{1}{3}}(v+2)^{\frac{1}{3}} \right),$$

3. on the chart $(V_{div}^-, (u, v, w))$, the linear projection $\pi(u, v, w) = v$ is every where transverse to the levels of the foliation $\sigma^* \mathcal{F}$ which corresponds simply to the graphs of the multivalued functions

$$v \mapsto (u, w) = \left(st^{-\frac{1}{3}}v^{-\frac{1}{3}}(v-2)^{-\frac{1}{3}}, t^{\frac{1}{3}}v^{\frac{1}{3}}(v-2)^{\frac{1}{3}} \right).$$

In third step, we compute the different scaled variations of relatives cycles using the local expression of two first integrals F_1 and F_2 above near the singular points p_1, p_2 and p_3 . Recall that the scaled variation of a relative cycle $\delta(s)$ is given by

$$\mathcal{V}ar_{(s,\beta)} \delta(s) = \delta(se^{i\pi\beta}) - \delta(se^{-i\pi\beta}).$$

In the local chart $(V_{div}^+, (u, v, w))$, the restriction of the blown-up foliation $\sigma_1^* \mathcal{F}$ to the transversals sections $\Sigma_{div}^- = \{w = 1\}$ (near the point p_3) and $\Omega_+ = \{u = 1\}$ (near the point p_1) is given respectively by

$$\begin{aligned} F_1|_{\Sigma_{div}^-} &= \frac{1}{v} = t, & F_2|_{\Sigma_{div}^-} &= u = s, \\ F_1|_{\Omega_+} &= \frac{w^3}{v} = t, & F_2|_{\Omega_+} &= w = s. \end{aligned}$$

Let us fix $t \in [N, +\infty]$. We observe that the restriction of the foliation $\sigma_1^* \mathcal{F}$ to the transversal section $\Sigma_{div}^+ = \{w = 1\}$ is analytic with respect to s . Then, after taking an scaled variation with respect to s , the relative cycle $\delta_{div}^+(s, t)$ is replaced by a loop θ_1 , modulo homotopy, which consists of line segment $\ell_{31} = [p_3, p_1]$ connecting the Morse point p_3 with the point p_1 encircling the latter along a small counterclockwise circular arc α_1 and then returning along the segment $\ell_{13} = [p_1, p_3]$. The loop θ_1 can be moved along the complex curve $\{u = w = 0\}$. Then, we have

$$\mathcal{V}ar_{(s,1)} \delta_{div}^+(s, t) = \theta_1 = \ell_{31} \alpha_1 \ell_{13}.$$

The same computation of the scaled variation with respect to s for the relative cycle $\delta_{div}^-(s, t)$ gives us a loop θ_3 , modulo homotopy, which can be moved along the complex plane $\{u = w = 0\}$. The loop θ_3

consists of line segment $\ell_{32} = [p_3, p_2]$ connecting the point p_3 with the point p_2 encircling the latter along a small counterclockwise circular arc α_3 and then returning along the segment $\ell_{23} = [p_2, p_3]$. Then, we have

$$\mathcal{V}ar_{(s,1)}\delta_{div}^-(s,t) = \theta_3 = \ell_{32}\alpha_3\ell_{23}.$$

In the local chart $(V_{div}, (u, v, w))$, we define the transversal section $\Omega_+ = \{u = 1\}$ (resp $\Omega_+ = \{u = 1\}$) near p_1 (resp near p_2). The restriction of the foliation $\sigma_1^*\mathcal{F}$ to the transversal section Ω_+ is given by

$$F_1|_{\Omega_+} = \frac{w^3}{v} = t, \quad F_2|_{\Omega_+} = w = s.$$

On the second step let us fix $t \in [0, 2N]$. After taking an scaled variation with respect to s , the relative cycle $\delta_{div}(s, t)$ is replaced by a figure eight cycle which can be moved along the complex line $C_{div}^t = \{x = 0, G_1 = t\}$ of the foliation $\sigma_1^*\mathcal{F}$. This case is similar to the classical situation which is studied by Bobieński and Mardešić in [2].

Now using the analyticity of the lifting $\sigma^{-1}\mathcal{F}$ with respect to s , the scaled variation of the cycle of integration $\delta(s, t)$ with respect to s is equal to the scaled variation with respect to s of the following difference $\delta_{div}^+(s, t) - \delta_{div}^-(s, t)$ which is equal, modulo homotopy, to the cycle $\theta_1\theta_3^{-1}$, where θ_3^{-1} is the inverse of the loop θ_3 . Schematically, the loop $\theta_1\theta_3^{-1}$ is a figure eight cycle. \square

Remark 3.

- In the local chart $(V_{div}^+, (u, v, w))$ (resp $V_{div}^-, (u, v, w)$), the loop θ_1 (resp θ_3) generating the fundamental group of the complex plane $\{u = w = 0\} \setminus \{p_1\}$ (resp $\{u = w = 0\} \setminus \{p_2\}$) with base point p_3 .
- By the univalence of the blown-up one form $\sigma_1^*\eta_2$, we have

$$\mathcal{V}ar_{(t,\alpha)} \int_{\delta(s,t)} \sigma_1^*\eta_2 = \int_{\mathcal{V}ar_{(t,\alpha)}\delta(s,t)} \sigma_1^*\eta_2.$$

5 Proof of the Theorem

In this section we first take benefit from the blowing-up in the family to prove our principal theorem. the proof is analogous of the following :

Theorem 2. *There exists a bound of the number of zeros of the function $t \mapsto J(s, t)$, for $t \in [0, +\infty]$ and $s > 0$ sufficiently small. This bound is locally with respect to all parameters uniform, in particular with respect to s .*

Let $\beta = (\beta_1, \dots, \beta_{k+2})$ where $\beta_1 = 3, \beta_2 = -1, \beta_3 = -\alpha_1, \dots, \beta_{k+2} = -\alpha_k$. Let D_1 is slit annulus in the complex plane \mathbb{C}_t^* with boundary ∂D_1 . This boundary is decomposed as follows $\partial D_1 = C_{R_1} \cup C_{r_1} \cup C^\pm$, where $C_{R_1} = \{|t| = R_1, |\arg t| \leq \alpha\pi\}$, $C^\pm = \{r_1 < |t| < R_1, |\arg t| = \pm\alpha\}$ and $C_{r_1} = \{|t| = r_1, |\arg t| \leq \alpha\pi\}$.

Petrov's method gives us that the number of zeros $\#Z(J(s, t))$ of the function $J(s, t)$ in slit annulus D_1 is bounded by the increment of the argument of $J(s, t)$ along ∂D_1 divided by 2π i.e.

$$\begin{aligned} \#Z(J(s, t)|_{D_1}) &\leq \frac{1}{2\pi} \Delta \arg(J(s, t)|_{\partial D_1}) = \frac{1}{2\pi} \Delta \arg(J(s, t)|_{C_{R_1}}) \\ &\quad + \frac{1}{2\pi} \Delta \arg(J(s, t)|_{C^\pm}) + \frac{1}{2\pi} \Delta \arg(J(s, t)|_{C_{r_1}}) \end{aligned}$$

(A) The increment of argument $\Delta \arg(J(s, t)|_{C_{R_1}})$ is uniformly bounded by Gabrielov's theorem [6].

(B) We use the Schwartz's principle

$$\operatorname{Im}(J(s, t))|_{C^\pm} = \mp 2i \operatorname{Var}_{(t, \alpha)} J(s, t).$$

Thus, the increments of argument along segments C^\pm are bounded by zeros of the variation $\operatorname{Var}_{(t, \alpha)} J(s, t)$ on segment (r, R) . By identity (18), the function $\operatorname{Var}_{(t, \beta_i)} J(s, t)$ can be written as follows

$$\begin{aligned} \operatorname{Var}_{(t, \beta_i)} J(s, t) &= K(t^{\frac{\beta_1}{\beta_i}}, \dots, t^{\frac{\beta_{k+\mu}}{\beta_i}}, s; \log s) \\ &= K(e^{\frac{\beta_1}{\beta_i} \log t}, \dots, e^{\frac{\beta_{k+\mu}}{\beta_i} \log t}, e^{\log s}; \log s) \end{aligned}$$

where K is a meromorphic function. The function $\operatorname{Var}_{(t, \beta_i)} J(s, t)$ is logarithmico-analytic function of type 1 in the variable s (see [9]). Then, there exist a finite cover of $\mathbb{R}^{k+\mu+1} \times \mathbb{R}$ by a logarithmico-exponential cylinders, using Rolin-Lion's theorem [9], such that on each cylinder of this family we have

$$\operatorname{Var}_{(t, \beta_i)} J(s, t) = y_0^{r_0} y_1^{r_1} A(t) U(t, y_0, y_1),$$

with $y_0 = s - \theta_0(t)$, $y_1 = \log y_0 - \theta_1(t)$, where θ_0, θ_1, A are logarithmico-exponential functions and U is a logarithmico-exponential unity function. As the number of zeros of a logarithmico-exponential function is bounded, the number of zeros of $\operatorname{Var}_{(t, \beta_i)} J(s, t)$ is bounded.

(C) Finally, we estimate the increment of argument of J along the small arc C_{r_1} . Then, it is necessarily to study the increment of argument of the leading term of the function J at $t = 0$.

Lemma 1. *The increment of the argument of $J(s, t)$ along the small circular arc C_{r_1} can be estimated by the increment of the argument of a some meromorphic function $F(s, t)$.*

Proof. The problem of the estimation of the increment of the argument of $J(s, t)$ along the circular arc C_{r_1} consist that the principal part of the function J contains the term $\log s \rightarrow -\infty$ as $s \rightarrow 0$. To resolve this problem we make a blowing-up at the origin in the total space with coordinates (x, y, z) where

$$x = J_1(s, t), \quad y = J_2(s, t), \quad z = (\log s)^{-1}.$$

The function $J(s, t)$ can be rewritten as follows

$$J(s, t) = J_1(s, t) + J_2(s, t) \log s = ((\log s)^{-1} J_1(s, t) + J_2(s, t)) \log s = (zx + y)z^{-1}.$$

Thus, for $z^{-1} \in \mathbb{R}$ be sufficiently small, we have

$$\arg(J(s, t)) = \arg((zx + y)z^{-1}) = \arg(zx + y).$$

To estimate the increment of argument of $zx + y$ uniformly with respect to $s > 0$ we make a quasi-homogeneous blowing-up π_1 with weight $(\frac{1}{2}, 1, \frac{1}{2})$ of the polynomial $zx + y$ at $C_1 = \{x = y = z = 0\}$ (the centre of blowing-up). The explicit formula of the quasi-homogeneous blowing-up π_1 in the affine charts $T_1 = \{x \neq 0\}$, $T_2 = \{y \neq 0\}$ and $T_3 = \{z \neq 0\}$ is written respectively as

$$\begin{aligned} \pi_{11} : x &= \sqrt{x_1}, \quad y = y_1 x_1, \quad z = z_1 \sqrt{x_1}, \\ \pi_{12} : x &= x_2 \sqrt{y_2}, \quad y = y_2, \quad z = z_2 \sqrt{y_2}, \\ \pi_{13} : x &= x_3 \sqrt{z_3}, \quad y = y_3 z_3, \quad z = \sqrt{z_3}. \end{aligned}$$

The pull-back $\pi_1^*(zx + y)$ is given, in different charts, by

$$\begin{aligned} \pi_{11}^*(zx + y) &= x_1(z_1 + y_1) = d_1 P_1(x_1, y_1, z_1), \\ \pi_{12}^*(zx + y) &= y_2(x_2 z_2 + 1) = d_2 P_2(x_2, y_2, z_2), \\ \pi_{13}^*(zx + y) &= z_3(x_3 + y_3) = d_3 P_3(x_3, y_3, z_3). \end{aligned}$$

where $d_i = 0$ and $P_i = 0$ are equations of exceptional divisor and the strict transform of $zx + y = 0$ respectively.

Observe that $P_i = 0, i = 1, 3$ has not a normal crossing with the exceptional divisor $d_i = 0, i = 1, 3$. To resolve this problem we make a second blowing-up π_2 with centre a subvariety C_2 which is given, in different charts, as following:

1. In the chart T_1 , choose a local coordinate chart with coordinates (x_1, y_1, z_1) in which $C_2 = \{y_1 = z_1 = 0\}$. Then $\pi_2^{-1}(C_2)$ is covered by two coordinate charts U_{y_1} and U_{z_1} with coordinate $(\tilde{x}_1, \tilde{y}_1, \tilde{z}_1)$ where in y_1 -chart U_{y_1} the blowing-up π_2 is given by $x_1 = \tilde{x}_1, y_1 = \tilde{y}_1, z_1 = \tilde{z}_1 \tilde{y}_1$ and in z_1 -chart U_{z_1} the blowing-up π_2 is given by $x_1 = \tilde{x}_1, y_1 = \tilde{y}_1 \tilde{z}_1, z_1 = \tilde{z}_1$.
2. In the chart T_2 , the blowing-up π_2 is a biholomorphism (π_2 is a proper map).
3. In this chart T_3 , choose a local coordinate chart with coordinates (x_3, y_3, z_3) in which $C_2 = \{x_3 = y_3 = 0\}$. Then $\pi_2^{-1}(C_2)$ is covered by two coordinate charts U_{x_3} and U_{y_3} with coordinate $(\tilde{x}_3, \tilde{y}_3, \tilde{z}_3)$ where in x_3 -chart U_{x_3} the blowing-up π_2 is given by $x_3 = \tilde{x}_3, y_3 = \tilde{y}_3 \tilde{x}_3, z_3 = \tilde{z}_3$ and in y_3 -chart U_{y_3} the blowing-up π_2 is given by $x_3 = \tilde{x}_3 \tilde{y}_3, y_3 = \tilde{y}_3, z_3 = \tilde{z}_3$.

The pull-back $\pi_1^*(zx + y)$ is given, in different charts, by

- In the y_1 -chart U_{y_1} , the transformation of the pull-back $\pi_1^*(zx + y)$ by the blowing-up π_2 is given by

$$\pi_2^* \circ \pi_1^*(zx + y) = \pi_2^*(d_1 P_1(x_1, y_1, z_1)) = \tilde{x}_1 \tilde{y}_1 (\tilde{z}_1 + 1) \stackrel{0}{\approx} \tilde{x}_1 \tilde{y}_1 = J_2(s, t) = F(s, t).$$

- In the z_1 -chart U_{z_1} , the transformation of the pull-back $\pi_1^*(zx + y)$ by the blowing-up π_2 is given by

$$\pi_2^* \circ \pi_1^*(zx + y) = \pi_2^*(d_1 P_1(x_1, y_1, z_1)) = \tilde{z}_1 \tilde{x}_1 (\tilde{y}_1 + 1) \stackrel{0}{\approx} \tilde{x}_1 \tilde{z}_1 = (\log s)^{-1} J_1(s, t) = F(s, t).$$

- In the chart T_2 , we have

$$\pi_2^* \circ \pi_1^*(zx + y) = \pi_2^*(d_2 P_2(x_2, y_2, z_2)) = d_2 P_2(x_2, y_2, z_2) = (\log s)^{-1} J_1(s, t) + J_2(s, t) = F(s, t).$$

- In the x_3 -chart U_{x_3} , the transformation of the pull-back $\pi_1^*(zx + y)$ by the blowing-up π_2 is given by

$$\pi_2^* \circ \pi_1^*(zx + y) = \pi_2^*(d_3 P_3(x_3, y_3, z_3)) = \tilde{z}_3 \tilde{x}_3 (\tilde{y}_3 + 1) \stackrel{0}{\approx} \tilde{x}_3 \tilde{z}_3 = (\log s)^{-1} J_1(s, t) = F(s, t).$$

- In the y_3 -chart U_{y_3} , the transformation of the pull-back $\pi_1^*(zx + y)$ by the blowing-up π_2 is given by

$$\pi_2^* \circ \pi_1^*(zx + y) = \pi_2^*(d_3 P_3(x_3, y_3, z_3)) = \tilde{z}_3 \tilde{y}_3 (\tilde{x}_3 + 1) \stackrel{0}{\approx} \tilde{y}_3 \tilde{z}_3 = J_2(s, t) = F(s, t).$$

Finally, we distinguish three cases:

1. $\arg_{C_{r_1}} J(s, t) = \arg_{C_{r_1}} ((\log s)^{-1} J_1(s, t)) = \arg_{C_{r_1}} J_1(s, t), ((\log s)^{-1} \in \mathbb{R})$
2. $\arg_{C_{r_1}} J(s, t) = \arg_{C_{r_1}} J_2(s, t),$
3. In the chart T_2 , the function $F(s, t) = ((\log s)^{-1} J_1(s, t)) + J_2(s, t)$ is meromorphic.

□

Now we define the functional space \mathcal{P}_β which are formed of coefficients of the polynomials P_i of the Darboux first integral H , the coefficients of the polynomials R, S of the perturbative one forme η , exponents α_i and degrees $n_i = \deg P_i, n = \max(\deg R, \deg S)$. Consider the following finite dimensional functional space \mathcal{P}_β

$$\mathcal{P}_\beta(m_\beta, M_\beta; \beta_1, \dots, \beta_{k+2}) = \left\{ \sum_{j=1}^{k+2} \sum_{n, \ell} A_{j\ell n}(s) t^{\beta_j n} s^m \log^\ell(t) : \right. \\ \left. A_{j\ell n}(s) \in \mathbb{C}, m_\beta < A_{j\ell n} < M_\beta, 0 \leq \ell \leq k+1 \right\}.$$

For the first two cases, the function $J_i(s, t), i = 1, 2$ satisfies the following iterated variations equation with respect to t

$$\mathcal{V}ar_{(t, \beta_1)} \circ \dots \circ \mathcal{V}ar_{(t, \beta_{k+2})} J_i(s, t) = 0.$$

Thus, by Lemma 4.8 from [2], there exists a non zero leading term $P_{i\beta} \in \mathcal{P}_\beta$ of $J_i(s, t), i=1,2$ at $t = 0$ such that $|J_i(s, t) - P_{i\beta}(s, t)| = O(t^{\mu_1}), \mu_1 > 0$, uniformly in s . Moreover, the function $J_i(s, t), i = 1, 2$ satisfies the iterated variation equation

$$\mathcal{V}ar_{(s, 1)} J_i(s, t) = 0.$$

Thus, we have $J_i(s, t) = O(s^{\mu_2}), \mu_2 > 0$, uniformly in t .

For each element in the parameter space, we can choose the leading term of $P_{i\beta}$. The increment of argument of this leading term is bounded by a constant $C(M_\beta, k+2, \beta_{k+2})$. Since the leading term of $P_{i\beta}$ is also the leading term of $J_i(s, t)$, the limit $\lim_{r_1 \rightarrow 0} \Delta \arg(J_i(s, t)|_{C_{r_1}}) \leq C(M_\beta, k+2, \beta_{k+2})$.

In the chart T_2 , the function F is meromorphic. Thus, this function can be rewritten as following

$$F(s, t) = (\log s)^{-1} J_1(s, t) + J_2(s, t) = G(t^{\beta_1}, \dots, t^{\beta_k}, s, (\log s)^{-1})$$

where G is meromorphic function. The number $\#Z(G)$ of zeros of the function G is uniformly bounded. The latter claim is a direct application of fewnomials theory of Khovanskii [8]: since the functions $\epsilon_i(t) = t^{\beta_i}, \epsilon(s) = (\log s)^{-1}$ are Pfaffian functions (solutions of Pfaffian equations $t d\epsilon_i - \beta_i \epsilon_i dt = 0$ and $s d\epsilon + \epsilon^2 ds$, respectively), the upper bound for this number of zeros can be given, using Rolle-Khovanskii arguments of [7], in terms of the number of zeros of some polynomial and its derivatives. The latter are uniformly bounded by Gabrielov's theorem [6].

References

- [1] Bobieński, Marcin Pseudo-Abelian integrals along Darboux cycles *a codimension one case*. *J. Differential Equations* 246 (2009), no. 3, 1264-1273.
- [2] Bobieński, Marcin; Mardešić, Pavao *Pseudo-Abelian integrals along Darboux cycles*. *Proc. Lond. Math. Soc. (3)* 97 (2008), no. 3, 669-688.
- [3] Bobieński, Marcin; Mardešić, Pavao; Novikov, Dmitry *Pseudo-Abelian integrals: unfolding generic exponential*. *J. Differential Equations* 247 (2009), no. 12, 3357-3376.
- [4] Bobieński, Marcin; Mardešić, Pavao; Novikov, Dmitry *Pseudo-Abelian integrals on slow-fast Darboux systems*. *Ann. Inst. Fourier (Grenoble)*, 2013, to appear.
- [5] Braghtha Aymen (2013) *Les zéros des intégrales pseudo-abéliennes: cas non générique*. *PhD thesis. Université de Bourgogne: France*.
- [6] Gabrièlov, A. M *Projections of semianalytic sets. (Russian) Funktsional. Anal. i Priložen.* 2 1968 no. 4, 18-30

- [7] Khovanskii, A, G. *Fewnomials. Translations of Mathematical Monographs, 88. American Mathematical Society, Providence, RI, 1991. 139 pp.*
- [8] Khovanskii, A, G. *Real analytic manifolds with property of finiteness, and complex abelian integrals, Funktsional. Ana. i Prolizhen. 18 (2) (1984) 40-50.*
- [9] Lion, Jean Marie; Rolin, Jean Phillippe *Théorème de préparation pour les fonctions logarithmico-exponentielles. Annales de l'institut de Fourier, tome 47, n3 (1997) p.859-884.*

**Burgundy University, Burgundy Institute of Mathematics,
U.M.R. 5584 du C.N.R.S., B.P. 47870, 21078 Dijon
Cedex - France.
E-mail adress: aymenbraghtha@yahoo.fr**