



**HAL**  
open science

## A Dual Polybasic Motif Determines Phosphoinositide Binding and Regulation in the P2X Channel Family

Louis-Philippe Bernier, Dominique Blais, Eric Boué-Grabot, Philippe Séguéla

► **To cite this version:**

Louis-Philippe Bernier, Dominique Blais, Eric Boué-Grabot, Philippe Séguéla. A Dual Polybasic Motif Determines Phosphoinositide Binding and Regulation in the P2X Channel Family. PLoS ONE, 2012, 7 (7), p.e0040595. 10.1371/journal.pone.0040595 . hal-01146617

**HAL Id: hal-01146617**

**<https://hal.science/hal-01146617v1>**

Submitted on 28 Apr 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A Dual Polybasic Motif Determines Phosphoinositide Binding and Regulation in the P2X Channel Family

Louis-Philippe Bernier<sup>1</sup>, Dominique Blais<sup>1</sup>, Éric Boué-Grabot<sup>2</sup>, Philippe Séguéla<sup>1\*</sup>

<sup>1</sup> Department of Neurology and Neurosurgery, Montreal Neurological Institute, Alan Edwards Centre for Research on Pain, McGill University, Montréal, Québec, Canada, <sup>2</sup> Institut des Maladies Neurodégénératives, Université Bordeaux Segalen, CNRS UMR 5293, Bordeaux, France

## Abstract

Phosphoinositides modulate the function of several ion channels, including most ATP-gated P2X receptor channels in neurons and glia, but little is known about the underlying molecular mechanism. We identified a phosphoinositide-binding motif formed of two clusters of positively charged amino acids located on the P2X cytosolic C-terminal domain, proximal to the second transmembrane domain. For all known P2X subtypes, the specific arrangement of basic residues in these semi-conserved clusters determines their sensitivity to membrane phospholipids. Neutralization of these positive charges disrupts the functional properties of the prototypical phosphoinositide-binding P2X4 subtype, mimicking wortmannin-induced phosphoinositide depletion, whereas adding basic residues at homologous positions to the natively insensitive P2X5 subtype establishes *de novo* phosphoinositide-mediated regulation. Moreover, biochemical evidence of *in vitro* P2X subunit-phospholipid interaction and functional intracellular phosphoinositide-binding assays demonstrate that the dual polybasic cluster is necessary and sufficient for regulation of P2X signaling by phospholipids.

**Citation:** Bernier L-P, Blais D, Boué-Grabot É, Séguéla P (2012) A Dual Polybasic Motif Determines Phosphoinositide Binding and Regulation in the P2X Channel Family. PLoS ONE 7(7): e40595. doi:10.1371/journal.pone.0040595

**Editor:** Yoshiro Ishimaru, University of Tokyo, Japan

**Received:** March 14, 2012; **Accepted:** June 11, 2012; **Published:** July 11, 2012

**Copyright:** © 2012 Bernier et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** Supported by CIHR and NSERC operating grants (PS) and CNRS (EBG). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors declare that no competing interests exist.

\* E-mail: philippe.seguela@mcgill.ca

## Introduction

ATP-gated P2X receptor channels play significant roles in pain transduction, neuro-immune interactions and inflammatory response therefore understanding their regulation mechanisms is critical. Plasma membrane phosphoinositides (PIP<sub>n</sub>) are anionic phospholipids that act as functional regulators of many types of ion channels. They are necessary cofactors for activation or desensitization of various channels, including transient receptor potential (TRP) channels [1,2], inward rectifier K<sup>+</sup> (Kir) channels [3,4] and voltage-gated KCNQ channels [5]. Most ATP-gated P2X receptor subtypes are potentiated by intracellular PIP<sub>n</sub>. P2X1, P2X2, P2X3, P2X4 and P2X7 are functionally sensitive to PIP<sub>n</sub> [6,7,8,9], however, P2X5 was found to be PIP<sub>n</sub>-insensitive [10].

In P2X subunits, the few residues shown to be implicated in PIP<sub>n</sub>-mediated regulation are located in the C-terminal domain, which is also involved in subunit trafficking, phosphorylation, heteromerization and multi-receptor crosstalks [11,12,13,14].

Although no consensus PIP<sub>n</sub> binding site exists among membrane proteins, analysis of the PIP<sub>n</sub>-binding region of PH domain-containing proteins points to the necessary presence of basic amino acids interacting with the anionic headgroup of PIP<sub>n</sub> [15]. The identity of PIP<sub>n</sub>-binding domains in ion channels in particular has been even more elusive, with only a few putative residues identified as involved in the interaction, all on the cytoplasmic side of the membrane [for review [16]]. Although direct protein-phospholipid binding was demonstrated for several families of PIP<sub>n</sub>-sensitive channels, no common motif can predict an effective interaction.

Here, we demonstrate that P2X channel subunits bind to PIP<sub>n</sub> via two clusters of positively charged residues located in the proximal C-terminal domain. The specific arrangement of basic and acidic amino acids found in these semi-conserved clusters predicts the PIP<sub>n</sub>-sensitivity of all known P2X subtypes. By mutating the prototypical PIP<sub>n</sub>-sensitive P2X4 and PIP<sub>n</sub>-insensitive P2X5 subtypes, we provide functional and biochemical evidence that a dual cluster motif in the proximal C-terminal domain is necessary and sufficient for the regulation of P2X receptor channels by PIP<sub>n</sub>.

## Results

### A PIP<sub>n</sub>-binding Site in P2X C-terminal Domains


Several studies have demonstrated functional modulation of P2X receptors by PIP<sub>n</sub> as well as direct PIP<sub>n</sub> binding to the C-terminal domain of various P2X subunits (Figure 1A, left columns) [6,7,8,9,10,17]. P2X1, P2X2 and P2X4 subunits directly bind PIP<sub>n</sub>, whereas P2X3 and P2X5 do not. Hence, we analyzed their respective C-terminal sequences and found that PIP<sub>n</sub> binding correlates with the net positive charge of two polybasic amino acid clusters (Figure 1, shaded areas 1 and 2). P2X1/2/4 contain 6–7 basic residues (lysine, arginine or histidine) in these two clusters and a maximum of one negatively-charged residue (aspartic or glutamic acid). On the other hand, the non PIP<sub>n</sub>-binding P2X3 and P2X5 subunits contain 5 and 6 basic residues, respectively, but also 3 acidic residues disrupting the global positive charge of the clusters. We therefore hypothesized that the dual cluster's charge is responsible for the affinity of the P2X C-terminus to

PIP<sub>n</sub>. Interestingly, a conserved hydrophobic tyrosine residue is located between the two clusters, and could also be involved in the interaction. This cytosolic region located 3 residues from the second transmembrane domain (TM2) region lies in close proximity to the plasma membrane where electrostatic interactions with the negative phosphate head groups of the membrane-anchored PIP<sub>n</sub> can take place.

A closer look at the single residues that have, in this report or previously, been demonstrated to be involved in PIP<sub>n</sub>-mediated regulation shows they are all (except one in P2X7) located within the two polybasic clusters (Fig. 1B). No evidence of binding has been found on the N-terminal domain of any P2X subunit. The P2X6 subtype was excluded from our study since it does not form functional homomeric receptors [18].

### A Dual Polybasic Cluster Motif is Necessary for PIP<sub>n</sub>-binding and P2X4 Channel Regulation

We reported that P2X4 is a prototypical PIP<sub>n</sub>-dependent P2X subtype, being tightly regulated via direct binding to PIP<sub>2</sub> and PIP<sub>3</sub> [8]. We therefore aimed to neutralize the PIP<sub>n</sub>-binding site by mutating key lysine residues. We found that neutralizing the charge of either of the two clusters, by mutating lysines 362 and 363 or lysines 370 and 371 into neutral glutamines, leads to a loss of PIP<sub>n</sub> binding in an *in vitro* binding assay where a GST-fusion protein coding for a 16-amino acid sequence (Figure 2A,B) is applied to various PIP<sub>n</sub>. The lysine-to-glutamine mutations performed on residues 362 and 363 also induced significant changes in the P2X4 channel activity. Expressed in the *Xenopus* oocyte expression system, the P2X4 mutant with lower PIP<sub>n</sub>-



**Figure 1. The proximal C-terminal domain of P2X subunits contains a semi-conserved PIP<sub>n</sub>-binding motif.** A) Sequence alignment of rat P2X C-termini proximal to the TM2 domain showing the two polybasic clusters (shaded area, 1 and 2). The left column summarizes, for each subunit, the presence (+) or absence (-) of binding of the GST-fusion C-terminal domain to PIP<sub>n</sub> in PIP strip assays. The second column shows the presence (+) or absence (-) of modulation by PIP<sub>n</sub> in functional assays. Basic residues are shown in red and acidic residues in blue. B) Sequences showing residues that were reported (here or previously) to be involved in PIP<sub>n</sub> regulation. Basic residues in red, acidic residues in blue and an uncharged serine in green. C) Schematic representation of the topology of a P2X subunit showing binding of two positively charged amino acid clusters to membrane-bound PIP<sub>n</sub>.

doi:10.1371/journal.pone.0040595.g001

binding affinity displayed a stronger current rundown upon repeated ATP applications as well as slower activation and desensitization current phases (Figure 2C,D), all these effects mimicking those brought by pharmacological  $\text{PIP}_n$  depletion [8]. The K362Q-K363Q mutant receptor was more strongly inhibited by wortmannin-induced  $\text{PIP}_n$  depletion than the wild-type (WT) receptor (Figure 2E), due to its lower affinity to  $\text{PIP}_n$ . The mutation targetting the second basic cluster (K370Q-K371Q) could not be tested functionally as P2X4 channels with mutations on residue 371 are non-functional due to the role of conserved lysine 371 in receptor trafficking [11].

### P2X1 and P2X7 Binding to $\text{PIP}_n$ is Consistent with the Dual Polybasic Cluster Model

For the P2X1 subtype, the results that we have previously reported are consistent with our model, in that mutating the K359 residue in the first cluster suppressed *in vitro* binding to  $\text{PIP}_n$ , and induced a  $\text{PIP}_n$ -depleted like current phenotype [7]. To confirm that both clusters are involved in the interaction with  $\text{PIP}_n$ , we neutralized the charge in the second cluster via a lysine-to-glutamine mutation on residue 364. This also induced a loss of binding in a phospholipid strip assay (Figure 3A), confirming that both clusters are necessary for the P2X1 C-terminus to bind  $\text{PIP}_n$ .

The P2X7 subtype was also analyzed, but no direct binding was found in our biochemical binding assay using C-terminal peptides of various length (Figure 3B). The absence of binding is likely due to the presence of only one polybasic cluster in the P2X7 C-terminus. Nevertheless, a previous report demonstrates through a mutational study that specific amino acids are involved in  $\text{PIP}_n$  modulation of P2X7 [17], suggesting a more complex binding mechanism, likely due to the presence of an additional 18-residue long sequence between the cluster and TM2.

### Generation of De Novo $\text{PIP}_n$ -regulation in the P2X5 Subtype

To verify if the presence of C-terminal polybasic clusters is sufficient for  $\text{PIP}_n$  regulation of P2X receptor channels, we chose the natively  $\text{PIP}_n$ -insensitive P2X5 subunit and examined the effect of adding basic residues to its C-terminus (Figure 4A,B). In the phospholipid binding assay, adding positive charges to the cluster proximal to TM2 by mutating residues 365 and 366 induced binding to several  $\text{PIP}_n$ . Also, mutating a negatively-charged glutamic acid into a lysine in the second cluster enhanced binding, thereby showing that  $\text{PIP}_n$  binding can be obtained via negative-to-positive mutations in the twin clusters (Figure 4C). We then analyzed the functional effect of that mutation by recording from the S365K-E366Y-E374K mutant in the *Xenopus* oocyte expression system: the mutant P2X5 receptor generated currents ~15 times larger than the WT upon 10  $\mu\text{M}$  ATP activation. Adding basic residues to the first cluster only (S365K-E366Y) also led to significantly larger currents than the WT. Whereas WT P2X5 is unaffected by wortmannin treatment, both  $\text{PIP}_n$ -binding mutants were strongly inhibited by wortmannin-induced  $\text{PIP}_n$  depletion, suggesting that  $\text{PIP}_n$  binding is responsible for the current amplitude increase induced by the C-terminal mutations (Figure 4B,D). WT P2X5 channels display a marked current rundown upon repeated activation; such a feature was absent in the triple mutant, but could be restored after pharmacological  $\text{PIP}_n$  depletion (Figure 4F). Also, the activation rate of the P2X5 current was faster in the  $\text{PIP}_n$ -binding mutant than in the WT, as was the desensitization rate. Both properties were restored towards WT levels after wortmannin treatment, confirming the  $\text{PIP}_n$ -sensitivity of the mutant receptor channel.

The human P2X5 ortholog has an arginine residue on position 365 and has been shown to evoke currents of much larger amplitude than its rat homolog [19], we therefore verified if  $\text{PIP}_n$  play a role in this difference. Wortmannin-induced  $\text{PIP}_n$  depletion significantly reduced the hP2X5 current amplitude (Figure 4E), indicating that interspecies differences exist in terms of functional  $\text{PIP}_n$  regulation of P2X5 channels and confirming the importance of the positive charges found in the proximal polybasic cluster.


### P2X4 C-terminal Peptides Compete for Intracellular $\text{PIP}_n$ and Induce a $\text{PIP}_n$ -depletion Current Phenotype

To confirm that P2X C-terminal polybasic clusters bind to  $\text{PIP}_n$  in a cytoplasmic environment, we performed an intracellular  $\text{PIP}_n$ -binding competition assay. HEK293 cells transiently expressing P2X4 were recorded in whole-cell patch-clamp configuration, and various GST fusion proteins containing the P2X C-terminal peptides (16 amino acid-long, P2X4: C360-V375, P2X5: L361-V376) were added to the intracellular milieu. When  $\text{PIP}_n$ -binding P2X4 C-terminus peptides were introduced through the patch pipette, a strong rundown of the ATP-mediated P2X4 current as well as a strong decrease in desensitization rate were observed, suggesting that the P2X4 C-terminal peptide competes for intracellular  $\text{PIP}_n$  binding, inducing a  $\text{PIP}_n$ -depletion current phenotype (Figure 5A,C,D). Peptides coding for the P2X4 K362Q-K363Q mutant C-terminal domain did not induce any change in the current phenotype as compared to a control GST peptide injection, indicating an inability to bind intracellular  $\text{PIP}_n$ . Reciprocally, the WT P2X5 C-terminal peptide had no effect on both functional parameters measured (Figure 5B,C,D) due to its low  $\text{PIP}_n$ -binding affinity. Strong interactions between the P2X5 S365K-E366Y-E374K mutant peptides and  $\text{PIP}_n$  led to rundown and slower desensitization of the P2X4 currents.

## Discussion

Membrane  $\text{PIP}_n$  regulate the activity of a wide variety of ion channels, and the mechanism of interaction between these important membrane proteins and the anionic phospholipids draws lots of attention. We show here that the modulation of P2X channel function by  $\text{PIP}_n$  is predicted by the subunit's ability to bind to the negative inositol triphosphate head group of the lipid via two adjacent clusters of basic amino acids located on the C-terminal domain. Binding of  $\text{PIP}_n$  to the clusters present in P2X1, P2X2 and P2X4 likely leads to a conformational change in the C-terminus, a domain highly involved in functional regulation of the channel.

The P2X3, P2X5 and P2X7 subunits lack this microdomain and therefore do not directly interact with  $\text{PIP}_n$ . While the absence of this microdomain renders P2X5 channel activity insensitive to  $\text{PIP}_n$ , P2X3 and P2X7 are functionally modulated by  $\text{PIP}_n$  [6,17], strongly indicating an indirect regulation. A mechanism in which a  $\text{PIP}_n$ -binding partner protein acts as a regulatory subunit has been proposed for TRPV1, where *phosphoinositide interacting regulator of TRP* (Pirt) is necessary for  $\text{PIP}_n$ -mediated enhancement of the channel activity [20]. Pirt, a membrane protein which binds  $\text{PIP}_2$  via a cluster of basic residues on its C-terminus, also complexes with TRPV1 to link both molecules. A similar interaction was observed in the case of NMDA receptors, where the cytosolic tails of the NR1 or NR2B subunits bind  $\alpha$ -actinin, an actin-crosslinking protein.  $\alpha$ -actinin also binds membrane  $\text{PIP}_2$  and modifies the NMDA receptor's intracellular tail conformation to promote channel opening [21]. A similar mechanism could underlie the indirect  $\text{PIP}_n$ -dependent regulation of P2X3 and P2X7, but the nature of the partner involved remains to be elucidated.




**Figure 2. Requirement of two polybasic clusters in the PIP<sub>n</sub>-regulated P2X4 subtype.** A) Sequence of the P2X4 C-terminus showing lysine to glutamine mutations disrupting the positive charge of the first or second cluster (basic residues in red, acidic in blue, neutral mutations in grey). B) The GST construct containing the P2X4 C-terminal domain C360-V375 binds to several PIP<sub>n</sub> including PIP<sub>2</sub> and PIP<sub>3</sub>. Mutating the basic lysine residues K362 and K363, or K370 and K371 into neutral glutamine leads to a loss of binding to PIP<sub>n</sub> (n = 3–6). C) Representative ATP-activated P2X4 current traces obtained on P2X4-expressing *Xenopus* oocytes showing the slower activation and desensitization rates induced by the K362Q-K363Q mutation decreasing PIP<sub>n</sub>-binding affinity. D) Quantitative analysis of the functional changes induced by the K362Q-K363Q mutation on P2X4 current rundown (left), activation (middle) and desensitization (right). A larger rundown between agonist applications is observed with the mutant than with the WT (2<sup>nd</sup>/1<sup>st</sup> application: WT: 63.5 ± 3.8%, mutant: 50.4 ± 4.9%, n = 10–11). The mutant P2X4 channel shows a slower activation rate (10–90% rise time: WT: 0.67 ± 0.03 s, mutant: 0.80 ± 0.05 s, n = 70–80) and a slower desensitization rate (5-second decay %: WT: 55.6 ± 3.1%, mutant: 30.1 ± 2.6%, n = 55–61). E) Wortmannin-induced PIP<sub>n</sub> depletion leads to a stronger inhibition of P2X4 current amplitude in the K362Q-K363Q mutant than in WT (post/pre-treatment: WT: 61.3 ± 7.5%, mutant: 29.5 ± 4.6%, n = 30–50). \*: p < 0.05; \*\*\*: p < 0.001. doi:10.1371/journal.pone.0040595.g002

Interestingly, P2X7 forms a signalling complex with various proteins that includes  $\alpha$ -actinin [22], which could link the P2X7 C-terminal tail to PIP<sub>n</sub>.

Proteins bind PIP<sub>n</sub> via multiple contacts that require the contribution of multiple amino acids [16], hence we analyzed the general charge of the P2X C-terminal domain, and found that the

13-amino acid sequence containing both clusters has a predicted isoelectric point of 10.4 to 10.8 for P2X1/2/4/5, but that P2X3 and P2X7 have lower predicted values of 9.2 and 8.5, respectively. This could explain the lack of direct PIP<sub>n</sub> binding in these two subunits. However, PIP<sub>n</sub>-sensitive P2X1/2/4 and PIP<sub>n</sub>-insensitive P2X5 subunits have similar isoelectric points, suggesting that PIP<sub>n</sub>





**Figure 3. Requirement of two polybasic clusters for PIP<sub>n</sub>-binding in P2X1 and P2X7.** A) The GST construct containing the WT P2X1 C-terminal domain (L352-E378) (basic residues in red, acidic in blue, neutral mutations in grey) binds various PIP<sub>n</sub> on a phospholipid strip assay, whereas disrupting the positive charge of the first or second polybasic cluster with K359Q and K364Q mutations suppresses binding (n = 3). B) The absence of two polybasic clusters in the C-terminus of P2X7 prevents its binding to PIP<sub>n</sub> on a phospholipid strip assay (n = 3). Shown in grey boxes are various GST-fusion peptides generated.

doi:10.1371/journal.pone.0040595.g003

interaction is not only determined by the global charge of the structure, but depends on the specific spatial arrangement of charged amino acids. Although the C-terminal domains of P2X4 and P2X5 contain a similar number of basic residues, the presence of 3 acidic residues distributed within the two clusters of P2X5 disrupts the electrostatic pattern required for effective PIP<sub>n</sub> affinity.

While single cationic residues involved in subunit-PIP<sub>n</sub> interaction have been identified for most PIP<sub>n</sub>-dependent channels, clusters of basic residues were found in the TRP box of TRPM8 channels as well as in the carboxy-terminal tail of Kv7 channels [23,24]. It was also reported that a conserved sequence in the juxtamembrane C-terminus of Kir 1.1 and Kir 2.1 takes part in the protein-phospholipid interaction [25]. X-ray crystal structure modelling confirmed that the equivalent structure in Kir6.2 is part of the PIP<sub>2</sub>-binding pocket that includes 3 other basic residues [26]. Furthermore, the crystallization of Kir 2.2 and of GIRK2 in the presence of short-chain PIP<sub>2</sub> directly shows that PIP<sub>2</sub>-binding can induce significant conformational changes that modulate the channel function [27,28]. The recently published crystal structure of a truncated zebrafish P2X4.1 receptor [29] unfortunately does not contain the N- and C-terminal tails, therefore additional work will be needed to obtain structural information on this intracellular regulatory motif and on the exact conformational effect of PIP<sub>n</sub>-binding.


Another property common to the primary sequence of most known PIP<sub>n</sub>-binding pockets is the presence of at least one aromatic residue [15]. Interestingly, a fully conserved tyrosine residue lies between the two clusters forming the proposed P2X PIP<sub>n</sub>-binding motif. Mutating this residue on the P2X1 (Y363Q) and P2X4 (Y367Q) C-terminal sequence did not affect the binding to PIP<sub>n</sub> in *in vitro* overlay assays (data not shown). However, it may play a structural role in the secondary or tertiary conformation of the dual cluster domain, a possibility that could not be tested functionally as mutating this conserved residue induces a loss of function [11].

The gating mechanism underlying ion conduction in P2X channels has been extensively studied in recent years. It is believed

that the gate is located between residues 340 to 347 (nomenclature for zP2X4.1) in the TM2 domain, and that opening of the channel triggers rearrangement of the TM2 helices that reveals access to even deeper parts of the pore a few residues away from the cytoplasmic tail [29,30,31]. The PIP<sub>n</sub>-binding region therefore lies in close proximity to the gating machinery of P2X receptor channels, likely impacting on the open/closed transition through conformational changes.

Results obtained on the P2X4 and P2X5 subunits not only enhance our understanding of the PIP<sub>n</sub>-binding site of P2X receptors, but also demonstrate the importance of PIP<sub>n</sub> in the functional regulation of the channels. Disrupting the PIP<sub>n</sub> affinity of P2X4 led to major changes in the current phenotype, similar to what was seen in other PIP<sub>n</sub>-binding P2X subtypes [7,9]. Moreover, we were able for the first time to induce a PIP<sub>n</sub>-binding phenotype through single mutations in the otherwise PIP<sub>n</sub>-insensitive P2X5 subtype, demonstrating that the polybasic clusters motif is sufficient for PIP<sub>n</sub> binding and functional regulation. The high amplitude currents obtained with the gain-of-binding mutant suggest that the small size of currents mediated by the WT rat P2X5 in several expression systems [10,32] is due to its low PIP<sub>n</sub> affinity. Altogether, our results indicate that membrane PIP<sub>n</sub> contribute to the full expression of P2X receptor channel function.

Interestingly, it was shown that the human and chicken P2X5 receptors give rise to currents that are significantly larger, and desensitize faster than their rat counterpart [19,33]. Analysis of their C-terminal sequence shows that human P2X5 has a basic arginine residue on position 365, instead of a neutral serine found in the rat sequence, and that chicken P2X5 has a neutral asparagine on position 366, instead of a negatively-charged glutamic acid in the rat sequence. Our results demonstrate that a neutral-to-basic mutation on residue 365 and an acidic-to-neutral mutation on residue 366 induces a PIP<sub>n</sub>-binding phenotype in the rat P2X5. We also show that the human P2X5 channel is regulated by PIP<sub>n</sub> as pharmacological depletion of intracellular PIP<sub>n</sub> induced dramatic changes to its current phenotype. It has to be noted that on position 375, at the extremity of the second



**Figure 4. Mutations creating two polybasic clusters on the P2X5 C-terminus lead to  $PIP_n$  binding and a  $PIP_n$ -regulated current phenotype.** A) Sequence of the P2X5 C-terminus showing mutations adding positive charges to one (S365K-E366Y, SE→KY) or two clusters (S365K-E366Y-E374K, SEE→KYK) (basic residues in red, acidic in blue, neutral mutations in grey). B) Representative ATP-activated current traces recorded in *Xenopus* oocytes expressing P2X5 WT or SEE→KYK mutant in control and wortmannin conditions. C) The GST construct containing the WT P2X5

proximal C-terminal domain L361-V376 does not bind to PIP<sub>n</sub>. Adding positive residues to one amino acid cluster with the SE→KY mutation induces binding to several PIP<sub>n</sub>. Creating a second positive cluster with the SEE→KYK mutation increases PIP<sub>n</sub> binding (n=3–6). D) Quantitative data showing the SE→KY and SEE→KYK gain-of-binding mutations lead to increased current amplitude in rP2X5-expressing *Xenopus* oocytes (left graph; rP2X5 WT: 0.21±0.04 μA, SE→KY: 1.22±0.11 μA, SEE→KYK: 3.19±0.55 μA, n=13–25). The gain-of-binding mutants are sensitive to intracellular PIP<sub>n</sub> levels as wortmannin-induced PIP<sub>n</sub> depletion leads to a decrease in current amplitude (right graph, post/pre-treatment amplitude; WT: 107.5±21.8%, SE→KY: 50.3±10.7%, SEE→KYK: 40.9±8.4%, n=4–17). E) Human P2X5 channel currents are significantly inhibited by PIP<sub>n</sub> depletion (vehicle=4.90±0.79 μA; post-wortmannin=1.44±0.24 μA, n=7–10). F) Differences in current rundown, activation rate and desensitization rate between WT P2X5 and SEE→KYK mutant under control and PIP<sub>n</sub>-depletion conditions. The current rundown between successive applications measured with the WT P2X5 is prevented by gain-of-binding mutations, and is partially restored after a wortmannin treatment of the mutant (2<sup>nd</sup>/1<sup>st</sup> application: WT: 40.0±4.4%, mutant control: 113.9±8.0%, mutant wortmannin: 83.7±9.8%, n=5–10). The mutant P2X5 channel shows a faster current activation compared to WT, and it is slowed by PIP<sub>n</sub> depletion (10–90% rise time: WT: 3.77±0.69 s, mutant control: 0.86±0.08 s, mutant wortmannin: 2.31±0.25 s, n=7–8). The gain-of-binding P2X5 mutant current desensitizes faster than WT, and its desensitization rate is slowed by PIP<sub>n</sub> depletion (decay slope: WT: -0.001±0.006, mutant control: 0.31±0.09, mutant wortmannin: 0.07±0.02, n=7–8). \*: p<0.05; \*\*: p<0.01; \*\*\*: p<0.001.

doi:10.1371/journal.pone.0040595.g004


polybasic cluster, a lysine is found in the rat sequence instead of a glutamic acid in the human and chicken orthologs. This suggests that the first cluster plays a preponderant role in PIP<sub>n</sub> binding, in agreement with the gain-of-binding S365K-E366Y mutation performed on rat P2X5. Since mutations increasing the PIP<sub>n</sub>-binding affinity of P2X5 have a drastic effect on its ion channel function, it is likely that these differences in C-terminal sequence account for the high degree of variability of P2X5 phenotypes observed among vertebrate species.

Identification of the molecular determinants of PIP<sub>n</sub>-protein interactions in the P2X family confirmed the intrinsic and essential nature of PIP<sub>n</sub> regulation of P2X channel activity. Knowing that intracellular PIP<sub>n</sub> levels are controlled by a wide array of ubiquitous pathways such as G<sub>q</sub>-coupled receptor-induced phospholipase C hydrolysis of PIP<sub>2</sub> or receptor tyrosine kinase activation of PI3K, the P2X-PIP<sub>n</sub> regulatory mechanism is likely involved in multi-receptor crosstalks. Our predictive model unifies various data obtained on PIP<sub>n</sub>-regulation of P2X receptors in physiological and pathological contexts and also provides useful insights on PIP<sub>n</sub>-regulation mechanisms of other ion channels.

## Materials and Methods

### Two-electrode Voltage-clamp Recordings in *Xenopus* Oocytes

Oocytes were removed from *Xenopus laevis* frogs as described [8] before intranuclear microinjection of 1 ng plasmid DNA coding for rat P2X4 (WT or K362Q-K363Q), rat P2X5 (WT, S365K-E366Y or S365K-E366Y-E374K) or human P2X5 (kind gift from Dr Alan North). Oocytes were then incubated in Barth's solution containing 1.8 mM CaCl<sub>2</sub> at 19°C for 24 to 72 h before electrophysiological recordings. Two-electrode voltage-clamp recordings (V<sub>hold</sub> = -60 mV for P2X4, -120 mV for P2X5) were performed using glass pipettes (1–3 MΩ) filled with 3 M KCl solution. The external Ringer's solution, pH 7.4, contained (in mM): 115 NaCl, 5 NaOH, 2.5 KCl, 1.8 CaCl<sub>2</sub>, and 10 HEPES. Membrane currents were recorded using a Warner OC-725B amplifier (Warner Instruments) and digitized at 1 kHz. For PIP<sub>n</sub>-depletion experiments, oocytes were incubated in 10 μM wortmannin for 1–2 hours prior to recording. Each series of recordings consisted of three successive applications of ATP (10 or 100 μM), with a 4-minute wash in Ringer's solution between each application.



**Figure 5. P2X C-terminal peptides compete with P2X channels for binding to intracellular PIP<sub>n</sub>.** A) Representative traces of patch-clamp recordings showing that intracellular injection of the P2X4 C-terminal (-CT) peptide leads to a rundown of the P2X4 current in HEK293 cells by competing for intracellular PIP<sub>n</sub>. B) Intracellular injection of the P2X5-CT peptide does not affect the P2X5 current phenotype. C) Effect on P2X4 current amplitude of injection of peptides from the P2X4 WT, P2X4 K362Q-K363Q (2M), P2X5 WT or P2X5 S365K-E366Y-E374K (3M) C-terminus. D) Competition for PIP<sub>n</sub> binding from P2X4-CT or P2X5-3M-CT peptide injection leads to a slower desensitization of the P2X4 current. Values were normalized to the initial recording value obtained immediately after whole-cell configuration was obtained (n=4–5; \*, p<0.05; \*\*, p<0.01; \*\*\*, p<0.001, each group compared to control).

doi:10.1371/journal.pone.0040595.g005



## Whole-cell Patch-clamp Recordings on HEK293 Cells

HEK293 cells (ATCC) were cultured in Dulbecco's modified Eagle's medium and 10% heat-inactivated fetal bovine serum (Invitrogen) containing penicillin and streptomycin supplemented with G-418 (250 µg/ml). The cells were transiently co-transfected with pEGFP and pcDNA3-rP2X4 using Polyfect (Qiagen) according to the manufacturer's instructions. Transfected cells were used for electrophysiological recordings 48 h after transfection. Whole-cell recordings ( $V_{\text{hold}} = -60$  mV) were performed using pipettes filled with internal solution, pH 7.2, containing (in mM): 120 K-gluconate, 1 MgCl<sub>2</sub>, 5 EGTA and 10 HEPES. The recording solution, pH 7.4, comprised (in mM): 140 NaCl, 5 KCl, 2 CaCl<sub>2</sub>, 2 MgCl<sub>2</sub>, 10 HEPES and 10 glucose. GST-fusion peptides (GST control, P2X4-CT, P2X4 K362Q-K363Q-CT, P2X5-CT or P2X5 S365K-E366Y-E374K-CT) were dissolved in the internal pipette solution to a 1 µM concentration. Membrane currents were recorded using an Axopatch 200B amplifier and digitized at 500 Hz with a Digidata 1330 interface (Axon Instruments). Agonists were dissolved in recording solution and applied using a SF-77B fast perfusion system (Warner Instruments) at a rate of 1 ml/min.

## Site-directed Mutagenesis

Point mutations on pcDNA3-rP2X4 (K362Q-K363Q) and pcDNA3-rP2X5 (S365K-E366Y-E374K) were introduced using the QuikChange mutagenesis method (Stratagene).

## Phospholipid-binding Assay

Oligonucleotide-based sequences coding for P2X C-terminal sequences (P2X1: L352-E378, P2X4: C360-V375, P2X5: L361-V376, P2X7: P376-I391, P376-P396, A378-E393) were subcloned into the pGEX-2T vector for the production of GST fusion

proteins. Binding assays were conducted on phospholipid-coated hydrophobic membranes (PIP Strips<sup>TM</sup>, Echelon Biosciences). GST fusion proteins (1 µg/ml) were applied overnight, primary (mouse anti-GST, 1:1000) and secondary (goat anti-mouse HRP, 1:5000) antibodies were applied for one hour. All washes and protein or antibody incubations were done in TBS+T solution supplemented with 3% BSA. Bound proteins were detected by ECL (PerkinElmer).

## Data Analysis

Peak currents, defined as the maximal amplitude recorded during agonist application, were measured. For current rundown, the amplitude of the second response (after a 4-minute wash) was compared to the first and expressed as a percentage. For current kinetics, activation rate was measured as the rise time (in seconds) from 10% to 90% of the peak amplitude. For desensitization rate, the 5-second decay % was used for P2X4 and the decay slope was measured for P2X5. Data are presented as mean ± SEM. Statistical analyses for the difference in means were carried out using Student's *t* test for two unpaired groups, one-way or two-way ANOVA followed by a Bonferroni post-test.

## Acknowledgments

We thank Ariel Ase (Montreal Neurological Institute) for useful discussions during the preparation of this manuscript.

## Author Contributions

Conceived and designed the experiments: LPB EBG PS. Performed the experiments: LPB DB. Analyzed the data: LPB EBG PS. Contributed reagents/materials/analysis tools: PS. Wrote the paper: LPB EBG PS.

## References

- Rohacs T, Thyagarajan B, Lukacs V (2008) Phospholipase C mediated modulation of TRPV1 channels. *Mol Neurobiol* 37: 153–163.
- Nilius B, Mahieu F, Prenen J, Janssens A, Owsianik G, et al. (2006) The Ca<sup>2+</sup>-activated cation channel TRPM4 is regulated by phosphatidylinositol 4,5-bisphosphate. *EMBO J* 25: 467–478.
- Fan Z, Makielski JC (1997) Anionic phospholipids activate ATP-sensitive potassium channels. *J Biol Chem* 272: 5388–5395.
- Haider S, Tarasov AI, Craig TJ, Sansom MS, Ashcroft FM (2007) Identification of the PIP<sub>2</sub>-binding site on Kir6.2 by molecular modelling and functional analysis. *EMBO J* 26: 3749–3759.
- Suh BC, Hille B (2007) Regulation of KCNQ channels by manipulation of phosphoinositides. *J Physiol* 582: 911–916.
- Mo G, Bernier LP, Zhao Q, Chabot-Dore AJ, Ase AR, et al. (2009) Subtype-specific regulation of P2X3 and P2X2/3 receptors by phosphoinositides in peripheral nociceptors. *Mol Pain* 5: 47.
- Bernier LP, Ase AR, Tong X, Hamel E, Blais D, et al. (2008) Direct modulation of P2X1 receptor-channels by the lipid phosphatidylinositol 4,5-bisphosphate. *Mol Pharmacol* 74: 785–792.
- Bernier LP, Ase AR, Chevallier S, Blais D, Zhao Q, et al. (2008) Phosphoinositides regulate P2X4 ATP-gated channels through direct interactions. *J Neurosci* 28: 12938–12945.
- Fujiiwara Y, Kubo Y (2006) Regulation of the desensitization and ion selectivity of ATP-gated P2X2 channels by phosphoinositides. *J Physiol* 576: 135–149.
- Ase AR, Bernier LP, Blais D, Pankratov Y, Seguela P (2010) Modulation of heteromeric P2X1/5 receptors by phosphoinositides in astrocytes depends on the P2X1 subunit. *J Neurochem* 113: 1676–1684.
- Chaumont S, Jiang LH, Penna A, North RA, Rassendren F (2004) Identification of a trafficking motif involved in the stabilization and polarization of P2X receptors. *J Biol Chem* 279: 29628–29638.
- Koshimizu TA, Ueno S, Tanoue A, Yanagihara N, Stojilkovic SS, et al. (2002) Heteromultimerization modulates P2X receptor functions through participating extracellular and C-terminal subdomains. *J Biol Chem* 277: 46891–46899.
- D'Arco M, Giniatullin R, Leone V, Carloni P, Birsa N, et al. (2009) The C-terminal Src inhibitory kinase (Csk)-mediated tyrosine phosphorylation is a novel molecular mechanism to limit P2X3 receptor function in mouse sensory neurons. *J Biol Chem* 284: 21393–21401.
- Toulme E, Blais D, Leger C, Landry M, Garret M, et al. (2007) An intracellular motif of P2X(3) receptors is required for functional cross-talk with GABA(A) receptors in nociceptive DRG neurons. *J Neurochem* 102: 1357–1368.
- Rosenhouse-Dantsker A, Logothetis DE (2007) Molecular characteristics of phosphoinositide binding. *Pflügers Arch* 455: 45–53.
- Suh BC, Hille B (2008) PIP<sub>2</sub> is a necessary cofactor for ion channel function: how and why? *Annu Rev Biophys* 37: 175–195.
- Zhao Q, Yang M, Ting AT, Logothetis DE (2007) PIP<sub>2</sub> regulates the ionic current of P2X receptors and P2X(7) receptor-mediated cell death. *Channels (Austin)* 1: 46–55.
- North RA (2002) Molecular physiology of P2X receptors. *Physiol Rev* 82: 1013–1067.
- Bo X, Jiang LH, Wilson HL, Kim M, Burnstock G, et al. (2003) Pharmacological and biophysical properties of the human P2X5 receptor. *Mol Pharmacol* 63: 1407–1416.
- Kim AY, Tang Z, Liu Q, Patel KN, Maag D, et al. (2008) Pirt, a phosphoinositide-binding protein, functions as a regulatory subunit of TRPV1. *Cell* 133: 475–485.
- Michailidis IE, Helton TD, Petrou VI, Mirshahi T, Ehlers MD, et al. (2007) Phosphatidylinositol-4,5-bisphosphate regulates NMDA receptor activity through alpha-actinin. *J Neurosci* 27: 5523–5532.
- Kim M, Jiang LH, Wilson HL, North RA, Surprenant A (2001) Proteomic and functional evidence for a P2X7 receptor signalling complex. *EMBO J* 20: 6347–6358.
- Rohacs T, Lopes CM, Michailidis I, Logothetis DE (2005) PI(4,5)P<sub>2</sub> regulates the activation and desensitization of TRPM8 channels through the TRP domain. *Nat Neurosci* 8: 626–634.
- Hernandez CC, Zaika O, Shapiro MS (2008) A carboxy-terminal inter-helix linker as the site of phosphatidylinositol 4,5-bisphosphate action on Kv7 (M-type) K<sup>+</sup> channels. *J Gen Physiol* 132: 361–381.
- Logothetis DE, Jin T, Lupyan D, Rosenhouse-Dantsker A (2007) Phosphoinositide-mediated gating of inwardly rectifying K<sup>+</sup> channels. *Pflügers Arch* 455: 83–95.
- Stansfeld PJ, Hopkinson R, Ashcroft FM, Sansom MS (2009) PIP<sub>2</sub>-binding site in Kir channels: definition by multiscale biomolecular simulations. *Biochemistry* 48: 10926–10933.

27. Whorton MR, MacKinnon R (2011) Crystal structure of the mammalian GIRK2 K<sup>+</sup> channel and gating regulation by G proteins, PIP<sub>2</sub>, and sodium. *Cell* 147: 199–208.
28. Hansen SB, Tao X, MacKinnon R (2011) Structural basis of PIP<sub>2</sub> activation of the classical inward rectifier K<sup>+</sup> channel Kir2.2. *Nature* 477: 495–498.
29. Kawate T, Michel JC, Birdsong WT, Gouaux E (2009) Crystal structure of the ATP-gated P2X<sub>4</sub> ion channel in the closed state. *Nature* 460: 592–598.
30. Kracun S, Chaptal V, Abramson J, Khakh BS (2010) Gated access to the pore of a P2X receptor: structural implications for closed-open transitions. *J Biol Chem* 285: 10110–10121.
31. Li M, Kawate T, Silberberg SD, Swartz KJ (2010) Pore-opening mechanism in trimeric P2X receptor channels. *Nat Commun* 1: 44.
32. Collo G, North RA, Kawashima E, Merlo-Pich E, Neidhart S, et al. (1996) Cloning OF P2X<sub>5</sub> and P2X<sub>6</sub> receptors and the distribution and properties of an extended family of ATP-gated ion channels. *J Neurosci* 16: 2495–2507.
33. Ruppelt A, Ma W, Borchardt K, Silberberg SD, Soto F (2001) Genomic structure, developmental distribution and functional properties of the chicken P2X<sub>5</sub> receptor. *J Neurochem* 77: 1256–1265.