

Potential knowledge gain in large-scale simulations of forest carbon fluxes from remotely sensed biomass and height

Valentin Bellassen, Nicolas Delbart, G. Le Maire, S. Luyssaert, Philippe Ciais,

N. Viovy

▶ To cite this version:

Valentin Bellassen, Nicolas Delbart, G. Le Maire, S. Luyssaert, Philippe Ciais, et al.. Potential knowledge gain in large-scale simulations of forest carbon fluxes from remotely sensed biomass and height. Forest Ecology and Management, 2011, 261 (3), pp.515-530. 10.1016/j.foreco.2010.11.002 . hal-01149366

HAL Id: hal-01149366 https://hal.science/hal-01149366

Submitted on 2 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

sensed biomass and height	
Journal: Forest Ecology and Management, Elsevier, 2011, 261 (3), pp.515.	
(10.1016/j.foreco.2010.11.002).	
Authors: Bellassen V ¹ , Delbart N ¹ , Le Maire G ² , Luyssaert S ¹ , Ciais P ¹ , Viovy N ¹	
¹ Laboratoire des Sciences du Climat et de l'Environnement, Commissariat à l'énergie atomiq	lne
/ CEA-Orme des Merisiers / F-91191 Gif-sur-Yvette CEDEX / France	
² CIRAD, Persyst, UPR 80, s/c UMR Eco&Sols, 2 Place Viala - bât 12, 34060 Montpellier cedex	01,
France	

- Corresponding author: Bellassen V
- Phone: +33 1 69 08 31 01
- Fax: +33 1 69 08 30 73
- E-mail: valentin.bellassen@lsce.ipsl.fr

- Title: Potential knowledge gain in large-scale simulations of forest carbon fluxes from remotely
- and biomass and boigh

17 Abstract

18 Global Vegetation Models (GVMs) simulate CO₂, water and energy fluxes at large scales, 19 typically no smaller than 10 x 10 km. GVM simulations are thus expected to simulate the 20 average functioning, but not the local variability. The two main limiting factors in refining this 21 scale are 1) the scale at which the pedo-climatic inputs – temperature, precipitation, soil water 22 reserve, etc. – are available to drive models and 2) the lack of geospatial information on the 23 vegetation type and the age of forest stands. This study assesses how remotely sensed biomass 24 or stand height could help the new generation of GVMs, which explicitly represent forest age 25 structure and management, to better simulate this local variability. For the ORCHIDEE-FM 26 model, we find that a simple assimilation of biomass or height brings down the root mean 27 square error (RMSE) of some simulated carbon fluxes by 30-50%. Current error levels of remote 28 sensing estimates do not impact this improvement for large gross fluxes (e.g. terrestrial 29 ecosystem respiration), but they reduce the improvement of simulated net ecosystem 30 productivity, adding 13.5-21% of RMSE to assimilations using the *in situ* estimates. The data 31 assimilation under study is more effective to improve the simulation of respiration than the 32 simulation of photosynthesis. The assimilation of height or biomass in ORCHIDEE-FM enables 33 the correct retrieval of variables that are more difficult to measure over large areas, such as 34 stand age. A combined assimilation of biomass and net ecosystem productivity could possibly 35 enable the new generation of GVMs to retrieve other variables that are seldom measured, such 36 as soil carbon content.

Keywords: remote sensing; global vegetation model; ORCHIDEE; carbon; biomass; height

39 1 Introduction

40 Along with the growing concern about climate change, international agreements and regional 41 markets have appeared and now give an economical value to carbon. These developments have 42 naturally led to look at forest services from a carbon perspective: what is the carbon budget of 43 a country's forests and how will it evolve? How can forest management increase the carbon 44 balance of a forest stand and how can this mitigation effort be rewarded? To answer these 45 questions, policy makers and foresters have progressively turned to forest models. 46 Stand-scale growth-and-yield models – such as FAGACEES (Dhôte and Hervé, 2000), SILVA 47 (Pretzsch et al., 2002) or CO2FIX (Masera et al., 2003) – or process-based models – such as CASTANEA (Dufrene et al., 2005) or GRAECO (Porté, 1999) – offer trustworthy local simulations 48 49 of carbon stocks. Nevertheless, the former require a local site productivity index and the latter 50 tend to require an intensive local calibration, based on extensive field observations. It renders 51 both types of models very site-specific and unfit to simulate regional fluxes and stocks. Another 52 solution is to use generic global vegetation models such as Ecosytem Demography (Moorcroft 53 et al., 2001), Biome-BGC (Thornton et al., 2002), LPJ (Sitch et al., 2003) or ORCHIDEE (Krinner et 54 al., 2005). Although these GVMs tend to perform worse than well calibrated stand-scale models 55 on a site-by-site basis (Loustau *et al.*, 2005), they provide estimates of carbon stocks and fluxes 56 at regional scales, or where data is lacking for a local calibration. Their estimates of continental 57 carbon budgets are within the range given by other methods (Lindner et al., 2004), but 58 uncertainty remains high: for the carbon balance of European forests (EU25), the estimates of 59 three similar GVMs exhibit a 2-fold difference (Luyssaert *et al.*, 2010).

The bigger uncertainty of GVM estimates has two main causes: first the pedo-climatic inputs
driving the models are too coarse and do not capture local variations, which in turn can
generate errors due to the non-linear response of physiological processes to these inputs.
Second most GVMs do not explicitly simulate management (Le Quere *et al.*, 2009), and
therefore fail to correctly simulate age-dependent variables such as aboveground biomass. This
pitfall is being increasingly addressed (Zaehle *et al.*, 2006; Desai *et al.*, 2007; Sato *et al.*, 2007;
Bellassen *et al.*, 2010a).

67 In even-aged stands, local variations of soil fertility, climate, and stand age can be combined in the notion of "growth stage". Forest yield tables indeed show that most characteristics – stand 68 69 density, height, basal area, diameter, aboveground biomass, ... – of two stands with different 70 fertility classes follow the same evolution with age, albeit not at the same pace (Vannière, 71 1984; JRC, 2009). A 50 year old forest standing on a productive site may be considered to be at 72 the same "growth stage" than a 100 year old forest standing on an unfertile soil, as they have 73 similar biomass and height. Information about the "growth stage" of forest stands could be 74 used to account for sub-grid heterogeneity in GVM simulations and increase the match 75 between simulations and local measurements. Initializing a GVM with a spatially explicit map of 76 growth stage may improve simulations in two ways: 1) at the local level, information on growth 77 stage recreates some intra-pixel variability in simulations which may improve the fit to site data 78 and 2) at the continental level, a map of growth stage provides a spatially more precise 79 initialization than the current regional or national age averages of forest inventories. The

80 initialization of carbon pools has already been shown to play an important role in the ability of
81 GVMs to reproduce local flux data (Carvalhais *et al.*, 2010).

82 Both average stand height and aboveground biomass could be estimated at large scales with 83 active remote sensing techniques such as RADAR and LiDAR. These two variables can therefore 84 be used to initialize growth stage in a model simulation. P-band RADAR has been particularly 85 used for estimating biomass, as the reflected signal is more specific to the woody components 86 of trees, and therefore saturates at higher levels of biomass than other techniques (Le Toan et 87 al., 2008). The LiDAR signal is less specific and most often used to estimate canopy height 88 structure. Many studies however derive biomass estimates from LiDAR measurements, using 89 the allometric relationship between biomass and height (Lim and Treitz, 2004; Lefsky et al., 90 2005b; Stephens et al., 2007; Naesset, 2009). Except for a few data from the GLAS LiDAR 91 satellite (Lefsky et al., 2005a; Boudreau et al., 2008), all P-band RADAR and LiDAR studies relied 92 on airborne campaigns, and were therefore limited to the local scale. As the European Space 93 Agency is currently assessing the need for a P-band RADAR satellite (Le Toan et al., 2008) and a 94 LiDAR satellite (Durrieu, 2010), there is a pressing need to quantify the benefits of remotely 95 sensed biomass or height for the new generation GVMs which simulate different stand growth 96 stages.

97 To this aim the current study compares the standard version of the ORCHIDEE GVM, with
98 steady-state equilibrium forests, to a more recent version, ORCHIDEE-FM, that simulates forest
99 management and the resulting tree height and biomass in a generic – that is even-aged –
100 managed stand. The ability of ORCHIDEE-FM to simulate the growth stages of a forest stand is

101 checked with forest inventory plots. Then, pseudo-RADAR and LiDAR estimates of height and 102 biomass are generated, based on existing in situ measurements from forest inventories and a 103 global flux database. These pseudo-measurements are used to initialize ORCHIDEE-FM, and the 104 improvement brought by this basic data assimilation is quantified for the simulation of volume 105 increment, gross primary production (GPP), total ecosystem respiration (TER) and net 106 ecosystem productivity (NEP). These quantified improvements provide a first assessment of the 107 knowledge that could be gained from remotely sensed "growth stage" maps for large-scale 108 carbon and water flux estimates.

109 2 Material and Methods

110 **2.1** Field data

111 Two in situ datasets are used in this study: the French national forest inventory (IFN) plots in
112 two different regions, and the global flux and biometry database of Luyssaert *et al.* (2007).

113 **2.1.1** French national forest inventory (IFN)

114 The first in situ dataset used in this study comes from the French national forest inventory

115 (IFN). The IFN conducts yearly field measurement campaigns covering the entire French

116 metropolitan territory. A systematic inventory grid is visited and inventoried following the IFN

- 117 protocol (IFN, 2006): circumference at breast height, width of the last five rings, height and
- 118 species are recorded for a representative sample of trees. IFN allometric rules are used to
- 119 estimate commercial volume and volume increment. For even-aged stands, a few tree cores

sampling all growth rings are used to estimate stand age class, with age class widths between
10 and 20 years. Raw data for each plot is available on the IFN website (www.ifn.fr).
When necessary, estimated standing volume is converted to total aboveground biomass, using
the default branch ratio and carbon density parameters of ORCHIDEE-FM (Table 1).
Furthermore, commercial wood increment is used to estimate annual woody Net Primary
Productivity (NPP), using Eq. 1.

126
$$NPP_{woody} = (I + \varepsilon) \times (1 + T_b \times br \times V) \times BEF_i \times d_w \times d_c$$
 (1)

127 where NPP_{woody} is the annual woody NPP in gC m⁻² yr⁻¹, *I* is the estimated commercial wood 128 increment in $m^3 m^{-2} yr^{-1}$, T_b is the annual turnover of branches in yr^{-1} , br is the total branch ratio (no unit), V is the estimated standing commercial wood in $m^3 m^{-2}$, ε is the averaged increment 129 of trees that died over the last five years before measurement in m³ m⁻² yr⁻¹, *BEF_i* is the biomass 130 131 expansion factor for volume increment (no unit), d_w is the wood density in gDM m⁻³ (grams of 132 dry matter), and d_c is the carbon density in gC gDM⁻¹. At most a few percent of trees commonly 133 die over 5 years. They are usually smaller trees and not all die right before measurement date. ε 134 is therefore much smaller than the wood increment of trees which survived, and neglected in 135 the calculations. For parameter values, see Table 1.

For this study, the results of three campaigns – 2005, 2006 and 2007 – are pooled, and only
even-aged stands are used. To assess the performance of ORCHIDEE-FM, we selected regions
that filled the following criteria:

As REMO climate input data have a 0.25° x 0.25° resolution (see section 2.3.2) we
 selected grid points where climate does not vary strongly at the considered spatial scale

- so that a single ORCHIDEE-FM simulation should be representative of neighbouring IFN
 plots
- A second criteria is to have enough IFN plots within a 0.5° radius for statistical
 purposes.
- 145 Two highly forested regions fit these criteria for both broadleaves and needleleaves:
- southwestern and northeastern France. We therefore selected one southwestern gridpoint,
- 147 hereafter referred to as "Landes", and one northeastern gridpoint, hereafter referred to as
- 148 "Vosges", with respectively 215 and 324 IFN plots within a 0.5° radius of the grid point
- 149 centre. The characteristics of these two grid points are summarized in Table 2. The IFN
- dataset thus provides a high number of *in situ* estimates for age, average height, standing
- aboveground biomass, and woody NPP at various growth stages (stand ages vary from 2 to
- 152 200 years). However, it does not provide direct carbon fluxes measurements.
- 153 **2.1.2 Global dataset for carbon fluxes**

154 Luyssaert et al. (2007) compiled a dataset of carbon flux measurements on forest sites, heavily 155 building on the eddy covariance FLUXNET network (Baldocchi et al., 2001). This dataset also 156 gives additional site information, when available, regarding stand age, average height and 157 biomass. All managed sites in temperate and boreal biomes informed for stand age, average 158 height and aboveground biomass, and either GPP, TER or NEP were retained for this study, thus 159 reducing the database to a subsample of 31 sites. When several years of measurements were 160 available for a given variable, the average was retained. The resulting dataset, hereafter 161 referred to as the "global flux dataset", is presented in Appendix A.

162 Three peculiar site configurations are beyond the expected validity domain of ORCHIDEE-FM163 and therefore excluded for the quantification of performance improvement:

- Forest stands younger than 20 years (8 sites): in such young stands, NEP is expected to
- 165 be heavily influenced by the site history before stand establishment (eg. afforested
- 166 farmland or clearcut old stand) for which we have no information. This hypothesis is
- 167 tested by analyzing the difference in simulations with two extreme sets of initial
- 168 conditions: forest regrowth initial conditions corresponding to the clear-cut of a
- 169 mature forest and reforestation initial conditions corresponding to a cropland.
- Carbon sources (2 sites): our dataset only contains growing managed stands younger
- 171 than 103 years old. Such stands are not expected to be net sources of carbon for several
- 172 years in a row, unless management events such as heavy thinnings or clear-cuts take
- 173 place within the footprint of the flux tower. Again, the absence of information on such
- 174 heavy management events makes it hazardous to simulate these sites.
- Collelongo (1 site): GPP at this 103 years old Italian site is twice larger than TER, which
- 176 makes it an outlier in most analysis of the global flux database (Luyssaert *et al.*, 2009). In
- 177 our case, climate is a possible explanation as it exhibit strong local variations –
- 178 uncaptured by the 0.25° resolution forcing data in this part of Italy.
- 179 The resulting dataset of 20 sites is hereafter referred to as the "screened dataset".

180 **2.2** Generation of pseudo remote sensing data

181 The *in situ* estimates of average stand height and biomass from the two datasets are used to

182 construct a set of pseudo remote sensing estimates of these variables. To this end, a random

183 error is added to the *in situ* estimates of height and biomass. This random error is drawn from a 184 normal probability law centred on the in situ estimate and with a standard deviation equal to 185 the typical RMSE of LiDAR and P-band RADAR estimates of average height and biomass. This 186 implies three assumptions: in situ estimates are assumed to be perfect, remote sensing observations are assumed to be unbiased and their error is assumed to be independent of the 187 188 estimate value. The typical RMSE assigned to these pseudo-remote sensing observations are 189 averages of RMSE from relevant literature studies (see Table 3): the typical RMSE of LiDAR is 190 lower than that of RADAR for average height (1.66 vs 2.34 m) and higher for average biomass 191 (23.7 vs 18.5 tC ha⁻¹). The procedure of "pseudo data" generation is repeated 10 000 times – 192 procedure based on the Monte Carlo technique (Rubinstein and Kroese, 1981) - for each in situ 193 estimate in order to generate a representative pseudo-dataset.

194 **2.3 Model**

195 **2.3.1 ORCHIDEE and ORCHIDEE-FM**

The ORCHIDEE global vegetation model ("ORganizing Carbon and Hydrology In Dynamic
Ecosystems") is designed to operate from regional to global scales (Krinner *et al.*, 2005).
ORCHIDEE typically represents an average mature forest at steady-state equilibrium in a "bigleaf" approach. For a given climatology, it simulates the carbon, water and energy budget at the
pixel scale. For carbon, ORCHIDEE computes its assimilation (GPP), allocates photosynthates to
the different biomass compartments where they are respired or stored, and recycles carbon
through constant tree mortality and soil respiration. This "standard" version of ORCHIDEE

203 (V1.9.4) is intended to simulate forests that have reached a steady-state equilibrium between 204 growth and mortality. Its uses the allocation framework of Friedlingstein et al. (1999), and does 205 not simulate the nitrogen cycle, recently included in ORCHIDEE-N by Zaehle and Friend (2010). 206 This version of ORCHIDEE, hereafter referred to as the standard version, is intended to simulate 207 forests that have reached a steady-state equilibrium between growth and mortality. 208 The standard version does not represent important processes driving the evolution of stand 209 structure such as competition, forest management, or the age-limitation of NPP and is 210 therefore not suited to simulate managed forests, or forests recovering from past disturbance. 211 As a consequence of its formulation all carbon pools, including biomass, need to be put to 212 equilibrium before studying the effect of varying climate and CO₂ conditions. This equilibrium is 213 obtained by a "spin-up", that is an initial simulation which stops when carbon and water pools 214 are in equilibrium with a fixed climate which can take up to 10 000 years. 215 In order to simulate forest management, several processes have been added to the standard 216 version of ORCHIDEE, among which a Forest Management Module (FMM) inspired from the

stand-level model FAGACEES (Dhôte and Hervé, 2000). The key concept is to add to the
"average tree" representation of ORCHIDEE an explicit distribution of individual trees, which is
the basis for a process-based simulation of mortality. The aboveground "stand-scale" wood
increment simulated by ORCHIDEE is distributed among individual trees according to the rule of
Deleuze et al. (2004): the basal area of each individual trees grows proportionally to its
circumference. Tree mortality is then determined by the structure of the stand. Mortality due
to natural competition relies on the self-thinning rule of Reineke (1933) while another set of

224 rules drives the mortality processes due to human interventions such as thinnings or clearcuts. 225 Some other small refinements have been added to ORCHIDEE such as a height limitation on leaf 226 area index (LAI) and an age-related decline in photosynthesis efficiency. As a result, this new 227 version of the model, called ORCHIDEE-FM, is able to simulate the carbon budget and detailed 228 stand structure of forests of varying ages (Bellassen et al., 2010b). Its equations are fully 229 described in Bellassen et al. (2010a). For both versions of the model, the standard value of the 230 maximum rate of carboxylation, Vcmax, is set to the optimized values for 6 sites in France and 231 Germany found by Santaren (2006) for needleleaves and broadleaves, that is respectively 42.6 232 and 52.2 in μ mol m⁻² s⁻¹.

233 **2.3.2 Simulations**

234 2.3.2.1 Input data

For simulations at the sites of the global flux dataset, the climate forcing data comes from the 2.5° resolution NCEP – National Centers for Environmental Prediction – reanalysis, adjusted with the 0.5° CRU – Climate Research Unit – data for temperature and precipitation and interpolated to a 0.5° resolution (Kalnay *et al.*, 1996; Mitchell and Jones, 2005). As in most global simulations (Krinner *et al.*, 2005), the soil bucket is uniformly taken to be 2 m deep and its texture is evenly distributed between clay, sand and silt, corresponding to a uniform water holding capacity of 300 mm

242	For the IFN sites, we use the 0.25° resolution REMO reanalysis (Vetter et al., 2008), which
243	covers Europe. Maps of soil depth and texture were derived from FAO and IGBP products
244	(Vetter <i>et al.,</i> 2008).

245 *2.3.2.2 Model "spinup"*

246 Following a standard method in GVM modelling, a model "equilibrium spinup" is performed 247 before all simulations to define the initial conditions of subsequent simulations, in particular for soil carbon. For this "spinup", ORCHIDEE is repeatedly run for the climate of the 10 years 248 249 preceding stand establishment until all ecosystem carbon and water pools, including soil, reach 250 a steady-state equilibrium. For the ORCHIDEE-FM simulations, the runs start with a clear-cut: 251 stems are exported and all the remaining biomass – branches, roots and leaves – goes to the 252 litter pools, except for a small fraction corresponding to the initial stand structure assumed by 253 the model.

254 2.3.2.3 Simulation set up

255 The reference simulation – STD – uses the standard version of ORCHIDEE. This simulation is 256 compared to seven simulations using ORCHIDEE-FM, the age-explicit version of the model. For 257 each grid point, ORCHIDEE-FM is run 15 times, with plantation dates spanning 150 years before 258 the date of observation, in order to capture all the growth stages potentially existing on the grid 259 point. Out of these 15 runs, seven simulations are then selected for comparison with the 260 reference simulations: FMga, FMgb and FMgh, the simulations with respectively the closest age, 261 biomass and height to the actual *in situ* estimate. FM_{lb} and FM_{lh} are the simulations with the 262 respectively closest biomass and height to the pseudo-lidar measurement. FM_{rb} and FM_{rh} are

the simulations with respectively closest biomass and height to pseudo-radar measurement.
This selection is in fact a basic synthetic data assimilation procedure (Piao *et al.*, 2009) assuming
a perfect model and random errors only: the starting date of ORCHIDEE-FM simulations is
chosen among different realizations in order to minimize the absolute difference between a
simulated variable – age, biomass, or height – and its measured counterpart. The simulation
procedure is summarized in



- 270 Figure 1 and simulations names are summarized in Appendix E, together with other
- abbreviations.

272 **2.4** Model assessment and quantification of simulation improvement

273

275

2.4.1 Assessment of stand growth simulation

274 The expected improvement of ORCHIDEE-FM simulations relies on the assumed ability of the

model to correctly simulate the evolution of forest stand variables such as height and biomass

as a function of age. To check this ability, two tests are performed: first, the simulated

277 evolution of woody NPP with age is compared to the IFN plots estimates. Second, since stand

age, biomass and height are correlated with each other, ORCHIDEE-FM should be able to

279 retrieve stand age when biomass or height is assimilated. For simulations *FM*_{gb} and *FM*_{gh} where

stand age is adjusted in the simulations to minimize the error on respectively biomass and

height, the inferred stand age can be cross-validated by the *in situ* data on stand age.

282 **2.4.2** Quantification of simulation improvement

283 2.4.2.1 Quantified evaluation criteria

To quantify the improvement in simulations resulting from the use of ORCHIDEE-FM in conjunction with external estimates of biomass or height, we focus on four carbon flux variables available in the datasets: GPP, TER, NEP, and woody NPP, this last being the indirect result of the allocation of NPP between the different organs of trees. While thirteen model evaluation criteria are computed (Appendix B) for each simulation according to the recommendations of Willmott (1982), we especially focus our analysis on two of them: the root mean square error (RMSE) as an indicator of average simulation error, and the slope of the 291 linear regression between simulated and measured values (*a*), as an indicator of the model's292 ability to reproduce a trend observed in the data.

293 While it is technically feasible to assimilate biomass in the standard version, as in the CASA 294 steady-state equilibrium model (Carvalhais *et al.*, 2010), the constant mortality assumed by 295 ORCHIDEE, and the absence of wood removals, lead to a very fast equilibrium between NPP and 296 TER, and therefore a limited ability to take advantage of biomass data. We therefore chose to 297 discuss directly the relative merit of ORCHIDEE-FM with biomass or height assimilation against 298 the standard version without data assimilation.

299 2.4.2.2 Example of application: maps of NEP

300 Data-derived maps of height and volume in French forests can be obtained by a smooth 301 interpolation of IFN data (Bellassen et al., 2010b): each grid cell is attributed the averaged 302 height and standing volume over IFN plots within a 50 km radius from its centre. In order to 303 illustrate the potential of remote sensing data assimilation in ORCHIDEE-FM, three maps of 304 average simulated NEP in the 1990s are then produced for France. One represents the NEP 305 simulated by ORCHIDEE-FM without prior knowledge of the growth stage of French forests: all 306 French forests are assumed to belong to the 40-50 years age class. The two others - one 307 assimilating height and the other volume – show how the initial map can be refined with 308 information on growth stage. From 15 ORCHIDEE-FM simulations over France representing 309 stands aged between 0 and 150 years, each grid cell is attributed the NEP of the simulation with 310 closest height or biomass to the corresponding data-derived map.

311 **3 Results**

316

312 **3.1** Simulation of age-related trends by ORCHIDEE-FM

313 Whereas the woody NPP simulated by the standard version of ORCHIDEE is by definition

314 insensitive to age, ORCHIDEE-FM reproduces the observed downward trend in woody NPP for

315 both locations and both functional types



Figure 2). The age-related decline is however steeper in IFN estimates than in simulations. In particular, the simulated decline of woody NPP is several times smaller than the standard deviation of observations for any given age class. In terms of absolute values, measured and

simulated woody NPP are comparable for broadleaves, but tend to be overestimated by themodel for needleleaves.

322 **3.2** Age retrieval from assimilation of height or biomass data

326

The *in situ* estimate of stand age is correctly retrieved by assimilating biomass – FM_{gb} simulation – or height – FM_{gh} simulation –for both PFTs: the shape of the frequency distribution of age differences between simulations and observations is close to a zero-centered Gaussian



327 Figure 3). The spread of the distribution is larger toward positive differences, leading to a

- 328 positive mean age difference of 5 to 34 years between simulations and observations, for ages
- 329 measured between 10 and 250 years. This bias is comparable to the precision of the
- 330 measurement (the width of measured age classes varies between 10 and 20 years). The
- 331 narrower shape of height histograms indicates than assimilating average height in ORCHIDEE-
- 332 FM is slightly more discriminating than assimilating biomass.

333 **3.3** No quantified improvement in simulated woody NPP



334 The ability of ORCHIDEE-FM to simulate an age-related decline in woody NPP



Figure 2) does not translate in a quantified improvement in the fit to IFN-derived estimates



close to 0 (-0.03 – 0.6). The values of the thirteen Wilmott performance indexes are listed in
Appendix C.

345 **3.4** Improvement in simulated GPP, TER and NEP

346 **3.4.1** Simulation of carbon sources and sinks

- 347 Despite the expected presence of outliers in the global flux dataset, ORCHIDEE-FM FM_{ga}
- 348 simulation is better able than ORCHIDEE to reproduce the observed cross-sites gradient of



Measured NEP (gC/m2/yr)

349 NEP across sites (

351 measured at the very young site of Vancouver Island.

³⁵⁰ Figure 5). Interestingly, it is able to simulate the huge carbon source of -606 gC m⁻² yr⁻¹

352 The use of our "screened dataset" is justified by the exponential decrease over time of the

difference in simulated TER between reforested cropland and regrowing clearcut forests



354

353

Figure 6). This difference can reach 1 000 gC m⁻² yr⁻¹ for younger stands but is below 300 gC m⁻² yr⁻¹ for all the sites of the screened dataset, older than 20 years. These figures compare to a typical simulated interannual variability of 300 gC m⁻² yr⁻¹.

358 **3.4.2** Quantification of improvement in simulated carbon fluxes

The standard version of ORCHIDEE correctly reproduces the spatial gradient of GPP across the screened dataset, despite a systematic positive bias, but not that of TER (Figure 7). Due to its representation of stand growth, ORCHIDEE-FM improves the simulation of TER, despite a

362	systematic positive bias, and consequently the simulation of NEP. This improvement is
363	quantified in Figure 8: the RMSE is improved (i.e. reduced) by 40-50% for TER, and by 20-40%
364	for NEP. More importantly for NEP, ORCHIDEE-FM is able to reproduce the observed variability,
365	with slopes of data-simulations linear regressions between 0.8 and 1. This is a clear
366	improvement from the NEP simulated by standard version of ORCHIDEE, which correlates very
367	poorly with observations.
368	Whereas large gross fluxes (GPP, TER) are unaffected by an additional error on pseudo remote
369	sensing data (FM_{rb} , FM_{lb} , FM_{rh} and FM_{lh} simulations), the simulation of NEP deteriorates with
370	decreasing precision of the biomass data assimilated: the assimilation of <i>in situ</i> biomass (FM_{gb})
371	improves RMSE by 37% against only 23% for pseudo-lidar biomass (<i>FM</i> _{lb}), and 28% for pseudo-
372	radar biomass (<i>FM_{rb}</i>). The slope of the linear regression is also worsened from 1.01 to
373	respectively 0.81 and 0.86.
374	Assimilating height or biomass leads to broadly similar improvements in flux simulations.
375	Biomass assimilation nevertheless seems most beneficial: while height assimilation further
376	reduces RMSE for GPP and TER, this reduction comes at the price of a degraded linear
377	regression. For NEP, biomass assimilation brings the strongest improvement in both RMSE and
378	slope of linear regression.
379	3.4.3 Application: maps of NEP

Our simple assimilation procedure – using data-derived maps of height or biomass to select an
 age-class in ORCHIDEE-FM simulations – produces less uniform and likely more realistic maps of
 NEP for France (Figure 9). In particular, the older forests of central and north-eastern France are

detected from their higher height and biomass, leading to a lower simulated NEP than thesimulation without information on growth stage (Figure 9a).

385 **4 Discussion**

386 4.1 Flux simulation improvement delivered by height/biomass data

387 Our results show that a simple height or biomass data assimilation already yields a 30-50% 388 decrease in RMSE for both TER and NEP. The tested range of measurement errors (representing 389 Lidar or Radar errors) does not affect the improvement in large gross fluxes (GPP and TER). 390 However, the precision of NEP simulation is impacted: a 18.5 tC ha⁻¹ uncertainty on the 391 assimilated biomass data, typical of P-band radar measurements, increases the simulation error 392 by 20 gC m² yr⁻¹ (7.5% of observed average NEP), and a 23.6 tC ha⁻¹ uncertainty, typical of LiDAR 393 derived estimates, increases the simulation error by 31 gC m² yr⁻¹ (11.5% of observed average 394 NEP). Thus, a reduced error on remote sensing estimates of biomass would not be useful for 395 the simulation of large fluxes, but could further reduce the error on simulated NEP by 20-31 gC 396 m² yr⁻¹ down to 149 gC m² yr⁻¹, that is the error obtained by assimilating the in situ estimate.

397 These results are both site- and model-dependent. For instance, the strong influence of land-

use history on the NEP of younger stands limits the benefits of the assimilation procedure



399

398

Figure 6). More elaborate distinctions in the usefulness of the method could possibly be made
based on PFT or climate, but the small size of the screened dataset does not allow to draw
robust conclusions.

The error reduction in simulated fluxes could also be further improved by improvements in the structure and in the parameterisation of ORCHIDEE-FM. In particular, our results point to systematic positive biases in the simulation of GPP and TER which offset one another in the simulation of NEP. Zaehle *et al.* (2010) demonstrated that modelling the nitrogen cycle 408 unable to simulate the large positive NEP typical of growing forest sites. In the near future,

eliminates this systematic bias. This new version of ORCHIDEE, ORCHIDEE-N, remains however

409 when ORCHIDEE-FM is merged with ORCHIDEE-N, we can expect to reduce the bias in GPP and

410 TER, and further decrease the error on NEP by assimilating age, height or biomass.

407

411 Our assumption that remote sensing measurement error is independent on measured value 412 and normally distributed is not realistic. Remote sensing measurements of biomass for example 413 are known to carry a larger error for larger biomass values due to signal saturation (Le Toan et 414 al., 2008). We nevertheless opted for this simplistic approach of error modelling due to the lack 415 of quantification of these error patterns. An alternative to improving error modelling would be 416 to use real remote sensing data where it coincides with *in situ* measurements of carbon fluxes, 417 but this additional requirement would further reduce an already small dataset. 418 The simple data assimilation framework tested in this study could easily be applied at 419 continental scale, provided reliable biomass or height estimates are available. It would be most 420 meaningful in temperate and boreal regions, where carbon fluxes are most impacted by 421 management and age effects (Grant et al., 2007; Magnani et al., 2007; Nunery and Keeton, 422 2010): most temperate forests are thinned or harvested, and large fires are the dominant 423 source of disturbance in boreal forests. 424 These large-scale applications are very promising: TER is the component of NEP for which

425 existing GVM estimates are most uncertain (Mitchell *et al.*, 2009), and it is also the flux for

426 which simulation improvement is largest in our framework. This large improvement in TER

427 comes from the process-based mortality and wood removals simulated by ORCHIDEE-FM. For a

test broadleaf site in northeastern France, Bellassen *et al.* (2010a) showed that the delay in
litterfall and the wood removals were responsible for 60% and 40% of the sink over one forest
rotation respectively. The combination of these two processes allows the simulation of a
realistic disequilibrium in biomass and soil carbon.

432 **4.2** Scale issues in spatial heterogeneity

433 The ability of ORCHIDEE-FM to simulate correctly the across-site gradient in "growth stages" of 434 the global flux dataset does not yield a quantitative improvement in the simulation of woody 435 NPP for the IFN dataset. This difference points out the limits in the notion of "growth stage". 436 While it applies relatively well to some characteristics such as NEP (these characteristics 437 evolving similarly in all stands, albeit not at the same pace), other variables such as woody NPP 438 cannot be easily explained in this light only. Forests standing on poor soils for example will 439 never have strong woody NPP, no matter how long one waits. As illustrated in Figure 10, the 440 large variability in woody NPP for stands of similar ages highlights the importance of other 441 factors than growth stage: the growth stage-dependent allocation and photosynthesis 442 efficiency simulated in ORCHIDEE-FM cannot be expected to reproduce such a wide range of 443 observed values.



444 The same phenomenon explains that the spread of age deviation distributions presented in

445

452

446 Figure 3 is larger on the positive (younger ages) than on the negative (older ages) side. The 447 assimilation framework can go very far towards younger simulated ages in order to match a 448 small observed biomass, but the reverse is not true for high observed biomass: the old stands 449 of average productivity simulated by ORCHIDEE-FM cannot reach the biomass of some 450 observed high productive stands which are only slightly younger. This large residual heterogeneity is most likely explained by species or local fertility, which 451

could also be integrated in ORCHIDEE-FM. Model parameters in particular could be adapted to

distinguish between fast-growing and slow-growing species. The main limitation for these
developments is the ability to get proxies for these factors on a large scale. Fine resolution
maps of species or site productivity are even more difficult to obtain than equivalent maps of
height or biomass, although they may not be completely unmanageable (Nabuurs *et al.*, 2008).

457 4.3 Biomass vs. height

458 The application of our method on the global flux dataset points to biomass as a more suitable 459 candidate than average height for assimilation in ORCHIDEE-FM. The smaller RMSE obtained for 460 GPP and TER in the FM_{ah}, FM_{lh}, and FM_{rh} simulations are indeed misleading. They result from an 461 overestimate of stand age which activates age-related decline processes in ORCHIDEE-FM: 462 while the previously discussed positive biases in GPP and TER are consequently reduced, they 463 are probably not reduced for the correct reason since the RMSE in NEP is higher than for the 464 FM_{qb} simulation. As shown by Figure 11, the assimilation of height in ORCHIDEE-FM indeed 465 leads to an overestimate of stand age for other plant functional types than temperate 466 summergreen broadleaves and temperate evergreen needleleaves. This overestimate probably 467 comes from the height-circumference allometry and the self-thinning relationship of 468 ORCHIDEE-FM, which have only been tested rigorously for temperate summergreen 469 broadleaves and temperate evergreen needleaves.

The IFN dataset, which is restricted to these two plant functional types, provides a different
picture. The narrower and more centered distributions of age deviation for *FM_{gh}* simulations



473 Figure 3) point to height as a more useful variable for assimilation in ORCHIDEE-FM. Height is

474 indeed expected to be less sensitive than biomass to varying intensities of management.

475 These contradicting results make it difficult to draw a general and definitive conclusion on the

476 relative merits of height vs. biomass assimilation in ORCHIDEE-FM. While height seems

477 theoretically more promising, the allometric and self-thinning rules of ORCHIDEE-FM may not

478 be currently generic enough to make the best use of it.

479 **4.4 Retrieval of unmeasured variables**

480 Some variables, such as stand age or soil carbon content, are difficult to measure and therefore 481 seldom available at a fine resolution over large areas. Ours results show that ORCHIDEE-FM is 482 able to use biomass or height data, in addition to pedo-climatic conditions, to correctly retrieve 483 stand age. If such data were available over large areas, this simple assimilation method could 484 therefore produce a new set of stand age maps. The method is independent of the combination 485 of inventory data and remote sensing of disturbances used by Pan et al. (2010) to produce age 486 maps over north America, and the disagreements between the two would undoubtedly provide 487 useful insights on the strengths and weaknesses of both methods. 488 Estimates of soil carbon content could also be retrieved from ORCHIDEE-FM by assimilating

simultaneously biomass and NEP. Assuming that the model simulates correctly NPP and

490 litterfall once it has been initialised for biomass, the resulting discrepancy between measured

and simulated NEP would be due to a faulty soil carbon content, which could then be corrected

492 in the model to match the NEP measurements.

493 **5 Conclusion**

Large-scale information on biomass derived from remote sensing estimates would provide valuable constraints for the simulation of carbon fluxes in ORCHIDEE-FM: the RMSE of simulated NEP is decreased by up to 30% for a global flux dataset. Most importantly, this improvement results from the ability of ORCHIDEE-FM, initialized with the correct "growth 498 stage", to reproduce spatial gradients in NEP, an ability that is lacking in the standard steady-499 state equilibrium version of ORCHIDEE.

500 At a smaller spatial scale, where climate conditions are comparable, remotely sensed

501 information on "growth stage" does not bring a useful constraint on woody NPP, probably

502 because of the relatively higher importance of local factors such as soil fertility and species mix.

503 The notion of "growth stage" may also be less relevant for woody NPP than for NEP.

504 Nevertheless, our simple assimilation framework for height or biomass correctly retrieves stand

505 age, despite a large standard deviation.

506 The simulated error of pseudo remote sensing estimates of biomass or height does not impact

507 the improvement of large gross fluxes (GPP and TER). For simulated NEP however, this

additional source of uncertainty increases the total error by 13.5% and 21% for P-band radar

509 and lidar respectively.

510 Finally, while the results of our simple assimilation framework are promising, they represent

511 only a first assessment of the potential of large-scale data assimilation in DGVMs. New

512 developments in ORCHIDEE (Zaehle and Friend, 2010), and more refined assimilation

513 frameworks, including other remotely sensed variables such as leaf area index (Demarty et al.,

514 2007) or CO₂ concentration (Sarrat *et al.*, 2009), will doubtlessly optimize the use that

515 ORCHIDEE-FM can make of remote sensing estimates of biomass and height.

516 Acknowledgements

517 We want to acknowledge the contribution of Antoine Colin (IFN), without whom the work on

518 the dataset he manages would have been both impossible and meaningless. We also

- 519 appreciated the expert comments of Thuy Le Toan (CESBIO) and Patrick Chazette (LSCE) on
- 520 remote sensing estimates.
- 521 This work was made possible thanks to a research grant from the French ministry for research.

523 Appendixes

524 Appendix A. Subsample of the global carbon flux dataset (Luyssaert *et al.*, 2007) used in this

525 study ("global flux dataset")

ID	Site name	Latitude	Longitude	PFT^1	ASH ²	Biomass	GPP	RECO	NEP	YE ³	$\rm ME^4$	Age
					(m)	(gC m-2)	(gC m ⁻² yr ⁻¹)	(gC m ⁻² yr ⁻¹)	(gC m ⁻² yr ⁻¹)			(years)
1003	Collelongo	41.75 °N	13.75 °E	TeBS	19.0	9 860	1 127	591	583	1895	1998	103
1004	Prince_Albert_SSA_(SOAS)	53.75 °N	106.25 °W	BoBS	20.3	1 997	1 215	1 030	178	1928	1998	70
1006	Prince_Albert_SSA_(SOJP)	53.75 °N	104.75 °W	BoNE	13.0	1 997	690	638	35	1931	1999	68
1007	Thompson_NSA_(NYJP)	55.75 °N	98.25 °W	BoNE	5.0	3 090	960	550	410	1972	1994	22
1012	Bayreuth/Weiden_Brunnen	50.25 °N	11.75 °E	TeNE	36.0	9 267	1 303	1 334	-32	1954	1998	44
1014	Slash_pine_Florida_Mid	29.75 °N	82.25 °W	TeNE	10.0	2 285	2 762	2 087	589	1987	1999	12
1015	Slash_pine_Florida_old	29.75 °N	82.25 °W	TeNE	19.0	8 271	2 606	1 944	675	1973	1998	25
1089	Duke_Forest	35.75 °N	79.25 °W	TeNE	14.0	5 128	1 788	1 233	497	1982	2000	18
1092	Harvard	42.75 °N	72.25 °W	TeBS	25.0	9 900	1 287	1 058	202	1936	1997	61
1093	Walker_Branch	35.75 °N	84.25 °W	TeBS	25.3	5 715	1 690	1 335	514	1930	1997	67
1095	Flakaliden_C	64.25 °N	19.25 °E	BoNE	4.7	1 770	1 000	932	104	1960	2000	40
1096	Norunda	60.25 °N	17.25 °E	BoNE	28.0	11 135	1 312	1 404	-61	1900	1999	99
1097	Hyytiala	61.75 °N	24.25 °E	BoNE	15.5	5 900	1 012	782	233	1964	2000	36
1101	Willow_Creek	45.25 °N	90.25 °W	BoBS	24.0	7 490	1 165	835	289	1933	1999	66
1106	Morgan_Monroe	39.25 °N	86.25 °W	TeBS	26.5	8 720	1 452	1 163	279	1924	2000	76
1109	Le_Bray	44.75 °N	0.75 °W	TeNE	19.0	7 008	1 833	1 451	407	1969	2000	31
1110	Balmoral	42.75 °S	172.75 °E	TeNE	8.0	3 700	1 774	1 166	608	1987	1996	9
1154	Oak_ridge_liriodendron	35.75 °N	84.25 °W	TeBS	30.0	6 288	na	na	249	1918	2000	82
1168	Skyttorp2	60.25 °N	17.75 °E	BoNE	16.0	6 063	1 232	953	360	1970	2004	34
1169	Puechabon	43.75 °N	3.75 °E	TeBE	6.0	5 424	1 379	1071	309	1942	2002	60
1170	Dooary	52.75 °N	7.25 °W	TeNE	8.0	6 162	2 001	1 141	860	1989	2004	15
1178	Takayama	36.25 °N	137.25 °E	TeBS	20.0	13 488	1 050	833	217	1962	1997	35
1185	Hyytiala_12	61.75 °N	24.25 °E	BoNE	4.0	250	854	752	102	1991	2002	11
1246	Hyytiala_75	61.75 °N	24.25 °E	BoNE	25.0	6 700	918	566	352	1927	2001	74
1328	Skyttorp3	60.25 °N	17.75 °E	BoNE	18.0	7 310	na	na	370	1938	2002	64
1364	Espirra	38.75 °N	8.75 °W	TeBE	20.0	4 212	1 495	876	619	1991	2004	13
1378	Bartlett	44.25 °N	71.25 °W	TeBS	19.0	10 730	1 053	790	263	1925	2005	80
1482	Chibougamau_EOBS	49.75 °N	74.25 °W	BoNE	14.0	4 500	584	580	4	1909	2004	95
1507	Vancouver_Island_HDF00	49.75 °N	125.25 °W	TeNE	1.0	2 775	435	1041	-606	1999	2002	3
1508	Vancouver_Island_HDF88	49.75 °N	124.75 °W	TeNE	8.0	5 700	1 214	1 347	-133	1988	2002	14
1509	Vancouver_Island_DF49	49.75 °N	125.25 °W	TeNE	29.0	10 550	1 991	1 737	337	1949	2001	52

1. TeNE: temperate needleleaf evergreen, TeBE: temperate broadleaf evergreen, TeBS: temperate broadleaf summergreen, BoNE: boreal needleleaf evergreen, BoBS: boreal broadleaf summergreen

526 2. Average stand height / 2. Year of establishment / 3. Measurement year

Index	Name	Equation
OA	Average value of observations	$\frac{1}{N}\sum_{i}O_{i}$
SA	Average value of simulations	$\frac{1}{N}\sum_{i}S_{i}$
OS	Standard deviation of observations	$\frac{1}{N-1}\sum_{i}\left(O_{i}-OA\right)$
SS	Standard deviation of simulations	$\frac{1}{N-1}\sum_{i} (S_{i} - SA)$
N	Number of observations	
а	Slope of linear regression (simulations = f(observations))	
b	Intercept of linear regression (simulations = f(observations))	
MAE	Mean absolute error	$\frac{1}{N} \sum_{i} \left S_{i} - O_{i} \right $
RMSE	Root mean square error	$\sqrt{\frac{1}{N}\sum_{i}(S_{i}-O_{i})^{2}}$
RMSEs	Systematic root mean square error	$\sqrt{\frac{1}{N}}\sum_{i} (O_{i} - P_{i})^{2}$
RMSEu	Unsystematic root mean square error	$\sqrt{\frac{1}{N}\sum_{i}(S_{i}-P_{i})^{2}}$
d	Index of agreement	$1 - \frac{\sum_{i} (S_{i} - O_{i})^{2}}{\sum_{i} (S_{i} - OA_{i} + O_{i} - OA_{i})^{2}}$
r²	Square of Pearson's correlation coefficient	$\frac{\sum_{i} (S_{i} - SA)(O_{i} - OA)}{\sqrt{\sum_{i} (S_{i} - SA)^{2}} \times \sqrt{\sum_{i} (O_{i} - OA)^{2}}}$

528 Appendix B. Wilmott performance indexes (Willmott, 1982)

 $S_{1..N}$ are the N simulate values, $O_{1..N}$ are the N measured values, and $P_{1..N}$ are the values predicted by the linear regression: $\forall i, P_i = aO_i + b$

- 530 Appendix C. Wilmott performance indexes for the simulation of volume increment (IFN dataset)
- 531 Abbreviations for the names of performance indexes are given in Appendix B. For details on
- 532 simulations names, see part 2.3.2, Appendix E and





Woody NPP	- broadleaves	(gC m ⁻² yr ⁻¹)											
Simulation	obs average	sim average	obs sd	sim sd	N	а	b	MAE	RMSE	RMSEs	RMSEu	d	r2
STD	426.7	520.7	241.9	55.5	211	0.00	518.81	214.0	263.8	258.0	55.4	0.36	0.00
FM _{ga}	468.9	523.4	280.5	41.3	86	-0.02	533.51	234.1	292.9	290.1	40.6	0.20	0.02
FM _{gb}	426.7	534.1	241.9	38.6	211	-0.02	543.00	221.5	271.4	268.7	38.2	0.35	0.02
FM _{gh}	426.7	530.2	241.9	37.6	211	-0.01	535.58	219.3	268.0	265.4	37.4	0.34	0.01
FM _{Ib}	426.7	536.2	241.9	37.5	211	-0.02	545.14	222.9	272.1	269.6	37.1	0.35	0.02
FM _{rb}	426.7	533.8	241.9	39.0	211	-0.03	545.63	223.8	272.8	270.1	38.3	0.34	0.03
FM _{Ih}	426.7	530.1	241.9	38.5	211	-0.01	536.03	219.3	268.3	265.6	38.3	0.34	0.01
FM _{rh}	426.7	531.9	241.9	38.2	211	-0.02	538.57	220.2	269.4	266.7	37.9	0.34	0.01

Woody NPP	- needleleave	s (gC m ⁻² yr ⁻¹)											
Simulation	obs average	sim average	obs sd	sim sd	Ν	а	b	MAE	RMSE	RMSEs	RMSEu	d	r2
STD	481.5	650.9	219.1	71.0	328	0.05	624.71	228.4	276.4	267.4	69.9	0.48	0.03
FM _{ga}	483.1	617.1	193.8	70.4	172	0.05	592.82	189.1	237.7	227.3	69.5	0.46	0.02
FM _{gb}	481.5	648.2	219.1	72.8	328	0.06	620.94	225.2	274.8	265.3	71.6	0.48	0.03
FM _{gh}	481.5	640.5	219.1	70.8	328	0.06	613.97	220.6	269.9	260.8	69.7	0.48	0.03
FM _{Ib}	481.5	651.6	219.1	71.6	328	0.05	627.34	227.3	277.6	268.5	70.7	0.48	0.02
FM _{rb}	481.5	649.4	219.1	72.7	328	0.06	621.86	225.0	275.4	265.9	71.5	0.48	0.03
FM _{Ih}	481.5	642.1	219.1	71.1	328	0.05	616.99	222.3	271.5	262.3	70.0	0.47	0.03
FM _{rh}	481.5	641.8	219.1	70.6	328	0.06	615.26	220.7	270.6	261.6	69.4	0.48	0.03

- 538 Appendix D. Wilmott performance indexes for the simulation of GPP, TER and NPP (screened
- 539 global flux dataset)
- 540 Abbreviations for the names of performance indexes are given in Appendix B. For details on
- 541 simulations names, see part 2.3.2, Appendix E and



543 Figure 1.

GPP (gC m ⁻² yr ⁻¹)	OA	SA	OS	SS	Ν	а	b	MAE	RMSE	RMSEs	RMSEu	d	r2
STD	1296	1579	502	582	18	1.12	129	286	325	289	149	0.91	0.93
FM _{ga}	1296	1562	502	593	18	1.14	87	269	315	274	154	0.92	0.93
FM _{gb}	1296	1562	502	601	18	1.15	71	279	319	276	160	0.92	0.92
FM _{gh}	1296	1472	502	613	18	1.15	-22	219	274	190	197	0.94	0.89
FM _{lb}	1296	1544	502	596	18	1.13	73	269	310	258	167	0.92	0.91
FM _{rb}	1296	1552	502	597	18	1.14	75	273	313	265	164	0.92	0.92
FM _{lh}	1296	1475	502	619	18	1.17	-43	219	272	198	185	0.94	0.90
FM _{rh}	1296	1475	502	620	18	1.17	-45	220	272	198	185	0.94	0.90
TER (gC m ⁻² yr ⁻¹)	OA	SA	OS	SS	Ν	а	b	MAE	RMSE	RMSEs	RMSEu	d	r2
STD	1032	1493	403	569	18	1.31	142	462	520	477	207	0.76	0.86
FM _{ga}	1032	1288	403	425	18	0.96	302	263	311	257	176	0.86	0.82
FM _{gb}	1032	1324	403	423	18	0.96	333	292	336	292	166	0.85	0.84
FM _{gh}	1032	1269	403	412	18	0.94	296	250	285	239	156	0.88	0.85
FM _{lb}	1032	1331	403	443	18	0.99	310	301	354	301	187	0.84	0.81
FM _{rb}	1032	1330	403	436	18	0.98	321	300	350	299	183	0.84	0.82
FM _{lh}	1032	1262	403	416	18	0.95	278	241	278	231	155	0.89	0.85
FM _{rh}	1032	1261	403	416	18	0.95	277	241	278	230	155	0.89	0.85
NEP (gC m ⁻² yr ⁻¹)	OA	SA	OS	SS	Ν	а	b	MAE	RMSE	RMSEs	RMSEu	d	r2
STD	269	94	152	70	20	0.12	62	185	229	219	66	0.46	0.06
FM _{ga}	269	282	152	176	20	0.79	69	97	129	34	125	0.82	0.47
FM _{gb}	269	250	152	214	20	0.99	-17	110	149	19	148	0.81	0.50
FM _{gh}	269	213	152	255	20	0.98	-49	155	210	56	202	0.70	0.34
FM _{lb}	269	226	152	215	20	0.79	15	134	180	59	163	0.74	0.40
FM _{rb}	269	235	152	212	20	0.84	10	127	169	47	158	0.76	0.42
FM _{lh}	269	223	152	244	20	0.96	-34	146	194	47	189	0.73	0.37
FM _{rh}	269	223	152	244	20	0.96	-34	146	194	47	188	0.73	0.37

- 547 Appendix E. Abbreviations
- 548 **5.1.1.1.1.1 General terms**
- 549 FMM: Forest Management Module
- 550 GPP: Gross Primary Productivity
- 551 GVM: Global Vegetation Model
- 552 IFN: French National Forest Inventory
- 553 NEP: Net Ecosystem Productivity (a positive value indicates a carbon sink)
- 554 NPP: Net Primary Productivity
- 555 PFT: Plant Functional Type
- 556 RMSE: Root Mean Square Error
- 557 TER: Terrestrial Ecosystem Respiration
- 558 5.1.1.1.2 Simulations names
- 559 STD: Simulation using the standard version of ORCHIDEE, representing a forest stand at steady-
- 560 state equilibrium.
- 561 *FM*_{ga}: Simulation with closest age to the *in situ* estimate, selected from a set of ORCHIDEE-FM
- simulations separated by 10-years intervals.
- 563 FM_{gb}: Simulation with closest biomass to the in situ estimate, selected from a set of ORCHIDEE-
- 564 FM simulations separated by 10-years intervals.
- 565 *FM*_{gh}: Simulation with closest average stand height to the *in situ* estimate, selected from a set
- 566 of ORCHIDEE-FM simulations separated by 10-years intervals.

- 567 FM_{lb}: Simulation with closest biomass to the pseudo-lidar estimate, selected from a set of
- 568 ORCHIDEE-FM simulations separated by 10-years intervals.
- 569 FM_{rb}: Simulation with closest biomass to the pseudo-radar estimate, selected from a set of
- 570 ORCHIDEE-FM simulations separated by 10-years intervals.
- 571 *FM*_{lh}: Simulation with closest average stand height to the pseudo-lidar estimate, selected from
- a set of ORCHIDEE-FM simulations separated by 10-years intervals.
- 573 FM_{rh}: Simulation with closest average stand height to the pseudo-radar estimate, selected from
- a set of ORCHIDEE-FM simulations separated by 10-years intervals.

Tables

Parameter	Conifers	Broadleaves	Unit	Source
BEFi	1.1	1.2	no unit	(IPCC, 2003)
T _b	0.025	0.025	yr-1	(Bellassen <i>et al.,</i> 2010a)
br	0.25	0.38	no unit	(Bellassen <i>et al.,</i> 2010a)
d _c	0.5	0.5	gC gDM ⁻¹	(Bellassen <i>et al.,</i> 2010a)
d _w	0.4*10 ⁶	0.6*10 ⁶	gDM m ⁻³	(Bellassen <i>et al.,</i> 2010a)

578 Table 1. Parameter values

Name	Landes	Vosges
Latitude	43.875°N	48.125°N
Longitude	0.875°W	6.875°E
2001-2005 average temperature (°C) 2001-2005 average rainfall	14.3	9.7
(mm yr ⁻¹)	925	971
Number of broadleaf plots within a 0.5° radius	78	133
Number of needleleaf plots within a 0.5° radius	137	191
Average volume increment of neighbouring broadleaf plots (m ³ ha ⁻¹)	12.2	14.0
Average volume increment of neighbouring needleleaf plots (m ³ ha ⁻¹)	19.8	22.2
Average age of neighbouring broadleaf plots (years)	92	97
Average age of neighbouring needleleaf plots (years)	42	71

581 Table 2. Description of the IFN plots used in this study

Remote sensing technique	LiDAR (airborne)	P-band RADAR (airborne)
Average stand height		
RMSE (m)	1.66	2.34
Number of studies	5	3
References	(Nelson <i>et al.</i> , 2003; Balzter <i>et al.</i> , 2007a; Balzter <i>et al.</i> , 2007b; Stephens <i>et al.</i> , 2007; Breidenbach <i>et al.</i> , 2008)	(Neeff <i>et al.,</i> 2005; Dubois- Fernandez <i>et al.,</i> 2008; Hajnsek <i>et al.,</i> 2009)
Aboveground biomass		
RMSE (tC ha ⁻¹)	23.66	18.5
Number of studies	9	6
(Means <i>et al.</i> , 1999; Drake 2002; Lim and Treitz, 200 Lefsky <i>et al.</i> , 2005b; Watt References Haywood, 2006; Hyde <i>et</i> 2007; Stephens <i>et al.</i> , 200 Boudreau <i>et al.</i> , 2008; Luca <i>al.</i> , 2008)		(Neeff <i>et al.,</i> 2005; Hyde <i>et al.,</i> 2007; Saatchi <i>et al.,</i> 2007; Le Toan <i>et al.,</i> 2008)

583 Table 3. RMSE of LiDAR and P-band radar for height and biomass

584 Figure legends



585

586 Figure 1. Combination of *input data (italic)* and **models (bold)** used in the simulation

587 procedure

588 A typical "spinup" is used to generate intial conditions for both ORCHIDEE and

589 ORCHIDEE-FM simulations. Out of the fifteen ORCHIDEE-FM simulations, seven are

- 590 selected for each site according to their proximity to *in situ* or "pseudo remote sensing"
- 591 measurements, and are given a specific name (FM_{ga} , FM_{gb} , ...).
- 592





The black solid line and grey area respectively give the average and standard deviation of measured woody NPP in National Forest Inventory (NFI) broadleaf (a and b) or needleleaf (c and d) plots within a 50 km radius of the selected "Landes" (a and c) or "Vosges" (b and d) grid cell. Measurements are pooled per age class, and the resulting statistics per age class are smoothed using a "loess" algorithm (only age classes with 5 or more plots are retained). The large-dashed red curve and the small-dashed blue curve respectively give the wood increment in the *STD* and *FM*_{ga} simulations.

602

593





```
The difference between the age retrieved by biomass (FM_{gb} simulation, a and c) or
```

607 height (*FM*_{gh} simulation, b and d) assimilation in ORCHIDEE-FM and the *in situ* estimates

- 608 (IFN dataset) is presented as a frequency distribution for the 201 broadleaf plots (a and
- b) and the 328 needleleaf plots (c and d) of the combined "Landes" and "Vosges"
- 610 locations. A negative value indicates that the simulated age is higher than the
- 611 measurement.







- 619 horizontal dashed purple) assimilating biomass in ORCHIDEE-FM, and FM_{gh}, FM_{lh} and
- 620 FM_{rh} (diagonal dashed green) assimilating height in ORCHIDEE-FM. For a full explanation
- 621 of simulations names, see part 2.3.2, Appendix E and



623 Figure 1.



Measured NEP (gC/m2/yr)

625



627 The values of the *STD* simulation are shown in red, while those of the FM_{ga} simulation

are shown in blue. The red and blue lines represent the respective linear trends of these

629 plot series. A different font is used for each plant functional type, with the following

630 code: TeNE for tempereate needleleaf evergreen, TeBE for temperate broadleaf

631 evergreen, TeBS for temperate broadleaf summergreen, BoNE for boreal needleleaf

evergreen, and BoBS for boreal broadleaf summergreen.





636 In the "Forest spinup" case, the intial conditions of the simulation correspond to the

637 clear-cut of a mature forest whereas in the "Cropland spinup" case, the initial conditions

638 of the simulation correspond to a cropland.



Figure 7. Simulation of GPP, TER and NEP with age assimilation – screened dataset
The values of the *STD* simulation are shown in red, while those of the *FM_{ga}* simulation
are shown in blue. The red and blue lines represent the respective linear trends of these
plot series. A different font is used for each plant functional type, with the following
code: TeNE for tempereate needleleaf evergreen, TeBE for temperate broadleaf
evergreen, TeBS for temperate broadleaf summergreen, BoNE for boreal needleleaf
evergreen, and BoBS for boreal broadleaf summergreen.







of simulations names, see part 2.3.2, Appendix E and

Figure 9. Maps of simulated NEP assimilated data-derived maps of height and volume
The average NEP in the 1990s is obtained from a single simulation of ORCHIDEE-FM for
the 40-50 age class (a), by selecting for each grid cell the ORCHIDEE-FM age-class with

665 closest biomass to the IFN-derived map (b), or by selecting for each grid cell the

666 ORCHIDEE-FM age-class with closest height to the IFN-derived map (c).

667



668



670 For a given age class, the whisker plot represents successively the mean, 1st and 3rd

671 quartile, and extreme values within a distance of twice the interquartile from the box.



674 Figure 11. Age retrieval by assimilating biomass (a) or height (b) in ORCHIDEE-FM

- 675 simulations global flux dataset
- 676 The difference between the age retrieved by biomass (*FM*_{gb} simulation, a) or height
- 677 (*FM*_{gh} simulation, b) assimilation in ORCHIDEE-FM and the in situ estimates (global flux
- dataset) is presented as a frequency distribution for the 31 sites. A negative value
- 679 indicates that the simulated age is higher than the measurement.
- 680
- 681

Baldocchi, D., Falge, E., Gu, L.H., Olson, R., Hollinger, D., Running, S., Anthoni, P.,

- 683 Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X.H.,
- Malhi, Y., Meyers, T., Munger, W., Oechel, W., U, K.T.P., Pilegaard, K., Schmid, H.P.,
- Valentini, R., Verma, S., Vesala, T., Wilson, K., Wofsy, S., 2001. FLUXNET: A new tool to
- 686 study the temporal and spatial variability of ecosystem-scale carbon dioxide, water
- vapor, and energy flux densities. Bulletin of the American Meteorological Society 82,2415-2434.
- Balzter, H., Luckman, A., Skinner, L., Rowland, C., Dawson, T., 2007a. Observations of
- 690 forest stand top height and mean height from interferometric SAR and LiDAR over a

- 691 conifer plantation at Thetford Forest, UK. International Journal of Remote Sensing 28,
- 692 1173-1197.
- Balzter, H., Rowland, C.S., Saich, P., 2007b. Forest canopy height and carbon estimation
- at Monks Wood National Nature Reserve, UK, using dual-wavelength SAR
- 695 interferometry. Remote Sensing of Environment 108, 224-239.
- 696 Bellassen, V., Le Maire, G., Dhote, J.F., Viovy, N., Ciais, P., 2010a. Modeling forest
- 697 management within a global vegetation model Part 1: model structure and general
 698 behaviour. Ecological Modelling 221, 2458–2474.
- Bellassen, V., Le Maire, G., Guin, O., Dhote, J.F., Viovy, N., Ciais, P., 2010b. Modeling
- forest management within a global vegetation model Part 2: model validation from
 tree to continental scale. Ecological Modelling in press.
- 702 Boudreau, J., Nelson, R.F., Margolis, H.A., Beaudoin, A., Guindon, L., Kimes, D.S., 2008.
- Regional aboveground forest biomass using airborne and spaceborne LiDAR in Québec.
 Remote Sensing of Environment 112, 3876.
- 705 Breidenbach, J., Koch, B., Kandler, G., Kleusberg, A., 2008. Quantifying the influence of
- slope, aspect, crown shape and stem density on the estimation of tree height at plot
- 707 level using lidar and InSAR data. International Journal of Remote Sensing 29, 1511-1536.
- 708 Carvalhais, N., Reichstein, M., Ciais, P., Collatz, G.J., Mahecha, M., Montagnani, L.,
- Papale, D., Rambal, S., Seixas, J., 2010. Identification of Vegetation and Soil Carbon Pools
- out of Equilibrium in a Process Model via Eddy Covariance and Biometric Constraints.
- 711 Global Change Biology in press.
- Deleuze, C., Pain, O., Dhote, J.F., Herve, J.C., 2004. A flexible radial increment model for
 individual trees in pure even-aged stands. Annals of Forest Science 61, 327-335.
- Demarty, J., Chevallier, F., Friend, A.D., Viovy, N., Piao, S., Ciais, P., 2007. Assimilation of
- 715 global MODIS leaf area index retrievals within a terrestrial biosphere model.
- 716 Geophysical Research Letters 34, 6.
- 717 Desai, A.R., Moorcroft, P.R., Bolstad, P.V., Davis, K.J., 2007. Regional carbon fluxes from
- an observationally constrained dynamic ecosystem model: Impacts of disturbance, CO2
- 719 fertilization, and heterogeneous land cover. Journal of Geophysical Research-
- 720 Biogeosciences 112.
- 721 Dhôte, J.-F., Hervé, J.-C., 2000. Changements de productivité dans quatre forêts de
- 722 chênes sessiles depuis 1930 : une approche au niveau du peuplement. Ann. For. Sci. 57,723 651-680.
- 724 Drake, J.B., Dubayah, R.O., Knox, R.G., Clark, D.B., Blair, J.B., 2002. Sensitivity of large-
- footprint lidar to canopy structure and biomass in a neotropical rainforest. Remote
- 726 Sensing of Environment 81, 378-392.
- 727 Dubois-Fernandez, P.C., Souyris, J.C., Angelliaume, S., Garestier, F., 2008. The Compact
- 728 Polarimetry Alternative for Spaceborne SAR at Low Frequency. leee Transactions on
- 729 Geoscience and Remote Sensing 46, 3208-3222.
- 730 Dufrene, E., Davi, H., Francois, C., le Maire, G., Le Dantec, V., Granier, A., 2005.
- 731 Modelling carbon and water cycles in a beech forest Part I: Model description and
- vuncertainty analysis on modelled NEE. Ecological Modelling 185, 407-436.

- 733 Durrieu, S., 2010. Lidar for Earth and Forests. Proposal in response to the ESA Call for
- 734 Earth Explorer Opportunity Mission EE-8. CEMAGREF, Montpellier, France, 5 p.
- 735 Friedlingstein, P., Joel, G., Field, C.B., Fung, I.Y., 1999. Toward an allocation scheme for 736 global terrestrial carbon models. Global Change Biology 5, 755-770.
- 737
- Grant, R.F., Black, T.A., Humphreys, E.R., Morgenstern, K., 2007. Changes in net
- 738 ecosystem productivity with forest age following clearcutting of a coastal Douglas-fir
- 739 forest: testing a mathematical model with eddy covariance measurements along a 740 forest chronosequence. Tree Physiology 27, 115-131.
- 741 Hajnsek, I., Kugler, F., Lee, S.K., Papathanassiou, K.P., 2009. Tropical-Forest-Parameter
- 742 Estimation by Means of Pol-InSAR: The INDREX-II Campaign. leee Transactions on 743 Geoscience and Remote Sensing 47, 481-493.
- 744 Hyde, P., Nelson, R., Kimes, D., Levine, E., 2007. Exploring LIDAR-RaDAR synergy -
- 745 predicting aboveground biomass in a southwestern ponderosa pine forest using LiDAR,
- 746 SAR and InSAR. Remote Sensing of Environment 106, 28-38.
- 747 IFN, 2006. Observer la forêt française : mission première de l'IFN. L'IF, 12.
- 748 IPCC, 2003. Good Practice Guidance for Land-Use, Land-Use Change and Forestry.
- 749 Intergovernmental Panel on Climate Change, Kanagawa, Japan, 534 p.
- 750 JRC, 2009. European Forest Yield Table's database,
- 751 http://afoludata.jrc.ec.europa.eu/DS Free/abc intro.cfm.
- 752 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M.,
- 753 Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W.,
- 754 Janowiak, J., Mo, K.C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R.,
- 755 Joseph, D., 1996. The NCEP/NCAR 40-year reanalysis project. Bulletin of the American 756 Meteorological Society 77, 437-471.
- 757 Krinner, G., Viovy, N., de Noblet-Ducoudre, N., Ogee, J., Polcher, J., Friedlingstein, P.,
- 758 Ciais, P., Sitch, S., Prentice, I.C., 2005. A dynamic global vegetation model for studies of
- 759 the coupled atmosphere-biosphere system. Global Biogeochemical Cycles 19, 44.
- 760 Le Quere, C., Raupach, M.R., Canadell, J.G., Marland, G., Bopp, L., Ciais, P., Conway, T.J.,
- 761 Doney, S.C., Feely, R.A., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R.A., House,
- 762 J.I., Huntingford, C., Levy, P.E., Lomas, M.R., Majkut, J., Metzl, N., Ometto, J.P., Peters,
- 763 G.P., Prentice, I.C., Randerson, J.T., Running, S.W., Sarmiento, J.L., Schuster, U., Sitch, S.,
- 764 Takahashi, T., Viovy, N., van der Werf, G.R., Woodward, F.I., 2009. Trends in the sources 765 and sinks of carbon dioxide. Nature Geoscience 2, 831-836.
- 766 Le Toan, T., Baltzer, H., Paillou, P., Papathanassiou, K., Plummer, S., Quegan, S., Rocca,
- 767 F., Ulander, L., 2008. Candidate Earth Explorer Core Mission - Biomass. ESA, Noordwijk, 768 124 p.
- 769 Lefsky, M.A., Harding, D.J., Keller, M., Cohen, W.B., Carabajal, C.C., Espirito-Santo, F.D.,
- 770 Hunter, M.O., de Oliveira, R., 2005a. Estimates of forest canopy height and aboveground 771 biomass using ICESat. Geophysical Research Letters 32.
- 772 Lefsky, M.A., Turner, D.P., Guzy, M., Cohen, W.B., 2005b. Combining lidar estimates of
- 773 aboveground biomass and Landsat estimates of stand age for spatially extensive
- 774 validation of modeled forest productivity. Remote Sensing of Environment 95, 549.

- Lim, K.S., Treitz, P.M., 2004. Estimation of above ground forest biomass from airborne
- discrete return laser scanner data using canopy-based quantile estimators. In, pp. 558-570.
- Lindner, M., Lucht, W., Bouriaud, O., Green, T., Janssens, I., 2004. Specific Study on
 Forest Greenhouse Gas Budget. CarboEurope-GHG, Jena, 62 p.
- 780 Loustau, D., Bosc, A., Colin, A., Ogee, J., Davi, H., Francois, C., Dufrene, E., Deque, M.,
- 781 Cloppet, E., Arrouays, D., Le Bas, C., Saby, N., Pignard, G., Hamza, N., Granier, A., Breda,
- 782 N., Ciais, P., Viovy, N., Delage, F., 2005. Modeling climate change effects on the potential
- production of French plains forests at the sub-regional level. Tree Physiology 25, 813-823.
- Lucas, R.M., Lee, A.C., Bunting, P.J., 2008. Retrieving forest biomass through integrationof CASI and LiDAR data. In, pp. 1553-1577.
- 787 Luyssaert, S., Ciais, P., Piao, S.L., Schulze, E.D., Jung, M., Zaehle, S., Schelhaas, M.J.,
- 788 Reichstein, M., Churkina, G., Papale, D., Abril, G., Beer, C., Grace, J., Loustau, D.,
- 789 Matteucci, G., Magnani, F., Nabuurs, G.J., Verbeeck, H., Sulkava, M., van der Werf, G.R.,
- Janssens, I.A., Team, C.-I.S., 2010. The European carbon balance. Part 3: forests. Global
 Change Biology 16, 1429-1450.
- 792 Luyssaert, S., Inglima, I., Jung, M., Richardson, A.D., Reichsteins, M., Papale, D., Piao,
- 793 S.L., Schulzes, E.D., Wingate, L., Matteucci, G., Aragao, L., Aubinet, M., Beers, C.,
- 794 Bernhoffer, C., Black, K.G., Bonal, D., Bonnefond, J.M., Chambers, J., Ciais, P., Cook, B.,
- 795 Davis, K.J., Dolman, A.J., Gielen, B., Goulden, M., Grace, J., Granier, A., Grelle, A., Griffis,
- T., Grunwald, T., Guidolotti, G., Hanson, P.J., Harding, R., Hollinger, D.Y., Hutyra, L.R.,
- 797 Kolar, P., Kruijt, B., Kutsch, W., Lagergren, F., Laurila, T., Law, B.E., Le Maire, G., Lindroth,
- A., Loustau, D., Malhi, Y., Mateus, J., Migliavacca, M., Misson, L., Montagnani, L.,
- 799 Moncrieff, J., Moors, E., Munger, J.W., Nikinmaa, E., Ollinger, S.V., Pita, G., Rebmann, C.,
- 800 Roupsard, O., Saigusa, N., Sanz, M.J., Seufert, G., Sierra, C., Smith, M.L., Tang, J.,
- Valentini, R., Vesala, T., Janssens, I.A., 2007. CO2 balance of boreal, temperate, and
- tropical forests derived from a global database. Global Change Biology 13, 2509-2537.
- 803 Luyssaert, S., Reichstein, M., Schulze, E.D., Janssens, I.A., Law, B.E., Papale, D., Dragoni,
- 804 D., Goulden, M.L., Granier, A., Kutsch, W.L., Linder, S., Matteucci, G., Moors, E., Munger,
- J.W., Pilegaard, K., Saunders, M., Falge, E.M., 2009. Toward a consistency cross-check of

806 eddy covariance flux-based and biometric estimates of ecosystem carbon balance.
807 Global Biogeochemical Cycles 23

- 807 Global Biogeochemical Cycles 23.
- 808 Magnani, F., Mencuccini, M., Borghetti, M., Berbigier, P., Berninger, F., Delzon, S.,
- 809 Grelle, A., Hari, P., Jarvis, P.G., Kolari, P., Kowalski, A.S., Lankreijer, H., Law, B.E.,
- Lindroth, A., Loustau, D., Manca, G., Moncrieff, J.B., Rayment, M., Tedeschi, V.,
- 811 Valentini, R., Grace, J., 2007. The human footprint in the carbon cycle of temperate and
- 812 boreal forests. Nature 447, 848-850.
- 813 Masera, O.R., Garza-Caligaris, J.F., Kanninen, M., Karjalainen, T., Liski, J., Nabuurs, G.J.,
- 814 Pussinen, A., de Jong, B.H.J., Mohren, G.M.J., Tz, 2003. Modeling carbon sequestration
- 815 in afforestation, agroforestry and forest management projects: the CO2FIX V.2
- 816 approach. Ecological Modelling 164, 177-199.

- 817 Means, J.E., Acker, S.A., Harding, D.J., Blair, J.B., Lefsky, M.A., Cohen, W.B., Harmon,
- 818 M.E., McKee, W.A., 1999. Use of large-footprint scanning airborne lidar to estimate
- 819 forest stand characteristics in the Western Cascades of Oregon. Remote Sensing of

820 Environment 67, 298-308.

- 821 Mitchell, S., Beven, K., Freer, J., 2009. Multiple sources of predictive uncertainty in
- modeled estimates of net ecosystem CO2 exchange. Ecological Modelling 220, 3259-3270.
- 824 Mitchell, T.D., Jones, P.D., 2005. An improved method of constructing a database of
- 825 monthly climate observations and associated high-resolution grids. International Journal 826 of Climatology 25, 693-712.
- 827 Moorcroft, P.R., Hurtt, G.C., Pacala, S.W., 2001. A method for scaling vegetation
- dynamics: The ecosystem demography model (ED). Ecological Monographs 71, 557-585.
- Nabuurs, G.J., Hengeveld, G., Heidema, N., Brus, D., Goedhart, P., Walvoort, D., van den
- 830 Wyngaert, I., van der Werf, B., Tröltzsch, K., Lindner, M., Zanchi, G., Gallaun, H.,
- Schwaiger, H., Teobaldelli, M., Seufert, G., Kenter, B., 2008. Mapping the continent: High
- resolution forest resource analyses of European forests. The European Carbon Balance Research Highlights 2008. CarboEurope-IP (eds.), 4 p.
- 834 Naesset, E., 2009. Effects of different sensors, flying altitudes, and pulse repetition
- frequencies on forest canopy metrics and biophysical stand properties derived from
- small-footprint airborne laser data. Remote Sensing of Environment 113, 148-159.
- 837 Neeff, T., Dutra, L.V., dos Santos, J.R., Freitas, C.D., Araujo, L.S., 2005. Tropical forest
- 838 measurement by interferometric height modeling and P-band radar backscatter. Forest839 Science 51, 585-594.
- Nelson, R.F., Parker, G.G., Hom, M., 2003. A portable airborne laser system for forest
 inventory. American Society for Photogrammetry and Remote Sensing, Bethesda, USA, 7
- 842

p.

- 843 Nunery, J.S., Keeton, W.S., 2010. Forest carbon storage in the northeastern United
- States: Net effects of harvesting frequency, post-harvest retention, and wood products.
 Forest Ecology and Management 259, 1363-1375.
- 846 Pan, Y., Chen, J.M., Birdsey, R., McCullough, K., He, L., Deng, F., 2010. Age structure and 847 disturbance logacy of North American forests. Biogeosciences Discuss. 7, 979, 1020
- disturbance legacy of North American forests. Biogeosciences Discuss. 7, 979-1020.
- Piao, S.L., Fang, J.Y., Ciais, P., Peylin, P., Huang, Y., Sitch, S., Wang, T., 2009. The carbon
 balance of terrestrial ecosystems in China. Nature 458, 1009-U1082.
- 850 Porté, A., 1999. Modélisation des effets du bilan hydrique sur la production primaire et
- la croissance d'un couvert de pin maritime (Pinus pinaster Ait.) en lande humide,
- 852 University of Paris XI, Orsay, France, 140 p.
- 853 Pretzsch, H., Biber, P., Durský, J., 2002. The single tree-based stand simulator SILVA:
- 854 construction, application and evaluation. Forest Ecology and Management 162, 3.
- 855 Reineke, L.H., 1933. Perfecting a stand-density index for even-aged forests. Journal of
- Agricultural Research 46, 627-638.
- 857 Rubinstein, R.Y., Kroese, D.P., 1981. Simulation and the Monte Carlo Method. Wiley,
- 858 New York, USAp.

- 859 Saatchi, S., Halligan, K., Despain, D.G., Crabtree, R.L., 2007. Estimation of forest fuel load
- from radar remote sensing. leee Transactions on Geoscience and Remote Sensing 45,1726-1740.
- 862 Santaren, D., 2006. Optimisation des paramètres du modèle de biosphère ORCHIDEE à
- 863 partir de mesures sur site des flux de carbone, d'eau et d'énergie, Université Versailles
- 864 Saint-Quentin, Versailles, 190 p.
- 865 Sarrat, C., Noilhan, J., Lacarrere, P., Ceschia, E., Ciais, P., Dolman, A.J., Elbers, J.A.,
- 866 Gerbig, C., Gioli, B., Lauvaux, T., Miglietta, F., Neininger, B., Ramonet, M., Vellinga, O.,
- Bonnefond, J.M., 2009. Mesoscale modelling of the CO2 interactions between the
 surface and the atmosphere applied to the April 2007 CERES field experiment.
- 869 Biogeosciences 6, 633-646.
- 870 Sato, H., Itoh, A., Kohyama, T., 2007. SEIB-DGVM: A new dynamic global vegetation
- 871 model using a spatially explicit individual-based approach. Ecological Modelling 200,872 279-307.
- Sitch, S., Smith, B., Prentice, I.C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J.O., Levis,
- 874 S., Lucht, W., Sykes, M.T., Thonicke, K., Venevsky, S., 2003. Evaluation of ecosystem
- dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global
 vegetation model. Global Change Biology 9, 161-185.
- Stephens, P.R., Watt, P.J., Loubser, D., Haywood, A., Kimberley, M.O., 2007. Estimation
 of carbon stocks in New Zealand planted forests using airborne scanning LiDAR. IAPRS
 36, 389-394.
- 880 Thornton, P.E., Law, B.E., Gholz, H.L., Clark, K.L., Falge, E., Ellsworth, D.S., Golstein, A.H., 881 Mansan, B.K., Hallinger, D., Falk, M., Chan, L., Sparke, J.D., 2002, Madaling and
- Monson, R.K., Hollinger, D., Falk, M., Chen, J., Sparks, J.P., 2002. Modeling and
- 882 measuring the effects of disturbance history and climate on carbon and water budgets
- in evergreen needleleaf forests. Agricultural and Forest Meteorology 113, 185-222.
- Vannière, B., 1984. Tables de production pour les forêts françaises. Ecole Nationale du
 Génie Rural, des Eaux et des Forêts, Nancy, 160 p.
- 886 Vetter, M., Churkina, G., Jung, M., Reichstein, M., Zaehle, S., Bondeau, A., Chen, Y., Ciais,
- 887 P., Feser, F., Freibauer, A., Geyer, R., Jones, C., Papale, D., Tenhunen, J., Tomelleri, E.,
- 888 Trusilova, K., Viovy, N., Heimann, M., 2008. Analyzing the causes and spatial pattern of
- the European 2003 carbon flux anomaly using seven models. Biogeosciences 5, 561-583.
- 890 Watt, P.J., Haywood, A., 2006. Evaluation of airborne scanning LiDAR generated data as
- input into biomass/carbon models. No. 38A08068, Ministry for the Environment,
- 892 Wellington, New Zealand, 34 p.
- Willmott, C.J., 1982. Some Comments on the Evaluation of Model Performance. Bulletinof the American Meteorological Society 63, 1309-1313.
- Zaehle, S., Friend, A.D., 2010. Carbon and nitrogen cycle dynamics in the O-CN land
- 896 surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter
- 897 estimates. Global Biogeochemical Cycles 24.
- Zaehle, S., Friend, A.D., Friedlingstein, P., Dentener, F., Peylin, P., Schulz, M., 2010.
- 899 Carbon and nitrogen cycle dynamics in the O-CN land surface model: 2. Role of the

- 202 Zaehle, S., Sitch, S., Prentice, I.C., Liski, J., Cramer, W., Erhard, M., Hickler, T., Smith, B.,
- 903 2006. The importance of age-related decline in forest NPP for modeling regional carbon
- 904 balances. Ecological Applications 16, 1555-1574.
- 905