

Multi-criteria spatialization of soil organic carbon sequestration potential from agricultural intensification in Senegal

Valentin Bellassen, Raphaël J. Manlay, J.P. Chéry, V. Gitz, A. Touré, M. Bernoux, J.L. Chotte

► To cite this version:

Valentin Bellassen, Raphaël J. Manlay, J.P. Chéry, V. Gitz, A. Touré, et al.. Multi-criteria spatialization of soil organic carbon sequestration potential from agricultural intensification in Senegal. Climatic Change, 2010, 98 (1-2), pp.213-243. 10.1007/s10584-009-9635-x . hal-01149374

HAL Id: hal-01149374 https://hal.science/hal-01149374

Submitted on 18 Oct 2023 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- 1 **Running head**: Soil organic carbon sequestration in Senegal
- 2 **Title**: Multi-criteria spatialization of soil organic carbon sequestration potential from
- 3 agricultural intensification in Senegal
- 4 **Authors**: Valentin Bellassen^{1,2}, Raphaël J. Manlay^{1,2}, Jean-Pierre Chéry³, Vincent Gitz⁴,
- 5 Assize Touré⁵, Martial Bernoux¹, Jean-Luc Chotte¹
- 6 Journal: Climatic Change
- 7 Year: 2010

8 Affiliations:

- 9 1. UR SeqBio, Institut de Recherche pour le Développement (IRD)
- 10 2. Ecole Nationale du Génie Rural, des Eaux et des Forêts (AgroParisTech–ENGREF)
- 11 3. UMR TETIS, Ecole Nationale du Génie Rural, des Eaux et des Forêts
- 12 (AgroParisTech-ENGREF)
- 13 4. Centre International de Recherche sur l'Environnement et le Développement (CIRED)
- 14 5. Centre de Suivi Ecologique (CSE)
- 15 **Full address for correspondence**:
- 16 Valentin Bellassen
- 17 LSCE-Orme, Bât. 712,
- 18 Orme des Merisiers,
- 19 F-91191 GIF-SUR-YVETTE CEDEX
- 20 FRANCE
- 21 Tel: + 33 1 58 50 19 75 / + 33 1 69 08 31 01
- 22 Fax: + 33 1 69 08 77 16
- 23 E-mail address: vbella@lsce.ipsl.fr
- 24
- 25 26

27 Abstract

On the eve of the 15th climate negotiations conference in Copenhagen, the pressure to assess all climate mitigation options is mounting. In this study, a bio-physic model and a socio-economic model were designed and coupled to assess the carbon sequestration potential of agricultural intensification in Senegal. The biophysical model is a multiple linear regression, calibrated and tested on a dataset of long-term agricultural trials established in West Africa. The socio-economic model integrates both financial and environmental costs related to considered practice changes. Both models are spatially explicit and the resulting 35 spatial patterns were computed and displayed over Senegal with a geographic information system. The national potential from large-scale intensification was assessed at 0.65-0.83 MtC. 36 37 With regards to local-scaled intensification as local projects, the most profitable areas were 38 identified in agricultural expansion regions (especially Casamance), while the areas that meet 39 the current financial additionality criteria of the Clean Development Mechanism were located 40 in the northern part of the Peanut Basin. Using the current relevant mode of carbon valuation 41 (Certified Emission Reductions), environmental benefits are small compared to financial 42 benefits. This picture is radically changed if "avoided deforestation", a likely consequence of 43 agricultural intensification, is accounted for as the greenhouse gases sink capacity of projects 44 increases by an average of a hundred-fold over Senegal.

45

46 **1 Introduction**

47 The amount of soil organic carbon (SOC) results from an equilibrium between the 48 inputs and outputs to the system, which are driven by various parameters of natural or human origins (Schlesinger et al. 2000). In West African agro-ecosystems, this equilibrium is put in 49 50 jeopardy as few inputs are available to compensate for harvested biomass, a major output. 51 According to Sanchez (1997), the soils of sub-Saharan Africa would have undergone an average depletion of 660 kg of nitrogen (N) ha⁻¹, 75 kg of phosphorus (P) ha⁻¹, and 450 kg of 52 53 potassium (K) ha⁻¹ over the 1970-2000 period, together with emitting 27 Mt of carbon (C) per year (with an additional 18 MtC y^{-1} lost by erosion and therefore assumed not to be emitted). 54

Soil carbon loss potentially contributes to another environmental problem: global 55 56 warming. In Senegal, 32% of greenhouse gases (GHG) emissions come from the agricultural 57 sector (Kante 1997). Since agriculture is part of the problem, it could also be part of the 58 solution to global warming, if agricultural soil carbon stocks are managed to be increased: 59 Parton et al. (2004) estimated that 50% of the carbon storage potential of Senegal (as defined 60 by Bernoux et al. (2006), "carbon storage" refers to the change in carbon stored in the system, whereas "carbon sequestration" includes all indirect emissions related to the practice that 61 62 "stores" carbon) would lie in improved practices in agriculture (that is mainly increased 63 fallowing time combined with manure application). The entry into force of the Kyoto Protocol 64 to the United Nations Framework Convention on Climate Change (UNFCCC) in 2005 has created an economic value for carbon emitted or stored, paving the way for an internalisation 65 66 of this potentially important economic externality, if climate change damages are expected to 67 be high. Carbon emitters may therefore consider African agriculture as an opportunity to get 68 carbon credits, in case this activity becomes eligible to the Kyoto framework, or to any other

69 frameworks rewarding carbon sequestration. Different agricultural practices (amount of 70 inputs, length of fallow, agroforestry, ...) result in different stocks of carbon in the soil 71 (Vagen et al. 2005). Considering the prevailing pressure on land in West Africa due to food 72 demand, tenure rights and fire risks, it is particularly relevant to focus on agricultural 73 intensification as a change of practice that is not likely to meet significant economic or social 74 barriers.

Fallowing, manuring and crop residue recycling are traditional practices that sustain soil fertility in West Africa (Kowal and Kassam 1978). The increased or stabilized yields provided by such practices partly rely on the maintaining of a threshold level of soil organic matter (SOM) (Pieri 1992; Pieri 1995). Moreover, several studies have shown that a synergistic effect exists between mineral fertilizers and organic amendments in Sub-Saharan Africa, that leads both to higher yields and SOC content (Pieri 1992; Badiane et al. 2001; Bationo and Buerkert 2001; Palm et al. 2001; Vagen et al. 2005).

These on-field results could seem counter-intuitive at first sight because microbial activity and therefore SOC mineralization is enhanced with increasing mineral form of nitrogen (Corbeels et al. 2006). But nitrogen has the capacity to cycle fairly rapidly between the inorganic and organic forms within the soils and therefore may gave different result than the most expected. Moreover, some recent developments in modelling and laboratory experiments point to more complex mechanisms that could explain field

The aim of the present study was to assess the carbon sequestration potential of the cropped soils of Senegal under a scenario of agricultural intensification. This was done by developing, and coupling, two models: a biophysical model to predict the quantity of carbon sequestrated by a given practice, and a socio-economic model to assess the net present value (NPV)¹ of these practices. This coupled approach is indeed necessary for two reasons:

93 - the pedo-climatic conditions that are suitable for carbon storage may not coincide
94 with areas where the corresponding practices are socio-economically applicable;

eligibility to carbon valuation often includes specific criteria. For example, in the
Clean Development Mechanism (CDM) framework (UNFCCC 2003), Certified
Emission Reductions (CERs) are only granted where their value tips the balance to
make the project profitable (financial additionality criteria). On the other hand, the
Global Environmental Fund (GEF) has an internal policy of lending money only for
projects estimated to cost less than 10 US\$ tCeq⁻¹. In such circumstances, project-

¹ The Net Present Value (NPV) of a given practice is its cumulated value over a given number of years, each year's value being actualized by a discount rate.

101 designers will be interested to find where such economically constrained projects may 102 be located. To increase the intelligibility of this work for such project-designers, we 103 purposely used the vocabulary of the CDM framework, and paid a special attention to 104 its eligibility criteria (financial and environmental additionality of the project, "sound 105 reporting, monitoring and verification", ...).

5 6

106 This kind of coupled approach has already been explored locally in the Old Peanut 107 Basin (Tschakert 2004), but not at the country scale in Senegal. Another peculiarity of the 108 present study is that the biophysical model, though simple, has been both calibrated and tested 109 on a set of long-term local agronomic trials. This information is therefore complementary to 110 the one given by more general models such as Century (Parton et al. 2004), as the model 111 narrow range of applicability compared to mechanistic models such as Century is balanced by 112 its specificity to the studied country (Senegal). Indeed, even the adapted version of Century 113 used by Tschakert (2004) in the Old Peanut Basin yields a surprisingly long time-scale for 114 carbon dynamics, and Cerri et al. (2004) highlighted the inadequacy of Century for short-term 115 changes prediction in the Amazonian context.

Recently, a system called GEFSOC (Easter et al. 2007; Milne et al. 2007) set out a 116 117 framework to make regional scale assessment of SOC stocks and change using site scale 118 experimental data, regional GIS databases and simulation models using Century and Roth-C. 119 But the GEFSOC system need in it first stage a model evaluation, based on long term field 120 experiment, of both Century and Roth-C for the range of the different land uses and covering 121 all the conditions (soil type, climate...) found in the studied area. This first stage could be 122 difficult to implement with a large number of climate-soil-management combinations (Cerri 123 et al. 2007). More over this first step is important as it could lead to under or overestimation 124 of the prediction if not perform adequately, and there as there are no automatic, built-in 125 functions for uncertainty analysis in the GEFSOC system this may furnish results with high 126 level of uncertainties (Easter et al. 2007). In our study we decided for a more robust and 127 simple procedure adapted to the available information.

Finally, the coupling of this model to a Geographic Information System (GIS) produces spatialized information, which is essential for many purposes. One example is the aforementioned CDM framework in which project perimeter identification is one of the crucial steps.

133 **2** Material and Methods

134 2.1 Study site

135 Most of Senegalese soils are ferruginous and very sandy, especially those in cultivated 136 areas: 85% of total cropped soils contain less than 20% of clay (Stancioff et al. 1986; DAAC 137 2002). As opposed to clayey soils where the SOM pool in mostly clay fraction, sandy soils 138 have been shown to be particularly sensitive to organic amendments (Feller et al. 1991). On 139 the other hand, the total SOM pool of non-degraded sandy soils tends too be smaller than that 140 of similar clayey soils, which renders their storage potential smaller (Jones 1973; Feller 141 1995). Another important pattern driving carbon storage is that of rainfall distribution which follows a southward gradient from 200 mm y⁻¹ in the North to 1400 mm y⁻¹ in the South 142 143 (Figure 1), with rain falling during a single rainy season of 3 to 6 months (Kelly et al. 1996; 144 Worldclim 2000).

145 As mentioned in introduction, agriculture and livestock breeding closely interact: 146 during the beginning of the dry season, livestock is left on fields during the day to graze on 147 crop residues and deposit manure (Manlay et al. 2004). Only later in the season when residues 148 are exhausted, or during the wet season when animals are excluded from cropped lands, do 149 they go find food on marginal lands. Moreover, during all the dry season, animals are 150 corralled at night on fields close to human dwellings, so that they fertilize them and so that 151 livestock theft can be prevented. This close kind of interaction is made possible either by 152 farmer ownership of livestock, or through grazing contracts with neighbouring herders 153 (Powell and Williams 1995; Manlay et al. 2004).

154 Senegal has also hosted many research projects, which yielded large sets of field data. 155 The project "Sequestration Of Carbon in Soil Organic Matter" co-financed by the USAID and 156 the USGS recently achieved its research, which led to acquisition of spatial data, and the 157 publication of a special issue of the Journal of Arid Environments (2004, Vol. 59). The 158

current study is a preliminary contribution to CBP:MMM ("Carbon benefits project,

159 monitoring measuring and modelling"), the second phase of the GEFSOC project, currently

160 under examination by the GEF executive board, and as such benefited from the active

161 collaboration of the Centre de Suivi Ecologique in Dakar.

162 Senegal has ratified both the United Nations Framework Convention on Climate 163 Change (UNFCCC) and the Kyoto Protocol. Moreover, it is fairly well rated by *Transparency* 164 International and COFACE for its political stability (COFACE 2006; Transparency International 2006). This set of political and scientific reasons makes it a potentially attractivecountry for environmental projects, such as CDM or carbon sequestration projects.

167

2.2 Agricultural intensification scenarios

168

2.2.1 Manure availability

169 For the organic inputs of the scenarios, only manure was taken into account, to be 170 consistent with the calibration data of the model. The pool of animals which manure could be 171 tapped through the traditional agriculture-livestock interaction was conservatively limited to 172 those located within 10km of a cultivated area, thereby excluding migrating herds. This 173 computation leaves out 45% of total livestock. Livestock density was assumed to be uniform 174 over the area suitable for livestock grazing (croplands, shrublands and savannas), and could 175 therefore be computed from departmental statistics and a land-cover map (DAAC 2002; 176 Ministère de l'Agriculture et de l'Hydraulique 2004). This pool of animals was then converted 177 into a manure pool using the daily excretion figures of Fernandez-Rivera et al. (1995). The 178 resulting figures for manure availability are consistent with previous assessments at national 179 or local scale (Fernandez-Rivera et al. 1995; Breman 1998).

180

2.2.2 Description of scenarios

181 Three scenarios were elaborated, and then compared to a baseline scenario with the 182 models designed in this study, in order to assess the opportunities for carbon sequestration by 183 intensifying agriculture:

Baseline scenario (Business as Usual or BAU): The baseline scenario considered was the continuation of current practices as described in part 2.1. With regards to mineral fertilizers, it assumes that none is used: the data gathered by the International Fertilizer Association tend to bolster up this assumption (International Fertilizer Industry Association 2003).

189 The collecting efficiency of the available manure pool for current practices was set at 190 50% during the dry season (as animals are corralled at night and also deposit manure on fields 191 during the beginning of the season), and at 0% during the wet season (Fernandez-Rivera et al. 192 1995). This is again a conservative approximation, as deposited manure is not as efficient as 193 manure collected in stables and manually spread (McIntire and Powell 1994), the latter kind 194 being the one used for model calibration. The length of the dry season necessary for this 195 computation was assumed to vary linearly with annual rainfall average from 3 months in 196 Northern Senegal to 6 months in Southern Senegal.

197 Dry season stalling (LOWINT): The LOWINT scenario represents a first possibility of 198 intensification at national scale. Animals are stalled during the dry season, which is translated 199 in the model into an improvement of collecting efficiency from 50% to 95% during this 200 period. Crops receive the amount of mineral fertilization recommended by agronomic 201 research (CIRAD et al. 2004). Since these recommendations are crop-specific, a crop 202 repartition pattern was required. As in Liu et al. (2004), the share of each crop on each grid 203 element was set at the crop's regional share of cultivated land in 2002-2003 (source: Senegal, 204 DSDIA-DAPS-MAE). Only the main crops were taken into account, and rice was excluded. 205 Rice is indeed so different from other crops in the way it is grown that this study's model is 206 probably irrelevant for it. Therefore, the parts of Senegal classified as rice-growing regions by 207 Le Fur (2000) were excluded from the analysis.

Wet season stalling (HIGHINT): The HIGHINT scenario represents a higher degree of intensification at national scale. Crops receive the same amount of mineral fertilizers as for LOWINT, but animals are stalled all year long, which leads to a 95% collecting efficiency even during the wet season. This scenario could be less plausible than LOWINT, depending on local labour availability: the labour needed for animal stalling (bringing food, cleaning the stalls, spreading the manure, ...) is indeed usually less readily available during the wet season, as crops are grown and require a lot of labour (Dugué et al. 2004).

215 Project-based intensification (MAXINT): The MAXINT scenario is set to assess the 216 opportunities arising from a maximal degree of intensification that can only be realized on a 217 local, project-based basis, on farms well-endowed in livestock. It assumes that any quantity of 218 inputs needed can be brought to the field, notwithstanding the regional availability of such 219 resources or the special needs of the regionally dominant crop. The system considered is a groundnut/millet rotation which receives recommended amount of all inputs: 5 tDM ha⁻¹ yr⁻¹ 220 of manure (Bosma et al. 1995), 25 kgN ha⁻¹ vr⁻¹, and 11.5 kgP ha⁻¹ vr⁻¹ (CIRAD et al. 2004). 221 222 Unlike the two other intensification scenarios, it is therefore unfit for a national assessment, 223 but allows the comparison of all sites on an identical basis.

224

2.3 The biophysical model: a statistical model of soil carbon change

225

2.3.1 Carbon storage dynamics

To comply with CDM framework, the model must quantify the final difference in SOC between several management techniques only, and therefore it needs not focus on the dynamics that lead to this final difference. However, to select data representing this final difference and therefore suitable to calibrate the model, a hypothesis on the time to reachequilibrium in topsoil SOC was needed.

231 Several studies (Bationo et al. 1995; Tiessen et al. 1998) suggest that following a 232 change in land-use the soil surface layer reaches a new equilibrium in SOC after 3-5 years in 233 West African sandy soils. However, other studies assume a much longer timeframe of 25-50 234 years (Batjes 2001; Tschakert 2004). Among the few long-term trials that fuel the dataset 235 gathered for this study and where SOC had been measured overtime with the same method, 236 the 5-year dynamics hypothesis was verified in Thilmakha (Senegal) and Bambey (Senegal), but not in Bebedjia (Tchad). Bebedjia is very rainy (around 1100 mm y^{-1}), and slightly more 237 clayey (10% clay in the surface horizon) than the first two locations. Therefore, for the 238 239 modelling purposes of this study, the assumption that the equilibrium for sub-surface SOC 240 was reached after five years of treatment was retained for most of Senegal, but slightly 241 modified for the most n environmental and treatment variables, the general linear model does 242 not strictly apply, forbidding the use of the AIC (StatSoft 2006). A better suited mixed linear 243 model could not be applied either, due to the limited number of repetition within certain sites. 244 It is however well established that soil clay content and annual rainfall play a major role in the 245 SOM dynamics of rainy areas. There, model outputs on total SOC change were still 246 considered to be true, but the equilibrium was assumed to be reached only as fast as a capped 247 maximum speed would allow. This difference in the timeframe of carbon dynamics based on rainfall is consistent with the one observed by Elberling et al. (2003). 248

The storage speed cap was set at $0.5 \text{ tC ha}^{-1} \text{ yr}^{-1}$, which is approximately the storage speed of the best management practice in the Peanut Basin as computed with the Century model in Tschakert (2004). It is also the value given by the IPCC calculator (IPCC 1996) for Senegal, and by Ringius (2002) for West Africa.

253

2.3.2 Data collection

An extensive literature survey was undertaken to gather calibration data from long-term agronomic trials conducted in Senegal, and in other West African sites with edaphic conditions similar to those existing in Senegal. Several criteria limited the amount of data retained for the calibration:

At least 4 years of treatment: as the model is geared to predict final SOC stocks
 changes, calibration data must correspond to the final state of the system that is
 measured for its SOC. Therefore, consistently with the aforementioned hypothesis,
 only the studies where treatments had been maintained for at least 4 years were used

262 for model calibration (for most of them, treatments were actually maintained for more263 than 10 years).

- 264 Minimal data requirement: as in most meta-analysis (Jones 1973), the original studies _ 265 used here had different methodologies. The most problematic here was the potentially 266 important discrepancies for SOC content measurements arising from use of varying 267 methods (Pansu and Gautheyrou 2006). To alleviate any such analytical bias, data were expressed as the difference in SOC content (ΔC) between each treatment and its 268 control (which receives no input whatsoever). Only variations in the 0-20 cm layer 269 have been considered², since the influence of management below this depth has been 270 found to be of small magnitude (Tiessen et al. 1998; Manlay et al. 2002; Elberling et 271 272 al. 2003). The standard of the International Panel on Climate Change (IPCC) is 30 cm, 273 but it may not be relevant for carbon sequestration projects in agricultural soils as the 274 costs of monitoring the 20-30 cm depth may override the potential increase in SOC stocks there, and since only shallow tillage is practised. Finally, the availability of a 275 276 control, and of a minimal set of environmental parameters such as average rainfall and soil texture, also limited the number of studies eligible to the analysis. 277
- 278 No crop residue return or fallow: The only kinds of organic amendments retained _ 279 were manure and compost. The C:N ratio of crop residues is indeed usually much 280 higher than compost or manure, which implies notable differences in decomposition 281 dynamics, and therefore in their effects on SOC and crop yields (Allard et al. 1983; De 282 Ridder and Van Keulen 1990; McIntire and Powell 1994). For this reason, treatments 283 with crop residue return were excluded from the analysis. Fallow periods, which 284 contribute to soil restoration through increased root density and residue return, were 285 excluded for the same reasons. These exclusions were also reasonable from a socio-286 economic point of view: current pressure on land tends to shorten fallow length and 287 the pressure on organic resources greatly limit the possibility of crop residue recycling 288 (De Ridder and Van Keulen 1990).
- These conditions restricted the number of suitable studies for model calibration and testing to the ones indicated in Annex 1. A last restriction was applied by removing two outliers from Saria, which huge organic inputs (40 tDM ha⁻¹ y⁻¹) risked to bias the multiple regression (StatSoft 2006).

 $^{^{2}}$ For the same reason of data disparity, some readjustments (such depth weighed averages for the layers measured down to 20 cm) were sometimes necessary to obtain a standard depth of 20 cm for all data.

2.3.3 Model construction

Four management variables were available for all sites as potential explanatory variables for the effect of treatment (average yearly quantity of N, P, K and organic inputs applied over the last 5 years of treatment). A preliminary step-wise method of model selection, with the Akaike Information Criterion (AIC) as the selection criteria (Akaike 1973; StatSoft 2006), showed that only N, P and organic inputs were to be retained for their explicative power on ΔC .

300 The selection of environmental explanatory variables could not be so straightforward: due to a 301 split-plot kind of difference between stocks (Jones 1973; Kowal and Kassam 1978; Feller et 302 al. 1991; Feller 1995; Batjes 2001). These two variables were therefore retained in the model 303 as interacting with organic matter inputs. Other environmental variables, such as clay 304 mineralogy, soil thermic and water regimes, vegetation features and intensity of landscape 305 anthropisation that regulates OM availability and biological activity, also influence SOM dynamics. Nevertheless, their influence or variation across Senegal is weaker than that of 306 307 rainfall and clay content, or is correlated to these, and in any case, no spatially explicit data is 308 available at the scale of Senegal. The structure of the biophysical model, with the level of 309 intervention of each parameter, is summarized on

310 Figure 2.

311 Since this regression was based on clay contents ranging 3-15%, the model was not 312 deemed relevant for clay contents exceeding 20%, where SOC dynamics could be 313 significantly altered. Therefore, areas with such soil features (making up 15% of total 314 croplands in Senegal) were excluded from the analysis.

315 The final results could then be converted into SOC stocks changes using the same 316 depth as considered by the model (0-20 cm), and an average bulk density of 1.52 kg dm⁻³ 317 (average obtained over the same dataset that was used for model calibration). Assuming no 318 spatial variation of bulk density is a rough simplification, which is unavoidable as no spatially 319 explicit database currently exists for bulk density of soils over Senegal. However its impact is 320 likely to be limited, since the range of values for the clay content of the soils considered is 321 narrow, and since texture is a significant driver of soil density (Bernoux et al. 1998). For the 322 national potential for carbon storage, that is an overall estimate of carbon storage in Senegal 323 through agricultural intensification, the LOWINT and HIGHINT scenarios were processed 324 only on agricultural land as determined by MODIS data (DAAC 2002).

The rainfall dataset used to run the biophysical model is the yearly average between 1950-2000. This range is thought to be representative of the region, being wider than the drought that lasted from the 1960s to the beginning of the 1980s. Nevertheless, more recent
averages, or even forecasts such as the climate change scenario available in Hulme et al.
(2001) or in IPCC (2006), could be more relevant depending on the exact objectives of the
model runs.

The clay content dataset used to run the biophysical model was derived from Stancioff *et al.* (1986), taking the clay content of the barycentre of each texture class as a representative value for the class.

334

2.3.4 Model test

To assess the overall error of this statistical model, seven studies taking place in Senegal were retained. Their management parameters were entered into the model, together with the average rainfall corresponding to the study's period, and the clay content coming from the map. The model output was then compared with the actual measured values of SOC content change.

340

2.4 Socio-economic model

The socio-economic model makes a cost-benefit analysis of the agricultural intensification scenarios, taking into account both financial and environmental costs. As appropriate to this study, environmental costs accounted for were restricted to GHG emissions. The data on prices were updated for inflation when necessary (Diarisso 2004). All the intervening factors of the analysis are summarized on

346 Figure 3 and detailed hereafter.

347

2.4.1 Cost of transportation of mineral fertilizers

• Financial cost:

In Senegal, most mineral fertilizers are put together in Mbao (close to Dakar) by the *Industries Chimiques du Sénégal (ICS)*, and then distributed to local Senegalese agricultural markets by trucks (Industries Chimiques du Sénégal 2006). ICS uses the local phosphate deposits of Taïba, and imports nitrogen fertilizers and potash through the Dakar harbour.

For both financial and transportation costs, a map representing the distance of each grid element to Dakar by road was required. The itinerary chosen from each point to Dakar was computed using the following hypothesis:

- fertilizers are delivered from Dakar to agricultural markets by the fastest road;

357 - farmers from any grid element in Senegal then use the fastest road to fetch fertilizers
358 at the closest market.

- no transportation itinerary goes through Gambia, as long waiting times and passage fees to cross the two borders and the Gambia River are prohibitive.

The fastest roads were computed by attributing an average speed of 80 km h⁻¹ to journeys on paved roads, 40 km h⁻¹ to journeys on laterite roads, 20 km h⁻¹ to journeys on tracks, and 5 km h⁻¹ to journeys off tracks. These average speeds are broad assumptions, but they generated realistic itineraries. For all these computations, as for all the steps of this study that required a GIS, the ArcGis 9.0 software was used. The grid resolution used for cell-bycell model runs was 1 km².

For any point in Senegal, the financial cost of mineral fertilizers was therefore assumed to be the sum of its price in Dakar (*Source: Ministère de l'Agriculture et de l'Hydraulique - unpublished data, 2005*) and of the transportation price to the field. For this, the average price per ton \times kilometer was set at the same level as in Mali (UNCTAD 2000). It seemed relevant to Senegal as gasoline prices were comparable in the two neighbouring countries.

373 •

Environmental costs

374 For valuation of environmental externalities (costs of emissions), three sources of 375 emissions were identified: manufacturing (Vlek et al. 2004), sea transportation to Dakar 376 (ADEME and MIES 2007), and road transportation to the field (ADEME and MIES 2007). 377 As in N'Guessan (2003), a 25 tons truck was used as the reference, and its emissions were 378 increased by 25% from the original figures to account for the generally older trucks and ill-379 maintained roads of Senegal (Fall 1998). As Europe is by far the most important trade partner 380 of Senegal with more than 50% of Senegalese imports coming from Europe (Diarisso 2004), 381 the standard maritime route length used for these calculations was 4604 km, corresponding to 382 the standard maritime route from Hamburg to Dakar (Dataloy 2006).

383

2.4.2 Cost of application of mineral fertilizers

- Financial costs:
- 385

The labour cost of applying fertilizers were those estimated by Badiane *et al.* (2001).

• Environmental costs:

387 To compute the additional N_2O emissions resulting from the application of mineral 388 fertilizers, the IPCC methodology for national GHG inventories was applied (IPCC 2006).

389

2.4.3 Cost of manure collection and application

• Financial costs:

The collection and application of manure from stables was estimated at US\$ 1.42 per ton of manure (Badiane et al. 2001), and the transfer of fodder to stables at US\$ 24.86 per ton of manure (CIRAD et al. 2004).

394

395

• Environmental costs:

396 Emissions of CH₄ or CO₂ resulting from manure decomposition were assumed to be 397 the same whether livestock is stalled or not, and therefore did not influence the model. A 398 decrease in N₂O emissions from cattle manure when it is manually collected and spread 399 instead of being directly deposited on field by animals was accounted for, following the 400 standard IPCC methodology (IPCC 2006). The emissions from urine were overlooked, 401 thereby making the implicit assumption that its decomposition is independent of its 402 management (either directly deposited in the field or stored in stables and manually spread). 403 This assumption is probably reasonable since most of the nitrogen in urine is volatilized into 404 ammoniac notwithstanding its management (Fernandez-Rivera et al. 1995).

405

2.4.4 Additional revenue from increase in crop yields

406 Another important component of the cost-benefit analysis is the increased revenue 407 from improved yields. This requires (i) to evaluate how crop yield evolves under the different 408 scenarios, and (ii) to specify hypothesis on crop prices.

409

• Modelling of crop yield change:

410 As for SOC change, we used a statistical approach to model relative change in crop 411 yield. The same dataset as for SOC change prediction was used to calibrate a statistical 412 model. As the average yields of each treatment were not always available, the dataset was 413 here further restricted to 42 points coming from 12 different studies. The model building 414 methodology was the same as for the biophysical model, which again excluded K inputs from 415 explicative variables. There, such parameters as clay content and rainfall were not retained, as 416 they did not improve the model. This tends to show that, although clay content and rainfall 417 may play a major role in explaining absolute crop yield and absolute crop yield change, they 418 do not significantly influence relative crop yield change. However, as in McIntire and Powell 419 (1995), the inclusion of the square of some treatment parameters effectively improved the 420 model's precision, and was therefore retained to yield the model equation.

421 Regional statistics (source: Senegal, DSDIA-DAPS-MAE) were retained for the 422 reference yield from which the model computes yield changes (which imply both changed 423 benefits as the crops are sold, and changed costs as the harvesting time is changed in 424 proportion to the quantity of matter to be harvested). Spatialized crop yields were therefore425 obtained through the following equation:

426 $Y = Y_{R} \times \frac{100 + (\Delta Y_{T} - \Delta Y_{BAU})}{100}$

427 where Y is the yield (in kg ha⁻¹ y⁻¹), Y_R is the average regional yield (in kg ha⁻¹ y⁻¹), ΔY_T and 428 ΔY_{BAU} are the yield changes (in %) from a no-input treatment respectively due to the inputs in 429 the new practice and in the baseline scenarios, as computed by the statistical model.

430 • Crop prices:

For the producer prices of crops, as they were not directly available from any official and easily accessible source (Senegal is surprisingly not listed in the FAO statistical database on agricultural commodity prices), price ratios between crops were assumed to be the same as in Mali (FAO 2006), and these ratios were then combined with current market prices for millet and groundnut in Senegal (D. Baggio, pers. comm.) to get the producer price of each crop.

- 437 2.5 Model coupling
- 438

2.5.1 Hypothesis on carbon prices

439 The biophysical and economic models were coupled by attributing the 2005 average price of CERs from registered candidate CDM projects (25.8 US\$ tCeq⁻¹) to sequestered 440 carbon (Rosenzweig and Forrister 2006). This financial retribution was assumed to be 441 442 available only five years after the beginning of the project. The CERs were therefore 443 accordingly discounted, using the same discount rate as Sankhayan and Hofstad (2001) for Southern Senegal (5 % y⁻¹). The variable that represented the overall potential was then the 444 change of NPV over five years (using the same 5 % y⁻¹ discount rate) between the MAXINT 445 446 scenario and the baseline (BAU scenario).

447

2.5.2 Accounting for "avoided deforestation", and valuing it

Agricultural intensification is likely to decrease pressure on natural ecosystems such as forests and savannas, since demand for agricultural expansion fuelled by population growth is a major driver of deforestation in Senegal (Wood et al. 2004). Therefore, it is relevant to take into account the environmental benefits resulting from this decreasing pressure, especially for national-scale scenarios, which could be used as the practical tools of policies that bestow compensation for reductions in deforestation rates, such as "Compensated Reductions" (Santilli et al. 2005). For this assessment, land vulnerable to deforestation was defined as the nonagricultural land in the eco-regions undergoing agricultural expansion (Tappan et al. 2004). This included 5 eco-regions (Agricultural Expansion, Casamance, Niayes, Saloum Agricultural, and Southern Pastoral), over which non-agricultural land was divided between forests (17%) and savannas (83%). The area-weighed average change in carbon stocks (soil and aboveground biomass) for these two land uses when converted into cropland was estimated to 37.3 tC ha⁻¹ from data collected in Senegal (Liu et al. 2004).

To quantify the impact of agricultural intensification on deforestation, the underlying hypothesis was that agricultural intensification would not impact food and cash demand, meaning that each ton of increased crop production due to agricultural intensification preserves the amount of land necessary to produce one ton of crops in areas vulnerable to deforestation (1.6 ha). The result was then introduced into the cost-benefit analysis for the MAXINT scenario with the same valuation method for carbon as described in 2.4.

- 469 **3 Results**
- 470 3.1 Biophysical model
- 471

472 A multiple linear regression on the variables selected by the method exposed in 2.3.3 473 yielded the following equation:

3.1.1 Parameterization of equation for soil carbon stock change

- 474 $\Delta C = \text{organic inputs} \times (-0.56 0.036 \times \text{clay} + 0.0015 \times \text{rain} + 0.011 \times \text{P} 0.0031 \times \text{N}) + 0.023 \times \text{P} 0.0076 \times \text{N}$ (1) 475 with ΔC in mgC g⁻¹, organic inputs in tDM ha⁻¹ y⁻¹, rain in mm y⁻¹, P (P inputs) in kgP ha⁻¹ y⁻¹ 476 and N (N inputs) in kgN ha⁻¹ y⁻¹. This statistical model significantly explained data variations 477 (R² = 0.68***; n = 64) and the standard error was 0.87 mg g⁻¹ (1.32 tC ha⁻¹ when converted 478 into stocks).
- 479

3.1.2 Carbon storage found under the different scenarios

480 For the MAXINT scenario, the potential for carbon storage follows roughly the 481 southward rainfall gradient, with positive values for most of the country (

Figure 4a). The regional averages range from -3.2 to 15.6 tC/ha (Table 1), with highest increases in absolute amounts in Casamance (Ziguinchor & Kolda) while highest benefits of carbon sequestration on a surface basis were obtained in the Kolda and Tambacounda regions.

For the LOWINT and HIGHINT scenarios, the magnitude of SOC change is lower than for the MAXINT scenario. This way, the national potential for carbon storage was estimated to 0.648 MtC and 0.832 MtC for the LOWINT and HIGHINT (Table 2) scenarios, respectively.

489 *3.1.3 Model test*

490 Tested on an independent dataset over seven Senegalese sites, the predicted 491 values for SOC changes were not significantly different from the field measurements (p-value 492 = 0.13; n = 18). Nevertheless, the residual error was important (3.6 mg g⁻¹; 5.5 tC ha⁻¹ when 493 converted into stocks).

494

3.1.4 Sensitivity analysis to clay content

The biophysical model is quite sensitive to clay for the considered scenarios: a sensitivity analysis performed over a hundred points randomly selected on the map shows an average 92% of variation in SOC change for the MAXINT scenario when clay content is uniformly increased by 3 g 100g⁻¹ soil.

3.2 Cost-Benefit Analysis

500

3.2.1 Parameterization of the equation for yield increase

501 A multiple linear regression on the variables selected by the method exposed in 2.4.4 502 yielded the following equation:

503 $\Delta Y = (0.0424 - 0.00106 \times P) \times P + 0.00886 \times N + 0.00869 \times OI^2 + 0.0370 \times P \times OI - 0.00978 \times N \times OI$ (2) 504 with ΔY the difference in yield compared to a no-input scenario in %, OI the organic inputs in 505 tDM ha⁻¹ y⁻¹, P (P inputs) in kgP ha⁻¹ y⁻¹ and N (N inputs) in kgN ha⁻¹ y⁻¹. This statistical 506 model significantly explained the data variations (R² = 0.66***; n = 41) and the standard error 507 was 74%.

508 This error was increased to 94% when tested on an independent dataset over seven 509 Senegalese sites. The seven sites, though offering a good spatial coverage, all received the 510 same level of inputs (72.7 kgN ha⁻¹ y⁻¹ / 12.8 kgP ha⁻¹ y⁻¹ / 34.5 kgK ha⁻¹ y⁻¹ / 1.5 tDM ha⁻¹ y⁻¹ 511 of manure) which represents a potentially important shortcoming.

512

3.2.2 Cost-Benefit analysis for the different scenarios

513 The cost-benefits analysis for the scenarios was positive for most of currently 514 cultivated areas (except for the Northern Peanut Basin), even without accounting for carbon 515 (Figure 5c,d). Results for changes in greenhouse gas (GHG) balances (including all GHG and 516 SOC stock changes), are highly dependent on the considered scenario: the local, project-based 517 MAXINT scenario, yielded net sequestration for all areas where cultivation is practicable 518 (Central and Southern Senegal), whereas the regionally realistic HIGHINT scenario yielded 519 net emissions for most of these areas, except for the Old Peanut Basin and Western 520 Casamance (Figure 5a,b & Table 1).

521 3.3 <u>Model coupling</u>

522 When carbon is accounted as a potential source of revenue according to the method 523 previously described, the change of NPV of a farm over five years, due to the MAXINT 524 scenario, becomes positive in a limited area of the Old Peanut Basin (Figure 6).

525 3.4 <u>Avoided deforestation</u>

526 For both types of scenarios (regional-scaled such as LOWINT and HIGHINT, or local, 527 project-based such as MAXINT), the environmental benefits are low on a hectare basis, and 528 therefore unlikely to sweep the board. However, when the carbon sequestered through 529 "avoided deforestation" is accounted for, this pattern changes radically. For the MAXINT 530 scenario, the change -on a per hectare basis- in GHG balance compared to the baseline

- 531 becomes positive everywhere in Senegal, and its average over Senegal increases to 120 times
- the average without accounting for deforestation (Table 1 and Figure 7). For the HIGHINT
- 533 scenario the net fixation of GHG would increase by 161 folds (Table 2).

535 **4** Discussion

544

536 4.1 The limited national potential of increased fertilization for in-situ soil C accretion The average changes in SOC stocks in regions of the Old Peanut Basin such as Kaolack, 537 Fatick or Diourbel (1.78-3.89 tC ha⁻¹; Table 2), are close the 3.17 tC ha⁻¹ obtained by 538 539 Tschakert (2004) with the Century model for a similar treatment in this region. The North-540 South gradient in carbon storage potential for the MAXINT scenario (Figure 6a) is also consistent with the difference between these figures, and higher SOC increases $(+9.9 \text{ tC ha}^{-1})$ 541 542 measured by Manlay et al. (2002) in the Kolda region (for an administrative map of Senegal, 543 see Annex 2). These increases are important compared to the existing SOC stocks of 15-30 tC ha⁻¹ in the 0-40 cm layer (Woomer et al. 2004).

545 The present estimates of the national potential of carbon accretion of 0.65-0.83 MtC 546 for scenarios realistic at this scale (LOWINT and HIGHINT) would represent only 7 and 9% 547 of Senegal's 1994 GHG emissions from activities others than land-use change and forestry, 548 and 11 and 14% of GHG emissions from all activities, respectively (UNFCCC 2005 549 according to which land-use change and forestry were net C sinks in 1994). These scenarios 550 would also barely change the national SOC stocks estimated at 561 MtC by Henry et al. 551 (2009) for the 0-30 cm layer.

552 Estimates for C sequestration for the HIGHINT scenario would be even more modest 553 since it would represent 3 % of Senegal's 1994 GHG emissions from activities others than 554 land-use change and forestry, and 5% of GHG emissions from all activities. Estimates for C 555 accretion are more difficult to relate with the estimate of Senegal's potential of carbon storage 556 made by Parton et al. (2004) under their "improved management practices" scenario. Their 557 116 MtC potential indeed includes the potential gains from the biomass compartment, and 558 also those resulting from improved management of pastures and forests. This wide difference 559 is nevertheless surprising since these authors estimate that agriculture would account for 50% 560 of this potential, and that in the Intensive Agricultural Region, 90% of the potential comes 561 from increased SOC. We found several explanations for this discrepancy. The main one is 562 that the storage potential in our scenario forbids dramatic changes in management practices: it 563 limits the increase in manure application to the gains in manure from improved collecting 564 efficiency where Parton et al. (2004) assume a doubling of fallow time, coupled with the application of manure and household wastes (unknown quantities). Their estimate of an 565 average potential storage of 17.1 tC ha⁻¹ in the Intensive Agricultural Region is indeed higher 566 567 than the value given by Tschakert (2004) for an "optimal" intensification, which requires among others 9 tons of manure per hectare, which is already six times the maximal manure available at regional scale in our HIGHINT scenario. For such scenarios with higher inputs in manure such as the MAXINT scenario, our model indeed yields much higher estimates.

The second source of difference comes from the figures for agricultural cover: as the present model is spatially explicit, it is bound to use spatialized data (DAAC 2002) which adds up to 17% of total Senegal under cropping, whereas Parton *et al.* (2004) use the estimate of 21% from Tappan *et al.* (2004), which is more specific to Senegal, but unfortunately nonspatialized. Moreover, the range of application was limited here to sandy soils (clay content < 20%), which excluded 15% of total cropland. Overall, the total surface of cropland on which their estimate is based is therefore 45% higher than the one used in the present study.

578

4.2 Avoided deforestation: a key element to improved the C balance of fertilization

579 The model runs for carbon balances are widely modified when accounting for 580 deforestation (Table 1; Table 2). In this case estimates for C sequestration for the HIGHINT 581 scenario would rise to 561% of Senegal's 1994 GHG emissions from activities others than 582 land-use change and forestry, and to 872% of GHG emissions from all activities (UNFCCC 2005). This is consistent with the results of Liu et al. (2004), who found that of the 583 52 MgC ha⁻¹ of carbon stocks lost between 1900 and 2000 in south Central Senegal, 88% 584 585 came from live biomass, and only 12% from litter and topsoil SOC (the 0-20 cm depth was 586 the only soil compartment included in the calculations). Such results are consistent with the 587 recent regain of interest for the issue of carbon emissions from deforestation (The World 588 Bank 2008; UNFCCC 2008).

589

4.3 Cost-benefit analysis and carbon sequestration (net of additional GHG emissions)

590

4.3.1 Profitability of carbon sequestration

The models predict the intensification scenarios to be profitable in almost all Central and Southern Senegal (Figure 5c,d). However, they also show that intensification does not always coincide with carbon sequestration: for the HIGHINT scenario, the GHG balance is lower (more emissions) than the baseline for all areas but the central Senegal (which has low emission levels due to its proximity to Dakar) and Ziguinchor and Kolda due to their high potential for carbon storage). In Eastern Senegal and most of southern Senegal, emissions due to fertilizer use override gains from carbon storage (Figure 5a).

598 Facing such positive figures for profitability, one could wonder why farmers do not 599 intensify. The reason probably dwells in the time-discrepancy between the time when money 600 is needed to buy fertilizer, and the time when benefits from yield increases are reaped. In a 601 context of lacking credit possibilities, and chaotic supply, this has been shown to be an 602 important obstacle (Vlek 1990; Tschakert 2004). Climate variability seriously affects the 603 impact of fertilization on crop yields, and may also be part of the answer (see 4.4.4).

One could therefore look at the income generated by carbon retribution as a source of savings that that would enable farmers to buy fertilizers at the beginning of the growing season. However as Tschakert (2004), in the present study these retributions are insufficient since the yearly gains from carbon never compensate for even the lowest yearly price of fertilizers as predicted by the model (US\$ 28 ha⁻¹ y⁻¹, data not shown).

609

4.3.2 Transportation costs

Transportation costs would be all the more relevant for Casamance where their high level significantly increases the local price of fertilizers (from US\$ 30 ha⁻¹ y⁻¹ close to Dakar to US\$ 41 ha⁻¹ y⁻¹ in Western Casamance for recommended amounts of fertilizer). This is why, as highlighted in other studies about Africa or Indonesia, a priority for better access to fertilizers is the development of better transportation infrastructures (Vlek 1990; Isherwood 2000; Smaling et al. 2006).

616 A potential pitfall in these estimates of transportation costs is that they are independent 617 of the total flow of fertilizer. Possible efficiency gains linked to large increases in fertilizer 618 flow (through improvements in the road network infrastructure for example) are thereby ruled 619 out.

620

4.4 Methodological considerations

621

4.4.1 Resolution of spatial data

622 The sources of spatial data are of different resolutions, but those that would have to be 623 refined in priority may depend on the kind of scenario. For national-scale assessments, the "socio-economic" data such as the desirable quantity of fertilizer inputs are generally given at 624 625 a much coarser resolution than the "environmental" data (region or department versus square 626 kilometre). The priority would therefore be to refine the scenarios, in particular by accounting 627 for rice cultivation and fallowing, and by taking better care of the feasibility of each scenario. 628 Indeed, fodder and labour availability for animal stalling during an entire season was taken for granted here as grazing animals are only able to consume 20% of residues (the rest being 629 630 trampled on, polluted by excrements, burnt, or carried away by termites) (Bosma et al. 1999) and as stabled animals eat less (more than 33% so according to CIRAD et al. 2004). However,
local patterns could countervail these broad generalities.

633 On the contrary, for evaluations of local project feasibility, it is assumed that such 634 "socio-economic" data could be accurately forecast as project managers know what they 635 intend to do with the land. Clay content however still varies at very small scales due to 636 erosion, parent material, and deposition (Cerri et al. 2004). Therefore, refining first the clay 637 content map would therefore be most advisable.

638

4.4.2 Organic matter displacement

An important hypothesis of this paper is that the organic matter used to increase SOC
in agricultural fields would not have also done so in their original location. Otherwise, the
environmental benefit to the atmosphere would be null.

642 For manure naturally deposited on pastures, there are reasons to think this hypothesis 643 is valid (Poulton 1996), even if no experimental demonstration in the West African context 644 has been identified to support it. Indeed, manure patchily deposited by grazing animals is 645 probably much less efficiently incorporated to the soil than manure collected from stables and 646 evenly spread over the field (Bationo et al. 1995). Also, organic inputs would probably be 647 more protected from decomposition in depleted agricultural soils than in pastures where the 648 threshold of SOC protection by clays may already be attained (Breman 1998; Six et al. 2002). 649 Finally, manure has been shown to actually increase SOC content only if associated with mineral inputs in South Senegal (Manlay et al. 2002), something unlikely to happen in 650 651 pastures unless they are also fertilized with synthetic inputs.

652

4.4.3 Yield model

The model used to compute yield increases is very simple. Whereas more complex models can offer better predictions (Christianson et al. 1990; Aune and Lal 1995; McIntire and Powell 1995), they are often too demanding in parameters to be used on such a broad scale as this study's and in developing countries were data is scarce. Therefore, many well described effects of treatments or the environment on yields are either very summarily or not at all included in this model.

Nevertheless, it is satisfying to verify that where Baidu-Forson and Bationo (1992) equate a 17.5 kgP ha⁻¹ treatment to a 8.7 kgP.ha⁻¹ treatment complemented by 5 tDM ha⁻¹ of manure in terms of yields, the model predicted similar yield increases of respectively 95% and 91% for these two treatments compared to a no-input treatment.

4.4.4 The risk issue

664 The Net Present Value of a practice is not the only parameter that will determine its 665 chances of adoption (Tschakert 2004). As underscored by Bartel (2004), the most important 666 for smallholders may be to minimize risk before maximising profits. It was therefore 667 interesting for us to assess the risk associated to each practice. Badiane et al. (2001) found 668 that the combination of organic and mineral inputs decreased the risk coming from yield 669 variability. Therefore, the different scenarios are likely to decrease risk in the areas where 670 they increase the NPV, but some critical elements were lacking to perform a thorough risk 671 analysis.

672 Another risk is that of non-permanence of carbon storage, although this risk falls on 673 the eventual recipient of CERs, who is not necessarily the farmer himself. Although part of it 674 is included in the diminished price of the CERs themselves compared to other carbon credits 675 (Rosenzweig and Forrister 2006), the market is probably not able to cope with all risks, 676 especially such complex ones as an eventual large scale SOC mineralization due to increased 677 temperature (Ringius 2002; Bellamy et al. 2005) in the most vulnerable continent to climate 678 change (Boko et al. 2007). On the other hand, as mentioned by Olsson and Ardo (2002), the 679 risk of future release of sequestered carbon is less important in agriculture than in forestry 680 since (1) SOM preservation is the interest of the farmer, as it contributes to soil fertility, and 681 (2) it is less subject to external hazards such as fire.

682

4.4.5 Model imprecision

683 The standard errors of both SOC and yield prediction models, as assessed on our limited independent dataset, are quite high (± 5.5 tC ha⁻¹ for SOC change, and $\pm 94\%$ for yield 684 685 increase). This could be expected as it includes both the inherent imprecision of the statistical model and the one arising from cartographic data, especially high for clay content which 686 687 varies on a much smaller scale than the map is able to represent (Cerri et al. 2004). Due to the 688 non-homogeneous distribution of test points both spatially over Senegal, and qualitatively 689 over the environmental parameters (clay content and rainfall), these error estimates have to be 690 considered with caution.

691 Overall, the model error seems nevertheless low enough to provide broad indications 692 on the most adequate locations for carbon sequestration projects. It would be interesting to 693 perform the same kind of analysis with some more standard models such as Century (Parton 694 et al. 1987). Indeed, it is difficult to get a sense of the overall precision of these from existing 695 studies: Cerri *et al.* (2004) obtained a standard error as low as $\pm 17\%$ for forest conversion to pasture in Brazil, but this does not include the uncertainty stemming from the use of mapped
data. For more diverse and more specific kinds of managements such as those studied here,
the standard error is often left untested (Tschakert 2004; Woomer et al. 2004).

699

4.4.6 Time window considered in the cost-benefit analysis

The time window considered for the cost-benefit analysis is five years, which makes sense as a minimal time to get credits from carbon sequestration projects. But while future financial benefits after five years may be irrelevant due to high discount rates, this may not be easily replicable for the environmental balance: indeed, the intensification treatments would have to be maintained overtime, thereby emitting always more greenhouse gas for the same amount of SOC storage, unless integrated, nutrient-conservative management strategies are identified and adopted.

708 **5** Conclusion

709 This study shows that the potential for carbon storage through the proposed scenarios 710 applicable at national scale is relatively low (0.65-0.83 MtC) compared to previous estimates. 711 Large inputs of manure can change the picture (MAXINT scenario) and make it coincide 712 better with existing local estimates, but these would not be available at a larger scale than a 713 few well-endowed farms, unless livestock resources are increased. And in that case, the 714 methane emissions from the additional livestock would probably override the amounts of 715 carbon stored in the soil. However, accounting for "avoided deforestation" as a likely side-716 effect of agricultural intensification completely changes the picture, with carbon balances 717 positive everywhere, and amounting to about a hundred-fold increase compared to the 718 original figures, thus representing a major source of revenue with the proposed valuation 719 method.

Through this valuation method, the internalisation of a restricted set of environmental costs and benefits, namely GHG emissions, was attempted. But the considered scenarios are likely to impact other aspects of the environment: mineral fertilizer application carries the risk of water table pollution and soil acidification, and manure application can help prevent soil drought, soil acidification, and erosion. The further internalisation of these aspects would be of great interest.

The means of reducing greenhouse gas emissions explored in this study's scenarios are far from being comprehensive. At least three complementary ways of sequestering carbon would deserve an explicit role in the models and scenarios: agroforestry, intercropping and no tillage cropping schemes. Other more indirect ways could also be considered, such as methane capture from manure decomposition in stables. All these unexplored potentials would have to be analysed in depth before any definitive conclusion on the carbon sequestration potential of agriculture as a whole can be drawn for Senegal.

Inclusion of considerations out of the Kyoto protocol scope such as resource use
efficiency (carbon intensity), per capita emissions or historical responsibility could also
broaden the view on how Senegalese farming systems interact with global change.

736

737 Acknowledgments

The authors would like to thank Alioune Ka and Jacques-André Ndione from the Centre de
Suivi Ecologique (Dakar), whose active collaboration and goodwill made this work possible.
For their more punctual but nonetheless essential contributions to this work, the authors are

- 741 very grateful to Vincent Eschenbrenner, Francis Ganry, Robert Oliver, David Baggio, Marion
- 742 Houles, Amadou Dieye, and Gray Tappan.
- 743 This work received financial support from the Institut de Recherche pour le Développement
- 744 (IRD), the Ecole Nationale du Génie Rural, des Eaux et des Forêts (AgroParisTech-
- 745 ENGREF), the Centre International de Recherche sur l'Environnement et le Développement
- 746 (CIRED), and the Centre de Suivi Ecologique (CSE). It was implemented within the "Carbon
- benefits project, monitoring measuring and modelling" project of the Global Environmental
- Fund³.
- 749

³ Still in the pipeline on 02/12/2008

Annex 1 – Sources used for building the soil carbon accretion and yield increase models (summary
 table)

Study	Site	Use	Number of	Yearly rainfall	Clay content
			data points	(mm y ⁻¹)	(%)
(Fernandes 1999)	Bambey, Senegal	Calibration	2	446	4
(Pieri 1979; Pieri 1992)	Bambey, Senegal	Calibration	1	412	3
(Richard and Djoulet 1985)	Bebedjia, Chad (Trial N)	Calibration	3	1120	9
(Richard and Djoulet 1985)	Bebedjia, Chad (Trial R – Serial 64)	Calibration	3	1120	10
(Richard and Djoulet 1985)	Bebedjia, Chad (Trial R – Serial 65)	Calibration	3	1120	10
(Richard and Djoulet 1985)	Bebedjia, Chad (Trial S1 – Serial 68)	Calibration	5	1120	11
(Richard and Djoulet 1985)	Bebedjia, Chad (Trial X – Serial 1968)	Calibration	6	1120	10
(Sarr 1981)	Nioro du Rip, Senegal	Calibration	1	718	11
(Rabot 1984)	Nioro du Rip, Senegal	Calibration	1	718	9
(Bationo et al. 1993; Bationo and Buerkert 2001)	Sadoré, Niger	Calibration	1	560	3
(Agbenin and Goladi 1997)	Samaru, Nigeria	Calibration	5	1062	10
(Jones 1971)	Samaru, Nigeria (DNPK	Calibration	S	1062	10
(Jones 1971)	Samaru, Nigeria (MVR trial)	Calibration	3	1062	10
(Jones 1971)	Samaru, Nigeria (PFM trial)	Calibration	3	1062	10
(Singh and Balasubramanian 1979), (Ogunwole and	Samaru, Nigeria	Calibration			
Ogunleye 2005)			5	1062	10
(Hien 2004)	Saria, Burkina Faso (EEC trial)	Calibration	7	794	14
(Pieri 1992; Hien 2004)	Saria, Burkina Faso (EFF trial –	Calibration			
	monoculture)		3	765	14
(Pichot et al. 1981)	Saria, Burkina Faso (EFF trial –	Calibration			
	monoculture)	Tash	4	/65	14
(Feller et al. 1981)	Bambey, Senegal	Test	7	412	3,3*
(Rabot 1984)	Missirah, Senegal	Test	1	800	23,1*
(Manlay et al. 2002)	Sare Yorobana, Senegal	Test	5	960	3,3*
(Rabot 1984)	Sinthiou, Senegal	Test	1	900	9,5*
(Badiane 1993)	Thilmakha, Senegal	Test	1	363	3,3*
(Cissé 1986; Pieri 1992)	Thilmakha, Senegal	Test	2	344	3,3*
(Rabot 1984)	Velingara, Senegal	Test	1	1150	9,5*

752 * Not a direct measurement. Clay content estimated from the clay content map of Senegal

- Annex 2. Administrative map of Senegal (source: http://www.culture.gouv.sn/)
- 755 See figures file for the map

757	Bibliography
758	ADEME and MIES (2007) Guide des facteurs d'émission - Version 5.0. In: Bilan Carbone
759	Agbenin, J. O. and J. T. Goladi (1997) Carbon, nitrogen and phosphorus dynamics under continuous cultivation as influenced by farmyard
760	manure and inorganic fertilizers in the savanna of northern Nigeria. Agriculture Ecosystems & Environment 63(1): 17-24
761	Akaike, H. (1973) Information theory and an extension of the maximum likelihood principle. Paper presented at Second International
762	Symposium on Information Theory, Budapest, 1973
763	Allard, J. L., Y. Bertheau, et al. (1983) Ressources en résidus de récolte et potentialités pour le biogaz au Sénégal. Agronomie Tropicale
764	38(3): p.213-221
765	Aune, J. B. and R. Lal (1995) The Tropical Soil Productivity Calculator - A Model for Assessing Effects of Soil Management on
766	Productivity. In: R. Lal and B. A. Stewart (ed) Soil Management: Experimental Basis for Sustainability and Envrironmental
767	Quality.
768	Badiane, A., A. Faye, et al. (2001) Use of compost and mineral fertilizers for millet production by farmers in the semiarid region of Senegal.
769	Biological Agriculture and Horticulture 19(3): 219-230
770	Badiane, N. (1993) Le statut organique d'un sol sableux de la zone Centre-Nord du Sénégal. Sciences Agronomiques. Nancy, Institut
771	National Polytechnique de Lorraine
772	Baidu-Forson, J. and A. Bationo (1992) An economic evaluation of a long-term experiment on phosphorus and manure amendments to sandy
773	Sahelian soils: using a stochastic dominance model. Fertilizer Research 33: 193-202
774	Bartel, P. (2004) Soil carbon sequestration and its role in economic development: a donor perspective. Journal of Arid Environments 59(3):
775	643-644
776	Bationo, A. and A. Buerkert (2001) Soil organic management for sustainable land use in Sudano-Sahelian West Africa. Nutrient Cycling in
777	Agroecosystems 61: 131-142
778	Bationo, A., A. Buerkert, et al. (1995) A critical review of crop-residue use as soil amendment in the West African semi-arid tropics. In: J.
779	M. Powell, S. Fernandez Rivera, T. O. Williams and C. Renard (ed) Livestock and sustainable nutrient cycling in mixed farming
780	systems of sub-saharan Africa. ILCA, Addis Ababa
781	Bationo, A., C. B. Christianson, et al. (1993) The Effect of Crop Residue and Fertilizer Use on Pearl-Millet Yields in Niger. Fertilizer
782	Research 34(3): 251-258
783	Batjes, N. H. (2001) Options for increasing carbon sequestration in West African soils: An exploratory study with special focus on Senegal.
784	Land Degradation & Development 12(2): 131-142
785	Bellamy, P. H., P. J. Loveland, et al. (2005) Carbon losses from all soils across England and Wales 1978-2003. Nature 437(7056): 245-248
786	Bernoux, M., D. Arrouays, et al. (1998) Bulk densities of Brazilian Amazon soils related to other soil properties. Soil Science Society of
787	America Journal 62(3): 743-749
788	Bernoux, M., F. C., et al. (2006) Soil carbon sequestration. In: E. Roose, R. Lal, F. C., B. Barthès and B. A. Stewart (ed) Erosion & Carbon
789	Dynamics. CRC
790	Boko, M., I. Niang, et al. (2007) Africa. In: M. L. Parry, O. F. Canziani, J. P. Palutikof, P. J. van der Linden and C. E. Hanson (ed) Climate
791	Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the
792	Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge UK

- Bosma, R., M. Bengaly, et al. (1995) Pour un système durable de production au Mali-Sud : accroître le rôle des ruminants dans le maintien
 de la matière organique des sols. In: J. M. Powell, S. Fernandez Rivera, T. O. Williams and C. Renard (ed) Livestock and
- 795 sustainable nutrient cycling in mixed farming systems of sub-saharan Africa. ILCA, Addis Ababa
- Bosma, R. H., M. Bos, et al. (1999) The promising impact of ley introduction and herd expansion on soil organic matter content in southern
 Mali. Agricultural Systems 62(1): 1-15
- Breman, H. (1998) L'intensification agricole au Sahel : vouloir c'est pouvoir ! In: H. Breman (ed) L'intensification agricole au Sahel.
 Karthala, Paris
- 800 Cerri, C. E. P., M. Easter, et al. (2007) Predicted soil organic carbon stocks and changes in the Brazilian Amazon between 2000 and 2030.
 801 Agriculture, Ecosystems and Environment 122(1): 58-72
- 802 Cerri, C. E. P., K. Paustian, et al. (2004) Combining Soil C and N Spatial Variability and Modeling Approaches for Measuring and
 803 Monitoring Soil Carbon Sequestration. Environmental Management 33(1): 274-288
- 804 Cerri, C. E. P., K. Paustian, et al. (2004) Modeling changes in soil organic matter in Amazon forest to pasture conversion with the Century
 805 model. Global Change Biology 10(5): 815-832
- 806 Christianson, C. B., A. Bationo, et al. (1990) Fate and efficiency of N fertilizers applied to pearl millet in Niger. Plant and Soil 125: 221-231
- 807 CIRAD, GRET, et al. (2004) Mémento de l'Agronome. CIRAD, Paris
- 808 Cissé, L. (1986) Etude des effets d'apports de matière orgnique sur les bilans hydriques et minéraux et la production de mil et de l'arachide
 809 sur un sol sableux dégradé du Centre-Nord du Sénégal. Nancy, Institut National Polytechnique de Lorraine
- 810 COFACE (2006) Country Risk Outlook 2006. In. COFACE, East Windsor, NJ (USA)
- 811 Corbeels, M., E. Scopel, et al. (2006) Soil carbon storage potential of direct seeding mulch-based cropping systems in the Cerrados of Brazil.
 812 Global Change Biology 12: 1773-1787
- 813 DAAC (2002) MODIS/Terra Land Cover Type 96-Day L3 Global 1km ISIN Grid. http://edcdaac.usgs.gov/modis/mod12q1.asp. Cited
- 814 Dataloy (2006) Dataloy website. <u>http://www.dataloy.com/</u>. Cited 18 May 2006
- Be Ridder, N. and H. Van Keulen (1990) Somme aspects of the role of organic matter in sustainable intensified arable farming systems in the
 West African semi-arid-tropics (SAT). Fertilizer Research 26(1-3): p.299-310
- 817 Diarisso, S. (2004) Situation économique et sociale du Sénégal Edition 2004. In. MEF/DPS, Dakar
- Bugué, P., E. Vall, et al. (2004) Evolution des relations entre l'agriculture et l'élevage dans les savanes d'Afrique de l'Ouest et du Centre.
 Oléagineux, Corps Gras, Lipides 11(4): 268-76
- Easter, M., K. Paustian, et al. (2007) The GEFSOC soil carbon modelling system: a tool for conducting regional-scale soil carbon inventories
 and assessing the impacts of land use change on soil carbon. Agriculture Ecosystems and Environment 122(1): 13-25
- 822 Elberling, B., A. Toure, et al. (2003) Changes in soil organic matter following groundnut-millet cropping at three locations in semi-arid
- 823 Senegal, West Africa. Agriculture Ecosystems and Environment 96(1-3): 37-47
- 824 Fall, E. H. B. (1998) Etude des impacts des activités du PST II sur l'environnement. In. W. Bank. World Bank, Washington
- 825 FAO (2006) FAOSTAT data. http://faostat.fao.org/. Cited February 2006
- 826 Feller, C. (1995) La matière organique dans les sols tropicaux à argile 1:1 Recherche de compartiments organiques fonctionnels Une
- 827 approche granulométrique. Institut de géologie. Strasbourg, Université Louis Pasteur
- 828 Feller, C., E. Fritsch, et al. (1991) Effet de la texture sur le stockage et la dynamique des matières organiques dans quelques sols ferrugineux
- 829 et ferrallitiques (Afrique de l'Ouest, en particulier). Cahiers de l'ORSTOM, série Pédologie 26(1): 25-36

830 Feller, C., F. Ganry, et al. (1981) Décomposition et humidification des résidus végétaux dans un agro-système tropical. I. Influence d'une 831 fertilisation azotée (urée) et d'un amendement organique (compost) sur la répartition du carbone et de l'azote dans différents 832 compartiments d'un sol sableux. II. Décomposition des résidus végétaux (compost) pendant une saison des pluies dans un sol 833 sableux. Agronomie Tropicale 36(1): p.9-25 834 Fernandes, P. (1999) Effet des modes de gestion des terres sur les termes et l'évolution de leur bilan organique et biologique : conséquences 835 pour une production céréalière soutenue - Cas du Sénégal. Sciences Agronomiques. Nancy, Institut National Polytechnique de 836 Lorraine 837 Fernandez-Rivera, S., T. O. Williams, et al. (1995) Faecal excretion by ruminants and manure availability for crop production in semi-arid 838 West Africa. In: J. M. Powell, S. Fernandez Rivera, T. O. Williams and C. Renard (ed) Livestock and sustainable nutrient cycling 839 in mixed farming systems of sub-saharan Africa. ILCA, Addis Ababa 840 Henry, M., R. Valentini, et al. (2009) Soil carbon stocks in ecoregions of Africa. Biogeosciences Discussions 6: 797-823 841 Hien, E. (2004) Dynamique du carbone dans un Acrisol ferrique du Centre Ouest Burkina : Influence des pratiques culuturales sur le stock et 842 la qualité de la matière organique. Ecole Doctorale Biologie Intégrative. Montpellier, Ecole Nationale Supérieure Agronomique 843 de Montpellier 844 Hulme, M., R. Doherty, et al. (2001) African climate change: 1900-2100. Climate Research 17(2): 145-168 845 Industries Chimiques du Sénégal (2006) ICS website. http://www.ics.sn/. Cited 12 Jun 2006 846 International Fertilizer Industry Association (2003) IFA website. http://www.fertilizer.org/ifa/. Cited 6 May 2006 847 IPCC (1996) Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories - Workbook (Volume 2). In: 848 IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories. In. IPCC 849 IPCC (2006) The IPCC Data Distribution Centre. http://ipcc-ddc.cru.uea.ac.uk/. Cited 5 May 2006 850 Isherwood, K. F. (2000) Mineral fertilizer distribution and the environment. In. IFA and UNEP, Paris 851 Jones, M. J. (1971) The maintenance of soil organic matter under continuous cultivation at Samaru, Nigeria. Journal of Agricultural Science 852 77: 473-482 853 Jones, M. J. (1973) The organic matter content of the savanna soils of West Africa. Journal of Soil Science 24(1): 42-53 854 Kante, B. (1997) Communication initiale du Sénégal à la Convention-Cadre des Nations-Unies sur les Changements Climatiques 855 (CCNUCC). In. Ministère de la protection de la nature - Direction de l'environnement, Dakar 856 Kelly, V., B. Diagana, et al. (1996) Cash Crop and Foodgrain Productivity in Senegal: Historical View, New Survey Evidence, and Policy 857 Implications. In. USAID 858 Kowal, J. M. and A. H. Kassam (1978) Agricultural Ecology of Savanna - A Study of West Africa. Clarendon Press, Oxford 859 Le Fur, A. (2000) Espace agricole. La construction de l'espace sénégalais depuis l'indépendance, 1960-2000. M.-C. Cormier, C. Gueye, A. 860 Lericollais and S. M. Seck. Bondy, IRD 861 Liu, S., M. Kaire, et al. (2004) Impacts of land use and climate change on carbon dynamics in south-central Senegal. Journal of Arid 862 Environments 59(3): 583-604 863 Manlay, R. J., J.-L. Chotte, et al. (2002) Carbon, nitrogen and phosphorus allocation in agro-ecosystems of a West African savanna III. Plant 864 and soil components under continuous cultivation. Agriculture Ecosystems & Environment 88(3): 249-269 865 Manlay, R. J., A. Ickowicz, et al. (2004) Spatial carbon, nitrogen and phosphorus budget in a village of the West African savanna - II. 866 Element flows and functioning of a mixed-farming system. Agricultural Systems 79(1): 83-107

- 867 McIntire, J. and J. M. Powell (1994) African semi-arid tropical agriculture cannot grow without external inputs. Paper presented at Livestock 868 and Sustainable Nutrient Cycling in Mixed Farming Systems of Sub-Saharan Africa. Proceedings of an International Conference, 869 Addis Ababa, Ethiopia, 1994 870 McIntire, J. and J. M. Powell (1995) African semi-arid tropical agriculture cannot grow without external inputs. In: J. M. Powell, S. 871 Fernandez Rivera, T. O. Williams and C. Renard (ed) Livestock and sustainable nutrient cycling in mixed farming systems of 872 sub-saharan Africa. ILCA, Addis Ababa 873 Milne, E., R. Al-Adamat, et al. (2007) National and sub national assessments of soil organic carbon stocks and changes: the GEFSOC 874 modelling system. Agriculture Ecosystems and Environment 122(1): 3-12 875 Ministère de l'Agriculture et de l'Hydraulique (2004) Rapport annuel 2004 - Direction de l'élevage. In. Senegal. Ministère de l'Agriculture et 876 de l'Hydraulique, Dakar 877 N'Guessan, N. (2003) Amélioration du transport de transit en Afrique de l'Ouest. In: Première session du comité intergouvernemental 878 préparatoire de la conférence ministérielle internationale sur la coopération en transport de transit. UNCTAD, New York
- 879 Ogunwole, J. O. and P. O. Ogunleye (2005) Influence of long-term application of organic and mineral fertilizers on quality of a savanna
 880 Alfisol. Journal of Sustainable Agriculture 26(3): 5-14
- 881 Olsson, L. and J. Ardo (2002) Soil carbon sequestration in degraded semiarid agro-ecosystems Perils and Potentials. Ambio 31(6): 471-477
- Palm, C. A., K. E. Giller, et al. (2001) Management of organic matter in the tropics: translating theory into practice. Nutrient Cycling in
 Agroecosystems 61: 63-75
- Pansu, M. and J. Gautheyrou (2006) Handbook of Soil Analysis. Mineralogical, Organic and Inorganic Methods. Springer, Berlin,
 Heidelberg, New York
- Parton, W., G. Tappan, et al. (2004) Ecological impact of historical and future land-use patterns in Senegal. Journal of Arid Environments
 59(3): 605-623
- Parton, W. J., D. S. Schimel, et al. (1987) Analysis of factors controlling soil organic matter levels in Great Plains grasslands. Soil Science
 Society of America Journal 51(5): 1173-1179
- Pichot, J.-P., M. P. Sedogo, et al. (1981) Evolution de la fertilité d'un sol ferrugineux tropical sous l'influence de fumures minérales et
 organiques. Agronomie Tropicale 36(2): p.122-133
- 892 Pieri, C. (1979) Etude de la composition de la solution d'un sol sableux cultivé au Sénégal à l'aide de capteurs en céramique poreuse.
 893 Agronomie Tropicale 34(1): p.9-22
- 894 Pieri, C. (1992) Fertility of soils A future for farming in the West African savannah. Springer-Verlag, Berlin
- Pieri, C. (1995) Long term soil management experiments in semiarid francophone Africa(ed) In : Soil management. Experimental basis for
 substainability and environmental quality = [Gestion du sol. Base expérimentale pour une durabilité et une qualité de
- 897 l'environnement]/Lal R., Stewart B.A. Boca Raton : CRC Press, 1995. (Advances in Soil Science, ISSN 0176-9340).
- 898 Poulton, P. (1996). In: D. S. Powlson (ed) Evaluation of Soil Organic Matter Models Using Existing Long-Term Datasets. Springer-Verlag,
 899 Berlin
- 900 Powell, J. M. and T. O. Williams (1995) An overview of mixed farming systems in sub-Saharan Africa. In: J. M. Powell, S. Fernandez
- Rivera, T. O. Williams and C. Renard (ed) Livestock and sustainable nutrient cycling in mixed farming systems of sub-saharan
 Africa, ILCA, Addis Ababa
- 903 Rabot, C. (1984) Vingt années de successions de cultures dans la moitié Sud du Sénégal Impacts écologiques. Ecologie Tropicale.
- 904 Montpellier, Université des sciences et techniques du Languedoc

905	Richard, L. and B. Djoulet (1985) La fertilité des sols et son évolution en zone cotonnière du Tchad. In: In : Coton et Fibres Tropicales.
906	Série Documents, Etudes et Synthèses = ISSN 0010-9711 (1985)n°6
907	Ringius, L. (2002) Soil carbon sequestration and the CDM: Opportunities and challenges for Africa. Climatic Change 54(4): 471-495
908	Rosenzweig, R. and D. Forrister (2006) State and Trends of the Carbon Market 2006 - Summary. In: Carbon Finance Business. W. Bank.
909	Natsource LLC, Washington
910	Sanchez, P. A. (1997) Soil Fertility Replenishement in Africa: An Investment in Natural Ressource Capital. In: Buresh (ed) Replenishing
911	Soil Fertility in Africa. American Society of Agronomy and Soil Science Society of America, Madison
912	Sankhayan, P. L. and O. Hofstad (2001) A village-level economic model of land clearing, grazing, and wood harvesting for sub-Saharan
913	Africa: with a case study in southern Senegal. Ecological Economics 38(3): 423-440
914	Santilli, M., P. Moutinho, et al. (2005) Tropical deforestation and the Kyoto Protocol. Climatic Change 71(3): 267-276
915	Sarr, P. L. (1981) Analyse des effets induits par l'intensification des cultures sur quelques caractéristiques physico-chimiques d'un sol
916	ferrugineux tropical du Sénégal (Nioro du Rip). Agronomie-Pédologie. Montpellier, Université des sciences et techniques du
917	Languedoc
918	Schlesinger, W. H., J. Palmer Winkler, et al. (2000) Soils and the global carbon cycle. In: T. M. L. Wigley and D. S. Schimel (ed) The
919	Carbon Cycle. Cambridge University Press, Cambridge
920	Singh, L. and V. Balasubramanian (1979) Effects of continuous fertilizer use on a ferruginous soil (haplustalf) in Nigeria. Experimental
921	Agriculture 15: 257-265
922	Six, J., R. T. Conant, et al. (2002) Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils. Plant and Soil
923	241(2): 155-176
924	Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer
924 925	Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006
924 925 926	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie,
924 925 926 927	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings
924 925 926 927 928	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006
924 925 926 927 928 929	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. <u>http://www.statsoft.com/textbook/stathome.html</u>. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462
 924 925 926 927 928 929 930 	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. <u>http://www.statsoft.com/textbook/stathome.html</u>. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank
 924 925 926 927 928 929 930 931 	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. <u>http://www.statsoft.com/textbook/stathome.html</u>. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117
 924 925 926 927 928 929 930 931 932 	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. <u>http://www.statsoft.com/textbook/stathome.html</u>. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005.
924 925 926 927 928 929 930 931 932 933	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. http://www.transparency.org/policy_research/surveys_indices/cpi/2005. Cited 4 Apr 2006
924 925 926 927 928 929 930 931 932 933 934	Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. <u>http://www.statsoft.com/textbook/stathome.html</u> . Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. <u>http://www.transparency.org/policy_research/surveys_indices/cpi/2005</u> . Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic
 924 925 926 927 928 929 930 931 932 933 934 935 	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria, 2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. http://www.transparency.org/policy_research/surveys_indices/cpi/2005. Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic Change 67(2-3): 273-290
 924 925 926 927 928 929 930 931 932 933 934 935 936 	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. http://www.transparency.org/policy research/surveys indices/cpi/2005. Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic Change 67(2-3): 273-290 Tschakert, P. (2004) The costs of soil carbon sequestration: an economic analysis for small-scale farming systems in Senegal. Agricultural
924 925 926 927 928 929 930 931 932 933 934 935 936 937	Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. <u>http://www.statsoft.com/textbook/stathome.html</u> . Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. <u>http://www.transparency.org/policy_research/surveys_indices/cpi/2005</u> . Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic Change 67(2-3): 273-290 Tschakert, P. (2004) The costs of soil carbon sequestration: an economic analysis for small-scale farming systems in Senegal. Agricultural Systems 81: 227–253
924 925 926 927 928 929 930 931 932 933 934 935 936 937 938	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria, 2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. http://www.transparency.org/policy_research/surveys_indices/cpi/2005. Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic Change 67(2-3): 273-290 Tschakert, P. (2004) The costs of soil carbon sequestration: an economic analysis for small-scale farming systems in Senegal. Agricultural Systems 81: 227-253 UNCTAD (2000) Guide d'investissement au Mali. http://www.unctad.org/fr/docs/poiteiim24.fr.pdf. Cited 28 Mar 2006
924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria, 2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. http://www.transparency.org/policy_research/surveys_indices/cpi/2005. Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic Change 67(2-3): 273-290 Tschakert, P. (2004) The costs of soil carbon sequestration: an economic analysis for small-scale farming systems in Senegal. Agricultural Systems 81: 227–253 UNCTAD (2000) Guide d'investissement au Mali. http://www.unctad.org/fr/docs/poiteiitm24.fr.pdf. Cited 28 Mar 2006 UNFCCC (2003) Modalities and procedures for afforestation and reforestation project activities under the clean development mechanism in
924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 930	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria,2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et telédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. http://www.transparency.org/policy_research/surveys_indices/cpi/2005. Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic Change 67(2-3): 273-290 Tschakert, P. (2004) The costs of soil carbon sequestration: an economic analysis for small-scale farming systems in Senegal. Agricultural systems 81: 227-253 UNCTAD (2000) Guide d'investissement au Mali. http://www.unctad.org/fr/docs/poiteiitm24.fr.pdf. Cited 28 Mar 2006 UNFCCC (2003) Modalities and procedures for afforestation and reforestation project activities under the clean development mechanism in the first commitment period of the Kyoto Protocol. 19CP9/FCCC/CP/2003/6/Add2. Bonn
924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 939 939 940 941	 Smaling, E., M. Touré, et al. (2006) Fertilizer Use and the Envrionment in Africa: Friends of Foes. Paper presented at African Fertilizer Summit, Abuja, Nigeria, 2006 Stancioff, A., M. Staljanssens, et al. (1986) Cartographie et télédétection des ressources de la république du Sénégal - Etude de la géologie, de l'hydrologie, des sols, de la végétation et des potentiels d'utilisation des sols. USAID, Brookings StatSoft, I. (2006) Electronic Statistics Textbook. http://www.statsoft.com/textbook/stathome.html. Cited 21 Feb 2006 Tappan, G. G., M. Sall, et al. (2004) Ecoregions and land cover trends in Senegal. Journal of Arid Environments 59(3): 427-462 The World Bank (2008) Forest Carbon Partnership Facility Information Memorandum - June 13, 2008, The World Bank Tiessen, H., C. Feller, et al. (1998) Carbon sequestration and turnover in semiarid savannas and dry forest. Climatic Change 40(1): 105-117 Transparency International (2006) Corruption Perceptions Index 2005. http://www.transparency.org/policy_research/surveys_indices/cpi/2005. Cited 4 Apr 2006 Tschakert, P. (2004) Carbon for farmers: Assessing the potential for soil carbon sequestration in the Old Peanut Basin of Senegal. Climatic Change 67(2-3): 273-290 Tschakert, P. (2004) The costs of soil carbon sequestration: an economic analysis for small-scale farming systems in Senegal. Agricultural Systems 81: 227-253 UNCTAD (2000) Guide d'investissement au Mali. http://www.unctad.org/fr/docs/poiteiitm24.fr.pdf. Cited 28 Mar 2006 UNFCCC (2003) Modalities and procedures for afforestation and reforestation project activities under the clean development mechanism in the first commitment period of the Kyoto Protocol. 19CP9/FCC/CP/2003/6/Add2. Bonn UNFCCC (2005) Sixth compilation and synthesis of initial national communications from Parties not included in Annex I to the Convention.

943	UNFCCC (2008) Report of the Conference of the Parties serving as the meeting of the Parties to the Kyoto Protocol on its third session, held
944	in Bali from 3 to 15 December 2007. FCCC/KP/CMP/2007/9. United Nations Office at Geneva, Geneva
945	Vagen, T. G., R. Lal, et al. (2005) Soil carbon sequestration in sub-Saharan Africa: A review. Land Degradation and Development 16(1): 53-
946	71
947	Vlek, P. L. G. (1990) The role of fertilizers in sustaining agriculture in sub-saharan Africa. Fertilizer Research 26(1-3): p.327-339
948	Vlek, P. L. G., G. Rodriguez-Kuhl, et al. (2004) Energy use and CO ₂ production in tropical agriculture and means and strategies for reduction
949	or mitigation. Environment, Development and Sustainability 6: 213-233
950	Wood, E. C., G. G. Tappan, et al. (2004) Understanding the drivers of agricultural land use change in south-central Senegal. Journal of Arid
951	Environments 59(3): 565-582
952	Woomer, P. L., L. L. Tieszen, et al. (2004) Land use change and terrestrial carbon stocks in Senegal. Journal of Arid Environments 59(3):
953	625-642
954	Worldclim (2000) Précipitations totales annuelles au Sénégal. Catalogue : Coordination Régionale (France). ROSELT. Berkeley, Berkeley,
955	University of California
956	
957	
958	

- 959 Figures legends
- 960 Figure 1: Clay content and average yearly rainfall (1950-2000) in Senegal (Stancioff et al.
- 961 1986; Worldclim 2000)





963







967 Figure 3: Factors intervening in the socio-economic model



969

970 Figure 4: Change in Soil Organic Carbon stocks for: (a) the MAXINT scenario; (b) the

971 LOWINT scenario;(c) the HIGHINT scenario

972 * White areas within national boundaries are either clayey soils (> 20% clay), rocks, or wetlands, over which the

973 model was not run; the negative values for Northern Senegal should be taken with caution due to increasing

974 imprecision of the model at low rainfall averages (extra- rather than inter-polations).

975 ** White areas within national boundaries correspond either to the model restriction previously mentioned, or to

976 non agricultural land-uses as identified by (DAAC 2002).



- 978
- 979 Figure 5: Results of model coupling for two scenarios: (a) Change in greenhouse gas (GHG)
- 980 balance for the HIGHINT scenario; (b) Change in GHG balance for the MAXINT scenario;
- 981 (c) Change in financial benefits for the HIGHINT scenario; (d) Change in financial benefits
- 982 for the MAXINT scenario
- 983 * Negative figures indicate net releases of GHG; the change in GHG balance includes emissions from fertilizer
- 984 manufacture, transportation, and use, emissions from manure use, and SOC changes



a. Change in GHG balance for the HIGHINT scenario*

b. Change in GHG balance for the MAXINT scenario*



- 986
- 987 Figure 6: Valuation of potential carbon sequestration at US\$ 25.8 tCeq⁻¹ for the MAXINT
- 988 scenario:
- 989 (a) Change in NPV over five years for the MAXINT scenario; (b) Profitability switch with
- 990 carbon valuation for the MAXINT scenario



- 991 992
- 993 Figure 7: Results from model coupling with accounting "avoided deforestation" for the
- 994 MAXINT scenario: (a) Change in carbon balance; (b) Change in NPV over five years



997 Tables

Region	Percent of Average change		Average	e GHG balance	Average change	Average change in NPV over five years with C valuation (\$/ha)	
	region subject to analysis (%)	in SOC stocks (tC/ha)	C stocks (tCeq/ha/y) including "avoided in financ (tC/ha) deforestation" (tCeq/ha/y) balance (\$/ha/		in financial balance (\$/ha/y)		
Dakar	59	-0.64	-0.07	5.61	3.2	6.1	
Diourbel	100	1.78	0.31	9.15	-7.3	-0.8	
Fatick	69	3.89	0.46	14.29	22.1	141	
Kaolack	71	3.17	0.33	19.35	58.8	285.4	
Kolda	64	12.75	0.56	28.99	139.3	654.1	
Louga	97	-1.05	-0.24	4.24	-25.1	-131.5	
Matam	55	0.01	-0.08	4.99	20	77.9	
Saint-Louis	73	-3.18	-0.58	2.47	-43, 6	-245.3	
Tambacounda	47	5.91	0.43	22.93	116.4	543.3	
Thiès	90	0.39	0.05	5.58	-15.3	-60.7	
Ziguinchor	44	15.64	0.58	20.23	65	337.7	
Senegal	65	3.03	0.11	13.27	40.1	183.4	

999 Table 1: Change in soil carbon sequestration amounts and benefits across regions of Senegal,

1000 in the MAXINT scenario with respect to the BAU scenario (NPV: Net Present Value)

1001

998

Region	Part of region	Carbon storage		GHG balance				
	subject to analysis			without accounting for		with accounting for		
	(%)			"avoided deforestation"		"avoided deforestation"		
	-	Average (tC/ha)	Total (tC)	Average	Total	Average	Total	
				(tCeq/ha/y)	(tCeq/y)	(tCeq/ha/y)	(tCeq/y)	
Dakar	27.58	0.71	10 553	0.10	1 503	7.06	104 495	
Diourbel	82.38	0.30	123 905	0.01	4 855	4.86	2 019 220	
Fatick	54.61	0.35	149 539	0.02	7 631	5.00	2 149 470	
Kaolack	30.94	0.43	200 442	0.04	17 572	4.93	2 280 350	
Kolda	0.93	0.28	5 492	-0.02	-458	5.22	102 899	
Louga	16.18	0.46	187 749	0.06	22 921	4.38	1 798 800	
Matam	0.01	0.03	5	-0.07	-14	4.97	995	
Saint-Louis	0.03	0.44	264	0.05	29	4.37	2 625	
Tambacounda	0.09	0.01	55	-0.07	-374	4.97	26 353	
Thiès	50.48	0.32	106 519	0.02	6 061	4.81	1 614 680	
Ziguinchor	10.00	0.64	48 023	0.07	5 201	4.84	361 393	
Senegal	10.99	0.38	832 546	0.03	64 930	4.82	10 461 280	

1003 Table 2: Change in soil carbon storage amounts in Senegal in the HIGHINT scenario with

1004 respect to the BAU scenario (for the grid elements where storage is positive)

1005

1002