



HAL
open science

Compactified universal jacobian and the double ramification cycle

Bashar Dudin

► **To cite this version:**

Bashar Dudin. Compactified universal jacobian and the double ramification cycle. 2015. hal-01150627

HAL Id: hal-01150627

<https://hal.science/hal-01150627>

Preprint submitted on 11 May 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMPACTIFIED UNIVERSAL JACOBIAN AND THE DOUBLE RAMIFICATION CYCLE

BASHAR DUDIN

ABSTRACT. Using the compactified universal jacobian $\overline{\mathcal{J}}_{g,n}$ over the moduli space of stable marked curves $\overline{\mathcal{M}}_{g,n}$, we give an expression in terms of natural classes of the zero section of $\overline{\mathcal{J}}_{g,n}$ in the (rational) Chow ring of $\overline{\mathcal{J}}_{g,n}$. After extending variants of the Abel-Jacobi map to a locus containing curves of treelike type we give a formula for the pullback of the said zero section along these extensions.

0. INTRODUCTION

0.1. The Eliashberg problem. Let $\underline{\tau} = (\tau_1, \dots, \tau_n)$ be an integer valued non-zero n -tuple having 0 sum entries. The double ramification cycle $[\mathrm{DR}]_{g,\underline{\tau}}$ in the moduli space of (smooth) genus g n -marked curves $\mathcal{M}_{g,n}$ is defined by the locus $\mathrm{DR}_{g,\underline{\tau}}$ of marked curves (X, \underline{p}) , for $\underline{p} = (p_1, \dots, p_n)$, such that

$$\mathcal{O}_X \left(\sum_{i=1}^n \tau_i p_i \right) \simeq \mathcal{O}_X.$$

This is precisely the locus of genus g curves having a dominant map to the projective line, such that the fiber of ∞ is given by the p_i points with positive τ_i , the fiber of 0 is given by the p_i points with negative τ_i and ramification index at p_i is $|\tau_i|$. The Eliashberg problem is to find compactifications of $\mathrm{DR}_{g,\underline{\tau}}$ in the moduli space of genus g stable n -marked curves $\overline{\mathcal{M}}_{g,n}$ where to write down the closure of $\mathrm{DR}_{g,\underline{\tau}}$ in terms of tautological classes of $\overline{\mathcal{M}}_{g,n}$.

There has been mainly two approaches to tackle the previous question, at least on partial compactifications of $\mathrm{DR}_{g,\underline{\tau}}$. One is to define an extension $[\mathrm{DR}]_{g,\underline{\tau}}^{GW}$ of $[\mathrm{DR}]_{g,\underline{\tau}}$ over the whole $\overline{\mathcal{M}}_{g,n}$ as the pushforward of the virtual fundamental class of the moduli space of relative stable maps to a rubber \mathbb{P}^1 , by the forgetful map to $\overline{\mathcal{M}}_{g,n}$. This is the point of view of [CMW12], it gives rise to a formula for $[\mathrm{DR}]_{g,\underline{\tau}}^{GW}$ over the locus of curves having rational tails. The other approach, on which we build up, is to look for extensions of the Abel-Jacobi map \underline{a} from $\mathcal{M}_{g,n}$ to the universal family \mathcal{X}_g over the moduli space of polarized abelian varieties \mathcal{A}_g , sending a curve (X, \underline{p}) to $(\mathrm{Jac}(X), \mathcal{O}_X(\sum_i \tau_i p_i))$, and compute the pullback of the zero section of \mathcal{X}_g along these extensions. By using the extension of \underline{a} to the locus $\mathcal{M}_{g,n}^{ct}$ of curves of compact type R. Hain computed in [Hai13] the class of an extension of $[\mathrm{DR}]_{g,\underline{\tau}}$ in terms of tautological classes over $\mathcal{M}_{g,n}^{ct}$. By [MW13] this class happens to be equal to $[\mathrm{DR}]_{g,\underline{\tau}}^{GW}$ when restricted to curves of compact

Date: May 11, 2015.

2010 Mathematics Subject Classification. 14H10, 14H40, 14H60.

The author was supported by the FCT research project *Geometry of moduli spaces of curves and abelian varieties* EXPL/MAT-GEO/1168/2013.

type. The starting point of Hain's formula is the elegant relation

$$(0.1) \quad [\mathcal{Z}_g] = \frac{[T]^g}{g!}$$

expressing the zero cycle of \mathcal{X}_g in terms of the universal symmetric theta divisor T trivialized along the zero section. The pullback of T to $\mathcal{M}_{g,n}^{ct}$ is then given by

$$(0.2) \quad \underline{\mathbf{a}}^*[T] = -\frac{1}{4} \sum_{\substack{0 \leq h \leq g \\ AC\{1, \dots, n\} \\ 1 \leq h+|A| \leq g+n-1}} \left(\sum_{i \in A} \tau_i \right)^2 \delta_{h,A},$$

where $\delta_{0,i} = -\psi_i$ and $\delta_{h,A} = \delta_{g-h,A^c}$ are boundary divisors for $(h, A) \neq (0, \{i\})$ or $(g, \{1, \dots, n\} - \{i\})$. Formula 0.2 was later proven using simpler techniques in [GZ14a]. Formula 0.1 was extended by S. Grushevsky and D. Zakharov in [GZ14b] to the partial compactification of \mathcal{A}_g by rank 1 semi-abelian varieties. This gave rise to an extension of the formula of $\underline{\mathbf{a}}^*[\mathcal{Z}_g]$ to the case of stable marked curves having at most one non-separating node. Recently A. Pixton conjectured a formula for $[\mathrm{DR}]_{g,\underline{\tau}}^{GW}$ on $\overline{\mathcal{M}}_{g,n}$.

The previous questions were considered by Hain in a more general setting. Let k be an integer and $\underline{\tau}$ an integer valued n -tuple, such that $\sum_i \tau_i = k(2g-2)$. Write $\mathrm{DR}_{g,\underline{\tau}}^k$ for the locus of curves (X, \underline{p}) in $\mathcal{M}_{g,n}$ satisfying

$$\mathcal{O}_X \left(\sum_{i=1}^n \tau_i p_i \right) = \omega_X^{\otimes k}$$

where ω_X is the dualizing sheaf of X —in the smooth case it is just the sheaf of regular differential forms. From a modular point of view this is about looking at curves together with (rational) multiples of canonical divisors having a given specific form. Hain gives equivalent formulas to 0.2 for any k over $\mathcal{M}_{g,n}^{ct}$ and Pixton's conjectures extend to these cases.

0.2. Content of the present paper. Rather than looking at the whole moduli space \mathcal{A}_g , we limit ourselves to the case of the universal jacobian $\mathcal{J}_{g,n}$ over $\mathcal{M}_{g,n}$. A point in $\mathcal{J}_{g,n}$ is simply a tuple (X, \underline{p}, L) where (X, \underline{p}) is a smooth marked curve and L is a degree 0 line bundle on X . For each k and $\underline{\tau}$ the Abel-Jacobi map $\mathbf{a}_k : \mathcal{M}_{g,n} \rightarrow \mathcal{J}_{g,n}$ sending (X, \underline{p}) on $(X, \underline{p}, \mathcal{O}_X(\sum_i \tau_i p_i) \otimes \omega_X^{-k})$ gives a section of the forgetful map $\epsilon_n : \mathcal{J}_{g,n} \rightarrow \mathcal{M}_{g,n}$.

A standard way to look for compactifications of $\mathcal{J}_{g,n}$ is by allowing points (X, \underline{p}, L) where L is a simple torsion-free sheaf of rank 1 on the stable marked curve (X, \underline{p}) . To get a proper stack one has to make assumptions on the multidegree of L . This is done by choosing a polarisation on $\overline{\mathcal{M}}_{g,n}$, imposing a stability condition on each torsion-free sheaf L , see section 1 for more details. In our case, the natural polarisation is the one giving an extension of the zero section $\mathcal{Z}_{g,n}$ to the corresponding compactification $\overline{\mathcal{J}}_{g,n}$. In section 2 we describe $\mathcal{Z}_{g,n}$ as the degeneracy locus of the connection morphism $\phi_*(\mathcal{L} \otimes \mathcal{O}_{D_n}) \rightarrow \mathrm{R}^1 \phi_* \mathcal{L}(-D_n)$ where $\phi : \overline{\mathcal{J}}_{g,n|1} \rightarrow \overline{\mathcal{J}}_{g,n}$ is the universal curve over $\overline{\mathcal{J}}_{g,n}$, \mathcal{L} is the universal quasi-stable torsion-free sheaf over $\overline{\mathcal{J}}_{g,n}$ (a sheaf over $\overline{\mathcal{J}}_{g,n|1}$ in fact) and D_n the n -th marked section. Using Thom-Porteous formula we get the relation

$$(2.1) \quad [\mathcal{Z}_{g,n}] = \left\{ \exp \left(\sum_{s \geq 1} (-1)^s (s-1)! \{ \phi_*(\mathrm{ch}(\mathcal{L}) \mathrm{Td}^\vee(\Omega_\phi) \} \}_s \right) \right\}_g$$

where Ω_ϕ is the relative sheaf of differential forms over $\overline{\mathcal{J}}_{g,n}$ and $\{\bullet\}_\ell$ the degree ℓ part of \bullet .

The usefulness of formula 2.1 lies in the fact that it is straightforward to pullback classes obtained by pushing forward Chern classes of \mathfrak{L} and Ω_ϕ . To give a formula for an extension of $[\mathrm{DR}]_{g,\underline{\tau}}^k$ is thus only about extending \mathbf{a}_k . Let $\pi : \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ be the universal curve over $\overline{\mathcal{M}}_{g,n}$ and D_i the section of π corresponding to the i -th marked point. To extend \mathbf{a}_k means giving a quasi-stable line bundle over a locus of $\overline{\mathcal{M}}_{g,n}$ whose restriction to $\mathcal{M}_{g,n}$ is given by $\mathcal{O}(\sum_i \tau_i D_i)$. Let $\overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}}$ be the union of the locus of curves of treelike type and the one of curves (X, \underline{p}) satisfying, for each union of irreducible components $Z \subset X$

$$(3.1) \quad \sum_{p_i \in Z} \tau_i \geq k \deg(\omega_{X|Z}) - \frac{|Z \cap \overline{X} - \overline{Z}|}{2},$$

with strict inequality whenever $p_1 \in Z$ and $\emptyset \subsetneq Z \subsetneq X$. The complement of $\overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}}$ in $\overline{\mathcal{M}}_{g,n}$ is the closure of the locus of curves having topological type a two vertex loopless graph, which doesn't satisfy 3.1. By twisting $\mathcal{O}(\sum_i \tau_i D_i)$ we show in section 3 that the line bundle

$$(3.5) \quad \mathcal{L}(\underline{\tau}, k) = \mathcal{O}\left(\sum_{i=1}^n \tau_i D_i\right) \otimes \omega^{-k} \otimes \mathcal{O}\left(\sum_{\substack{0 \leq h \leq \lfloor g/2 \rfloor \\ AC\{1, \dots, n\} \\ 2 \leq |A| + h \leq g + n - 2}} [k(1-2h) + \sum_{i \in A} \tau_i] \delta_{h, A \cup \{n+1\}}\right).$$

is a quasi-stable line bundle over $\overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}}$. The Abel-Jacobi map does thus extend to maps $\bar{\mathbf{a}}_k$ defined over $\overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}}$. Obstructions to extend yet further \mathbf{a}_k maps are shortly discussed in section 3. Pulling back 2.1 along $\bar{\mathbf{a}}_k$ we get the formula

$$(4.3) \quad \bar{\mathbf{a}}_k^*[\mathcal{Z}_{g,n}] = \left\{ \exp\left(\sum_{s \geq 1} (-1)^s (s-1)! \{\pi_* (\mathrm{ch}(\mathcal{L}(\underline{\tau}, k)) \mathrm{Td}^\vee(\Omega_\pi))\}_s\right) \right\}_g.$$

It agrees with Hain's formula for curves of compact type but we don't have a direct proof of this fact. The right hand side of 4.3 is an explicit degree g polynomial in tautological classes on $\overline{\mathcal{M}}_{g,n}$. It is unfortunately much less transparent than the formula given by Hain and comparison with the extension given by Grushevsky and Zakharov is not clear.

The starting point of the previous result lies in the fact we're able to write down the zero cycle of the compactified jacobian in terms of its universal sheaf, as well as sheaves coming from $\overline{\mathcal{M}}_{g,n}$. The pullback along extensions of Abel-Jacobi maps is then straightforward. Following the same line of thought, we write down the class of the universal theta divisor $[\Theta]$ (trivialized along the zero section) in terms of \mathfrak{L} , in this case we have that

$$(2.10) \quad [\Theta] = -\phi_* \left(\frac{c_1(\mathfrak{L})^2}{2} \right).$$

A simple computation in section 4 gives yet another way of computing the pullback of $[\Theta]$ along $\bar{\mathbf{a}}_k$.

Lastly, in section 4.3, we briefly explain how to adapt our strategy to compute the pullback of the universal theta divisor of the degree $g-1$ compactified jacobian along the Abel-Jacobi map. We get back the class computed in [GZ14a] up to the generic vanishing of the theta function along the locus of irreducible curves

in $\overline{\mathcal{M}}_{g,n}$. This also recovers the formula given by F. Müller in [Mül13], which computes the closure in $\overline{\mathcal{M}}_{g,n}$ of the locus

$$\mathcal{D}_g = \left\{ (X, \underline{p}) \mid h^0(\mathcal{O}_X(\sum_i \tau_i p_i)) \geq 1 \right\}$$

for smooth marked curves X , $\sum_i \tau_i = g - 1$ and at least one τ_i is negative.

Acknowledgements. I'm grateful to Margarida Melo and Filippo Viviani for presenting me the questions studied here, for many useful discussions and for reading and commenting on early versions of the present paper.

1. COMPACTIFIED UNIVERSAL JACOBIAN OVER $\overline{\mathcal{M}}_{g,n}$

We review the definition of compactified jacobians following [Est01]. Points of any such compactification are given by stable marked curves endowed with simple rank 1 torsion-free sheaves satisfying a semi-stability condition. The first appearance of such compactifications is due to R. Pandharipande in [Pan96] for slope semi-stability, in the more general case of vector bundles. This last point of view, in the case of line bundles, is equivalent to the original approach by L. Caporaso in [Cap94]. Both were compactifications of the universal jacobian over the moduli space of stable curves $\overline{\mathcal{M}}_g$. The corresponding construction for marked stable curves was studied by M. Melo in [Mel11]. The equivalence between Pandharipande's and Caporaso's approaches were first proven in [Pan96]. An extension to the marked case and more general stability conditions can be found in [EP14]. In the case at hand—the case of compactified universal jacobians—our main reference is [Mel].

Convention 1.1. All our schemes are schemes over \mathbb{C} . We fix integers $d \in \mathbb{Z}$, $g \in \mathbb{N}$ and $n \in \mathbb{N}^*$ such that $2g + 2 + n > 0$.

Let $\tilde{\mathcal{J}}_{g,n}^d$ be the stackification of the prestack whose S -sections are given by couples $(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{L})$ consisting of a relative stable curve f of genus g , disjoint étale relative Cartier divisors $\underline{D} = (D_1, \dots, D_n)$ in the smooth locus of f and a coherent sheaf \mathcal{L} over \mathcal{X} which is flat over S and has simple torsion-free degree d geometric fibers. The Cartier divisors \underline{D} are called markings or marked points of f and are assumed to be defined by images of sections of f . Morphisms from $(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{L})$ on $(\psi : \mathcal{Y} \rightarrow T, \underline{E}, \mathcal{M})$ over $\alpha : S \rightarrow T$ are given by S -morphisms $\beta : \mathcal{X} \rightarrow \alpha^* \mathcal{Y}$ commuting with the markings and for which there is an invertible sheaf \mathcal{N} on S and an isomorphism $\mu : \mathcal{L} \simeq \beta^* \mathcal{M} \otimes f^* \mathcal{N}$.

Theorem 1.1 ([Mel]). *The stack $\tilde{\mathcal{J}}_{g,n}^d$ is a smooth irreducible DM-stack of dimension $4g - 3 + n$. It is representable over $\overline{\mathcal{M}}_{g,n}$ and satisfies the existence part of the relative criterion for properness.*

The previous theorem says that to compactify the open substack $\mathcal{J}_{g,n}^d$, corresponding to smooth underlying curves, one has to look for separated universally closed substacks of $\tilde{\mathcal{J}}_{g,n}^d$.

Let X be a stable marked curve of genus g over \mathbb{C} or, more generally, over an algebraically closed field. By a subcurve of X we shall always mean a *closed* subcurve of X . A polarization of degree d on X is a locally free \mathcal{O}_X -module P of rank $r > 0$ and degree $r(d + 1 - g)$. Given any subcurve $Y \subset X$ we write

$$q_Y(P) = \frac{\deg P|_Y}{r} + \frac{\deg \omega_{X|Y}}{2}$$

where ω_X is the canonical sheaf on X .

Definition 1.1. Let P be a polarization a degree d on X and $x \in X$ a smooth point. Given a closed subcurve Y of X we write κ_Y for the number of points in the intersection of Y with the closure of its complement in X . Given a line bundle L on X we write $\deg_Y L$ for the degree of $L|_Y$ modulo torsion. A torsion-free sheaf L of rank 1 on X of degree d is said to be

- (1) P -semistable if for all non-empty subcurves $Y \subsetneq X$, $\deg_Y L \geq q_Y - \frac{\kappa_Y}{2}$
- (2) P -stable if inequalities in 1. are strict
- (3) (P, x) -quasistable (q-stable for short) if (1) is verified and is strict whenever $x \in Y$.

By a polarization on $\overline{\mathcal{M}}_{g,n}$ we understand a polarization on the universal curve over $\overline{\mathcal{M}}_{g,n}$. This means a locally free sheaf \mathcal{P} on $\overline{\mathcal{M}}_{g,n+1}$ of rank r and degree $r(d+1-g)$.

Fix a polarization \mathcal{P} on $\overline{\mathcal{M}}_{g,n}$ and a section τ of $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$. Write $\overline{\mathcal{J}}_{g,n}^{d,\mathcal{P},ss}$, $\overline{\mathcal{J}}_{g,n}^{d,\mathcal{P},s}$ and $\overline{\mathcal{J}}_{g,n}^{d,\mathcal{P},\tau}$ for the substacks of $\overline{\mathcal{J}}_{g,n}^d$ given by triples $(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{L})$ for which \mathcal{L} has respectively P -semistable, P -stable and (P, τ) -quasistable simple torsion-free geometric fibers of degree d .

Theorem 1.2 ([Mel]). *$\overline{\mathcal{J}}_{g,n}^{d,\bullet}$ are smooth irreducible DM-stacks of dimension $4g-3+n$ that are representable and of finite type over $\overline{\mathcal{M}}_{g,n}$. Furthermore*

- (1) $\overline{\mathcal{J}}_{g,n}^{d,\mathcal{P},ss}$ is universally closed
- (2) $\overline{\mathcal{J}}_{g,n}^{d,\mathcal{P},s}$ is separated
- (3) $\overline{\mathcal{J}}_{g,n}^{d,\mathcal{P},\tau}$ is proper and has a projective coarse moduli space.

In the following, we will mainly be interested in the compactification $\overline{\mathcal{J}}_{g,n}$ of $\mathcal{J}_{g,n}$ given by the polarization $\omega^{-1} \oplus \mathcal{O}$ and q-stable torsion-free rank 1 sheaves. For clarity the section τ defining q-stability is D_1 , the relative étale Cartier divisor defined by the first marking. We limit ourselves to this case from now on.

Let $\overline{\mathcal{J}}_{g,n|1}$ be the stack whose S -sections are given by sections $(f : \mathcal{X} \rightarrow S, \underline{D}, D_{n+1}, \mathcal{L})$ where D_{n+1} is an extra section. There are no assumptions on D_{n+1} ; it can intersect other marked sections or the singular locus of f . The universal family over $\overline{\mathcal{J}}_{g,n}$ is given by the map $\phi : \overline{\mathcal{J}}_{g,n|1} \rightarrow \overline{\mathcal{J}}_{g,n}$ forgetting the extra section. It comes with a universal torsion free sheaf \mathcal{L} of rank 1 given over an S -section $(f : \mathcal{X} \rightarrow S, \underline{D}, D_{n+1}, \mathcal{L})$ by the pullback of \mathcal{L} along D_{n+1} . One has also another forgetful map $\epsilon_{n|1} : \overline{\mathcal{J}}_{g,n|1} \rightarrow \overline{\mathcal{M}}_{g,n+1}$ sending $(f : \mathcal{X} \rightarrow S, \underline{D}, D_{n+1}, \mathcal{L})$ on $(f : \mathcal{X} \rightarrow S, \underline{D}, D_{n+1})$. The stack of stable n -marked curves having an extra section is isomorphic to the one of $n+1$ -marked stable curves. We shall constantly make this identification. This discussion fits into the picture

$$(1.1) \quad \begin{array}{ccc} \mathcal{L} & \longrightarrow & \overline{\mathcal{J}}_{g,n|1} \xrightarrow{\phi} \overline{\mathcal{J}}_{g,n} \\ & & \downarrow \epsilon_{n|1} \quad \downarrow \epsilon_n \\ & & \overline{\mathcal{M}}_{g,n+1} \xrightarrow{\pi} \overline{\mathcal{M}}_{g,n} \end{array}$$

where the square diagram is commutative but generally not cartesian.

2. THE ZERO CYCLE OF $\overline{\mathcal{J}}_{g,n}$ AND CLASS OF THE THETA DIVISOR

2.1. The zero cycle of $\overline{\mathcal{J}}_{g,n}$. A line bundle L on a stable curve X is semistable for $\omega^{-1} \oplus \mathcal{O}$ if and only if for each proper subcurve $Y \subset X$ the inequality

$$\deg_Y L \geq -\frac{\kappa_Y}{2}$$

holds. In particular the trivial line bundle \mathcal{O}_X is stable and thus q -stable. The zero section of the universal Jacobian over $\mathcal{M}_{g,n}$ does therefore extend to $\overline{\mathcal{J}}_{g,n}$, its image is simply denoted by $\mathcal{Z}_{g,n}$. It is a smooth irreducible substack of $\overline{\mathcal{J}}_{g,n}$ of codimension g , its cycle in the Chow ring of $\overline{\mathcal{J}}_{g,n}$ is written $[\mathcal{Z}_{g,n}]$. We will be expressing this cycle in terms of “natural” cycles on $\overline{\mathcal{J}}_{g,n}$. Namely the Chern characters of the universal torsion-free sheaf \mathcal{L} , the relative dualizing sheaf of $\overline{\mathcal{J}}_{g,n|1} \rightarrow \overline{\mathcal{J}}_{g,n}$ and boundary classes. We first start by a lemma on the space of global sections of L .

Lemma 2.1. *For semistable L we have that $h^0(L) \leq 1$ with equality if and only if L is locally free of $\underline{0}$ multidegree.*

Proof. Let $\nu : X^\nu \rightarrow X$ be the normalisation of X . The natural map $L \rightarrow \nu_*\nu^*L$ induces an injective map $H^0(L) \rightarrow H^0(\nu^*L)$. Recall that

$$\deg(L) = \deg(\nu^*L) + t_L$$

where t_L is the number of nodal points of X where L is not free. In particular, when L is not free the total degree of ν^*L is negative (by definition the total degree of L is 0).

There is nothing to prove when X has only one irreducible component. Assume that X has at least two such components and L is not of multidegree $\underline{0}$. Notice that this is automatically the case if L is not free. One can write $X = Z \cup Z^c$ for Z an irreducible component of X such that $L|_Z$ has negative degree and Z^c is the closure of the complement of Z in X . Let X_j for $j = 1, \dots, m$ be the connected components of Z^c and write $E = Z \cap Z^c$, $E_j = E \cap X_j$. We have that

$$H^0(X, L) = H^0\left(Z^c, L\left(\sum_{x \in E} -x\right)\right) = \bigoplus_j H^0\left(X_j, L\left(\sum_{x \in E_j} -x\right)\right).$$

Now the semistability condition says

$$\deg_{X_j} L \leq \frac{|E_j|}{2}.$$

The connectedness of X insures $L(\sum_{x \in E_j} -x)$ has negative degree on X_j . Using iteratively semistability inequalities we get that $h^0(L) = 0$. We’ve thus shown that if L is not of multidegree $\underline{0}$ then $h^0(L) = 0$. Since L cannot be of multidegree $\underline{0}$ if non-free we have just shown our claim. \square

Corollary 2.2. *For semistable L on X , $h^0(L) \neq 0$ if and only if $L \simeq \mathcal{O}_X$.*

Proof. By the previous lemma L has to be of multidegree $\underline{0}$. If a global section of L vanishes at some point it vanishes on the irreducible component containing it. Since X is connected it has to identically vanish on X . When $h^0(L) \neq 0$ there is thus a section that doesn’t vanish anywhere on X . \square

The substack of $\overline{\mathcal{J}}_{g,n}$ given by the zero section of $\overline{\mathcal{J}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ is formally defined by S -points $(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{L})$ in $\overline{\mathcal{J}}_{g,n}$ such that $\mathcal{L} \simeq f^*\mathcal{M}$ where \mathcal{M} is an invertible sheaf on S . By base change in cohomology this is equivalently given by the S -points in $\overline{\mathcal{J}}_{g,n}$ such that $\mathcal{L}_s \simeq \mathcal{O}_{\mathcal{X}_s}$ for every fiber over $s \in S$. In the light of the previous lemma $\mathcal{Z}_{g,n}$ is thus given by S -points $(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{L})$ such that $h^0(\mathcal{L}_s) \geq 1$ for every $s \in S$.

We describe $\mathcal{Z}_{g,n}$ as the degeneracy locus of a map between vector bundles on $\overline{\mathcal{J}}_{g,n}$. Let $(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{L})$ be an S -point in $\overline{\mathcal{J}}_{g,n}$. Pick one of the marked

sections, say D_n , and consider the short exact sequence of sheaves over \mathcal{X}

$$0 \longrightarrow \mathcal{L}(-D_n) \longrightarrow \mathcal{L} \longrightarrow \mathcal{O}_{D_n} \otimes \mathcal{L} \longrightarrow 0$$

Its direct image by f gives the long exact sequence

$$0 \longrightarrow f_*\mathcal{L} \longrightarrow \mathcal{O}_{D_n} \otimes \mathcal{L} \xrightarrow{\nabla} R^1 f_*\mathcal{L}(-D_n) \longrightarrow R^1 f_*\mathcal{L} \longrightarrow 0$$

The map ∇ is a map between two locally free bundles over S . The source is of rank 1 and the target of rank g . The degeneracy locus of ∇ is defined by points $s \in S$ such that $\text{rank}(\nabla \otimes k(s)) \leq 0$. Because of commutation to base change of both source and target of ∇ the tensor product $\nabla_s = \nabla \otimes k(s)$ is the one appearing in the long exact sequence in cohomology

$$0 \longrightarrow H^0(\mathcal{L}_s) \longrightarrow H^0(\mathcal{O}_{D_n} \otimes \mathcal{L}_s) \xrightarrow{\nabla_s} H^1(\mathcal{L}_s(-D_n)) \longrightarrow H^1(\mathcal{L}_s) \longrightarrow 0.$$

The degeneracy locus of ∇ is thus defined by the set of points $s \in S$ such that $H^0(\mathcal{L}_s) \geq 1$ and this is precisely $\mathcal{Z}_{g,n}$.

Theorem 2.3. *Let $\phi : \overline{\mathcal{J}}_{g,n|1} \rightarrow \overline{\mathcal{J}}_{g,n}$ be the universal curve over $\overline{\mathcal{J}}_{g,n}$ and \mathfrak{L} the universal torsion-free sheaf over $\overline{\mathcal{J}}_{g,n|1}$. The zero cycle $[\mathcal{Z}_{g,n}]$ is given by*

$$(2.1) \quad [\mathcal{Z}_{g,n}] = \left\{ \exp \left(\sum_{s \geq 1} (-1)^s (s-1)! \{ \phi_*(\text{ch}(\mathfrak{L}) \text{Td}^\vee(\Omega_\phi)) \}_s \right) \right\}_g$$

where Ω_ϕ is the sheaf of relative differentials of $\overline{\mathcal{J}}_{g,n|1}$ over $\overline{\mathcal{J}}_{g,n}$ and $\{\bullet\}_\ell$ is the degree ℓ part of \bullet .

Proof. The zero locus $\mathcal{Z}_{g,n}$ is the degeneracy locus of ∇ . Since $\overline{\mathcal{J}}_{g,n}$ is smooth and $\mathcal{Z}_{g,n}$ of the expected codimension g we get the relation

$$[\mathcal{Z}_{g,n}] = c_g(R^1 \phi_* \mathfrak{L}(-D_n) - \phi_*(\mathcal{O}_{D_n} \otimes \mathfrak{L}))$$

out of the Thom-Porteous formula [Ful98]. The standard relation between Chern classes and Chern characters gives

$$[\mathcal{Z}_{g,n}] = \left\{ \exp \left(\sum_{s \geq 1} (-1)^{s-1} (s-1)! \left[\text{ch}_s(R^1 \phi_* \mathfrak{L}(-D_n)) - \text{ch}_s(\phi_* \mathcal{O}_{D_n} \otimes \mathfrak{L}) \right] \right) \right\}_g$$

The exact sequence

$$0 \longrightarrow \phi_* \mathfrak{L} \longrightarrow \phi_*(\mathcal{O}_{D_n} \otimes \mathfrak{L}) \longrightarrow R^1 \phi_* \mathfrak{L}(-D_n) \longrightarrow R^1 \phi_* \mathfrak{L} \longrightarrow 0$$

and the additivity of the Chern character in exact sequences says

$$[\mathcal{Z}_{g,n}] = \left\{ \exp \left(\sum_{s \geq 1} (-1)^s (s-1)! \text{ch}_s(\phi_* \mathfrak{L}) \right) \right\}_g.$$

The desired formula is obtained using the GRR formula. \square

Previously known expressions of 2.1 on partial compactifications of $\mathcal{J}_{g,n}$ were based on relation 0.1, which is only valid over curves of compact type. Correction terms were computed in [GZ14b] to extend the latter relation to the locus of curves having at most 1 non-separating node. The advantage of the present formula, from a modular point of view, is that \mathfrak{L} is much easier to deal with than the theta divisor. This will be put in practice in section 4. We identify in the following section the term in 2.1 coming from the class of the universal theta divisor $[\Theta]$, trivialized along the zero section.

2.2. Relating the theta divisor to \mathfrak{L} . Marked points do not play a role in this section. The corresponding compactification $\overline{\mathcal{J}}_g^{d,\mathcal{P}}$ of the universal jacobian \mathcal{J}_g over (unmarked) stable curves is given by semi-stable torsion-free sheaves with respect to a given polarisation \mathcal{P} over $\overline{\mathcal{M}}_g$, see section 1 for details. We will only be concerned by the degree 0 and $g-1$ compactified universal jacobians $\overline{\mathcal{J}}_g$ and $\overline{\mathcal{J}}_g^{g-1}$; the former has polarisation $\omega^{-1} \otimes \mathcal{O}$ while the latter has trivial polarisation \mathcal{O} . The compactified universal jacobian $\overline{\mathcal{J}}_g^{g-1}$ comes with a canonical theta divisor Θ_{g-1} set theoretically given by

$$(2.2) \quad \Theta_{g-1} = \{(X, L) \mid h^0(L) \geq 1\}.$$

We have the relation

$$(2.3) \quad -\Theta_{g-1} = d_\pi(\mathcal{L}_{g-1})$$

where $d_\pi(\bullet)$ is the determinant in cohomology $\det(\pi_!\bullet)$, see for instance [ACG11, 13]. This description can be found in [Ale04] and [Cap09]. We question how to pullback Θ_{g-1} on $\overline{\mathcal{J}}_g$.

For the degree 0 jacobian of a single smooth curve X , defining a theta divisor is about pulling back Θ_{g-1} along an isomorphism $-\otimes \xi : \text{Jac}(X) \simeq \text{Jac}^{g-1}(X)$ for a degree $g-1$ line bundle ξ . A line bundle on $\text{Jac}(X)$ is said to be symmetric if it is invariant by pullback along the involution $L \mapsto L^\vee$. Using the set-theoretic description 2.2 one can show that the pullback of Θ_{g-1} along $-\otimes \xi$ is symmetric if and only if ξ is a theta characteristic, i.e. if and only if $\xi^{\otimes 2} \simeq \omega$. There is no canonical choice of a theta characteristic for a given curve, there is thus no hope to define a symmetric theta divisor this way on \mathcal{J}_g . There is, however, a roundabout way to partially tackle this question: the pullback of the rational equivalence class of $2\Theta_{g-1}$ by $-\otimes \xi$ does not depend on the theta characteristic ξ . Indeed, two theta characteristics ξ and ξ' differ by a root of the trivial line bundle $\eta = \xi \otimes \xi'^{-1}$. Using the theorem of the square [Mum08, II. 6.] we have that

$$(2.4) \quad 2\Theta_{g-1} \sim 2t_\eta \Theta_{g-1},$$

where t_η is translation by η on the Picard group of $\text{Jac}^{g-1}(X)$. Pulling back 2.4 along $-\otimes \xi$ we get the desired relation between the pullbacks along $-\otimes \xi$ and $-\otimes \xi'$ of $2\Theta_{g-1}$.

We should thus be able to pullback $2\Theta_{g-1}$ to \mathcal{J}_g . The point is that there is no global choice of a theta characteristic on \mathcal{M}_g , one has to take in account all such possible choices. Assume for now $g \geq 2$ and let \mathcal{S}_g^2 be the moduli space of smooth genus g spin curves, see for instance [Cor89]. A point of \mathcal{S}_g^2 is a couple (X, ξ) where X is a marked curve of genus g and ξ a line bundle of degree $g-1$ together with an isomorphism $\alpha : \xi^{\otimes 2} \simeq \omega_X$. The moduli space \mathcal{S}_g^2 comes with a forgetful map to \mathcal{M}_g only keeping the underlying marked curve, and a universal line bundle Ξ on the universal curve such that $\Xi^{\otimes 2} \simeq \omega$. On the fiber product $\mathcal{S}_g^2 \times_{\mathcal{M}_g} \mathcal{J}_g$ we have two maps

$$(2.5) \quad \mathcal{J}_g \xleftarrow{\beta} \mathcal{S}_g^2 \times_{\mathcal{M}_g} \mathcal{J}_g \xrightarrow{\alpha} \mathcal{J}_g^{g-1}$$

where β is the projection on the second factor and α sends $(f : \mathcal{X} \rightarrow S, \mathcal{L}, \xi)$ on $(f : \mathcal{X} \rightarrow S, \mathcal{L} \otimes \xi)$. By the standard theory of spin curves, the map β is finite of degree 2^{2g} . Write $\langle \bullet, - \rangle$ for the Deligne pairing of two line bundles \bullet and $-$ on

the universal curves over \mathcal{J}_g^{g-1} or \mathcal{J}_g . Using 2.3 we have that

$$-2\alpha^*\Theta_{g-1} = d_\phi(\beta^*\mathcal{L} \otimes \Xi)^{\otimes 2}.$$

Using GRR [ACG11, XII 6. 5.31], we can write

$$\begin{aligned} -2\alpha^*\Theta_{g-1} &= \langle \beta^*\mathcal{L} \otimes \Xi, \beta^*\mathcal{L} \otimes \Xi^{-1} \rangle d_\phi(\mathcal{O})^{\otimes 2} \\ &= \langle \beta^*\mathcal{L}, \beta^*\mathcal{L} \rangle \langle \beta^*\mathcal{L}, \Xi \rangle \langle \beta^*\mathcal{L}, \Xi^{-1} \rangle \langle \Xi, \Xi^{-1} \rangle d_\phi(\mathcal{O})^{\otimes 2}. \end{aligned}$$

Since $\langle \bullet, \Xi^{-1} \rangle = \langle \bullet, \Xi \rangle^{-1}$ we get that

$$-2\alpha^*\Theta_{g-1} = \langle \beta^*\mathcal{L}, \beta^*\mathcal{L} \rangle \langle \Xi, \Xi \rangle^{-1} d_\phi(\mathcal{O})^{\otimes 2}.$$

One can check by looking at the Chern classes in the group of relative Cartier divisors of $\mathcal{S}_g^2 \times_{\mathcal{M}_g} \mathcal{J}_g$ that

$$c_1 \langle \Xi, \Xi^{-1} \rangle = -\phi_* \left(\frac{c_1(\omega)^2}{4} \right) = -3\lambda_1 = c_1(d_\phi(\mathcal{O})^{-\otimes 3}).$$

The middle equality comes from the Mumford relation over \mathcal{M}_g [ACG11, XII 7.3] and λ_1 is the first Chern class of the pullback to $\mathcal{S}_g^2 \times_{\mathcal{M}_g} \mathcal{J}_g$ of the Hodge bundle over \mathcal{M}_g . Putting the previous computations together we get that

$$(2.6) \quad -2\alpha^*\Theta_{g-1} = \beta^* (\langle \mathcal{L}, \mathcal{L} \rangle d_\phi(\mathcal{O})^{-1}).$$

Remark 1. Computations in the genus 1 case are similar. One has to be careful though, in order to work with Deligne-Mumford stacks one has to stick to working with $\mathcal{M}_{1,1}$. We get the same expression as 2.6 but over $\mathcal{M}_{1,1}$.

Remark 2. We extend the Deligne pairing to the case of \mathcal{L} and \mathcal{L}^\vee over $\overline{\mathcal{J}}_g$ using the relation

$$\langle L, M \rangle = d_\phi(L \otimes M) d_\phi(L)^{-1} d_\phi(M)^{-1} d_\phi(\mathcal{O})$$

for L and M invertible sheaves or in $\{\mathcal{L}^{\otimes m}, \mathcal{L}^{\vee \otimes m} \mid m \in \mathbb{N}\}$. All standard computational properties including the Serre duality extend to this case. A way to check this fact is by recalling that $\overline{\mathcal{J}}_g^d$ is isomorphic to the compactification by Caporaso of the universal jacobian using quasi-stable curves. The pullback of \mathcal{L} to the latter compactification is a well-behaved invertible sheaf. See [EP14] for details. One can thus check the standard properties extend as claimed by pullback on Caporaso's compactification using semi-stable line bundles on quasi-stable curves.

Definition 2.1. A squared theta divisor over $\overline{\mathcal{J}}_g$ is a divisor given on each geometric fiber over \mathcal{M}_g^{ct} as the pullback of $2\Theta_{g-1}$ along a theta characteristic. It is said to be

- trivialized along the zero section if its pullback along the zero section $\overline{\mathcal{M}}_g \rightarrow \overline{\mathcal{J}}_g$ is trivial
- symmetric if it is invariant under the involution of the Picard group of $\overline{\mathcal{J}}_{g,1}$ sending \mathcal{L} on \mathcal{L}^\vee .

In case of genus 1 the definition makes sense replacing $\overline{\mathcal{J}}_g$ and $\overline{\mathcal{M}}_g$ respectively by $\overline{\mathcal{J}}_{1,1}$ and $\overline{\mathcal{M}}_{1,1}$.

Proposition 2.4. *There is a unique symmetric squared theta divisor 2Θ over $\overline{\mathcal{J}}_g$ which is trivialized along the zero section. It is then given by the formula*

$$(2.7) \quad 2\Theta = \langle \mathcal{L}, \mathcal{L} \rangle^{-1}.$$

Proof. The fact the right-hand side of 2.7 is symmetric comes from the standard properties of the Deligne pairing in the case of torsion free sheaves as mentioned in remark 2.

Over \mathcal{M}_g the pullback of $2\Theta_{g-1}$ is given by formula 2.6. The line bundle $\langle \mathcal{L}, \mathcal{L} \rangle^{-1} d_\phi(\mathcal{O})$ is thus a squared theta divisor over \mathcal{J}_g . Its pullback along the zero section is given by $d_\phi(\mathcal{O})$. It is enough to get the desired formula over \mathcal{J}_g . To extend it on \mathcal{J}_g^{ct} on only needs to check formula 2.7 on the locus of curves having two components. Let $\iota_h : \mathcal{J}_{h,x} \times \mathcal{J}_{g-h,y} \rightarrow \mathcal{J}_g^{ct}$ be the gluing maps identifying x and y and write p_i for the first and second projections from $\mathcal{J}_{h,x} \times \mathcal{J}_{g-h,y}$ respectively on $\mathcal{J}_{h,x}$ and $\mathcal{J}_{g-h,y}$. The universal curve over $\mathcal{J}_{h,x} \times \mathcal{J}_{g-h,y}$ is the disjoint union

$$(2.8) \quad (\mathcal{J}_{h,x|1} \times \mathcal{J}_{g-h,y}) \bigsqcup (\mathcal{J}_{h,x} \times \mathcal{J}_{g-h,y|1})$$

with the obvious maps ϕ and $\iota_{h|1}$ respectively to $\mathcal{J}_{h,x} \times \mathcal{J}_{g-h,y}$ and \mathcal{J}_g^{ct} . Write p_h and p_{g-h} for the maps whose domain is 2.8 and only keeping $\mathcal{J}_{h,x|1}$ and $\mathcal{J}_{g-h,y|1}$. We have that $\phi \circ p_h = p_1$ and $\phi \circ p_{g-h} = p_2$. Using the simple relation $\iota_{h|1}^* \mathcal{L} = p_h^* \mathcal{L} + p_{g-h}^* \mathcal{L}$ on the corresponding universal curves we get that

$$(2.9) \quad \begin{aligned} \iota_h^* \langle \mathcal{L}, \mathcal{L} \rangle &= \langle p_h^* \mathcal{L}, p_h^* \mathcal{L} \rangle \langle p_h^* \mathcal{L}, p_{g-h}^* \mathcal{L} \rangle^{\otimes 2} \langle p_{g-h}^* \mathcal{L}, p_{g-h}^* \mathcal{L} \rangle \\ &= \langle p_h^* \mathcal{L}, p_h^* \mathcal{L} \rangle \langle p_{g-h}^* \mathcal{L}, p_{g-h}^* \mathcal{L} \rangle. \end{aligned}$$

The middle term vanishes because $p_h^* \mathcal{L}$ and $p_{g-h}^* \mathcal{L}$ have disjoint supports. Each term of the right-hand side of 2.9 respectively correspond to the pullbacks $-2p_1^* \Theta$ and $-2p_2^* \Theta$ because of the previous work on the smooth case. Now, the equality

$$\iota_h^* \langle \mathcal{L}, \mathcal{L} \rangle = -2\Theta$$

is just about rewriting the fact that the theta divisor of a decomposable jacobian is the union of pullbacks along the projection to each factor.

Uniqueness is a consequence of the work of Melo and Viviani [MV14] on the Picard group of $\overline{\mathcal{J}}_g$. Using our notation theorem B of *loc. cit.* states that the Picard group of $\overline{\mathcal{J}}_g$ is freely generated by $\langle \mathcal{L}, \mathcal{L} \rangle d_\phi(\mathcal{O})^{\otimes 2}$, $d_\phi(\omega)$ and line bundles obtained by pullback from $\overline{\mathcal{M}}_g$. Thus, the only extension of 2Θ to $\overline{\mathcal{J}}_g$ that is trivialized along the zero section is the one given by formula 2.7. \square

Corollary 2.5. *Let $[\Theta]$ be half the class of the symmetric squared theta divisor trivialized along the zero section in the (rational) Chow ring of \mathcal{J}_g . We then have*

$$(2.10) \quad [\Theta] = -\phi_* \left(\frac{c_1(\mathcal{L})^2}{2} \right).$$

Proof. This is a direct computation of first Chern classes of both sides of 2.7. \square

Remark 3. In the case of marked underlying curves, the theta divisor is obtained by pullback along the forgetful morphism $\overline{\mathcal{J}}_{g,n} \rightarrow \overline{\mathcal{J}}_g$ in the case of genus $g \geq 2$ and by pullback along $\overline{\mathcal{J}}_{1,n} \rightarrow \overline{\mathcal{J}}_{1,1}$ for the genus 1 case. Formula 2.7 does still hold in these cases.

Putting back relation 2.10 into the expression of the zero cycle 2.1 one can rewrite 2.1 in the form

$$[\mathcal{Z}_{g,n}] = \left\{ \exp\left(-\phi_*\left(\frac{c_1(\mathcal{L})^2}{2}\right)\right) \times \exp\left(\frac{c_1(\mathcal{L})c_1(\omega_\phi)}{2} - \frac{B_2}{2}(c_1(\omega_\phi)^2 + \Delta)\right) \right. \\ \left. \times \exp\left(\sum_{s \geq 2} (-1)^s (s-1)! \left(\phi_*(\text{ch}(\mathcal{L}) \text{Td}^\vee(\Omega_\phi))\right)_s\right) \right\}_g$$

where ω_ϕ is the dualizing sheaf of ϕ , Δ the singular locus in the fibers of ϕ . This gives

$$(2.11) \quad [\mathcal{Z}_{g,n}] = \sum_{a+b=g} \frac{[\Theta]^a}{a!} P_b$$

where P_b is a polynomial in pushforward of classes on $\overline{\mathcal{J}}_{g,n|1}$ lying in the degree b part of the Chow ring. It is not clear how to write P_b as a polynomial in an elegant minimal set of classes on $\overline{\mathcal{J}}_{g,n}$, the main reason being that not much is known about the intersection theory of $\overline{\mathcal{J}}_{g,n}$. Relation 0.1 (pulled back to the universal jacobian) does however imply that on the locus $\mathcal{M}_{g,n}^{ct}$ of curves of compact type we have

$$(2.12) \quad \sum_{a+b=g} \frac{[\Theta]^a}{a!} P_b = \frac{[\Theta]^g}{g!}.$$

3. PARTIAL EXTENSION OF ABEL-JACOBI MAPS

Let $\underline{\tau} = (\tau_1, \dots, \tau_n)$ be a non-zero n -tuple of integer entries having zero sum. The forgetful map $\epsilon_n : \mathcal{J}_{g,n} \rightarrow \mathcal{M}_{g,n}$ sending an S -point of $\mathcal{J}_{g,n}$ on the underlying marked stable curve has a section \mathbf{a} -known as the Abel-Jacobi map-sending an S -point $(f : \mathcal{X} \rightarrow S, \underline{D})$ on $(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{O}_{\mathcal{X}}(\sum_i \tau_i D_i))$. More generally given an integer $k \in \mathbb{Z}$ and an n -tuple of integers (τ_1, \dots, τ_n) having sum equal to $k(2g-2)$ one can define the section of ϵ_n

$$\mathbf{a}_k(f : \mathcal{X} \rightarrow S, \underline{D}) = \left(f : \mathcal{X} \rightarrow S, \underline{D}, \mathcal{O}_{\mathcal{X}}\left(\sum_i \tau_i D_i\right) \otimes \omega^{-k}\right).$$

These Abel-Jacobi maps can be extended to partial compactifications of $\mathcal{M}_{g,n}$. The extension to curves of compact type is standard and the one to curves with at most 1 non-separating node was described in [GZ14b]. These extensions appear to give partial answers to the Eliashberg problem: express the pullback of $[\mathcal{Z}_{g,n}]$ along extensions of \mathbf{a}_0 to partial compactifications of $\mathcal{M}_{g,n}$.

We will extend the \mathbf{a}_k s yet further. We start first by noticing there is an open substack of $\overline{\mathcal{M}}_{g,n}$ where \mathbf{a}_k makes sense without any change. We then review the extension to the case of curves of treelike type. The indeterminacy locus of \mathbf{a}_k s is contained in the complement of the union of both previous loci.

3.1. The locus of balanced curves. Given a family of curves $f : \mathcal{X} \rightarrow S$ and a line bundle \mathcal{L} over \mathcal{X} the set of points $s \in S$ where \mathcal{L}_s is q -stable in the sense of definition 1.1 is open in S , see for instance [Est01, 1.4].

Definition 3.1. A curve (X, \underline{p}) in $\overline{\mathcal{M}}_{g,n}$ is said to be $(\underline{\tau}, k)$ -balanced if for each subcurve

$$(3.1) \quad Z \subset X, \quad \sum_{p_i \in Z} \tau_i \geq k \deg(\omega_{X|Z}) - \frac{kZ}{2}$$

with strict inequality whenever $p_1 \in Z$.

This condition is the one for the q-stability of $\mathcal{O}_X(\sum_{i=1}^n \tau_i p_i) \otimes \omega_X^{-k}$. It defines an open substack $\mathcal{B}_{g,n}^{k,\underline{\tau}}$ of $\overline{\mathcal{M}}_{g,n}$. It is clear that the section \mathbf{a}_k extends to $\mathcal{B}_{g,n}^{k,\underline{\tau}}$. The extension is denoted by \mathbf{a}_k as well.

3.2. Further extensions and twistors. Over the complement of $\mathcal{B}_{g,n}^{k,\underline{\tau}}$ the line bundle $\mathcal{O}(\sum_i \tau_i D_i) \otimes \omega^{-k}$ is no more q-stable; the section \mathbf{a}_k does not make sense anymore as it is. Notice however that any extension of \mathbf{a}_k to a point in $\overline{\mathcal{M}}_{g,n} - \mathcal{B}_{g,n}^{k,\underline{\tau}}$ gives a point in the closure $\overline{\mathbf{a}_k \mathcal{B}_{g,n}^{k,\underline{\tau}}}$. Such a point in the image appears as the special fiber of a family $(f : \mathcal{X} \rightarrow \text{Spec}(R), \underline{D}, \mathcal{L})$ where R is a DVR and \mathcal{L} is a q-stable torsion-free sheaf whose restriction to the generic fiber \mathcal{X}_η is given by $\mathcal{O}_{\mathcal{X}_\eta}(\sum_i \tau_i D_{i,\eta}) \otimes \omega_{\mathcal{X}_\eta}^{-k}$. Properness of $\overline{\mathcal{F}}_{g,n}$ says that \mathcal{L} is unique up to tensor product with an invertible sheaf coming from the base. We shortly recall facts about twistors to make clear how to extend \mathbf{a}_k on a bigger substack in $\overline{\mathcal{M}}_{g,n}$. Basic facts about twistors can be found in [Cap12].

Let $(f : \mathcal{X} \rightarrow T, \underline{D})$ be a regular model over a DVR R having smooth generic fiber \mathcal{X}_η and central fiber X . Write X_v for the irreducible component of X corresponding to the vertex $v \in V$ of the dual graph of X . Over the generic smooth fiber the line $\mathcal{L}_\eta = \mathcal{O}_{\mathcal{X}_\eta}(\sum_i \tau_i D_i) \otimes \omega^{-k}$ is q-stable. The valuative criterion of properness for $\overline{\mathcal{F}}_{g,n}$ implies \mathcal{L}_η extends uniquely (up to tensor product by a pullback from the base) into a q-stable line bundle \mathcal{L} over \mathcal{X} . We are going to build such limits for treelike curves by tensoring the line bundle $\mathcal{O}_{\mathcal{X}}(\sum_i \tau_i D_i) \otimes \omega^{-k}$ with suitable twistors over \mathcal{X} . A treelike curve is a stable curve such that each edge joining two different vertices of its dual graph is separating, i.e. taking out that edge gives a graph with an extra connected component.

Recall that a twistor is a line bundle over X of the form $\mathcal{O}_{\mathcal{X}}(\sum_v \gamma_v X_v) \otimes \mathcal{O}_X$. The set of twistors Tw_f acts faithfully on $\mathcal{O}_{\mathcal{X}}(\sum_i \tau_i D_i) \otimes \omega^{-k}$ by tensor product.

Lemma 3.1. *If X is treelike then any q-stable line bundle L on X has $\underline{0}$ multidegree.*

Proof. If X has only one irreducible component there is nothing to prove. Assume X has at least 2 components. Let v be a leaf of the dual graph Γ_X and X_v the corresponding irreducible component. The q-stability condition states

$$-\frac{1}{2} \leq \deg_{X_v} L \leq \frac{1}{2}$$

and thus L has degree 0 on X_v . Write X' for $X_v^c = \overline{X - X_v}$. Taking any subcurve $Z = Y \cup X_v$ we have the inequality

$$-\frac{\kappa_Z}{2} \leq \deg_Z L \leq \frac{\kappa_Z}{2}.$$

The middle term is in fact equal to $\deg_Y L$ and $\kappa_Z = \kappa_{Y|X'}$, the number of intersection points of Y with Y^c in X' . We thus get the inequality

$$-\frac{\kappa_Y}{2} \leq \deg_Y L \leq \frac{\kappa_Y}{2}$$

in X' . Since this condition is harder than the one for Y in X , it is enough to check the multidegree of L on X' is 0. Since X' has less components than X the result is obtained by induction. \square

Remark 4. In general a q-stable line bundle is not of multidegree $\underline{0}$. The simplest example comes from a banana curve.

Lemma 3.2. *Assume X is treelike. There is then a unique q -stable line bundle in the orbit of $\mathcal{O}_{\mathcal{X}}(\sum_i \tau_i D_i) \otimes \omega^{-k}$ under the action of Tw_f .*

Proof. The point is to prove that, given a line bundle \mathcal{L} on \mathcal{X} , there is a twister sending $\mathcal{L}|_X$ to a multidegree $\underline{0}$ line bundle on X . A short answer would be to recall that the class group of a treelike graph is trivial, see for instance [Cap12]. We shall however go more into details in order to keep track of how to get the desired twister.

Let Γ be the dual graph of X and V the set of vertices of Γ . The irreducible component of X corresponding to a vertex $v \in V$ is denoted X_v . Let L be the restriction $\mathcal{L}|_X$ and write $(m_v^0)_{v \in V}$ for the multidegree of L on $X = X_0$. Since Γ is treelike there is a filtration covering Γ

$$\Gamma = \Gamma_0 \supsetneq \Gamma_1 \supsetneq \cdots \supsetneq \Gamma_{|V|-1} \supsetneq \emptyset$$

built up by choosing leaves v_r of the loopless graph obtained out of Γ_r by contracting possible loops and satisfying

$$\Gamma_{r+1} = \Gamma_r - v_r.$$

The right hand side of the previous equality is short for the subgraph of Γ_r generated by all vertices except v_r . It does contain all edges that are not half-edges of v_r or in geometric edges incident to v_r . For each v_r let B_{v_r} be the branch of Γ out of v_r ; removing the unique edge linking v_r to Γ_{r+1} , B_{v_r} is the component containing v_r . Define on \mathcal{X} the following sequence $(\mathcal{L}_r)_{0 \leq r \leq |V|-1}$ of line bundles

$$(3.2) \quad \begin{aligned} (1) \quad & \mathcal{L}_0 = \mathcal{L} \\ (2) \quad & \text{let } L_r = \mathcal{L}_r|_X \text{ and write } (m_v^r)_{v \in V} \text{ for the multidegree of } L_r, \text{ then} \\ & \forall r \geq 0, \quad \mathcal{L}_{r+1} = \mathcal{L}_r \left(m_{v_r}^r \sum_{w \in B_{v_r}} X_w \right). \end{aligned}$$

Each L_r is obtained out of $L = L_0$ by tensor product with a given twister. Let F_r be the set of vertices of Γ not in Γ_r . This is a strictly increasing finite collection of sets.

Claim. For all $j \geq r$, L_j has zero degree on each component X_v for $v \in F_r$.

Notice first that each $v \in F_r$ is of the form v_ℓ for some $\ell \leq r$. It is thus enough to show that for all $j \geq r$, L_j has degree 0 on v_r . The claim is clear if $r = 0$, assume $r \geq 1$. The degree of L_r on v_r is given by

$$(3.3) \quad \deg(L_r|_{X_{v_r}}) = \underbrace{\deg(L_{r-1}|_{X_{v_{r-1}}})}_{(I)} + \underbrace{\deg\left(\mathcal{O}_{\mathcal{X}}\left(m_{v_{r-1}}^{r-1} \sum_{w \in B_{v_{r-1}}} X_w\right)\right)|_{X_{v_{r-1}}}}_{(II)}.$$

By definition (I) is equal to $m_{v_{r-1}}^{r-1}$. The term (II) is equal to

$$m_{v_{r-1}}^{r-1} (\text{val}(v_r) - 1) - m_{v_{r-1}}^{r-1} \text{val}(v_r) = -m_{v_{r-1}}^{r-1}.$$

The first term of the left hand side is the contribution of neighboring vertices of v_{r-1} while the second one comes from the self-intersection of $X_{v_{r-1}}$. The total sum (I) + (II) is thus 0 as claimed. Now, for any $j > r$ we have that

$$\deg(L_j|_{X_{v_r}}) = \underbrace{\deg(L_r|_{X_{v_r}})}_{=0} + \sum_{r < \ell \leq j} \deg\left(\mathcal{O}_{\mathcal{X}}\left(m_{v_r}^r \sum_{w \in B_{v_r}} X_w\right)\right)|_{X_{v_r}}.$$

If $v_r \notin B_{v_\ell}$ for some ℓ then the corresponding term in the right hand sum is clearly 0. Otherwise we have that

$$\deg \left(\mathcal{O}_X(m_{v_r}^r \sum_{w \in B_\ell} X_w)_{|X_{v_r}} \right) = m_{v_r}^r \text{val}(v_r) - m_{v_r}^r \text{val}(v_r) = 0$$

where the first right hand term comes from contributions of neighbors of v_r , the second from the self intersection of X_{v_r} .

Our claim is proved. We have shown that we have an increasing filtration by F_r s that covers V and on which $\mathcal{L}_r|_X$ has $\underline{0}$ multidegree. There is therefore a line bundle \mathcal{L}_∞ whose restriction to X has multidegree $\underline{0}$. \square

Remark 5. Each component of the multidegree $(m_v^r)_{v \in V}$ of L_r can be written explicitly in terms of the combinatorics of Γ . We have already seen during the previous proof that given $v \in F_r$

$$\forall j \geq r, \quad m_v^j = \deg(L_j|_{X_v}) = 0.$$

In general we have the relation

$$\deg(L_j|_{X_v}) = \deg(L_{j-1}|_{X_v}) + \deg \left(\mathcal{O}_X(m_{v_{j-1}}^{j-1} \sum_{w \in B_{v_j}} X_w)_{|X_v} \right)$$

which can also be written

$$m_v^j = m_v^{j-1} + \deg \left(\mathcal{O}_X(m_{v_{j-1}}^{j-1} \sum_{w \in B_{v_j}} X_w)_{|X_v} \right).$$

When $j < r$, the vertex v doesn't appear in $B_{v_{j-1}}$. The second term of the previous right hand side is 0 unless v_j is adjacent to v , in which case it is 1. Let $\chi_{v_{j-1}}$ be the function whose value is 0 on a vertex v if v_{j-1} is adjacent to v and 1 otherwise. We thus have that

$$m_v^j = m_v^{j-1} + \chi_{v_{j-1}}(v) m_{v_{j-1}}^{j-1}.$$

An inductive argument thus gives

$$m_v^j = \sum_{\ell < j} \chi_{v_\ell}(v) m_{v_\ell}^\ell = \sum_{w \in B_v} m_w.$$

In particular one can write \mathcal{L}_{r+1} as

$$(3.4) \quad \mathcal{L}_{r+1} = \mathcal{L}_r \left(\left(\sum_{w \in B_{v_r}} m_w \right) \sum_{w \in B_{v_r}} X_w \right).$$

3.3. Defining a partial extension. Let $\overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}}$ be the open substack union of the locus of curves of treelike type and the one of $(\underline{\tau}, k)$ -balanced ones $\mathcal{B}_{g,n}^{k,\underline{\tau}}$. This open substack can be described as the complement in $\overline{\mathcal{M}}_{g,n}$ of the following closed locus: let $\mathcal{I}_{g,n}^{k,\underline{\tau}}$ be the closure in $\overline{\mathcal{M}}_{g,n}$ of curves having topological type a two vertex loopless graph, which is not $(\underline{\tau}, k)$ -balanced; then $\overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}} = \overline{\mathcal{M}}_{g,n} - \mathcal{I}_{g,n}^{k,\underline{\tau}}$. Write $\overline{\mathcal{J}}_{g,n}^{k,\underline{\tau}}$ for the substack of $\overline{\mathcal{J}}_{g,n}$ obtained by pullback along $\overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}} \subset \overline{\mathcal{M}}_{g,n}$.

Recall that in $\overline{\mathcal{M}}_{g,n+1}$ the divisor $\delta_{h,A}$ is the closure of stable marked curves of topological type a graph having only one edge and two vertices, one of which is of genus h and whose set of legs (marked points) is A . The divisor D_i corresponding to the image of the sections of $\pi : \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ is simply $\delta_{0,\{n+1,i\}}$.

Proposition 3.3. *Let $\mathcal{L}(\underline{\tau}, k)$ be the line bundle on $\overline{\mathcal{M}}_{g,n+1}$ given by*

$$(3.5) \quad \mathcal{L}(\underline{\tau}, k) = \mathcal{O}\left(\sum_{i=1}^n \tau_i D_i\right) \otimes \omega^{-k} \otimes \mathcal{O}\left(\sum_{\substack{0 \leq h \leq \lfloor g/2 \rfloor \\ A \subset \{1, \dots, n\} \\ 2 \leq |A| + h \leq g + n - 2}} [k(1-2h) + \sum_{i \in A} \tau_i] \delta_{h, A \cup \{n+1\}}\right).$$

Then $\mathcal{L}(\underline{\tau}, k)$ defines a section $\bar{\mathbf{a}}_k : \overline{\mathcal{M}}_{g,n}^{k,\underline{\tau}} \rightarrow \overline{\mathcal{J}}_{g,n}^{k,\underline{\tau}}$ extending \mathbf{a}_k .

Proof. We need to check that along fibers of π the line bundle $\mathcal{L}(\underline{\tau}, k)$ is q-stable, i.e. has multidegree $\underline{0}$. Let (X, \underline{p}) be a treelike stable n -marked curve over a point. Write Γ for the dual graph of X and V for its set of vertices. Let X_v be the irreducible component of X corresponding to $v \in V$ and L_v the set of marked points of X_v .

Choose a regular smoothing $f : \mathcal{X} \rightarrow S$ of X and let D_i be the corresponding marked point of f . If X is a point in $\delta_{h,A} \subset \overline{\mathcal{M}}_{g,n}$ write $\Lambda_{h,A}$ for the subgraph of Γ that comes from a specialization of the genus h A -marked vertex of the generic point of $\delta_{h,A}$. The pull back of $\delta_{h, A \cup \{n+1\}}$ over f gives the vertical divisor

$$\Xi_{h,A} = \sum_{v \in \Lambda_{h,A}} X_v.$$

The $\delta_{h,A}$ contribution in 3.5 is given by

$$(3.6) \quad [k(1-2h) + \sum_{i \in A} \tau_i] \Xi_{h,A}.$$

Now, writing $\hat{g}(v) = g(v) + b_1(v)$ where $g(v)$ is the genus of v (its weight in the dual graph) and $b_1(v)$ is the number of loops v has, we have that

$$(3.7) \quad k(1-2h) + \sum_{i \in A} \tau_i = \sum_{v \in \Lambda_{h,A}} (k(2-2\hat{g}(v) + \kappa_v) + \sum_{i \in A \cap L_v} \tau_i).$$

When X is $(\underline{\tau}, k)$ -balanced 3.7 is 0 by assumption. Thus $\mathcal{L}(\underline{\tau}, k)$ is the desired line bundle in the $(\underline{\tau}, k)$ -balanced case. When X is treelike notice that each subgraph $\Lambda_{h,A}$ is a branch of Γ . The vertex of that branch corresponds to the nodal point linking the subcurve of X in $\overline{\mathcal{M}}_{h,A}$ to the one in $\overline{\mathcal{M}}_{g-h, \mathbb{C}A}$. One can order $\delta_{h, A \cup \{n+1\}}$ s in such a way that complements of $\Lambda_{h,A}$ s give a decreasing finite cover of Γ as in the setting we started with in the proof of lemma 3.2. From 3.7 and 3.6 the contribution of $\delta_{h, A \cup \{n+1\}}$ is given by

$$\mathcal{O}_{\mathcal{X}} \left(\left[\sum_{v \in \Lambda_{h,A}} (k(2-2\hat{g}(v) + \kappa_v) + \sum_{i \in A \cap L_v} \tau_i) \right] \sum_{v \in \Lambda_{h,A}} X_v \right)$$

By 3.4 this is precisely the contribution we need to build up a q-stable line bundle out of $\mathcal{O}_{\mathcal{X}}(\sum_{i=1}^n \tau_i D_i) \otimes \omega^{-k}$ by tensoring with twistors. \square

3.4. Obstructions to extend Abel-Jacobi maps yet further. There are two obstructions to yet further extend the Abel-Jacobi maps. The first is a failure of the approach in its present shape; one can obtain too many q-stable limits out of a given line bundle by tensoring with twistors. More details on this feature can be found in [Cap08, 4]. The second is of more geometric nature.

Combinatorial obstruction. An example of the first phenomenon is given by the stable curve (X, \underline{p}) having dual graph a cycle with 3 vertices. Write X_1, X_2 and X_3 for the three irreducible components of X and assume $p_1 \in X_1$. The two multidegrees $(0, -1, 1)$ and $(1, 0, -1)$ are q-stable. Assuming that $(f : \mathcal{X} \rightarrow S, \underline{D})$ is a regular one parameter smoothing of (X, \underline{p}) , then one can go from the first to the second multidegree by tensoring \mathcal{X} with the twister $\mathcal{O}_{\mathcal{X}}(X_2)$. There is thus no unique twister on \mathcal{X} sending $\mathcal{O}_{\mathcal{X}}(\sum_i \tau_i D_i)$ on a q-stable line bundle.

Geometry. The previous phenomenon doesn't happen for banana curves, which are obtained by identifying two distinct points on a smooth curves with two others from another curve. Given such a curve (X, \underline{p}) , there are only two possible q-stable multidegrees given by $(0, 0)$ and $(1, -1)$, the first coordinate corresponding to the component marked by p_1 . Given a regular one parameter smoothing $(f : \mathcal{X} \rightarrow S, \underline{D})$ there is only one twister getting $\mathcal{O}_{\mathcal{X}}(\sum_i \tau_i D_i)$ to a q-stable line bundle having either degree $(0, 0)$ or $(1, -1)$. The expected multidegree is $(0, 0)$ if the multidegree of the line bundle $\mathcal{O}_X(\sum_i \tau_i p_i)$ is even and $(1, -1)$ if it is odd. Despite this pleasant feature, this construction cannot be globalized to the universal family. The reason is that the corresponding twister would have to be trivial over the locus of irreducible curves and thus only have support over the one of banana curves. This locus is of codimension 2 in $\overline{\mathcal{M}}_{g,n+1}$ and cannot correspond to a divisor on $\overline{\mathcal{M}}_{g,n+1}$. This strongly suggests that one needs to blow up $\overline{\mathcal{M}}_{g,n}$ along the locus of none $(\underline{\tau}, k)$ -stable banana curves to be able to extend $\overline{\mathbf{a}}_k$ yet further.

Among the possible pathological phenomena we could have faced, the one where there is no q-stable limit that is a line bundle does not occur. This is a general fact.

Lemma 3.4. *Let (X, \underline{p}) be a marked stable curve, $f : \mathcal{X} \rightarrow S$ a one parameter smoothing of X and \mathcal{L} a line bundle on \mathcal{X} , with $L = \mathcal{L}|_X$. Then there is always a q-stable line bundle on X coming from an element in the orbit of \mathcal{L} under the action of Tw_f .*

Proof. By [Cap08, 4.12] we can assume the restriction L of \mathcal{L} on the central fiber is semistable. Let $Y \subset X$ be the irreducible component of X containing p_1 . If L is semistable then

$$\deg_Y L = -\frac{\kappa_Y}{2}.$$

Let Z_1, \dots, Z_m be the connected components of Y^c . We have that $\kappa_Y = \kappa_{Z_1} + \dots + \kappa_{Z_m}$. Using the fact $\deg_{Y^c} L = -\deg_Y L$ and the stability condition on each Z_j we get that

$$\deg_{Z_j} L = \frac{\kappa_{Z_j}}{2}.$$

Tensoring \mathcal{L} by the twister given by $\mathcal{O}_{\mathcal{X}}(Y)$ we get a line bundle L' equal to L on components of X not neighboring Y and satisfying

$$\deg_Y L' = \frac{\kappa_Y}{2} \quad \text{and} \quad \deg_{Z_j} L' = -\frac{\kappa_{Z_j}}{2}.$$

It is enough to claim L' is q-stable. □

4. PULLBACKS ALONG ABEL-JACOBI MAPS

4.1. **Pulling back the zero section.** Our goal is to compute the pullbacks of the class of the zero section $[\mathcal{Z}_{g,n}]$ in $\overline{\mathcal{J}}_{g,n}^{k,\mathcal{I}}$ along $\bar{\mathbf{a}}_k$ s. Recall that

$$(2.1) \quad [\mathcal{Z}_{g,n}] = \left\{ \exp \left(\sum_{s \geq 1} (-1)^s (s-1)! \{ \phi_* (\text{ch}(\mathcal{L}) \text{Td}^\vee(\Omega_\phi)) \}_s \right) \right\}_g$$

where $\{\bullet\}_\ell$ is the degree ℓ part of \bullet . One can picture the involved data by looking at the commutative diagram

$$(4.1) \quad \begin{array}{ccc} \mathcal{L} & \longrightarrow & \overline{\mathcal{J}}_{g,n|1}^{k,\mathcal{I}} \xrightarrow{\epsilon_{n|1}} \overline{\mathcal{M}}_{g,n+1}^{k,\mathcal{I}} \longrightarrow \mathcal{L}(\mathcal{I}, k) \\ & & \downarrow \phi \qquad \qquad \downarrow \pi \\ & & \overline{\mathcal{J}}_{g,n}^{k,\mathcal{I}} \xrightarrow{\epsilon_n} \overline{\mathcal{M}}_{g,n}^{k,\mathcal{I}} \\ & & \longleftarrow \bar{\mathbf{a}}_k \end{array}$$

Notice that we have a tautological lift $\hat{\mathbf{a}}_k : \overline{\mathcal{M}}_{g,n+1}^{k,\mathcal{I}} \rightarrow \overline{\mathcal{J}}_{g,n|1}^{k,\mathcal{I}}$ simply given by sending an S -section $(f : \mathcal{X} \rightarrow S, \underline{D}, D_{n+1})$ on $(f : \mathcal{X} \rightarrow S, \underline{D}, D_{n+1}, \mathcal{L}(\mathcal{I}, k))$. The corresponding S -object in $\overline{\mathcal{J}}_{g,n}^{k,\mathcal{I}}$ is obtained by simply forgetting the extra section D_{n+1} . We can thus complete diagram 4.1 to get

$$(4.2) \quad \begin{array}{ccc} \mathcal{L} & \longrightarrow & \overline{\mathcal{J}}_{g,n|1}^{k,\mathcal{I}} \xrightarrow{\epsilon_{n|1}} \overline{\mathcal{M}}_{g,n+1}^{k,\mathcal{I}} \longrightarrow \mathcal{L}(\mathcal{I}, k) \\ & & \downarrow \phi \qquad \qquad \downarrow \pi \\ & & \overline{\mathcal{J}}_{g,n}^{k,\mathcal{I}} \xrightarrow{\epsilon_n} \overline{\mathcal{M}}_{g,n}^{k,\mathcal{I}} \\ & & \longleftarrow \bar{\mathbf{a}}_k \end{array}$$

Lemma 4.1. *On the level of Chow groups we have the relation $\bar{\mathbf{a}}_k^* \phi_* = \pi_* \hat{\mathbf{a}}_k^*$.*

Proof. This is a standard argument. It is straightforward to check that the red part of diagram 4.2 is in fact cartesian. Now given an element α in the Chow group of $\overline{\mathcal{J}}_{g,n|1}^{k,\mathcal{I}}$. We have that

$$\bar{\mathbf{a}}_k^* \phi_*(\alpha) = \epsilon_{n*}(\phi_*(\alpha) \cap [\mathcal{Z}_{g,n}]) = \epsilon_{n*} \phi_*(\alpha \cap \phi^*[\mathcal{Z}_{g,n}])$$

since ϕ is flat we get that

$$\bar{\mathbf{a}}_k^* \phi_*(\alpha) = \pi_* \epsilon_{n|1*}(\alpha \cap [\phi^{-1} \mathcal{Z}_{g,n}])$$

which, because the red part of diagram 4.2 is cartesian, gives

$$\bar{\mathbf{a}}_k^* \phi_*(\alpha) = \pi_* \hat{\mathbf{a}}_k^*(\alpha).$$

□

Theorem 4.2. *The pullbacks $\bar{\mathbf{a}}_k^*[\mathcal{Z}_{g,n}]$ are given over $\overline{\mathcal{M}}_{g,n}^{k,\mathcal{I}}$ by the relation*

$$(4.3) \quad \bar{\mathbf{a}}_k^*[\mathcal{Z}_{g,n}] = \left\{ \exp \left(\sum_{s \geq 1} (-1)^s (s-1)! \{ \pi_* (\text{ch}(\mathcal{L}(\mathcal{I}, k)) \text{Td}^\vee(\Omega_\pi)) \}_s \right) \right\}_g$$

where $\{\bullet\}_\ell$ is the degree ℓ part of \bullet .

Proof. This is a direct consequence of the previous lemma including the fact the red part of 4.2 is cartesian. □

Remark 6. Formula 4.3 is an explicit polynomial in the tautological classes of $\overline{\mathcal{M}}_{g,n}$. Unfortunately, it is not clear how to get back Hain's formula or its extension in [GZ14b] by directly manipulating 4.3.

4.2. Pullback of the theta divisor to $\overline{\mathcal{M}}_{g,n}$. We follow notation of [ACG11, 17]. Write \tilde{K} for the Chern class of ω and ψ_i for the Chern class of the cotangent bundle along the marked section D_i . The ψ_i classes can be described as push-forwards under π by the relation $\psi_i = -\pi_*(D_i^2)$. Using relation 2.10 we have that

$$(4.4) \quad \bar{a}_k^*[\Theta] = -\pi_* \left(\frac{c_1(\mathcal{L}(\underline{t}, k))^2}{2} \right)$$

Assume that each time a subset A of $\{1, \dots, n\}$ indexes a sum we have that $2 \leq |A| + h \leq g + n - 2$. Unravelling the right hand side of 4.4 we have

$$\begin{aligned} \bar{a}_k^*[\Theta] = -\pi_* \left[\right. & \frac{1}{2} \sum_{i=1}^n \tau_i^2 D_i^2 \\ & + \sum_{\substack{0 \leq h \leq \lfloor g/2 \rfloor \\ A \subset \{1, \dots, n\}}} \sum_{i=1}^n \tau_i (\gamma_h + \sum_{j \in A} \tau_j) D_i \delta_{h, A \cup \{n+1\}} \\ & + \frac{1}{2} \sum_{\substack{0 \leq h, l \leq \lfloor g/2 \rfloor \\ A, B \subset \{1, \dots, n\}}} (\gamma_h + \sum_{j \in A} \tau_j) (\gamma_l + \sum_{j \in B} \tau_j) \delta_{h, A \cup \{n+1\}} \delta_{l, B \cup \{n+1\}} \\ & \left. + \frac{k^2 \tilde{K}^2}{2} - k \sum_{i=1}^n \tau_i D_i \tilde{K} - k \sum_{\substack{0 \leq h \leq \lfloor g/2 \rfloor \\ A \subset \{1, \dots, n\}}} (\gamma_h + \sum_{i \in A} \tau_i) \tilde{K} \delta_{h, A \cup \{n+1\}} \right]. \end{aligned}$$

where γ_h stands for $k(1 - 2h)$. Taking into account that

- $\tilde{K} D_i = -D_i^2$
- $\pi_*(D_i \delta_{h, A \cup \{n+1\}})$ is zero unless $i \in A$, in which case it is $\delta_{h, A}$
- $\pi_*(\delta_{h, A \cup \{n+1\}} \delta_{l, B \cup \{n+1\}})$ is zero unless $A = B$ and $h = l$ in which case it is equal to $-\delta_{h, A}$
- $\pi_*(\tilde{K} \delta_{h, A \cup \{n+1\}}) = (2h - 1) \delta_{h, A}$

we get the expression

$$(4.5) \quad \bar{a}_k^*[\Theta] = \sum_{i=1}^n \left(\frac{\tau_i^2}{2} + k\tau_i \right) \psi_i - \frac{k^2}{2} \tilde{\kappa}_1 - \frac{1}{2} \sum_{\substack{0 \leq h \leq \lfloor g/2 \rfloor \\ A \subset \{1, \dots, n\}}} \left(k(1 - 2h) + \sum_{i \in A} \tau_i \right)^2 \delta_{h, A}$$

where for each A we assume $2 \leq |A| + h \leq g + n - 2$. For $k = 0$ rearranging terms we recover the relation 0.2. More generally, we recover back Hain's formula.

4.3. Pulling back the theta divisor in degree $g - 1$. We briefly show how to adapt the present strategy to compute the pullback of the theta divisor on the compactification $\overline{\mathcal{J}}_{g,n}^{g-1}$ of the degree $g - 1$ universal jacobian with canonical trivial polarization \mathcal{O} . A point of $\overline{\mathcal{J}}_{g,n}^{g-1}$ is a tuple (X, \underline{p}, L) where (X, \underline{p}) is a stable n -marked curve and L is a q -stable torsion-free rank 1 sheaf on \tilde{X} . The compactification $\overline{\mathcal{J}}_{g,n}^{g-1}$ has a well defined theta divisor given by

$$(4.6) \quad \Theta_{g-1} = \{(X, L) \mid h^0(L) \geq 1\}.$$

See for instance [Ale04] or [Cap09]. In particular one can write down the class of the theta divisor as

$$(4.7) \quad -[\Theta_{g-1}] = \phi_* \left(\frac{c_1(\mathfrak{L}_{g-1})^2}{2} \right) - \phi_* \left(\frac{c_1(\mathfrak{L}_{g-1})c_1(\omega)}{2} \right) + \lambda_1$$

where $\phi : \overline{\mathcal{J}}_{g,n|1}^{g-1} \rightarrow \overline{\mathcal{J}}_{g,n}^{g-1}$ is the universal curve and \mathfrak{L}_{g-1} the universal torsion-free sheaf of rank 1 over $\overline{\mathcal{J}}_{g,n|1}^{g-1}$. Given an n -tuple of integers $\underline{\tau} = (\tau_1, \dots, \tau_n)$ such that $\sum_i \tau_i = g-1$ we get on $\mathcal{M}_{g,n}$ a section $\mathbf{a}^{g-1} : \mathcal{M}_{g,n} \rightarrow \overline{\mathcal{J}}_{g,n}^{g-1}$ sending points (X, \underline{p}) on $(X, \underline{p}, \mathcal{O}_X(\sum_i \tau_i p_i))$. To pullback the theta divisor on $\overline{\mathcal{J}}_{g,n}^{g-1}$ it is enough to extend \mathbf{a}^{g-1} to codimension 1 loci given by stable curves having 1 node. The Abel-Jacobi map \mathbf{a}^{g-1} extends as it is to the case of irreducible curves. For a curve of compact type (X, \underline{p}) notice that a q -stable line bundle L is of multidegree

$$\deg_{X_v} L = g(v) - \chi_{p_1^c}(v)$$

where $\chi_{p_1^c}(v)$ is 0 if $p_1 \notin X_v$ and 1 otherwise. The multidegree of L does thus only depend on (X, \underline{p}) . Twisting the line bundle $\mathcal{O}(\sum_i \tau_i D_i)$ over the locus $\overline{\mathcal{M}}_{g,n}^{\leq 1}$ of stable curves having at most one node we get that

$$\mathcal{O}\left(\sum_i \tau_i D_i\right) \otimes \mathcal{O}\left(\sum_{\substack{0 \leq h \leq \lfloor g/2 \rfloor \\ A \subset \{1, \dots, n\} \\ 2 \leq |A| \leq g+n-2}} \left(\sum_{i \in A} \tau_i - h + \chi_{A^c}(D_1)\right) \delta_{h, A \cup \{n+1\}}\right)$$

is a q -stable line bundle. It does thus define an extension $\bar{\mathbf{a}}^{g-1}$ to $\overline{\mathcal{M}}_{g,n}^{\leq 1}$. Pulling formula 4.7 along $\bar{\mathbf{a}}^{g-1}$ and using the same relations on tautological classes as the previous section we get that

$$(4.8) \quad \bar{\mathbf{a}}^{g-1*}[\Theta_{g-1}] = \sum_{i=1}^n \frac{\tau_i(\tau_i + 1)}{2} \psi_i - \lambda_1 - \sum_{\substack{0 \leq h \leq \lfloor g/2 \rfloor \\ A \subset \{1, \dots, n\}}} \left(\sum_{i \in A} \tau_i - h\right) \left(\sum_{i \in A} \tau_i - h + 1\right) \frac{\delta_{h,A}}{2}$$

where $2 \leq |A| + h \leq g+n-2$. Notice that this class differs from the one computed by Grushevsky and Zakharov in [GZ14a] by $\delta_{irr}/8$. The reason is that the theta divisor here is not given by the vanishing of a theta function but using description 4.6. As is the case in [GZ14a], formula 4.8 generalizes a formula given by Müller in [Mül13]. We shortly recall their arguments. The locus $\overline{\mathcal{D}}_g$ computed by Müller is the closure in $\overline{\mathcal{M}}_{g,n}$ of

$$\mathcal{D}_g = \left\{ (X, \underline{p}) \mid h^0\left(\mathcal{O}_X\left(\sum_i \tau_i p_i\right)\right) \geq 1 \right\}$$

for smooth X and at least one negative τ_i . The locus \mathcal{D}_g is given by the pullback $(\mathbf{a}^{g-1})^* \Theta_{g-1}$ on $\mathcal{M}_{g,n}$. Over the boundary locus $\mathcal{M}_{h,A,x} \times \mathcal{M}_{g-h,A^c,y}$ the Abel-Jacobi map $\bar{\mathbf{a}}^{g-1}$ sends a curve $X = X_h \cup X_{g-h}$ on the line bundle

$$\left(\mathcal{O}_{X_h} \left(\sum_{i \in A} \tau_i D_i - \left(\sum_{i \in A} \tau_i - h + 1 \right) x \right), \mathcal{O}_{X_{g-h}} \left(\sum_{i \in A^c} \tau_i D_i + \left(\sum_{i \in A} \tau_i - h \right) y \right) \right).$$

If any of the entries of the first factor is negative the image of $\mathcal{M}_{h,A,x} \times \mathcal{M}_{g-h,A^c,y}$ by $\bar{\mathbf{a}}^{g-1}$ doesn't lie in the support of Θ_{g-1} . The pullback of Θ_{g-1} over this locus is in a codimension 2 locus of $\overline{\mathcal{M}}_{g,n}$. The divisors $\bar{\mathbf{a}}^{g-1*}[\Theta_{g-1}]$ and $[\overline{\mathcal{D}}_g]$ are thus equal on each such locus. When all entries are positive $\bar{\mathbf{a}}^{g-1}$ sends $\mathcal{M}_{h,A,x} \times \mathcal{M}_{g-h,A^c,y}$ into the support of Θ_{g-1} . To get $[\overline{\mathcal{D}}_g]$ one needs to correct

$\bar{a}^{g-1}[\Theta_{g-1}]$ by subtracting the multiplicity of \bar{a}^{g-1} along the previous locus. By the Riemann singularity theorem we get

$$[\bar{\mathcal{D}}_g] = \bar{a}^{g-1*}[\Theta_{g-1}] - \sum_{(h,A) \in E_+} \left(h - \sum_{i \in A} \tau_i \right) \delta_{h,A}$$

where E_+ is the subset of couples (h, A) having the usual constraints as in 4.8, for each $i \in A$, τ_i is positive and $h \geq \sum_{i \in A} \tau_i$.

REFERENCES

- [ACG11] E. Arbarello, M. Cornalba, and P. A. Griffiths. *Geometry of Algebraic Curves vol II*. Springer, 2011.
- [Ale04] Valery Alexeev. Compactified Jacobians and Torelli map. *Publ. Res. Inst. Math. Sci.*, 40(4):1241–1265, 2004.
- [Cap94] Lucia Caporaso. A compactification of the universal picard variety over the moduli space of stable curves. *J. Amer. Math. Soc.*, 7:589–660, 1994.
- [Cap08] Lucia Caporaso. Néron models and compactified Picard schemes over the moduli stack of stable curves. *Amer. J. Math.*, 130(1):1–47, 2008.
- [Cap09] Lucia Caporaso. Geometry of the theta divisor of a compactified Jacobian. *J. Eur. Math. Soc. (JEMS)*, 11(6):1385–1427, 2009.
- [Cap12] Lucia Caporaso. Compactified Jacobians of Néron type. *Atti Accad. Naz. Lincei, Cl. Sci. Fis. Mat. Nat., IX. Ser., Rend. Lincei, Mat. Appl.*, 23(2):213–227, 2012.
- [CMW12] Renzo Cavalieri, Steffen Marcus, and Jonathan Wise. Polynomial families of tautological classes on $\mathcal{M}_{g,n}^{\text{st}}$. *J. Pure Appl. Algebra*, 216(4):950–981, 2012.
- [Cor89] Maurizio Cornalba. Moduli of curves and theta-characteristics. In *Lectures on Riemann surfaces (Trieste, 1987)*, pages 560–589. World Sci. Publ., Teaneck, NJ, 1989.
- [EP14] E. Esteves and M. Pacini. Semistable modifications of families of curves and compactified Jacobians. *ArXiv e-prints*, June 2014.
- [Est01] Eduardo Esteves. Compactifying the relative Jacobian over families of reduced curves. *Trans. Am. Math. Soc.*, 353(8):3045–3095, 2001.
- [Ful98] W. Fulton. *Intersection theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete 3.Folge. A series of modern surveys in mathematics. Springer-Verlag, 1998.
- [GZ14a] Samuel Grushevsky and Dmitry Zakharov. The double ramification cycle and the theta divisor. *Proc. Am. Math. Soc.*, 142(12):4053–4064, 2014.
- [GZ14b] Samuel Grushevsky and Dmitry Zakharov. The zero section of the universal semiabelian variety and the double ramification cycle. *Duke Math. J.*, 163(5):953–982, 2014.
- [Hai13] Richard Hain. Normal functions and the geometry of moduli spaces of curves. In *Handbook of moduli. Vol. I*, volume 24 of *Adv. Lect. Math. (ALM)*, pages 527–578. Int. Press, Somerville, MA, 2013.
- [Mel] Margarida Melo. Compactifications of the universal jacobian over curves with marked points. In preparatoion.
- [Mel11] Margarida Melo. Compactified Picard stacks over the moduli stack of stable curves with marked points. *Adv. Math.*, 226(1):727–763, 2011.
- [Mül13] Fabian Müller. The pullback of a theta divisor to $\overline{\mathcal{M}}_{g,n}$. *Math. Nachr.*, 286(11-12):1255–1266, 2013.
- [Mum08] David Mumford. *Abelian varieties*, volume 5 of *Tata Institute of Fundamental Research Studies in Mathematics*. Published for the Tata Institute of Fundamental Research, Bombay; by Hindustan Book Agency, New Delhi, 2008. With appendices by C. P. Ramanujam and Yuri Manin, Corrected reprint of the second (1974) edition.
- [MV14] Margarida Melo and Filippo Viviani. The Picard group of the compactified universal Jacobian. *Doc. Math., J. DMV*, 19:457–507, 2014.
- [MW13] S. Marcus and J. Wise. Stable maps to rational curves and the relative Jacobian. *ArXiv e-prints*, October 2013.
- [Pan96] Rahul Pandharipande. A compactification over $\overline{\mathcal{M}}_g$ of the universal moduli space of slope-semistable vector bundles. *J. Amer. Math. Soc.*, 9(2):425–471, 1996.

BASHAR DUDIN, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DE COIMBRA, APARTADO 3008, EC SANTA CRUZ, 3001 - 501 COIMBRA, PORTUGAL
E-mail address: dudin@mat.uc.pt