



HAL
open science

Counterflow in a doubly superfluid mixture of Bosons and Fermions

F Chevy

► **To cite this version:**

F Chevy. Counterflow in a doubly superfluid mixture of Bosons and Fermions. *Physical Review A: Atomic, molecular, and optical physics* [1990-2015], 2015, 91, pp.063606. <hal-01152336>

HAL Id: hal-01152336

<https://hal.science/hal-01152336v1>

Submitted on 15 May 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Counterflow in a doubly superfluid mixture of Bosons and Fermions

F. Chevy¹

¹*Laboratoire Kastler Brossel, ENS-PSL Research University,
CNRS, UPMC-Sorbonne Universités, Collège de France*

In this article, we calculate the friction between two counter-flowing bosonic and fermionic superfluids. In the limit where the boson-boson and boson-fermion interactions can be treated within the mean-field approximation, we show that the force can be related to the dynamical structure factor of the fermionic component. Finally, we provide asymptotic expressions for weakly and strongly attractive fermions and show that the damping rate obeys simple scaling laws close to the critical velocity.

PACS numbers: 67.85.-d, 67.10.-j, 67.85.Pq, 67.85.De

I. INTRODUCTION

The onset of frictionless flow in quantum fluids is probably one of the most intriguing macroscopic manifestations of quantum mechanics. Recent experiments on Bose-Fermi superfluid mixtures gave a new twist to this old question by probing the critical velocity of a superfluid counterflow [1]. When two miscible superfluids flow through each-other, Castin *et al.* suggested a generalization of the celebrated Landau criterion where superfluidity is destroyed by the shedding of a pair of elementary excitations in the two systems [2]. Later-on, this scenario was supported by the study of the lifetime of the quasi-particles [3, 4] or by the calculation of the hydrodynamic spectrum [5].

The results put forward in [2] were based on heuristic arguments and were focusing on the velocity threshold above which the counter-flow is damped. In this work, we provide a full microscopic treatment of the friction in a superfluid counterflow and we determine the explicit velocity dependence of the damping force above the critical velocity. Assuming that the boson-boson and boson-fermion interactions can be treated within the mean-field approximation, we show that the force can be related to the dynamic structure factor of the fermionic superfluid. Although the general expression of the structure factor of an attractive Fermi gas is not known exactly in the crossover between the Bardeen-Cooper-Schrieffer (BCS) and molecular Bose-Einstein Condensate (BEC) regimes, we provide asymptotic expressions in the limits of weak and strong interactions where the fermionic component behaves respectively as an ideal Fermi gas and an hydrodynamic Bose-Einstein condensate of dimers [6]. We find that close to the critical velocity, the force obeys simple scaling laws, with exponent depending on the value of the fermion-fermion scattering length.

II. GENERAL SETTING

We consider a mixture between a fermionic superfluid and a weakly interacting Bose-Einstein condensate moving at velocities \mathbf{V}_f and \mathbf{V}_b respectively. The Hamiltonian

of the system can be written as

$$\hat{H} = \hat{H}_b + \hat{H}_f + \hat{H}_{bf}, \quad (1)$$

where $\hat{H}_{b,f}$ are the Hamiltonians of the bosons and the fermions and \hat{H}_{bf} describes the coupling between the two species. In the mean-field approximation, we have

$$\hat{H}_{bf} = g_{bf} \int d^3\mathbf{r} \sum_{\sigma} \hat{\varphi}(\mathbf{r})^{\dagger} \hat{\varphi}(\mathbf{r}) \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}) \hat{\psi}_{\sigma}(\mathbf{r}), \quad (2)$$

where $\hat{\psi}_{\sigma}$ is the field operator of spin σ fermions, and $\hat{\varphi}$ that of the bosons. Assuming that the BEC can also be described within the mean-field approximation, then the bosonic field operator can be expanded over the Bogoliubov modes

$$\hat{\varphi}(\mathbf{r}) = e^{i\mathbf{k}_b \cdot \mathbf{r}} \left[\sqrt{n_b} + \frac{1}{\sqrt{\Omega}} \sum_{\mathbf{q}} \left(u_{\mathbf{q}} \hat{b}_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{r}} - v_{\mathbf{q}} \hat{b}_{\mathbf{q}}^{\dagger} e^{-i\mathbf{q} \cdot \mathbf{r}} \right) \right], \quad (3)$$

where Ω is the quantization volume, $\hbar\mathbf{k}_b = m\mathbf{V}_b$, n_b is the density of bosons and the $(u_{\mathbf{q}}, v_{\mathbf{q}})$ are the usual Bogoliubov coefficients. The exponential prefactor describes the Galilean boost associated with the motion of the condensate (see Appendix A). In addition, due to the motion of the BEC, the energies of the Bogoliubov modes are shifted by Doppler effect and their spectrum is given by

$$E_{\mathbf{q},b} = E_{\mathbf{q},b}^{(0)} + \hbar\mathbf{q} \cdot \mathbf{V}_b, \quad (4)$$

where $E_{\mathbf{q},b}^{(0)} = \hbar c_b q \sqrt{1 + q^2 \xi_b^2 / 2}$ is the Bogoliubov spectrum (here c_b is the sound velocity and ξ_b is the healing length) [7].

Expanding \hat{H}_{bf} to first order in fluctuations of the bosonic field, we have

$$\hat{H}_{bf} = \hat{H}_{bf}^{(0)} + \hat{H}_{bf}^{(1)} + \hat{H}_{bf}^{(1)\dagger} + \dots \quad (5)$$

with

$$\hat{H}_{bf}^{(0)} = g_{bf} n_b \int d^3\mathbf{r} \sum_{\sigma} \hat{\psi}_{\sigma}^{\dagger} \hat{\psi}_{\sigma}(\mathbf{r}), \quad (6)$$

and

$$\widehat{H}_{\text{bf}}^{(1)} = g_{\text{bf}} \sqrt{\frac{n_b}{\Omega}} \int d^3 \mathbf{r} \sum_{\mathbf{q}, \sigma} (u_{\mathbf{q}} - v_{\mathbf{q}}) e^{-i\mathbf{q} \cdot \mathbf{r}} \widehat{\psi}_{\sigma}^{\dagger}(\mathbf{r}) \widehat{\psi}_{\sigma}(\mathbf{r}) \widehat{b}_{\mathbf{q}}^{\dagger}. \quad (7)$$

$\widehat{H}_{\text{bf}}^{(0)}$ being proportional to the number of fermions, it is a constant of motion, and can therefore be gauged away. $\widehat{H}_{\text{bf}}^{(1)}$ and $\widehat{H}_{\text{bf}}^{(1)\dagger}$ respectively create and destroy Bogoliubov excitations in the BEC. In the weak coupling limit we can expand the damping rate in power of g_{bf} . To first order, only $\widehat{H}_{\text{bf}}^{(1)}$ plays a role and using Fermi's Golden Rule the rate of creation of excitations of momentum $\hbar \mathbf{q}$ in the BEC and in an eigenstate $|\alpha_f\rangle$ of the fermionic Hamiltonian \widehat{H}_f is given by

$$\Gamma(\alpha, \mathbf{q}) = \frac{g_{\text{bf}}^2 n_b}{\Omega \hbar} (u_{\mathbf{q}} - v_{\mathbf{q}})^2 \delta(E_{\alpha, f} + E_{\mathbf{q}, b}) \times \left| \langle \alpha_f | \int d^3 \mathbf{r} \sum_{\sigma} \widehat{\psi}_{\sigma}^{\dagger} \widehat{\psi}_{\sigma}(\mathbf{r}) e^{-i\mathbf{q} \cdot \mathbf{r}} | 0_f \rangle \right|^2, \quad (8)$$

where $E_{\alpha, f}$ is the energy of the state $|\alpha_f\rangle$ relatively to that of the ground state $|0_f\rangle$ of the moving fermionic superfluid. Since the creation of a Bogoliubov excitation imparts a momentum $\hbar \mathbf{q}$ to the BEC, the force acting on the bosons can be written as

$$\mathbf{F} = \sum_{\alpha, \mathbf{q}} \hbar \mathbf{q} \Gamma(\alpha, \mathbf{q}) \quad (9)$$

$$= 2\pi \frac{g_{\text{bf}}^2 n_b}{\Omega} \sum_{\mathbf{q}} \mathbf{q} (u_{\mathbf{q}} - v_{\mathbf{q}})^2 S(-\mathbf{q}, -E_{\mathbf{q}, b}), \quad (10)$$

where we introduced the dynamical structure factor of the Fermi gas

$$S(\mathbf{q}, E) = \sum_{\alpha} \delta(E - E_{\alpha, f}) \left| \langle \alpha_f | \int d^3 \mathbf{r} \sum_{\sigma} \widehat{\psi}_{\sigma}^{\dagger} \widehat{\psi}_{\sigma}(\mathbf{r}) e^{i\mathbf{q} \cdot \mathbf{r}} | 0_f \rangle \right|^2 \quad (11)$$

Note that the expression for the force is galilean invariant and depends only on the relative velocity $\mathbf{V}_b - \mathbf{V}_f$. Indeed, one can show that the dynamical structure factor S of a system moving at velocity \mathbf{V}_f is related to the one of a system at rest (denoted as S_{st}) by

$$S(\mathbf{q}, E) = S_{\text{st}}(\mathbf{q}, E - \hbar \mathbf{q} \cdot \mathbf{V}_f) \quad (12)$$

(this result is a direct consequence of Eq. (A12) giving the energy in the moving frame, and the fact that the operator $\int d^3 \mathbf{r} \sum_{\sigma} \widehat{\psi}_{\sigma}^{\dagger} \widehat{\psi}_{\sigma}(\mathbf{r}) e^{i\mathbf{q} \cdot \mathbf{r}}$ gives a kick $\hbar \mathbf{q}$ to the system). Inserting this expression in Eq. (10) and using the expression for the Bogoliubov spectrum of a moving BEC, Eq. (4) we have for the force

$$\mathbf{F} = 2\pi \frac{g_{\text{bf}}^2 n_b}{\Omega} \sum_{\mathbf{q}} \mathbf{q} (u_{\mathbf{q}} - v_{\mathbf{q}})^2 S_{\text{st}}(-\mathbf{q}, -E_{\mathbf{q}, b}^{(0)} - \hbar \mathbf{q} \cdot \mathbf{V}), \quad (13)$$

with $\mathbf{V} = \mathbf{V}_f - \mathbf{V}_b$. Finally, using the explicit form of the Bogoliubov coefficients, we have

$$(u_{\mathbf{q}} - v_{\mathbf{q}})^2 = \frac{\varepsilon_{\mathbf{q}, b}}{E_{\mathbf{q}, b}^{(0)}} \quad (14)$$

with $\varepsilon_{\mathbf{q}, b} = \hbar^2 q^2 / 2m_b$, hence

$$\mathbf{F} = 2\pi \frac{g_{\text{bf}}^2 n_b}{\Omega} \sum_{\mathbf{q}} \mathbf{q} \frac{\varepsilon_{\mathbf{q}, b}}{E_{\mathbf{q}, b}^{(0)}} S_{\text{st}}(-\mathbf{q}, -E_{\mathbf{q}, b}^{(0)} - \hbar \mathbf{q} \cdot \mathbf{V}). \quad (15)$$

Eq. (15) is the main result of this paper and shows that the friction between the two superfluids is directly related to the dynamic structure factor of the fermionic component of the system. The dynamical structure factor describes the response of a given system to a sinusoidal perturbation and can be related to two-body correlations. In strongly correlated gases, it was measured using Bragg spectroscopy [8] and was used to determine the value of Tan's contact parameter describing the tail of the momentum distribution [9].

The properties of strongly correlated Fermi systems are notoriously hard to calculate, and approximate expression for the dynamic structure factor have been obtained using the approximate methods [10, 11]. However, its general properties can be summarized on the sketch of Fig. 1. At low momenta, the excitation spectrum is dominated by a phonon branch $E = \hbar c_f k$, with c_f the sound velocity of the fermionic superfluid. Since the ground state defines the energy zero, S_{st} vanishes for $E < 0$. However, the presence of a pairing gap implies of a region where S_{st} vanishes at positive energies. Landau's critical velocity v_L is defined as the highest value of v such that $S_{\text{st}}(E, q)$ vanishes for all $E \leq vq$ (see Fig. 1). In the BEC regime of the BEC-BCS crossover, we have $v_L = c_F$, while in the BCS regime, Landau's velocity is driven by pair breaking excitations and we have $v_L \ll c_f$.

These general features implies that if for all \mathbf{q} we have $-E_{\mathbf{q}, b}^{(0)} - \hbar \mathbf{q} \cdot \mathbf{V} < qv_L$, the force is zero. To get some damping, there must be values of \mathbf{q} for which $-E_{\mathbf{q}, b}^{(0)} - \hbar \mathbf{q} \cdot \mathbf{V} \geq \hbar qv_L$. Using the general Landau argument, we get that $|\mathbf{V}| \geq v_L + \min_{\mathbf{q}} (E_{\mathbf{q}, b}^{(0)} / \hbar q) = v_L + c_b$. Note that this bound sets a necessary, but not sufficient, condition on the relative velocity. As a consequence $v_L + c_b$ is only a lower bound for the real critical velocity.

Another consequence of this inequality is that the values of q contributing to the integral are bounded. Indeed, from the inequality $E \leq \hbar qv_L$, we get that

$$q\xi_b \leq \sqrt{2x(x+2)},$$

where $x_{c_b} = V - c_b - v_L$. In other words, when the relative velocity gets close to $c_b + v_L$, the damping is dominated by long wave-length excitations.

In what follows, we compute the force in the deep BEC and BCS limits where analytical expressions can be obtained.




FIG. 1: Sketch of the structure factor of an interacting Fermi gas: S_{st} vanishes outside the gray area and Δ is the excitation gap. Dotted line: low momentum phonon branch $E = \hbar c_f q$. Dashed line: Landau's critical velocity $E = \hbar v_L q$ below which the structure factor vanishes. Note that this sketch corresponds to a weakly interacting system where Landau's velocity is fixed by pair breaking, and $v_L < c_f$. In the molecular regime of the crossover, $v_L = c_f$ [2, 12].

III. BCS LIMIT

Let's consider first the deep BCS limit where the fermions can be described as an ideal gas. Since there is a vanishingly small gap in the fermionic spectrum the lower bound derived above is actually saturated and we have $V_c = c_b$. Working close to c_b , the relevant values of q much smaller than k_F , and we can therefore replace the expression of the dynamical structure factor by its low-energy approximation (see for instance [13], and Appendix B)

$$S_{\text{st}}(\mathbf{q}, E) = \frac{m_f^2 \Omega E}{2\pi^2 \hbar^4 q},$$

this replacement being valid as long as the condition

$$k_F \xi \ll \sqrt{2 \left(\frac{V^2 - c_b^2}{c_b^2} \right)}$$

is fulfilled.

Replacing the discrete sum over q by an integral, one gets

$$\frac{\mathbf{F}}{\Omega} = -\frac{2g_{\text{bf}}^2 n_b m_f^2}{(2\pi)^4 \hbar^4} \int d^3 \mathbf{q} \frac{\mathbf{q} \varepsilon_{\mathbf{b}, \mathbf{q}}}{q E_{\mathbf{b}, \mathbf{q}}^{(0)}} (E_{\mathbf{b}, \mathbf{q}}^{(0)} + \hbar \mathbf{q} \cdot \mathbf{V}) \Theta(-E_{\mathbf{b}, \mathbf{q}}^{(0)} - \hbar \mathbf{q} \cdot \mathbf{V}). \quad (16)$$

where Θ is Heaviside's step function. The integral can be computed analytically and taking $V' = V/c_b$, the force becomes

$$\frac{\mathbf{F}}{\Omega} = -\frac{g_{\text{bf}}^2 n_b m_f^2}{(2\pi)^3 m_b \hbar^2 \xi^5} I(V'), \quad (17)$$

with

$$I(V') = \frac{1}{105\sqrt{2}V'^2} \left[105V'^3 \sinh^{-1}(\sqrt{V'^2 - 1}) + \sqrt{V'^2 - 1} (6V'^6 - 39V'^4 - 80V'^2 + 8) \right]. \quad (18)$$

Close to $V' = 1$, this expression can be expanded as

$$I(V') \simeq \frac{128}{315} (V' - 1)^{9/2} + \dots \quad (19)$$

Even though this scaling is strictly-speaking obtained for a non-interacting Fermi gas, it can be extended to the case of a BEC moving inside an interacting Fermi gas in its normal phase when it can be described within the Fermi Liquid Theory. Indeed, close the critical velocity, the damping mechanism described involves only low energy excitations of the system which behave as fermionic quasiparticles with renormalized physical parameters. In particular, the scaling $S(q, E) \propto E/q$ is still valid [14].

IV. HYDRODYNAMIC LIMIT

In the BEC limit of the cross-over, the excitations of the fermionic superfluid are dominated by phonons of energy $E_{f, \mathbf{q}}$. In this case, the dynamical structure factor can be approximated by a Dirac function and we have in this case

$$S_{\text{st}}(\mathbf{q}, E) = \mathcal{A} \delta(E - E_{f, \mathbf{q}}^{(0)}) \quad (20)$$

where the normalization constant \mathcal{A} is fixed by the f-sum rule [7, 15] and is given by

$$\mathcal{A} = N_f \frac{\varepsilon_{\mathbf{q}, f}}{E_{f, \mathbf{q}}^{(0)}}. \quad (21)$$

In this limit, we obtain for the force

$$\mathbf{F} = 2\pi g_{\text{bf}}^2 n_b n_f \sum_{\mathbf{q}} \mathbf{q} \frac{\varepsilon_{\mathbf{q}, b} \varepsilon_{\mathbf{q}, f}}{E_{\mathbf{q}, b}^{(0)} E_{-\mathbf{q}, f}^{(0)}} \delta(E_{\mathbf{q}, b}^{(0)} + E_{-\mathbf{q}, f}^{(0)} + \hbar \mathbf{q} \cdot \mathbf{V}). \quad (22)$$

Using the usual Landau argument, we see that the force vanishes if the velocity is smaller than a critical velocity V_c given by

$$V_c = \min_{\mathbf{q}} \left(\frac{E_{\mathbf{q}, b}^{(0)} + E_{-\mathbf{q}, f}^{(0)}}{\hbar q} \right). \quad (23)$$

In case of linear dispersion relation, the critical velocity is simply

$$V_c = c_b + c_f, \quad (24)$$

where $c_{f,b}$ are the sound velocities of the fermions and the bosons.

To calculate explicitly the force, we assume that both the bosons and the fermions follow the Bogoliubov dispersion relation, an assumption valid in the far-BEC limit of the BEC-BCS crossover. Taking $\xi_{b,f}$ the healing lengths of the two superfluids, we have after performing the angular integral

$$\frac{F}{\Omega} = -\frac{g_{bf}^2 n_b n_f \hbar}{8\pi m_f m_b c_b c_f V} \times \int_0^{q_M} q^4 dq \frac{\left(\frac{c_b}{V} \sqrt{1 + q^2 \xi_b^2/2} + \frac{c_f}{V} \sqrt{1 + q^2 \xi_f^2/2} \right)}{\sqrt{(1 + q^2 \xi_b^2/2)(1 + q^2 \xi_f^2/2)}} \quad (25)$$

with q_M given by the condition

$$c_b \sqrt{1 + q_M^2 \xi_b^2/2} + c_f \sqrt{1 + q_M^2 \xi_f^2/2} = V \quad (26)$$

For $V \simeq V_c$, q_M is vanishingly small, and is given by

$$q_M \simeq 2 \sqrt{\frac{V - V_c}{c_b \xi_b^2 + c_f \xi_f^2}},$$

hence

$$\frac{F}{\Omega} \simeq -\frac{4g_{bf}^2 n_b n_f \hbar}{5\pi m_f m_b c_b c_f V_c} \left(\frac{V - V_c}{c_b \xi_b^2 + c_f \xi_f^2} \right)^{5/2} \quad (27)$$

V. CONCLUSION

In this work we have shown that the damping between two bosonic and fermionic superfluids could be related to the dynamical structure factor of the fermions. In the weakly and strongly attractive limits, we find a power law dependence of the damping force vs velocity close to the critical velocity. Like previous findings [3], we find that the exponent varies across the BEC-BCS crossover and is different from the scaling $F \propto (V' - 1)^2$ found for the drag of a particle in a Bose-Einstein condensate [16]. These different scalings highlight the many-body nature of the two counter-flowing systems and the role of the collective excitations in the damping mechanism. Our prediction can be directly tested by measuring the damping of the relative oscillations of two superfluids, as presented in [1] for a unitary fermions. In particular, comparison with Eq. 15 provides a quantitative test of theoretical predictions for the structure factor of a strongly correlated fermionic superfluid.

Acknowledgments

The authors thank X. Leyronas, C. Salomon, Y. Castin, I. Ferrier-Barbut, S. Laurent, M. Delehaye and S. Stringari for stimulating discussions. This work was supported by Région Île de France (IFRAF) and European Union (ERC grant ThermoDynaMix).

Appendix A: Bogoliubov Transformation in a Moving Frame

Consider an ensemble of N particles described by a state $|\psi\rangle$. The action of a velocity \mathbf{V} boost is described by the operator

$$\hat{U} = \exp \left(i \sum_{j=1}^N \mathbf{k} \cdot \hat{\mathbf{r}}_j \right), \quad (A1)$$

with $\mathbf{k} = m\mathbf{V}/\hbar$. In second quantization, this operator is represented by

$$\hat{U} = \exp \left(i \int d^3 \mathbf{r}' \hat{\varphi}(\mathbf{r}')^\dagger \hat{\varphi}(\mathbf{r}') \mathbf{k} \cdot \mathbf{r}' \right). \quad (A2)$$

Define $\hat{\varphi}'(\mathbf{r})$ the field operator in the moving frame. By definition, we must have for $|\psi'_{i=1,2}\rangle = \hat{U}|\psi_{i=1,2}\rangle$, $\langle \psi'_1 | \hat{\varphi}'(\mathbf{r}) | \psi'_2 \rangle = \langle \psi_1 | \hat{\varphi}(\mathbf{r}) | \psi_2 \rangle$, hence

$$\langle \psi_1 | \hat{U}^\dagger \hat{\varphi}'(\mathbf{r}) \hat{U} | \psi_2 \rangle = \langle \psi_1 | \hat{\varphi}(\mathbf{r}) | \psi_2 \rangle. \quad (A3)$$

This equation being valid for any $|\psi_i\rangle$, we must have $\hat{U}^\dagger \hat{\varphi}'(\mathbf{r}) \hat{U} = \hat{\varphi}(\mathbf{r})$ hence

$$\hat{\varphi}'(\mathbf{r}) = \hat{U} \hat{\varphi}(\mathbf{r}) \hat{U}^\dagger. \quad (A4)$$

$\hat{\varphi}'$ can be calculated using the Campbell-Hausdorff formula. Taking $\hat{U} = \exp(\hat{T})$, we have indeed

$$\hat{\varphi}'(\mathbf{r}) = \hat{\varphi}(\mathbf{r}) + [\hat{T}, \hat{\varphi}(\mathbf{r})] + \frac{1}{2} [\hat{T}, [\hat{T}, \hat{\varphi}(\mathbf{r})]] + \dots \quad (A5)$$

The commutator $[\hat{T}, \hat{\varphi}(\mathbf{r})]$ is readily calculated and yields $[\hat{T}, \hat{\varphi}(\mathbf{r})] = -i\mathbf{k} \cdot \mathbf{r} \hat{\varphi}(\mathbf{r})$. After resumming the Campbell-Hausdorff formula, we obtain

$$\hat{\varphi}'(\mathbf{r}) = e^{-i\mathbf{k} \cdot \mathbf{r}} \hat{\varphi}(\mathbf{r}). \quad (A6)$$

In the moving frame, the BEC is stationary and we can therefore decompose $\varphi'(\mathbf{r})$ using a Bogoliubov transform

$$\hat{\varphi}'(\mathbf{r}) = \sqrt{n_0} + \frac{1}{\sqrt{\Omega}} \sum_{\mathbf{q}} \left[u_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{r}} \hat{b}_{\mathbf{q}} - v_{\mathbf{q}} e^{-i\mathbf{q} \cdot \mathbf{r}} \hat{b}_{\mathbf{q}}^\dagger \right]. \quad (A7)$$

We thus have in the lab frame

$$\begin{aligned} \hat{\varphi}(\mathbf{r}) &= e^{i\mathbf{k} \cdot \mathbf{r}} \hat{\varphi}'(\mathbf{r}) \\ &= e^{i\mathbf{k} \cdot \mathbf{r}} \left[\sqrt{n_0} + \frac{1}{\sqrt{\Omega}} \sum_{\mathbf{q}} \left(u_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{r}} \hat{b}_{\mathbf{q}} - v_{\mathbf{q}} e^{-i\mathbf{q} \cdot \mathbf{r}} \hat{b}_{\mathbf{q}}^\dagger \right) \right] \end{aligned} \quad (A8)$$

The eigenenergies of the Bogoliubov excitations of the moving Bose-Einstein condensate are obtained by applying the Galilean boost operator \hat{U} to the eigenstates $|\beta\rangle_b$ of the condensate at rest. Taking $|\beta'\rangle_b = \hat{U}|\beta\rangle_b$, and consider the quantum many-body Hamiltonian

$$\begin{aligned} \hat{H}_b = & - \int \hat{\varphi}(\mathbf{r})^\dagger \frac{\hbar^2 \nabla^2}{2m} \hat{\varphi}(\mathbf{r}) d^3\mathbf{r} \\ & + \int V(\mathbf{r} - \mathbf{r}') \hat{\varphi}(\mathbf{r})^\dagger \hat{\varphi}(\mathbf{r}')^\dagger \hat{\varphi}(\mathbf{r}') \hat{\varphi}(\mathbf{r}) d^3\mathbf{r} d^3\mathbf{r}'. \end{aligned} \quad (\text{A10})$$

By using Eq. A6), we see readily that if $|\beta\rangle_b$ is a common eigenstate of \hat{H}_b and the momentum and particle number operators, then $|\beta'\rangle_b$ is an eigenstate of \hat{H}_b for the eigenvalue

$$E'_\beta = E_\beta + \hbar\mathbf{q}_\beta \cdot \mathbf{V} + N_b \frac{mV^2}{2}, \quad (\text{A11})$$

where $\hbar\mathbf{q}_\beta$ is the momentum of state $|\beta\rangle_b$. Compared to the energy E'_0 of the boosted ground-state Bose-Einstein condensate, the energy of the excitation $|\beta'\rangle_b$ is therefore

$$E'_\beta - E'_0 = E_\beta + \hbar\mathbf{q}_\beta \cdot \mathbf{V}, \quad (\text{A12})$$

and is simply Doppler-shifted.

Appendix B: Dynamic structure factor of the ideal Fermi gas.

In this appendix, we rederive the dynamical structure factor for zero-temperature, non-interacting fermions. A more general derivation including finite temperature effects can be found in [13].

The dynamic structure factor is given by the matrix elements of the excitation operator $\hat{F}_\mathbf{q} = \sum_\sigma \int d^3\mathbf{r} e^{i\mathbf{q}\cdot\mathbf{r}} \hat{\psi}_\sigma(\mathbf{r})^\dagger \hat{\psi}_\sigma(\mathbf{r})$. Expanding the field operators over the plane wave basis as $\hat{\psi}_\sigma(\mathbf{r}) = \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} \hat{c}_{\mathbf{k},\sigma} / \sqrt{\Omega}$, where Ω is a quantization volume, yields

$$\hat{F}_\mathbf{q} = \sum_{\mathbf{k},\sigma} \hat{c}_{\mathbf{k}+\mathbf{q},\sigma}^\dagger \hat{c}_{\mathbf{k},\sigma}.$$

In other words, the operator $\hat{F}_\mathbf{q}$ kicks one particles of the system and gives it a momentum \mathbf{q} . The states $|n\rangle$ excited by $\hat{F}_\mathbf{q}$ from the ground state $|0\rangle$ correspond to particle-hole pairs of momenta \mathbf{k} (hole) and $\mathbf{k} + \mathbf{q}$ (particle). The energy of such an excitation is $E_n = \varepsilon_{\mathbf{k}+\mathbf{q}} - \varepsilon_{\mathbf{k}} = \hbar^2 q^2 / 2m + \hbar^2 \mathbf{k} \cdot \mathbf{q} / m$ and we can write the structure factor as

$$S(\mathbf{q}, E) = 2 \int_{k < k_F, |\mathbf{k}+\mathbf{q}| > k_F} \frac{d^3\mathbf{k}\Omega}{(2\pi)^3} \delta\left(E - \frac{\hbar^2 q^2}{2m} - \frac{\hbar^2 k_z q}{m}\right),$$




FIG. 2: Construction of the dynamic structure factor in momentum space. The shaded area represents the Fermi sphere. Energy-momentum conservation fixes the initial k_z to a value K_z (see text). Since the initial state must lie inside the Fermi Sea, we must have $\kappa \leq \kappa_1 = \sqrt{k_F^2 - K_z^2}$. Moreover, the final momentum must be above the Fermi surface, hence $\kappa \geq \sqrt{k_F^2 + (K_z + q)^2}$. The thick solid line represents the momenta contributing to the dynamical structure factor. From top to bottom. Upper panel, case (1): $K_z + q$ larger than k_f . The final momentum is always above the Fermi surface. Middle panel, case 2: $K_z + q$ is lower than k_F , with $\kappa_2 \leq \kappa_1$. The permitted values of \mathbf{k} are located in the ring $\kappa_2 \leq \kappa \leq \kappa_1$. Lower panel, case 3. For $\kappa_2 \geq \kappa_1$, the contribution to the structure factor vanishes.

where the z axis is chosen along \mathbf{q} and where the factor 2 comes from the two spin states.

We perform first the integral over k_z . Take $K_z(\mathbf{q}, E) = (E - \frac{\hbar^2 q^2}{2m})m/\hbar^2 q$. The delta-function selects $k_z = K_z(\mathbf{q}, E)$ and taking $\boldsymbol{\kappa} = (k_x, k_y)$ we have

$$S(\mathbf{q}, E) = \frac{2m}{\hbar^2 q} \int_{|K_z| < k_F, |\mathbf{k} + \mathbf{q}| > k_F} \frac{d^2 \boldsymbol{\kappa} \Omega}{(2\pi)^3}. \quad (\text{B1})$$

The condition $|K_z| \leq k_F$ selects a band in the (q, E) plane. Indeed, we must have

$$-\frac{\hbar^2 k_F q}{m} \leq \left(E - \frac{\hbar^2 q^2}{2m} \right) \leq \frac{\hbar^2 k_F q}{m}.$$

The constraint $k_z = K_z$ selects a plane in initial-momentum space. Taking into account the fact that the initial momenta must lie within the Fermi Sea, we see that the transverse momentum $\boldsymbol{\kappa}$ lies within a disk of radius $\kappa_1 = \sqrt{k_F^2 - K_z^2}$. The set of allowed values for $\boldsymbol{\kappa}$ is further constrained by Pauli Blocking on the final states. Indeed, the final momentum must lie outside the Fermi Sea and as such it implies the condition $|\mathbf{k} + \mathbf{q}| \geq k_F$. We must then consider three cases (see also Fig. 2) :

1. If $K_z + q$ is larger than k_F , then for any \mathbf{k} such that $k_z = K_z$, the final momentum $\mathbf{k} + \mathbf{q}$ is outside the Fermi Sea. The constraint on the final state is thus always satisfied and does not have any effect on the calculation of the integral over $\boldsymbol{\kappa}$, which yields the area of the disk of radius κ_1 . We thus have

$$S(\mathbf{q}, E) = \frac{m\Omega\kappa_1^2}{4\pi^2\hbar^2 q}. \quad (\text{B2})$$

2. For $K_z + q \leq k_F$, final momenta corresponding to small values of κ (more precisely smaller than $\kappa_2 = \sqrt{k_F^2 - (K_z + q)^2}$) lie inside the Fermi Sea and are therefore forbidden by Pauli Principle. We have then

$$S(\mathbf{q}, E) = \frac{m\Omega(\kappa_1^2 - \kappa_2^2)}{4\pi^2\hbar^2 q} = \frac{m^2\Omega E}{2\pi^2\hbar^4 q}. \quad (\text{B3})$$

Take $k_F^3 = 3\pi^2 n_f$, where n_f is the total density of fermions, this expression can be recast as

$$S(\mathbf{q}, E) = \frac{3N_f}{8} \left(\frac{k_F}{q} \right) \frac{E}{E_F^2}, \quad (\text{B4})$$

with $N_f = n_f \Omega$ the total number of fermions.

3. Eq. B3 is valid as long as $\kappa_1 \geq \kappa_2$. Graphically, case (2) ends when $K_z + q = -K_z$, ie for $E = 0$.

For $E \leq 0$, the dynamic structure factor vanishes, as expected since the Fermi Sea is the ground state of the system.




FIG. 3: Dynamical structure factor of the ideal gas. The shaded area corresponds to the region where the structure factor takes finite values. The solid lines corresponds to condition $-k_F \leq K_z \leq k_F$. The dashed marks the limit between cases (1) and (2) and is associated with the condition $K_z + q = k_F$.

Finally, let's consider some simple limiting cases. For $q \geq 2k_F$, we are in case (1) (see Fig. 3), and $S(\mathbf{q}, E)$ is therefore given by

$$S(\mathbf{q}, E) = \frac{m\Omega}{4\pi^2\hbar^2 q} \left[k_F^2 - \frac{m^2}{\hbar^2 q^2} \left(E - \frac{\hbar^2 q^2}{2m} \right)^2 \right]. \quad (\text{B5})$$

In this regime, S is peaked around the energy $E = \varepsilon_{\mathbf{q}} = \hbar^2 q^2 / 2m$ with a half-width $\Delta E = \hbar^2 k_F q / m$. We note that $\Delta E / \varepsilon_{\mathbf{q}} = 2k_F / q \rightarrow 0$ for $q \rightarrow \infty$. In other words, we recover at large momenta the dynamic structure factor of a free particle.

For low values of q , the range of validity of case (1) is vanishingly small and the dynamic structure factor is mostly given by case (2). We therefore have

$$S(\mathbf{q}, E) = \frac{3N_f}{8} \left(\frac{k_F}{q} \right) \frac{E}{E_F^2}, \quad (\text{B6})$$

as long as $0 \leq E \leq 2E_F q / k_F$, and $S = 0$ otherwise.

One readily checks that both approximate expressions (B6) and (B5) satisfy the f-sum rule.

-
- [1] I Ferrier-Barbut, M. Delehaye, S. Laurent, A.T. Grier, M. Pierce, B.S Rem, F. Chevy, and C. Salomon. A mixture of Bose and Fermi superfluids. *Science*, 345:1035–1038, 2014.
- [2] Yvan Castin, Igor Ferrier-Barbut, and Christophe Salomon. The Landau critical velocity for a particle in a fermi superfluid. *Comptes Rendus Physique*, 16(2):241–253, 2015.
- [3] Wei Zheng and Hui Zhai. Quasiparticle lifetime in a mixture of Bose and fermi superfluids. *Phys. Rev. Lett.*, 113(26):265304, 2014.
- [4] J.J. Kinnunen and G.M. Bruun. Induced interactions in a superfluid Bose-Fermi mixture. *Phys. Rev. A*, 91(4):041605, 2015.
- [5] Abad, Marta, Recati, Alessio, Stringari, Sandro, and Chevy, Frédéric. Counter-flow instability of a quantum mixture of two superfluids. *Eur. Phys. J. D*, 69(5):126, 2015.
- [6] W. Zwerger, editor. *The BCS-BEC Crossover and the Unitary Fermi Gas*, volume 836 of *Lecture Notes in Physics*. Springer, Berlin, 2012.
- [7] L. Pitaevskii and S. Stringari. *Bose-Einstein Condensation*. Clarendon Press, 2003.
- [8] G. Veeravalli, E Kuhnle, P Dyke, and C.J. Vale. Bragg spectroscopy of a strongly interacting fermi gas. *Phys. Rev. Lett.*, 101(25):250403, 2008.
- [9] E. D. Kuhnle, H. Hu, X.-J. Liu, P. Dyke, M. Mark, P. D. Drummond, P. Hannaford, and C. J. Vale. Universal behavior of pair correlations in a strongly interacting fermi gas. *Phys. Rev. Lett.*, 105(7):070402, Aug 2010.
- [10] Anna Minguzzi, Gabriele Ferrari, and Yvan Castin. Dynamic structure factor of a superfluid fermi gas. *The European Physical Journal D-Atomic, Molecular, Optical and Plasma Physics*, 17(1):49–55, 2001.
- [11] H.P. Büchler, P. Zoller, and W. Zwerger. Spectroscopy of superfluid pairing in atomic fermi gases. *Physical Rev. Lett.*, 93(8):080401, 2004.
- [12] R. Combescot, M. Kagan, and S. Stringari. Collective mode of homogeneous superfluid Fermi gases in the BEC-BCS crossover. *Phys. Rev. A*, 74(4):042717, October 2006.
- [13] K Baerwinkel. Dynamic structure factor of ideal quantum systems. *Zeitschrift für Physik B Condensed Matter*, 12(4):287–291, 1971.
- [14] S. Stringari. Density and spin response function of a normal Fermi gas at unitarity. *Phys. Rev. Lett.*, 102(11):110406, 2009.
- [15] D. Pines and P. Nozières. *The Theory of Quantum Liquids, vol. I: Normal Fermi Liquids*. WA Benjamin, New York, 1966.
- [16] G.E. Astrakharchik and L.P. Pitaevskii. Motion of a heavy impurity through a Bose-Einstein condensate. *Phys. Rev. A*, 70(1):013608, 2004.