



HAL
open science

Acetylene (C₂H₂) and hydrogen cyanide (HCN) from IASI satellite observations: global distribution, validation, and comparison with model.

Valentin Duflot, Catherine Wespes, Lieven Clarisse, Daniel Hurtmans, Yasmine Ngadi, Nicolas Jones, Claire Paton-Walsh, Juliette Hadji-Lazaro, Corinne Vigouroux, Martine de Mazière, et al.

► To cite this version:

Valentin Duflot, Catherine Wespes, Lieven Clarisse, Daniel Hurtmans, Yasmine Ngadi, et al.. Acetylene (C₂H₂) and hydrogen cyanide (HCN) from IASI satellite observations: global distribution, validation, and comparison with model.. Atmospheric Chemistry and Physics Discussions, 2015, doi:10.5194/acpd-15-14357-2015. 10.5194/acpd-15-14357-2015 . hal-01154522

HAL Id: hal-01154522

<https://hal.science/hal-01154522>

Submitted on 3 Jun 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Global
measurements of
HCN and C₂H₂ from
IASI

V. Dufлот et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Acetylene (C₂H₂) and hydrogen cyanide (HCN) from IASI satellite observations: global distributions, validation, and comparison with model

V. Dufлот^{1,2}, C. Wespes¹, L. Clarisse¹, D. Hurtmans¹, Y. Ngadi¹, N. Jones³, C. Paton-Walsh³, J. Hadji-Lazaro⁴, C. Vigouroux⁵, M. De Mazière⁵, J.-M. Metzger⁶, E. Mahieu⁷, C. Servais⁷, F. Hase⁸, M. Schneider⁸, C. Clerbaux^{1,4}, and P.-F. Coheur¹

¹Spectroscopie de l'Atmosphère, Service de Chimie Quantique et Photophysique, Université Libre de Bruxelles (U.L.B.), 50 Av. F. D. Roosevelt, 1050, Brussels, Belgium

²Laboratoire de l'Atmosphère et des Cyclones (LACy), Université de la Réunion, UMR CNRS-Météo-France 8105, Saint-Denis de la Réunion, France

³School of Chemistry, University of Wollongong, Wollongong, New South Wales, Australia

⁴UPMC Université Paris 06; Université Versailles-St. Quentin; CNRS/INSU, LATMOS-IPSL, Paris, France

⁵Belgian Institute for Space Aeronomy (BIRA-IASB), 3, Av. Circulaire, 1180, Brussels, Belgium

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁶UMS3365 de l'OSU-Réunion, CNRS – Université de la Réunion, Saint Denis de la Réunion, France

⁷Institut d'Astrophysique et de Géophysique, Université de Liège, 17, Allée du 6 Août, 4000, Liège, Belgium

⁸Institute for Meteorology and Climate Research (IMK-ASF), Karlsruhe Institute of Technology, Karlsruhe, Germany.

Received: 13 April 2015 – Accepted: 20 April 2015 – Published: 21 May 2015

Correspondence to: V. Dufлот (valentin.dufлот@univ-reunion.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Global
measurements of
HCN and C₂H₂ from
IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We present global distributions of C_2H_2 and HCN total columns derived from the Infrared Atmospheric Sounding Interferometer (IASI). These distributions are obtained with a fast method allowing to retrieve C_2H_2 abundance globally with a 5% precision and HCN abundance in the tropical (subtropical) belt with a 10% (30%) precision. IASI data are compared for validation purposes with ground-based Fourier Transform Infrared (FTIR) spectrometer measurements at four selected stations. We show that there is an overall agreement between the ground-based and space measurements. Global C_2H_2 and subtropical HCN abundances retrieved from IASI spectra show the expected seasonality linked to variations in the anthropogenic emissions and seasonal biomass burning activity, as well as exceptional events, and are in good agreement with previous spaceborne studies. IASI measurements are also compared to the distributions from the Model for Ozone and Related Chemical Tracers, version 4 (MOZART-4). Seasonal cycles observed from satellite data are reasonably well reproduced by the model. However, the model seems to overestimate (underestimate) anthropogenic (biomass burning) emissions and a negative global mean bias of 1% (16%) of the model relative to the satellite observations was found for C_2H_2 (HCN).

1 Introduction

Hydrogen cyanide (HCN) and acetylene (or ethyne, C_2H_2) are ubiquitous atmospheric trace gases with medium lifetime, which are frequently used as indicators of combustion sources and as tracers for atmospheric transport and chemistry. For HCN, biomass burning is the primary source, followed by fossil fuel combustion and higher plants, bacteria and fungi (Cicerone and Zellner, 1983; Li et al., 2000), and its primary sink is thought to be ocean uptake (Li et al., 2000). For C_2H_2 , biofuel combustion is considered as the dominant source, followed by fossil fuel combustion and biomass burning

ACPD

15, 14357–14401, 2015

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Xiao et al., 2007). Reaction with hydroxyl radical (OH) is the main sink for C₂H₂, which may also act as a precursor of secondary organic aerosols (Volkamer et al., 2009).

With a tropospheric lifetime of 2–4 weeks for C₂H₂ (Logan et al., 1981) and 5–6 months for HCN (Li et al., 2000; Singh et al., 2003), these two species are interesting tracers for studying atmospheric transport. The study of the ratio C₂H₂/CO (carbon monoxide) can also help to estimate the age of emitted plumes (Xiao et al., 2007).

Long-term local measurements of HCN and C₂H₂ are sparse and mainly performed from ground-based Fourier transform infrared (FTIR) spectrometer at selected stations of the Network for the Detection of Atmospheric Composition Change (NDACC, <http://www.ndacc.org>) (Vigouroux et al., 2012, and references therein). Global distributions of HCN and C₂H₂ may thus help to reduce the uncertainties remaining with regard to the magnitude of their sources and sinks, as well as to their spatial distribution and seasonality in the atmosphere (Li et al., 2009; Parker et al., 2011).

Satellite sounders have provided considerable new information in the past years, with measurements from the Atmospheric Chemistry Experiment (ACE-FTS) (Lupu et al., 2009; Gonzalez Abad et al., 2011), the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) (Parker et al., 2011; Wiegele et al., 2012; Glatthor et al., 2015) and the Microwave Limb Sounder (MLS) (Pumphrey et al., 2011). These measurements were all made in limb geometry and consequently mostly in the upper troposphere or higher; also the spatial sampling from these instruments is limited, making it less well-suited when studying dynamical events on short time-scales.

Having a twice daily global coverage and a 12 km diameter footprint at nadir, the IASI infrared sounder (Clerbaux et al., 2009) aboard the MetOp-A satellite has the potential for providing measurements for these two species globally, and with higher spatial resolution and temporal sampling than what has been obtained up to now.

Previous studies have demonstrated that HCN and C₂H₂ can be observed with the IASI infrared nadir-looking hyperspectral sounder, e.g. in a specific biomass burning plume (Clarisse et al., 2011a), as well as in an anthropogenic pollution plume uplifted in the free troposphere (Clarisse et al., 2011b). More recently, Dufлот et al. (2013) have

by Rodgers (2000):

$$\hat{x} = x_0 + (K^T S_e^{\text{tot}-1} K)^{-1} K^T S_e^{\text{tot}-1} (y - F(x_0)) \quad (1)$$

where y is the spectral measurements, x_0 is the linearization point, F is the forward model (FM), S_e^{tot} is the covariance of the total error (random + systematic), and the Jacobian K is the derivative of the FM to the target species in a fixed atmosphere.

S_e^{tot} can be estimated considering an appropriate ensemble of N measured spectra which can be used to build up the total measurement error covariance S_y^{obs} :

$$S_e^{\text{tot}} \simeq \frac{1}{N-1} \sum_{j=1}^N (y_j - \bar{y})(y_j - \bar{y})^T = S_y^{\text{obs}} \quad (2)$$

where \bar{y} is the calculated mean spectrum for the ensemble.

To generate S_y^{obs} , we randomly chose 1 million spectra observed by IASI all over the world, above both land and sea, during the year 2009. Then, we applied a BTD test to remove the spectra contaminated by the target species. For HCN (C_2H_2), the wavenumbers 716.5 and 732 cm^{-1} (712.25 and 737.75 cm^{-1}) were used as reference channels and 712.5 cm^{-1} (730 cm^{-1}) was used as test channel (Fig. 1, middle panel). Given the medium lifetimes of the target species (few weeks for C_2H_2 to few months for HCN), and the limited accuracy of the BTD test due to the weak spectral signatures of the target species, it is likely that such randomly chosen and filtered spectra still contain a small amount of the target species whose signal may come out from the noise. This limitation decreases the sensitivity of the method, which is discussed in Sect. 2.2.3.

The spectral ranges considered to compute the S_y^{obs} matrices are 645–800 cm^{-1} for HCN and 645–845 cm^{-1} for C_2H_2 (Fig. 1, top panel). These ranges were chosen as they include parts of the spectrum which have a relatively strong signal from the target species but also from the main interfering species (CO_2 , H_2O and O_3 , Fig. 1, bottom panel) in order to maximize the contrast with the spectral background.

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Having calculated S_y^{obs} and \bar{y} , the HRI of a measured spectrum y can be defined as:

$$\text{HRI} = G(y - \bar{y}) \quad (3)$$

with G the measurement contribution function

$$G = (K^T S_y^{\text{obs}-1} K)^{-1} K^T S_y^{\text{obs}-1} \quad (4)$$

The HRI is a dimensionless scalar similar, other than units, to the apparent column retrieved in Walker et al. (2011). Unlike the optimal estimation method, no information about the vertical sensitivity can be extracted. Note also that the use of a fixed Jacobian to calculate HRI does not allow generating meaningful averaging kernels.

2.2.2 Conversion of HRI into total columns

Having calculated the matrices \mathbf{G} for HCN and C_2H_2 , each observed spectrum can be associated through Eq. (3) with a value of HRI for HCN (HRI_{HCN}) and C_2H_2 ($\text{HRI}_{\text{C}_2\text{H}_2}$). These HRIs are only metrics for determining whether levels of the gas are enhanced with respect to the climatological background over the vertical levels where the instrument is sensitive. For a given atmosphere atm, the main challenge is then to link the HRI to a column amount of the target molecule, i.e. to find $B_{\text{HCN}_{\text{atm}}}$ and $B_{\text{C}_2\text{H}_{2\text{atm}}}$ such as:

$$[X] = B_{X_{\text{atm}}} \text{HRI}_X \quad (5)$$

$[X]$ being the species abundance in molec cm^{-2} .

To determine these coefficients linking the HRIs to total column amounts, HCN and C_2H_2 profiles have been constructed, with enhanced concentrations of the species located in a 1 km thick layer, whose altitude is varied from the ground up to 30 km for HCN and up to 20 km for C_2H_2 (the choice of these maximum altitudes are made with respect to the Jacobians of the FM that are shown in Fig. 3 and commented

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in Sect. 2.2.3). Each of the constructed profile has been associated with a spectrum through the FM of *Atmosphit* considering standard absorption profiles. The associated values of HRI_{HCN} and $\text{HRI}_{\text{C}_2\text{H}_2}$ have then been computed for each of the simulated spectra. Figure 2 shows the look up tables (LUTs) of HRI_{HCN} (top) and $\text{HRI}_{\text{C}_2\text{H}_2}$ (bottom) as a function of the abundance of the target molecule and of the altitude of the polluted layer in a standard tropical modeled atmosphere (Anderson et al., 1986). Similar LUTs have been computed for standard temperate (US standard atmosphere) and polar (Anderson et al., 1986) atmospheres (data not shown). The satellite viewing angles were taken into account in the HRI calculation similarly to Van Damme et al. (2014). One can see that, for a given atmosphere and for a given altitude of the polluted layer, the abundances of both species linearly depend on the HRI value, which validates Eq. (5). For a given atmosphere atm and a given species X , the different values of B with respect to the altitude z of the polluted layer will be noted $b_{X_{\text{atm}}}(z)$ and $k_{X_{\text{atm}}}(z)$ in the following.

Figure 3 shows the normalised Jacobians of the FM for HCN and C₂H₂ averaged over the spectral ranges given in Sect. 2.2.1 (645–800 cm⁻¹ for HCN and 645–845 cm⁻¹ for C₂H₂) and for each of the three standard modeled atmospheres. These Jacobians express the sensitivity of the FM, i.e. both the radiative transfer model and IASI (through its instrumental function), to the target species abundance X in a fixed atmosphere atm:

$$K_{X_{\text{atm}}} = \left[\frac{\partial F_{\text{atm}}}{\partial X}(z_1) \dots \frac{\partial F_{\text{atm}}}{\partial X}(z_n) \right] = \left[k_{X_{\text{atm}}}(z_1) \dots k_{X_{\text{atm}}}(z_n) \right] \quad (6)$$

We then obtain the coefficients $B_{X_{\text{atm}}}$ by multiplying the $b_{X_{\text{atm}}}(z)$ by the value of the Jacobian at the altitude z :

$$B_{X_{\text{atm}}} = \sum_{i=1}^n \left(b_{X_{\text{atm}}}(z_i) \times k_{X_{\text{atm}}}(z_i) \right) \quad \text{with} \quad \sum_{i=1}^n k_{X_{\text{atm}}}(z_i) = 1 \quad (7)$$

Applying this method to the three standard modeled atmospheres (tropical, temperate and polar), we get a B_X value for each, which we have associated with the corresponding range of latitude ($[\pm 20^\circ]$, $[\pm 45^\circ : \pm 60^\circ]$, $[\pm 75^\circ : \pm 90^\circ]$, respectively), and linearly interpolated between. Figure 4 gives the resulting values of B_{HCN} (blue) and $B_{\text{C}_2\text{H}_2}$ (green) in function of the latitude.

2.2.3 Sensitivity and stability of the method

The sensitivity of the method can be assessed from the Jacobians presented in Fig. 3. For HCN, one can see that there is no sensitivity at the surface and above ~ 30 km, and the altitude of the sensitivity peak is located close to the tropopause at ~ 9 , ~ 11 and ~ 14 km for the polar, temperate and tropical atmospheres, respectively. For C_2H_2 , there is no sensitivity above ~ 20 km, and the maximum sensitivity is reached at ~ 8 , ~ 10 and ~ 11 km for the polar, temperate and tropical atmospheres, respectively.

The HRIs presented here above are sensitive to the abundance of the target species – this is what they are made for – and to their vertical distribution. However, the measured column amount may also depend on: (1) the proper suppression of the spectral background, (2) the conditions of thermal contrast with the surface (TC), and (3) the accuracy of the FM to simulate the spectra used to build up the LUTs. The latter was discussed already by Dufлот et al. (2013). In order to test the impact of the two first factors (spectral background suppression and TC) on the retrieved column amount, HCN and C_2H_2 profiles have been constructed with varying TC and concentrations of the interfering and target species. The TC is defined here as the difference between the skin (surface) temperature and that of the air at an altitude of 1.5 km. These variations in interfering species abundances and TC were considered to be independent and were taken within the range $\pm 2\%$ for CO_2 and $\pm 20\%$ for H_2O and O_3 , and in the range $\pm 10\text{K}$ for the TC. For a fixed column amount of the target species, the HRIs were compared one by one to a HRI corresponding to a standard spectrum (i.e. with background concentrations of the interfering species and a TC equal to zero) and if the

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



difference between the two HRIs was lower than 10 %, then this fixed abundance of the target species was tagged as detectable independently from the listed parameters.

The TC was found to be the major source of HRI variation for both target species, and a serious cause of limitation only for HCN. Figure 5 shows the variation of HRI_{HCN} caused by a TC equal to $\pm 10\text{K}$. One can see that HCN column amount can be detected with a variation due to the TC below 10 % when its abundance is higher than 0.28 , 1.2 and $1.6 \times 10^{16} \text{ molec cm}^{-2}$ for the tropical, temperate and polar atmospheres, respectively. This gives the stability thresholds above which HCN column amount can be measured with a 10 % confidence in the independence of the retrieval method to the atmospheric parameters. Consequently, as the stability thresholds of the method for HCN in temperate and polar atmospheres are too high (1.2 and $1.6 \times 10^{16} \text{ molec cm}^{-2}$, respectively) to allow the detection of HCN background abundances as compared to usual background column of typically $0.35 \times 10^{16} \text{ molec cm}^{-2}$ (Vigouroux et al., 2012; Dufлот et al., 2013), IASI HCN measurements have to be rejected in these two types of atmosphere, and considered in the tropical belt for values above $0.28 \times 10^{16} \text{ molec cm}^{-2}$. In order to broaden the exploitable latitude range, we take into account the IASI HCN measurements at subtropical latitudes with the same stability threshold ($0.28 \times 10^{16} \text{ molec cm}^{-2}$), assuming a 30 % confidence in the independence of the retrieval method to the atmospheric parameters – which is quite a prudent assumption. As a result, in the following, IASI HCN measurements is considered in the $\pm 35^\circ$ latitude band with a stability threshold of $0.28 \times 10^{16} \text{ molec cm}^{-2}$, and confidence in the stability of the method is 10 % at tropical latitudes ($[\pm 20^\circ]$) and 30 % at subtropical latitudes ($[\pm 35^\circ : \pm 20^\circ]$). Oppositely to HCN, for C_2H_2 , the variation of $HRI_{\text{C}_2\text{H}_2}$ due to varying TC was found to be lower than 5 % for every C_2H_2 abundances (data not shown). Consequently, in the following no IASI C_2H_2 measurements is rejected.

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

The goal of this section is to describe and evaluate the C₂H₂ and HCN total columns as measured by IASI. We first compare HCN and C₂H₂ total columns retrieved from IASI spectra and from ground-based FTIR spectra. We then depict the C₂H₂ and HCN total columns at global and regional scales. IASI global and regional distributions are finally compared with output from the Model for Ozone and Related Chemical Tracers, version 4 (MOZART-4) in order to evaluate the agreement between the model and the IASI distributions.

3.1 Comparison with ground-based observations

We compare in this section HCN and C₂H₂ total columns retrieved from IASI spectra and from ground-based FTIR spectra for the years 2008–2010 for four selected ground-based FTIR observation sites: Wollongong (34° S; 151° E; 30 m a.m.s.l.), Reunion Island (21° S; 55° E; 50 m a.m.s.l.), Izaña (28° N; 16° W; 2367 m a.m.s.l.) and Jungfrau-joch (46° N; 8° E; 3580 m a.m.s.l.) (Fig. 6). IASI cloudy spectra were removed from the data set using a 10 % contamination threshold on the cloud fraction in the pixel. As exposed in Sect. 2.2.3, errors in retrieved species abundances from IASI spectra due to variations in atmospheric parameters are 10 % at tropical latitudes ([±20°]) and 30 % at subtropical latitudes ([±35° : ±20°]) for HCN and 5 % for C₂H₂, and comparison with ground-based HCN measurements are only performed for tropical and subtropical sites (Reunion Island, Wollongong and Izaña).

Total errors for ground-based measurements at Reunion Island are 17 % for both species, total error for HCN ground-based measurements at Wollongong is 15 %, total error for HCN ground-based measurements at Izaña is 10 %, and total error for C₂H₂ ground-based measurements at Jungfrau-joch is 7 %. Detailed description of ground-based FTIR data set, retrieval method and error budget can be found in Vigouroux et al. (2012) for Reunion Island and in Mahieu et al. (2008) for Jungfrau-joch. However, at Reunion Island, the retrieval strategies have been slightly improved from Vigouroux

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al. (2012), mainly concerning the treatment of the interfering species, but the same spectral signatures are used. Izaña data set and error budget were obtained from the NDACC database (<ftp://ftp.cpc.ncep.noaa.gov/ndacc/station/izana/>). Wollongong data set and error budget were calculated by N. Jones from the University of Wollongong (N. Jones, personal communication, 2015).

Figure 7 shows the mean total column averaging kernels for the ground-based FTIR at each of the four sites. Similarly to IASI (Fig. 3), information content from ground-based instruments measurements is mostly in the middle-high troposphere for both species. The main difference can be observed for tropical C_2H_2 : while IASI Jacobian peaks at 10 km for C_2H_2 in a tropical atmosphere, ground-based FTIR averaging kernel peaks at 15 km for C_2H_2 at Reunion Island.

Figure 8 shows the comparison between the IASI and the ground-based measurements. IASI retrieved total columns were averaged on a daily basis and on a $1^\circ \times 1^\circ$ area around the observation sites. HCN retrieved abundances below 2.8×10^{15} molec cm^{-2} have been removed from both ground-based and space measurements to allow comparison of both datasets (cf. Sect. 2.2.3). One can see that there is an overall agreement between the IASI and the ground-based FTIR measurements considering the error bars. An important result from this study is that IASI seems to capture the seasonality in the two species in most of the cases. This is best seen by looking at the IASI monthly mean retrieved total columns (black circles and lines in Fig. 8). The scatter of the IASI daily mean measurements (red dots) are due to the averaging on a $1^\circ \times 1^\circ$ area around the observation sites.

At Reunion Island HCN and C_2H_2 peak in October-November and are related to the Southern Hemisphere biomass burning season (Vigouroux et al., 2012). We find maxima of around 12×10^{15} molec cm^{-2} for HCN and 10×10^{15} molec cm^{-2} for C_2H_2 . The seasonality and interannual variability matches very well that of the ground-based FTIR measurements for HCN (correlation coefficient of 0.81 for the entire daily mean dataset, and of 0.98 for the monthly mean data set) but with the IASI columns being biased high by 0.79×10^{15} molec cm^{-2} (17%). For C_2H_2 at Reunion Island, the sea-

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sonality and interannual variability matches reasonably well that of the ground-based measurements (correlation coefficient of 0.40 for the entire daily mean dataset, and of 0.72 for the monthly mean data set) but with the IASI columns being biased high by $1.10 \times 10^{15} \text{ molec cm}^{-2}$ (107%). Such a high bias between the two datasets could be due to the difference between space and ground-based instruments sensitivity (Figs. 3 and 7). One can also notice that the C₂H₂ and HCN peaks are higher in 2010. As South American biomass burning plumes are known to impact trace gases abundance above Reunion Island (Edwards et al., 2006a, b; Dufлот et al., 2010), these 2010 higher peaks are probably due to the 2010 great Amazonian fires (Lewis et al., 2011) influence.

At Wollongong HCN peaks also in October–November due to the Southern Hemisphere biomass burning season (Paton-Walsh et al., 2010). We find maxima of around $11 \times 10^{15} \text{ molec cm}^{-2}$ in October 2010, which is, similarly to Reunion Island, very likely to be a signature of the great Amazonian fires as South American biomass burning plumes are known to impact trace gases abundance above Australia (Edwards et al., 2006a, b). The seasonality and interannual variability matches well that of the ground-based FTIR measurements (correlation coefficient of 0.55 for the entire daily mean dataset, and of 0.83 for the monthly mean data set), with the IASI columns being biased low by $0.48 \times 10^{15} \text{ molec cm}^{-2}$ (10%).

At Izaña HCN peaks in May–July due to the biomass burning activity occurring in Northern America and Europe (Sancho et al., 1992). We find maxima of around $8 \times 10^{15} \text{ molec cm}^{-2}$. The seasonality and interannual variability matches poorly that of the ground-based FTIR measurements for HCN (correlation coefficient of 0.28 for the entire daily mean dataset, and of 0.64 for the monthly mean data set), with the IASI columns being biased high by $0.45 \times 10^{15} \text{ molec cm}^{-2}$ (11%). One can notice that HCN total columns as measured by ground-based FTIR are below the HCN stability threshold in boreal winter, which may result in erroneous IASI measurements (because unstable) and explain this poor match between the two datasets.

For C₂H₂ at the Jungfraujoch site, the agreement between IASI and the ground-based retrieved columns is good (correlation coefficient of 0.70 for the entire daily mean

dataset, and of 0.85 for the monthly mean data set), with the IASI columns being biased low by $0.15 \times 10^{15} \text{ molec cm}^{-2}$ (12%), opposite to the observations at Reunion. The larger columns observed in late winter are caused by the increased C_2H_2 lifetime in that season (caused by the seasonal change in OH abundance) (Zander et al., 1991), and we find corresponding maxima of up to $4 \times 10^{15} \text{ molec cm}^{-2}$.

3.2 IASI Global distributions

We focus in this section on the description of the C_2H_2 and HCN distributions retrieved from IASI spectra. For practical reasons, the figures used in this section also show simulated distributions that will be analyzed afterwards.

The left panels of Figs. 9 and 10 provide the seasonal global and subtropical distributions of C_2H_2 and HCN total columns, respectively, as measured by IASI and averaged over the years 2008 to 2010.

Looking at IASI measurements (Figs. 9 and 10 – left panels), one can notice the following main persisting features for both C_2H_2 and HCN:

- the hot spots mainly due to the biomass burning activity occurring in Africa and moving southward along the year (Sauvage et al., 2005; van der Werf et al., 2006);
- the hot spot located in South East Asia being likely a combination of biomass burning and anthropogenic activities;
- the transatlantic transport pathway linking the African west coast to the South American east coast and moving southward along the year (Edwards et al., 2003, 2006a, b; Glatthor et al., 2015).

The following seasonal features can also be observed:

- the transpacific transport pathway linking Eastern Asia to Western North America, especially in March-April-May (MAM) (Yienger et al., 2000);

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- the transport pathway from Southern Africa to Australia in June-July-August (JJA) and September-October-November (SON) (Annegarn et al., 2002; Edwards et al., 2006a, b);
- the transport pathway linking South America (especially Amazonia) to Southern Africa and Australia during the SON period (Edwards et al., 2006a, b; Glatthor et al., 2015);
- the transport of the northern African plume over southern Asia to as far as the eastern Pacific by the northern subtropical jet during the MAM period (Glatthor et al., 2015);
- the Asian monsoon anticyclone (AMA), which is the dominant circulation feature in the Indian-Asian upper troposphere-lower stratosphere (UTLS) region during the Asian summer monsoon, spanning South East Asia to the Middle East and flanked by the equatorial and sub-tropical jets (Hoskins and Rodwell, 1995). The AMA is a known region of persistent enhanced pollution in the upper troposphere, linked to rapid vertical transport of surface air from Asia, India, and Indonesia in deep convection, and confinement by the strong anticyclonic circulation (Randel et al., 2010). The enhanced abundance of C₂H₂ and HCN within the AMA in JJA observed by IASI is in accordance with previous studies (Park et al., 2008; Randel et al., 2010; Parker et al., 2011; Glatthor et al., 2015); however, one should keep in mind that this enhanced abundance measured by IASI is likely due to the combination of this pollution uplift and confinement with the higher sensitivity of the method in the upper troposphere (Fig. 2).

One can also notice the very good agreement between the seasonal HCN distributions shown in our Fig. 10 and the ones published recently in Glatthor et al. (2015, Fig. 3).

Figures 11 and 12 show the C₂H₂ and HCN total columns time series, respectively, as measured by IASI (red dots) with the associated SD (light red lines) for each of the zones defined in Fig. 6.

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In Northern America, Europe and Boreal Central Asia (Fig. 11 – Zones NAM, EUR and BCA), C₂H₂ peaks in late boreal winter due to the increased C₂H₂ lifetime as already noticed over Jungfrauoch (Fig. 8). The boreal summer 2008 California wildfires event (Gyawali et al., 2009) is clearly visible in the NAM plot, as well as the August 2009 Russian wildfires in the NAM, EUR and BCA plots (Parrington et al., 2012; R'honi et al., 2013).

In North Central America (Fig. 12 – Zone NCA), the annual HCN peak in April–June is driven by local fire activity (van der Werf et al., 2010).

In South America, Southern Africa and Australia (Figs. 11 and 12 – Zones SAM, SAF and AUS), the Southern Hemisphere biomass burning season clearly drives the C₂H₂ and HCN peaks in September–November each year. The signature of the great 2010 Amazonian fires (Lewis et al., 2011) is visible on each of the these three Zones, South American fire plumes being known to impact Southern Africa and Australia (Edwards et al., 2003, 2006a, b). The February 2009 Australian bush fires (Glatthor et al., 2013) are also noticeable on Zone AUS for both species.

In Northern Africa (Figs. 11 and 12 – Zone NAF), C₂H₂ and HCN peak in boreal winter because of the biomass burning activity occurring in the Zone, and peak also in boreal summer because of the European and South Mediterranean fires (Van der Werf et al., 2010).

In South East Asia (Figs. 11 and 12 – Zone SEA), the observed C₂H₂ and HCN peaks in July–September and January–March are due to local fire activity (Fortems-Cheiney et al., 2011; Magi et al., 2012). Additionally, the July–September peaks are also likely due to the combination of the pollution uplift and confinement within the AMA with the higher sensitivity of the method in the upper troposphere.

In Equatorial Asia (Figs. 11 and 12 – Zone EQA), local fire activity is visible in July–October, as well as the South East Asian fire activity in January–March (Fortems-Cheiney et al., 2011; Magi et al., 2012). The high biomass burning activity occurring in Indonesia from July to December 2009 (Yulianti et al., 2013; Hyer et al., 2013) is also clearly noticeable.

In Table 2, for C_2H_2 , the correlation coefficients are good (≥ 0.6) to very good (≥ 0.9) except for the zones SAM (South America), SEA (South East Asia) and EQA (Equatorial Asia). For HCN, the correlation coefficients are good (≥ 0.6) except for the zones NCA (North Central America), NAF (Northern Africa), SEA and EQA.

For South America (Zone SAM), correlation coefficient is not as good for C_2H_2 ($R = 0.54$) due to a backward shift of the species abundance peaks in years 2008 and 2009: in the model, this increase occurs from July to October while observations (and previous studies, e.g. van der Werf et al., 2010) show an increase from August to December. This backward shift is also visible for HCN (Fig. 12), but to a lesser extent.

For South East Asia and Equatorial Asia (Zones SEA and EQA), the low correlation coefficients (cf. Table 2) can be attributed to the difficulty of locating precisely with the model the intercontinental convergence zone (ITCZ) which drives the long-range transport of C_2H_2 and HCN-loaded plumes into the zone. Additionally, for Equatorial Asia, the too low fire emissions considered in the model for Indonesia from July to December 2009 may also be a cause for these low correlation coefficients.

For HCN in northern Africa (Zone NAF), correlation coefficient is very low ($R = 0.07$) because the model sets the abundance peaks around August while observations show peaks occurring around December, which is in accordance with previous studies (van der Werf et al., 2010). This inadequate timing for HCN in the model simulations could be due to an overestimation of the Southern African contribution to the Northern African loading and is visible on Fig. 10 (JJA).

4 Conclusions

We have presented a fast method to retrieve HCN and C_2H_2 total columns from IASI spectra. The sensitivity of this method to the two species is mostly in the mid-upper troposphere. With this method, C_2H_2 total columns can be retrieved globally with 5% precision, while HCN abundances can be retrieved for abundances greater than

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$0.28 \times 10^{16} \text{ molec cm}^{-2}$ with 10 % precision in the $\pm 20^\circ$ latitudinal band and with 30 % precision in the $[\pm 35^\circ : \pm 20^\circ]$ latitudinal band.

Total columns have been retrieved globally for a three year period and compared to routine FTIR measurements available at Reunion Island (HCN and C_2H_2), Wollongong (HCN), Jungfraujoch (C_2H_2), and Izaña (HCN). The comparison between IASI and FTIR retrieved total columns demonstrates the capabilities of IASI to capture the seasonality in HCN and C_2H_2 in most cases.

Global seasonal distributions, as well as regional time series of the total columns, have been shown for the two species. IASI is able to capture persisting, seasonal and exceptional features for both species, and the observed patterns are in a general good agreement with previous spaceborne studies (ACE-FTS and MIPAS).

The comparison between these observations and MOZART-4 simulations leads to the following conclusions: (i) the model is able to capture most of the hot spots and seasonal cycles, but not the exceptional events, (ii) the model seems to overestimate (underestimate) anthropogenic (biomass burning) emissions for both species, (iii) the model dynamical scheme and/or the modeled species lifetime could be improved to simulate the impact of the long range transport for these species.

Acknowledgements. IASI has been developed and built under the responsibility of the Centre National d'Etudes Spatiales (CNES, France). It is flown onboard the MetOp satellites as part of the EUMETSAT Polar System. The IASI L1 data are received through the EUMETSAT-Cast near real time data distribution service. Part of the research is supported by EUMETSAT through the O3SAF project. P. F. C. is Senior Research Associate and L.C. is Research Associate at the F.R.S.-FNRS. The research in Belgium was funded by the F.R.S.-FNRS, the Belgian Science Policy Office, and the European Space Agency (ESA Prodex arrangements and the AGACC-II project). The Australian Research Council has provided financial support over the years for the NDACC site at Wollongong, most recently as part of project DP110101948. The Liège team further acknowledges the Fédération Wallonie-Bruxelles for supporting travel costs to the Jungfraujoch station and wishes to thank the International Foundation High Altitude Research Stations Jungfraujoch and Gornergrat (HFSJG, Bern) for supporting the facilities needed to perform the observations. Since 1999, the Izaña FTIR activities have been

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



supported by different funding agencies: European Commission, European Research Council, European Space Agency, EUMETSAT, Deutsche Forschungsgemeinschaft, Deutsches Zentrum für Luft- und Raumfahrt, and the Ministerios de Ciencia e Innovación and Educación from Spain. The authors also wish to thank the French regional, national (INSU, CNRS), and international (NASA/GFSC) organizations for supporting the OPAR (Observatoire de Physique de l'Atmosphère de la Réunion) station. The NDACC is also acknowledged for providing consistent and well documented datasets.

References

- Anderson, G. P., Clough, S. A., Kneizys, F. X., Chetwynd, J. H., and Shettle, E. P.: AFGL Atmospheric Constituent Profiles (0–120 km), Environmental Research Papers no. 954, Air Force Geophysics Laboratory, Hanscom AFB Massachusetts, AFGL TR-86-0110, 1986.
- Annegarn, H. J., Otter L., Swap R. J., and Scholes R. J.: Southern Africa's ecosystem in a test-tube – a perspective on the Southern African Regional Science Initiative (SAFARI 2000), *S. Afr. J. Sci.*, 98, 111–113, 2002.
- Cicerone, R. J. and Zellner, R.: The atmospheric chemistry of hydrogen cyanide (HCN), *J. Geophys. Res.*, 88, 10689–10696, 1983.
- Clarisse, L., Coheur, P. F., Prata, A. J., Hurtmans, D., Razavi, A., Phulpin, T., Hadji-Lazaro, J., and Clerbaux, C.: Tracking and quantifying volcanic SO₂ with IASI, the September 2007 eruption at Jebel at Tair, *Atmos. Chem. Phys.*, 8, 7723–7734, doi:10.5194/acp-8-7723-2008, 2008.
- Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., and Coheur, P.-F.: Global ammonia distribution derived from infrared satellite observations, *Nat. Geosci.*, 2, 479–483, doi:10.1038/NGEO551, 2009.
- Clarisse, L., R'honi, Y., Coheur, P.-F., Hurtmans, D., and Clerbaux, C.: Thermal infrared nadir observations of 24 atmospheric gases, *Geophys. Res. Lett.*, 38, L10802, doi:10.1029/2011GL047271, 2011a.
- Clarisse, L., Formm, M., Ngadi, Y., Emmons, L., Clerbaux, C., Hurtmans, D., and Coheur, P. F.: Intercontinental transport of anthropogenic sulfur dioxide and other pollutants: an infrared remote sensing case study, *Geophys. Res. Lett.* 38, L19806, doi:10.1029/2011GL048976, 2011b.

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmos. Chem. Phys.*, 9, 6041–6054, doi:10.5194/acp-9-6041-2009, 2009.

5 Coheur, P.-F., Barret, B., Turquety, S., Hurtmans, D., Hadji-Lazaro, J., and Clerbaux, C.: Retrieval and characterization of ozone vertical profiles from a thermal infrared nadir sounder, *J. Geophys. Res.*, 110, D24303, doi:10.1029/2005JD005845, 2005.

Dufлот, V., Dils, B., Baray, J. L., De Mazière, M., Attié, J. L., Vanhaelewyn, G., Senten, C., Vigouroux, C., Clain, G., and Delmas, R.: Analysis of the origin of the distribution of
10 CO in the subtropical southern Indian Ocean in 2007, *J. Geophys. Res.*, 115, D22106, doi:10.1029/2010JD013994, 2010.

Dufлот, V., Hurtmans, D., Clarisse, L., R'honi, Y., Vigouroux, C., De Mazière, M., Mahieu, E., Servais, C., Clerbaux, C., and Coheur, P.-F.: Measurements of hydrogen cyanide (HCN) and acetylene (C₂H₂) from the Infrared Atmospheric Sounding Interferometer (IASI), *Atmos. Meas. Tech.*, 6, 917–925, doi:10.5194/amt-6-917-2013, 2013.

15 Edwards, D. P., Lamarque, J.-F., Attié, J.-L., Emmons, L. K., Richter, A., Cammas, J.-P., Gille, J. C., Francis, G. L., Deeter, M. N., Warner, J., Ziskin, D. C., Lyjak, L. V., Drummond, J. R., and Burrows, J. P.: Tropospheric ozone over the tropical Atlantic: a satellite perspective, *J. Geophys. Res.*, 108, 4237, doi:10.1029/2002JD002927, 2003.

20 Edwards, D. P., Emmons, L. K., Gille, J. C., Chu, A., Attié, J.-L., Giglio, L., Wood, S. W., Haywood, J., Deeter, M. N., Massie, S. T., Ziskin, D. C., and Drummond, J. R.: Satellite-observed pollution from Southern Hemisphere biomass burning, *J. Geophys. Res.*, 111, D14312, doi:10.1029/2005JD006655, 2006a.

25 Edwards, D. P., Emmons, L. K., Gille, J. C., Chu, A., Attié, J.-L., Giglio, L., Wood, S. W., Haywood, J., Deeter, M. N., Massie, S. T., Ziskin, D. C., and Drummond, J. R.: Satellite-observed pollution from Southern Hemisphere biomass burning, *J. Geophys. Res.*, 111, D14312, doi:10.1029/2005JD006655, 2006b.

30 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, doi:10.5194/gmd-3-43-2010, 2010a.

**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Emmons, L. K., Apel, E. C., Lamarque, J.-F., Hess, P. G., Avery, M., Blake, D., Brune, W., Campos, T., Crawford, J., DeCarlo, P. F., Hall, S., Heikes, B., Holloway, J., Jimenez, J. L., Knapp, D. J., Kok, G., Mena-Carrasco, M., Olson, J., O'Sullivan, D., Sachse, G., Walega, J., Weibring, P., Weinheimer, A., and Wiedinmyer, C.: Impact of Mexico City emissions on regional air quality from MOZART-4 simulations, *Atmos. Chem. Phys.*, 10, 6195–6212, doi:10.5194/acp-10-6195-2010, 2010b.

Fortems-Cheiney, A., Chevallier, F., Pison, I., Bousquet, P., Szopa, S., Deeter, M. N., and Clerbaux, C.: Ten years of CO emissions as seen from Measurements of Pollution in the Troposphere (MOPITT), *J. Geophys. Res.*, 116, D05304, doi:10.1029/2010JD014416, 2011.

Glatthor, N., von Clarmann, T., Stiller, G. P., Funke, B., Koukouli, M. E., Fischer, H., Grabowski, U., Höpfner, M., Kellmann, S., and Linden, A.: Large-scale upper tropospheric pollution observed by MIPAS HCN and C₂H₆ global distributions, *Atmos. Chem. Phys.*, 9, 9619–9634, doi:10.5194/acp-9-9619-2009, 2009.

Glatthor, N., Höpfner, M., Semeniuk, K., Lupu, A., Palmer, P. I., McConnell, J. C., Kaminski, J. W., von Clarmann, T., Stiller, G. P., Funke, B., Kellmann, S., Linden, A., and Wiegele, A.: The Australian bushfires of February 2009: MIPAS observations and GEM-AQ model results, *Atmos. Chem. Phys.*, 13, 1637–1658, doi:10.5194/acp-13-1637-2013, 2013.

Glatthor, N., Höpfner, M., Stiller, G. P., von Clarmann, T., Funke, B., Lossow, S., Eckert, E., Grabowski, U., Kellmann, S., Linden, A., A. Walker, K., and Wiegele, A.: Seasonal and inter-annual variations in HCN amounts in the upper troposphere and lower stratosphere observed by MIPAS, *Atmos. Chem. Phys.*, 15, 563–582, doi:10.5194/acp-15-563-2015, 2015.

González Abad, G., Allen, N. D. C., Bernath, P. F., Boone, C. D., McLeod, S. D., Manney, G. L., Toon, G. C., Carouge, C., Wang, Y., Wu, S., Barkley, M. P., Palmer, P. I., Xiao, Y., and Fu, T. M.: Ethane, ethyne and carbon monoxide concentrations in the upper troposphere and lower stratosphere from ACE and GEOS-Chem: a comparison study, *Atmos. Chem. Phys.*, 11, 9927–9941, doi:10.5194/acp-11-9927-2011, 2011.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.

Gyawali, M., Arnott, W. P., Lewis, K., and Moosmüller, H.: In situ aerosol optics in Reno, NV, USA during and after the summer 2008 California wildfires and the influence of absorbing

**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and non-absorbing organic coatings on spectral light absorption, *Atmos. Chem. Phys.*, 9, 8007–8015, doi:10.5194/acp-9-8007-2009, 2009.

Hoskins, B. J. and Rodwell, M. J.: A model of the Asian summer monsoon, part I: the global scale, *J. Atmos. Sci.*, 52, 1329–1340, 1995.

5 Hyer, E. J., Reid, J. S., Prins, E. M., Hoffman, J. P., Schmidt, C. C., Miettinen, J. I., and Giglio, L.: Patterns of fire activity over Indonesia and Malaysia from polar and geostationary satellite observations, *Atmos. Res.*, 122, 504–519, 2013.

Lamarque, J.-F., Hess, P., Emmons, L., Buja, L., Washington, W., and Granier, C.: Tropospheric ozone evolution between 1890 and 1990, *J. Geophys. Res.*, 110, D08304, doi:10.1029/2004JD005537, 2005.

10 Lewis, S. L., Brando, P. M., Phillips, O. L., van der Heijden, G. M. F., and Nepstad, D.: The 2010 Amazon drought, *Science*, 331, 554, doi:10.1126/science.1200807, 2011.

Li, Q., Jacob, D., Bey, I., Yantosca, R., Zhao, Y., Kondo, Y., and Notholt, J.: Atmospheric hydrogen cyanide (HCN): biomass burning source, ocean sink?, *Geophys. Res. Lett.*, 27, 357–360, 2000.

15 Li, Q., Palmer, P. I., Pumphrey, H. C., Bernath, P., and Mahieu, E.: What drives the observed variability of HCN in the troposphere and lower stratosphere?, *Atmos. Chem. Phys.*, 9, 8531–8543, doi:10.5194/acp-9-8531-2009, 2009.

Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: a global perspective, *J. Geophys. Res.*, 86, 7210–7254, 1981.

Lupu, A., Kaminski, J. W., Neary, L., McConnell, J. C., Toyota, K., Rinsland, C. P., Bernath, P. F., Walker, K. A., Boone, C. D., Nagahama, Y., and Suzuki, K.: Hydrogen cyanide in the upper troposphere: GEM-AQ simulation and comparison with ACE-FTS observations, *Atmos. Chem. Phys.*, 9, 4301–4313, doi:10.5194/acp-9-4301-2009, 2009.

25 Magi, B. I., Rabin, S., Shevliakova, E., and Pacala, S.: Separating agricultural and non-agricultural fire seasonality at regional scales, *Biogeosciences*, 9, 3003–3012, doi:10.5194/bg-9-3003-2012, 2012.

Mahieu, E., Duchatelet, P., Bernath, P. F., Boone, C. D., De Mazière, M., Demoulin, P., Rinsland, C. P., Servais, C., and Walker, K. A.: Retrievals of C₂H₂ from high-resolution FTIR solar spectra recorded at the Jungfraujoch station (46.5° N) and comparison with ACE-FTS observations, *Geophys. Res. Abstr.*, 10, available at: <http://hdl.handle.net/2268/15191> (last access: 21 May 2015), 2008.

**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mei, L., Xue, Y., de Leeuw, G., Guang, J., Wang, Y., Li, Y., Xu, H., Yang, L., Hou, T., He, X., Wu, C., Dong, J., and Chen, Z.: Integration of remote sensing data and surface observations to estimate the impact of the Russian wildfires over Europe and Asia during August 2010, *Biogeosciences*, 8, 3771–3791, doi:10.5194/bg-8-3771-2011, 2011.

5 Park, M., Randel, W. J., Emmons, L. K., Bernath, P. F., Walker, K. A., and Boone, C. D.: Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric Chemistry Experiment (ACE-FTS) data, *Atmos. Chem. Phys.*, 8, 757–764, doi:10.5194/acp-8-757-2008, 2008.

Parker, R. J., Remedios, J. J., Moore, D. P., and Kanawade, V. P.: Acetylene C₂H₂ retrievals from MIPAS data and regions of enhanced upper tropospheric concentrations in August 2003, *Atmos. Chem. Phys.*, 11, 10243–10257, doi:10.5194/acp-11-10243-2011, 2011.

10 Parrington, M., Palmer, P. I., Henze, D. K., Tarasick, D. W., Hyer, E. J., Owen, R. C., Helmig, D., Clerbaux, C., Bowman, K. W., Deeter, M. N., Barratt, E. M., Coheur, P.-F., Hurtmans, D., Jiang, Z., George, M., and Worden, J. R.: The influence of boreal biomass burning emissions on the distribution of tropospheric ozone over North America and the North Atlantic during 2010, *Atmos. Chem. Phys.*, 12, 2077–2098, doi:10.5194/acp-12-2077-2012, 2012.

Paton-Walsh, C., Deutscher, N. M., Griffith, D. W. T., Forgan, B. W., Wilson, S. R., Jones, N. B., and Edwards, D. P.: Trace gas emissions from savanna fires in northern Australia, *J. Geophys. Res.*, 115, D16314, doi:10.1029/2009JD013309, 2010.

20 Pfister, G. G., Emmons, L. K., Hess, P. G., Honrath, R., Lamarque, J.-F., Val Martin, M., Owen, R. C., Avery, M. A., Browell, E. V., Holloway, J. S., Nédélec, P., Purvis, R., Ryrson, T. B., Sachse, G. W., and Schlager, H.: Ozone production from the 2004 North American boreal fires, *J. Geophys. Res.*, 111, D24S07, doi:10.1029/2006JD007695, 2006.

Pfister, G. G., Emmons, L. K., Hess, P. G., Lamarque, J.-F., Thompson, A. M., and Yorks, J. E.: Analysis of the summer 2004 ozone budget over the United States using Intercontinental Transport Experiment Ozone-sonde Network Study (IONS) observations and Model of Ozone and Related Tracers (MOZART-4) simulations, *J. Geophys. Res.*, 113, D23306, doi:10.1029/2008JD010190, 2008.

25 Pfister, G. G., Parrish, D. D., Worden, H., Emmons, L. K., Edwards, D. P., Wiedinmyer, C., Diskin, G. S., Huey, G., Oltmans, S. J., Thouret, V., Weinheimer, A., and Wisthaler, A.: Characterizing summertime chemical boundary conditions for air masses entering the US West Coast, *Atmos. Chem. Phys.*, 11, 1769–1790, doi:10.5194/acp-11-1769-2011, 2011.

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pumphrey, H. C., Santee, M. L., Livesey, N. J., Schwartz, M. J., and Read, W. G.: Microwave Limb Sounder observations of biomassburning products from the Australian bush fires of February 2009, *Atmos. Chem. Phys.*, 11, 6285–6296, doi:10.5194/acp-11-6285-2011, 2011.
- Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., Walker, K. A., Boone, C., and Pumphrey, H.: Asian monsoon transport of pollution to the stratosphere, *Science*, 328, 611–613, doi:10.1126/science.1182274, 2010.
- R'Honi, Y., Clarisse, L., Clerbaux, C., Hurtmans, D., Dufлот, V., Turquety, S., Ngadi, Y., and Coheur, P.-F.: Exceptional emissions of NH₃ and HCOOH in the 2010 Russian wildfires, *Atmos. Chem. Phys.*, 13, 4171–4181, doi:10.5194/acp-13-4171-2013, 2013.
- Rodgers, C. D.: *Inverse Methods for Atmospheric Sounding: Theory and Practice*, Series on Atmospheric, Oceanic and Planetary Physics – Vol. 2, World Scientific Publishing CO, Singapore, 2000.
- Sancho, P., De La Cruz, J., Diaz, A., Martin, F., Hernandez, E., Valero, F., and Albarran, B.: A five-year climatology of back-trajectories from the Izaña baseline station, Tenerife, Canary Islands, *Atmos. Environ.*, 26A, 1081–1096, 1992.
- Sauvage, B., Thouret, V., Cammas, J.-P., Gheusi, F., Athier, G., and Nédélec, P.: Tropospheric ozone over Equatorial Africa: regional aspects from the MOZAIC data, *Atmos. Chem. Phys.*, 5, 311–335, doi:10.5194/acp-5-311-2005, 2005.
- Singh, H. B., Salas, L., Herlth, D., Kolyer, R., Czech, E., Viezee, W., Li, Q., Jacob, D. J., Blake, D., Sachse, G., Harward, C. N., Fuelberg, H., Kiley, C. M., Zhao, Y., and Kondo, Y.: In situ measurements of HCN and CH₃CN over the Pacific Ocean: sources, sinks and budgets, *J. Geophys. Res.*, 108, 8795, doi:10.1029/2002JD003006, 2003.
- Standard Atmosphere 1976, National Oceanic and Atmospheric Administration, National Aeronautics and Space Administration, U.S. Air Force, U.S. Government Printing Office, NOAA-S/T 76-1562, Washington, DC, 228 pp., 1976.
- Tilmes, S., Emmons, L. K., Law, K. S., Ancellet, G., Schlager, H., Paris, J.-D., Fuelberg, H. E., Streets, D. G., Wiedinmyer, C., Diskin, G. S., Kondo, Y., Holloway, J., Schwarz, J. P., Spackman, J. R., Campos, T., Nédélec, P., and Panchenko, M. V.: Source contributions to Northern Hemisphere CO and black carbon during spring and summer 2008 from POLARCAT and START08/preHIPPO observations and MOZART-4, *Atmos. Chem. Phys. Discuss.*, 11, 5935–5983, doi:10.5194/acpd-11-5935-2011, 2011.
- Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J., Erisman, J. W., and Coheur, P. F.: Global distributions, time series and error characterization

**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of atmospheric ammonia (NH₃) from IASI satellite observations, *Atmos. Chem. Phys.*, 14, 2905–2922, doi:10.5194/acp-14-2905-2014, 2014.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emission from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Vigouroux, C., Stavrakou, T., Whaley, C., Dils, B., Dufлот, V., Hermans, C., Kumps, N., Metzger, J.-M., Scolas, F., Vanhaelewyn, G., Müller, J.-F., Jones, D. B. A., Li, Q., and De Mazière, M.: FTIR time-series of biomass burning products (HCN, C₂H₆, C₂H₂, CH₃OH, and HCOOH) at Reunion Island (21° S, 55° E) and comparisons with model data, *Atmos. Chem. Phys.*, 12, 10367–10385, doi:10.5194/acp-12-10367-2012, 2012.

Volkamer, R., Ziemann, P. J., and Molina, M. J.: Secondary organic aerosol formation from acetylene (C₂H₂), seed effect on SOA yields due to organic photochemistry in the aerosol aqueous phase, *Atmos. Chem. Phys.*, 9, 1907–1928, doi:10.5194/acp-9-1907-2009, 2009.

Walker, J.C, Dudhia, A., and Carboni, E.: An effective method for the detection of trace species demonstrated using the MetOp Infrared Atmospheric Sounding Interferometer, *Atmos. Meas. Tech.*, 4, 1567–1580, doi:10.5194/amt-4-1567-2011, 2011.

Wespes, C., Emmons, L., Edwards, D. P., Hannigan, J., Hurtmans, D., Saunio, M., Coheur, P.-F., Clerbaux, C., Coffey, M. T., Batchelor, R. L., Lindenmaier, R., Strong, K., Weinheimer, A. J., Nowak, J. B., Ryerson, T. B., Crouse, J. D., and Wennberg, P. O.: Analysis of ozone and nitric acid in spring and summer Arctic pollution using aircraft, ground-based, satellite observations and MOZART-4 model: source attribution and partitioning, *Atmos. Chem. Phys.*, 12, 237–259, doi:10.5194/acp-12-237-2012, 2012.

Wiedinmyer, C., Quayle, B., Geron, C., Belote, A., McKenzie, D., Zhang, X., O'Neill, S., and Wynne, K.: Estimating emissions from fires in North America for air quality modeling, *Atmos. Environ.*, 40, 3419–3432, 2006.

Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, doi:10.5194/gmd-4-625-2011, 2011.

**Global
measurements of
HCN and C₂H₂ from
IASI**

V. Dufлот et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Wiegele, A., Glatthor, N., Höpfner, M., Grabowski, U., Kellmann, S., Linden, A., Stiller, G., and von Clarmann, T.: Global distributions of C₂H₆, C₂H₂, HCN, and PAN retrieved from MIPAS reduced spectral resolution measurements, *Atmos. Meas. Tech.*, 5, 723–734, doi:10.5194/amt-5-723-2012, 2012.
- 5 Xiao, Y., Jacob, D. J., and Turquety, S.: Atmospheric acetylene and its relationship with CO as an indicator of air mass age, *J. Geophys. Res.-Atmos.*, 112, D12305, doi:10.1029/2006JD008268, 2007.
- Yienger, J. J., M. Galanter, T. A. Holloway, M. J. Phadnis, S. K. Guttikunda, G. R. Carmichael, W. J. Moxim, and Levy II, H.: The episodic nature of air pollution transport from
10 Asia to North America, *J. Geophys. Res.*, 105, 2693126945, doi:10.1029/2000JD900309, 2000.
- Yulianti, N. and Hayasaka, H.: Recent active fires under El Niño conditions in Kalimantan, Indonesia, *Am. J. Plant Sci.*, 4, 685–696, 2013.
- Zander, R., Rinsland, C. P., Ehhalt, D. H., Rudolph, J., and Demoulin P. H.: Vertical column abundances and seasonal cycle of acetylene, C₂H₂, above the Jungfraujoch Station, derived
15 from IR solar observations, *J. Atm. Chem.*, 13, 359–372, 1991.
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131–5153,
20 doi:10.5194/acp-9-5131-2009, 2009.

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Global C₂H₂ and HCN emission sources (Tg(species) year⁻¹) during the period 2008–2010 from the dataset used in MOZART-4.

Sources/Year	C ₂ H ₂			HCN		
	2008	2009	2010	2008	2009	2010
Anthropogenic	3.37	3.37	3.37	1.67	1.67	1.67
Biomass Burning	0.64	0.71	0.83	1.38	1.33	1.58
Total	4.01	4.07	4.20	3.05	3.00	3.25

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Table 2. Correlation coefficients (R) and biases (Bias) between IASI observations and MOZART-4 simulations for each of the zones defined in Fig. 6.

Zones	C ₂ H ₂		HCN	
	R	Bias (%)	R	Bias (%)
NAM	0.93	47	–	–
NCA	–	–	0.44	–1
SAM	0.54	–50	0.76	–14
EUR	0.88	115	–	–
NAF	0.66	–35	0.07	–2
SAF	0.69	–67	0.86	–28
BCA	0.82	105	–	–
SEA	–0.31	–23	0.57	–4
EQA	0.45	–51	0.09	–27
AUS	0.65	–61	0.77	–26
Global	0.72	–1	0.69	–16

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 1. (Top) Simulated spectra in the region of the HCN ν_2 band and C_2H_2 ν_5 band. The green (brown) double sided arrow gives the spectral range used to compute the S_e matrices for HCN (C_2H_2). (Middle) Contributions of climatological background levels of HCN and C_2H_2 . (Bottom) Contribution of CO_2 (red line), O_3 (green line) and H_2O (blue line) to a simulated spectrum for background concentrations. Calculations have been made for the US Standard Atmosphere (US Government Printing Office, 1976) with CO_2 concentrations scaled to 390 ppmv.





Figure 2. Variations of the HRI with HCN (top) and C_2H_2 (bottom) column (molec cm^{-2}) integrated over the 1 km-thick polluted layer in a standard modeled subtropical atmosphere from forward model simulations. The colorscale gives the altitude of the polluted layer.

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. Normalised Jacobians of the forward model implemented in *Atmosphit* for HCN (solid lines) and C_2H_2 (dotted lines) and for the standard modeled polar (blue lines), temperate (green lines) and subtropical (red lines) atmospheres. These are averaged Jacobians over the spectral ranges $645\text{--}800\text{ cm}^{-1}$ for HCN and $645\text{--}845\text{ cm}^{-1}$ for C_2H_2 .

Global measurements of HCN and C_2H_2 from IASI

V. Dufлот et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.




Figure 4. Values of B_{HCN} (blue) and $B_{\text{C}_2\text{H}_2}$ (green) as a function of the latitude.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.




Figure 5. Variation of HRI_{HCN} caused by a TC equal to ± 10 K for the polar (solid line and circles), temperate (dashed line and squares) and tropical (dotted line and crosses) atmospheres.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.




Figure 6. Locations of the 4 ground-based FTIR measurements sites (Jungfrauoch, Izaña, Reunion Island and Wollongong) and map of the 10 regions used in this study: NAM: Northern America, NCA: North Central America, SAM: South America, EUR: Europe, NAF: Northern Africa, SAF: Southern Africa, BCA: Boreal Central Asia, SEA: South East Asia, EQA: Equatorial Asia, AUS: Australia.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.




Figure 7. Total column averaging kernels of ground-based FTIR in $\text{molec cm}^{-2}/\text{molec cm}^{-2}$ for C₂H₂ at Reunion Island (red stars and line), HCN at Reunion Island (black circles and line), HCN at Wollongong (green dots and line), HCN at Izaña (light blue diamonds and line) and C₂H₂ at Jungfrauoch (blue squares and line).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.




Figure 8. Time series of HCN (left panel) and C₂H₂ (right panel) measurements for Reunion Island (HCN and C₂H₂), Wollongong (HCN only), Izaña (HCN only), and Jungfraujoch (C₂H₂ only). IASI measurements are shown as daily and 1° × 1° means (red dots) with associated SDs (light red lines), and as monthly and 1° × 1° means (black circles and line) with associated SD (vertical black lines). Ground-based FTIR measurements are shown as daily means with associated total error by green crosses and lines. Correlation coefficients are given on each plot for daily means in red and for monthly means in black.

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.




Figure 9. Seasonal distribution of the C₂H₂ total column (in molec cm⁻²) as measured by IASI (left panel) and simulated by MOZART-4 (right panel) averaged over the years 2008 to 2010. The IASI global distributions are given with the same horizontal resolution as MOZART-4 (1.875° latitude × 2.5° longitude). DJF = December–January–February, MAM = March–April–May, JJA = June–July–August, SON = September–October–November.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 10. Same as Fig. 9 for HCN.

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 11. Evolution with time of the mean C₂H₂ total column (in molec cm⁻²) over the zones defined in Fig. 6 as measured by IASI (red dots) with associated SD (light red lines), and as simulated by MOZART-4 (black dots). Correlation coefficients (R) and biases (Bias) between IASI and MOZART-4 are given on each plot for daily means.

Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.




Figure 12. Same as Fig. 11 for HCN.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global measurements of HCN and C₂H₂ from IASI

V. Dufлот et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 13. C₂H₂ and HCN surface emission fluxes (kg m⁻² s⁻¹) averaged over the period 2008–2010 from the anthropogenic and fire emissions inventories used in MOZART-4.