



HAL
open science

Formation and properties of SiC nano-particles colloids in non polar liquids

Hamza Hajjaji, Sergei Alekseev, David Philippon, Gérard Guillot, Philippe Vergne, Jean-Marie Bluet

► **To cite this version:**

Hamza Hajjaji, Sergei Alekseev, David Philippon, Gérard Guillot, Philippe Vergne, et al.. Formation and properties of SiC nano-particles colloids in non polar liquids. E-MRS 2014, May 2014, Lille, France. hal-01154677

HAL Id: hal-01154677

<https://hal.science/hal-01154677>

Submitted on 22 May 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

E-MRS: European Materials Research Society-2014

Formation and properties of SiC nano-particles colloids in non polar liquids

Authors : Hamza Hajjaji 1, Sergei Alekseev 2, David Philippon 3, Gérard Guillot 1, Philippe Vergne 3, Jean-Marie Bluet 1

Affiliations :

1 Université de Lyon, CNRS, UMR 5270, INSA-Lyon, INL, F-69621 Villeurbanne, France

2 Chemistry Faculty, Kiev National Taras Shevchenko University, 01601 Kiev, Ukraine 3 Université de Lyon, CNRS, UMR5259, INSA-Lyon, LaMCoS, F-69621 Villeurbanne, France

Resume : The goal of this work is to form SiC nanoparticles colloids in non polar liquids toward temperature sensing application in lubricant. The formation of stable SiC colloids in polar liquids like water or ethanol has been demonstrated and explained by their surface chemistry resulting in a negative zeta potential [1, 2]. The strong fluorescence emission of these colloids led to application as a non toxic bio label [3]. Here, we will present surface functionalization of SiC nanoparticles by long chain amines (nonylamine, octadecylamine) in order to disperse them in non polar liquids like alkane or alkene. The SiC nanoparticles have been obtained by two different top-down technology : the first is electrochemical anodization of polycrystalline SiC in an hydrofluoric acid-ethanol mixture, the second is chemical etching of micrometer size SiC particles in a hydrofluoric -nitric acid mixture. Chemical (FTIR, Raman), structural (TEM, DLS) and optical characterization (fluorescence and fluorescence lifetime) of the resulting colloids will be presented.

[1] J. Botsoa, J.-M. Bluet, V. Lysenko et G. Guillot, J. of Appl. Phys. 102 (8), p. 083526 (2007).

[2] Yu. Zakharko, J. Botsoa, S. Alekseev, V. Lysenko, J.-M. Bluet, O. Marty, V. A. Skryshevsky, and G. Guillot, J. of Appl. Phys. 107, 013503 (2010).

[3] J. Botsoa, V.Lysenko, A. Geloën, O. Marty, J.M. Bluet and G. Guillot, Appl. Phys. Lett. 92, 173902 (2008).