



HAL
open science

BSDEs with diffusion constraint and viscous Hamilton-Jacobi equations with unbounded data

Andrea Cosso, Huyên Pham, Hao Xing

► **To cite this version:**

Andrea Cosso, Huyên Pham, Hao Xing. BSDEs with diffusion constraint and viscous Hamilton-Jacobi equations with unbounded data. 2015. hal-01154898v1

HAL Id: hal-01154898

<https://hal.science/hal-01154898v1>

Preprint submitted on 25 May 2015 (v1), last revised 7 Mar 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BSDEs with diffusion constraint and viscous Hamilton-Jacobi equations with unbounded data

Andrea COSSO*

Huy en PHAM†

Hao XING‡

May 25, 2015

Abstract

We provide a stochastic representation for a general class of viscous Hamilton-Jacobi (HJ) equations, which has convexity and superlinear nonlinearity in its gradient term, via a type of backward stochastic differential equation (BSDE) with constraint in the martingale part. We compare our result with the classical representation in terms of (super)quadratic BSDE, and show in particular that existence of a solution to the viscous HJ equation can be obtained under more general growth assumptions on the coefficients, including both unbounded diffusion coefficient and terminal data.

Keywords: Backward stochastic differential equation (BSDE), randomization, viscous Hamilton-Jacobi equation, deterministic KPZ equation, nonlinear Feynman-Kac formula.

AMS 2010 subject classification: 60H30, 35K58.

1 Introduction

Given $T \in (0, \infty)$ and $d \in \mathbb{N} \setminus \{0\}$, we consider the parabolic semilinear partial differential equation (PDE) of the form:

$$\left\{ \begin{array}{l} -\frac{\partial u}{\partial t}(t, x) - \frac{1}{2} \text{tr}[\sigma \sigma^\top(x) D_x^2 u(t, x)] - b(x) \cdot D_x u(t, x) \\ \quad + F(x, u(t, x), \varrho^\top(x) D_x u(t, x)) = 0, \quad (t, x) \in [0, T) \times \mathbb{R}^d, \\ u(T, x) = g(x), \quad x \in \mathbb{R}^d. \end{array} \right. \quad (1.1)$$

Here σ^\top and ϱ^\top stand for the transpose of σ and ϱ , $D_x u$ and $D_x^2 u$ are the gradient and the Hessian of u , respectively, and \cdot represents the inner product between two vectors in

*Laboratoire de Probabilit es et Mod es Al atoires, CNRS, UMR 7599, Universit  Paris Diderot, cosso@math.univ-paris-diderot.fr

†Laboratoire de Probabilit es et Mod es Al atoires, CNRS, UMR 7599, Universit  Paris Diderot, and CREST-ENSAE, pham@math.univ-paris-diderot.fr

‡Department of Statistics, London School of Economics and Political Science, h.xing@lse.ac.uk

\mathbb{R}^d . The functions $b: \mathbb{R}^d \rightarrow \mathbb{R}^d$, $\sigma, \varrho: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$, and $F: \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ are called coefficients of the equation, and $g: \mathbb{R}^d \rightarrow \mathbb{R}$ gives the terminal condition.

The focus of this paper is to provide a probabilistic representation for solutions to (1.1). This is a classical problem and the standard approach (when $\varrho = \sigma$) is to study a backward stochastic differential equation (BSDE) whose generator is given by F , and associated to a forward diffusion with coefficients b and σ , namely:

$$\begin{cases} X_s^{t,x} &= x + \int_t^s b(X_r^{t,x})dr + \int_t^s \sigma(X_r^{t,x})dW_r, & (t, x) \in [0, T] \times \mathbb{R}^d, t \leq s \leq T, \\ Y_s^{t,x} &= g(X_T^{t,x}) - \int_s^T F(X_r^{t,x}, Y_r^{t,x}, Z_r^{t,x})dr - \int_s^T Z_r^{t,x}dW_r, \end{cases} \quad (1.2)$$

where W is a d -dimensional Brownian motion. The forward backward SDE (1.2) is then formally connected to the PDE (1.1) by the nonlinear Feynman-Kac formula: $Y_t^{t,x} = u(t, x)$. Assuming that F is superlinear (which includes the quadratic and superquadratic case) and convex in its last argument (in which case, the PDE (1.1) is referred to as generalized *viscous Hamilton-Jacobi equation*), the existence and uniqueness of a solution to the BSDE (1.2) is a quite complicated issue, extensively studied in the literature, and this paper provides an alternative representation by considering furthermore the case when ϱ may be different from σ , without imposing any non degeneracy condition on ϱ, σ . Comparison to the literature will be presented in Remark 1.1 and in three examples below. To state our main result, we introduce the following assumptions which will be enforced throughout the paper.

Standing Assumption 1.1

(i) *There exists a constant $L_{b,\sigma,\varrho}$ such that*

$$|b(x) - b(x')| + \|\sigma(x) - \sigma(x')\| + \|\varrho(x) - \varrho(x')\| \leq L_{b,\sigma,\varrho}|x - x'|, \quad \text{for all } x, x' \in \mathbb{R}^d,$$

where $\|A\| = \sqrt{\text{tr}(AA^\top)}$ denotes the Frobenius norm of any matrix A .

(ii) *There exist constants $M_\varrho \geq 0$ and $p_\varrho \in [0, 1]$ such that*

$$\|\varrho(x)\| \leq M_\varrho(1 + |x|^{p_\varrho}), \quad \text{for all } x \in \mathbb{R}^d.$$

(iii) *There exists a constant L_F such that*

$$|F(x, y, z) - F(x, y', z)| \leq L_F|y - y'|, \quad \text{for any } y, y' \in \mathbb{R} \text{ and } (x, z) \in \mathbb{R}^d \times \mathbb{R}^d.$$

(iv) *The map $z \mapsto F(x, y, z)$ is convex. There exist constants $m_F, M_F \geq 0$, $q \geq p > 1$, $p_F \geq 0$, and $q_F \in [0, (1 - p_\varrho)q/(q - 1)]$ if $p_\varrho < 1$, or $q_F = 0$ if $p_\varrho = 1$, such that*

$$-m_F(1 + |x|^{p_F} - \frac{1}{p}|z|^p) \leq F(x, 0, z) \leq M_F(1 + |x|^{q_F} + \frac{1}{q}|z|^q),$$

for any $(x, z) \in \mathbb{R}^d \times \mathbb{R}^d$.

(v) *Define the convex conjugate (or Fenchel-Legendre transform) of F as*

$$f(x, a, y) = - \inf_{z \in \mathbb{R}^d} [a \cdot z + F(x, y, z)], \quad \text{for all } (x, a, y) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}. \quad (1.3)$$

The map $x \mapsto f(x, a, y)$ is continuous.

- (vi) The function g is continuous. There exist constants $m_g, M_g, q_g \geq 0$, and $p_g \in [0, (1 - p_\varrho)q/(q - 1))$ if $p_\varrho < 1$, or $p_g = 0$ if $p_\varrho = 1$, such that

$$-m_g(1 + |x|^{p_g}) \leq g(x) \leq M_g(1 + |x|^{q_g}), \quad \text{for all } x \in \mathbb{R}^d.$$

Let us now comment the above assumptions and their connection with related literature.

Remark 1.1 1. The condition $q \geq p > 1$ in Assumption 1.1 (iv) means that F has a superlinear growth in z . We also consider that F may have some polynomial growth condition on x , and we distinguish the growth coefficients p_F and q_F for the lower and upper bounds. Indeed, notice that no condition is required on p_F , while we impose some upper bound on q_F depending on q and the (sub)linear growth coefficient p_ϱ in Assumption 1.1 (ii). Observe that this upper bound $\bar{q}_F = (1 - p_\varrho)q/(q - 1)$ is decreasing with p_ϱ and q , with a limiting value equal to infinity when q goes to 1, and equal to $1 - p_\varrho$ when q goes to infinity; $1 - p_\varrho$ shrinks to zero (i.e. F is upper-bounded in x) when $p_\varrho = 1$ (i.e. ϱ satisfies a linear growth condition). Similarly, the terminal function g is enabled to satisfy polynomial growth with the same condition on p_g related to q and p_ϱ .

When $q = 2$, F has at most quadratic growth in z . A stochastic representation of (1.1) is given by the *quadratic* BSDE, which has been studied extensively, see [13, 4, 5, 9, 21, 2, 6, 1], amongst others. For example, in the Markovian case of [13], F and g are assumed to be bounded in x , i.e. $q_F = p_F = q_g = p_g = 0$, with $\varrho = \sigma$ satisfying a linear growth condition, i.e. $p_\varrho = 1$. This case is covered by our assumptions (actually, we only need to assume $q_F = p_g = 0$ when $p_\varrho = 1$, but no condition is required on p_F, q_g). In the Markovian setting of [5], [9] or [21], $\varrho = \sigma$ is assumed to be a bounded function, which corresponds to the case where $p_\varrho = 0$ in Assumption 1.1 (ii). Notice that we do not assume any non degeneracy condition on σ so that the case of time dependent coefficients as in [9] or [21], can be embedded in our framework by extending the spatial variables from x to (t, x) . When $p > 2$ in Assumption 1.1 (iv), F has *super-quadratic* growth in z , which has been studied in [3, 10, 11, 8, 21, 14]. We shall illustrate the scope of Assumption 1.1 in the next three examples. In particular, we compare the condition on q_F, p_g to existing results from both analytic and probabilistic aspects.

2. The case where $z \mapsto F(x, y, z)$ is *concave* can be deduced from the convex case. Indeed, set $\tilde{F}(x, y, z) = -F(x, -y, -z)$. Then \tilde{F} is convex in z and our result can be applied to \tilde{u} which solves (1.1) with \tilde{F} in place of F and $-g$ in place of g .

3. Assumption 1.1 (v) is satisfied under the following two sufficient conditions:

- The map $x \mapsto F(x, y, z)$ is continuous, uniformly with respect to y and z , i.e., for all $x \in \mathbb{R}^d$ and $\varepsilon > 0$, there exists $\delta = \delta(x, \varepsilon) > 0$ such that,

$$|F(x, y, z) - F(x', y, z)| \leq \varepsilon, \quad \text{for any } |x - x'| \leq \delta \text{ and } (y, z) \in \mathbb{R} \times \mathbb{R}^d.$$

- The map $(x, z) \mapsto F(x, y, z)$ is continuous for any $y \in \mathbb{R}$, and the optimizer $z^* = z^*(x, a, y)$ in (1.3) is continuous in x . \square

Example 1.1 (Generalized deterministic KPZ equation) Consider $F(z) = -\lambda|z|^q$ for some constants $\lambda > 0$ and $q > 1$. The equation (1.1), with $\varrho = \sigma$ as the identity matrix, is referred to as the *generalized deterministic KPZ equation* with the $q = 2$ case introduced by Kardar, Parisi, and Zhang in connection with the study of growing surfaces. In this case, $\tilde{F}(z) = \lambda|z|^q$ satisfies Assumption 1.1, in particular, $p_\varrho = 0$ in (ii) and $p_F = q_F = 0$ in (iv). This equation has been studied from the analytical point of view in, for example, [3], [10], and [11]. In particular, concerning Assumption 1.1 (iv), in [11, Theorem 2.6] $\max\{p_g, q_g\} < q/(q-1)$ is assumed, while here we only assume: $p_g < q/(q-1)$, but no restriction on q_g . Moreover, when $\max\{p_g, q_g\} = q/(q-1)$, a solution which explodes in finite time is presented in [11, Remark 4.6]. \square

Example 1.2 (BSDEs with superquadratic growth) Motivated by the previous example, Delbaen, Hu, and Bao [8] studied BSDEs whose generator has superquadratic growth in the “ Z -component”. A solution to this superquadratic BSDE provides a probabilistic representation for the solution of (1.1) (when $\varrho = \sigma$) with $q > 2$. In particular, given a forward process

$$dX_s = b(s, X_s)ds + \sigma dW_s,$$

with a Brownian motion W , a continuous differentiable function b with bounded derivative, and a constant matrix σ , consider the following BSDE

$$Y_s = g(X_T) - \int_s^T F(Z_u)du - \int_s^T Z_u dW_u,$$

where g is bounded and continuous, F is nonnegative, convex, and satisfies $F(0) = 0$ and $\overline{\lim}_{|z| \rightarrow \infty} \frac{F(z)}{|z|^2} = \infty$. Proposition 4.4 in [8] presents a solution to the previous superquadratic BSDE. This result has been extended in [21] and [14], where F can depend on x and y , without convexity assumption on z , and σ can be a deterministic function of time. In these cases, $p_\varrho = 0$, [14, Assumptions (B.1) and (TC.1)] implies that $\max\{p_F, q_F\} < q/(q-1)$ and $\max\{p_g, q_g\} < q/(q-1)$. When σ depends on x , only existence for small time is available, see [21, Proposition 3.1]. Compared to aforementioned works, our main result (see Theorem 1.1 below) provides an alternative representation for (1.1) when F is convex in z , allowing a more general setting with (sub)linear growth condition on ϱ , and restrictions on q_F, p_g only, but not on p_F, q_g (hence with both unbounded diffusion coefficient and terminal data), and including also the case where $\varrho = \sigma$ is bounded, i.e. $p_\varrho = 0$, and $q_F, p_g < q/(q-1)$. \square

Example 1.3 (Utility maximization) Our framework allows us to incorporate the two following financial applications.

(i) *Portfolio optimization.* Consider a factor model of a financial market with a risk free asset S^0 and risky assets $S = (S^1, \dots, S^n)$ with dynamics

$$dS_t^0 = S_t^0 r dt \quad \text{and} \quad dS_t = \text{diag}(S_t)[(r1_n + \mu(X_t))dt + a(X_t)dB_t], \quad (1.4)$$

where $r \in \mathbb{R}$, μ and a are measurable functions on \mathbb{R}^d , valued respectively on \mathbb{R}^n and $\mathbb{R}^{n \times n}$, such that aa^\top is invertible, $\text{diag}(S)$ is a diagonal matrix with elements of S on the diagonal, 1_n is a n -dimensional vector with every entry 1, and B is a n -dimensional Brownian motion.

We denote by $\lambda = a^\top(aa^\top)^{-1}\mu$, the so-called Sharpe ratio. In the dynamics (1.4), the factor X is a d -dimensional process governed by

$$dX_t = \beta(X_t)dt + \sigma(X_t)dW_t \quad (1.5)$$

where β and σ are Lipschitz functions on \mathbb{R}^d , valued respectively on \mathbb{R}^d and $\mathbb{R}^{d \times d}$, and W is a d -dimensional Brownian motion. The correlation between B and W is given by $d\langle W, B \rangle_t = \rho dt$ for some $\rho \in \mathbb{R}^{n \times d}$.

An agent with power utility $U(w) = w^\gamma/\gamma$ for $\gamma < 1$, $\gamma \neq 0$ invests in this market in a self-financing way in order to maximize her expected utility of terminal wealth at an investment horizon T . Let $v(t, w, x)$ be the value function of the investor and define the reduced value function u via $v(t, w, x) = (w^\gamma/\gamma)e^{u(t, x)}$. Then the following equation, satisfied by u , is of the same type as (1.1) with $\varrho = \sigma$ (see e.g. [16, Equation (2.14)]):

$$\begin{cases} -\frac{\partial u}{\partial t} - \frac{1}{2}\text{tr}[\sigma\sigma^\top D_x^2 u] - b \cdot D_x u + F(x, \sigma^\top D_x u) & = 0 \\ u(T, \cdot) & = 0, \end{cases}$$

where

$$\begin{aligned} b(x) &= \beta(x) + \frac{\gamma}{1-\gamma}\sigma(x)\rho^\top\lambda(x), & F(x, z) &= -\frac{1}{2}zMz^\top - h(x), \\ M &= \text{diag}(1_d) + \frac{\gamma}{1-\gamma}\rho^\top\rho, & h(x) &= \gamma r + \frac{1}{2}\frac{\gamma}{1-\gamma}|\lambda(x)|^2. \end{aligned}$$

We assume that b is Lipschitz. Moreover, note that M is positive definite. Hence $\tilde{F}(x, z) = -F(x, -z)$ satisfies Assumption 1.1 (iv) with $p = q = 2$, and whenever λ satisfies the condition:

$$-m_F(1 + |x|^{p_F}) \leq \gamma|\lambda|^2 \leq M_F(1 + |x|^{q_F}),$$

for some nonnegative constants m_F , M_F , p_F , and $q_F < 2(1 - p_\varrho)$. Hence, when $\gamma > 0$, such condition holds whenever λ satisfies a strict sub-linear growth condition; while for $\gamma < 0$ (the empirically relevant case in financial context), this holds once λ satisfies a polynomial growth condition. In particular, the second scenario includes the case where a constant, and $\mu(x)$ (thus $\lambda(x)$) is affine in x , as in the original Kim-Omberg model where X is an Ornstein-Uhlenbeck process with σ constant and $\beta(x)$ affine in x .

(ii) *Indifference pricing.* We consider a financial model as in (1.4)-(1.5), where the process X represents now the level of nontraded assets (e.g. volatility index, temperature), correlated with the traded assets of price S , for which the Sharpe ratio λ is assumed to be bounded. Given an European option written on the nontraded asset, with payoff $g(X_T)$ at maturity T , and following the indifference pricing criterion (see e.g. [15]), we consider the problem of an agent with exponential utility $U(w) = -e^{-\gamma w}$, $\gamma > 0$, who invests in the traded assets S up to T where he has to deliver the option $g(X_T)$. Let $v(t, w, x)$ be the value function of the agent, and define the reduced value function u via: $v(t, w, x) = U(w - u(t, x))$. Then, u satisfies an equation of type (1.1) with $\varrho = \sigma$:

$$\begin{cases} -\frac{\partial u}{\partial t} - \frac{1}{2}\text{tr}[\sigma\sigma^\top D_x^2 u] - b \cdot D_x u + F(x, \sigma^\top D_x u) & = 0 \\ u(T, \cdot) & = g, \end{cases}$$

where

$$b(x) = \beta(x) + \sigma(x)\rho^\top\lambda(x), \quad F(x, z) = -\frac{1}{2}z(\text{diag}(1_d) - \rho^\top\rho)z^\top + \gamma r + \frac{1}{2}\frac{|\lambda(x)|^2}{\gamma}.$$

Since λ is assumed to be bounded, $\tilde{F}(x, z) = -F(x, -z)$ satisfies Assumption 1.1 (iv) with $p = q = 2$, $p_F = q_F = 0$. Moreover, Assumptions 1.1 (ii) and (vi) enable us to consider unbounded diffusion coefficient σ and unbounded payoff function g , for example when X is governed by the “shifted” CEV model with $\sigma(x) = \sigma_0(\sigma_1 + x)^{p_\sigma}$, for some positive constants σ_0, σ_1 , and $p_\sigma \in [0, 1)$ (the introduction of the positive constant σ_1 ensures that σ is a Lipschitz function), and g satisfies the growth condition (recall Remark 1.1 **2.**):

$$-M_g(1 + |x|^{q_g}) \leq g(x) \leq m_g(1 + |x|^{p_g}),$$

for some nonnegative constants m_g, M_g, q_g , and $0 \leq p_g < 2(1 - p_\sigma)$. \square

Before presenting our main result, let us first present an equivalent formulation of (1.1) in terms of an Hamilton-Jacobi-Bellman (HJB) equation. Recall the convex conjugate function f in (1.3). By convexity of F in z , we then have the following duality relationship

$$F(x, y, z) = -\inf_{a \in \mathbb{R}^d} [a \cdot z + f(x, a, y)], \quad \text{for all } (x, y, z) \in \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d.$$

Therefore equation (1.1) can be rewritten as the following HJB equation:

$$\begin{cases} -\frac{\partial u}{\partial t}(t, x) - \inf_{a \in \mathbb{R}^d} [\mathcal{L}^a u(t, x) + f(x, a, u(t, x))] = 0, & (t, x) \in [0, T) \times \mathbb{R}^d, \\ u(T, x) = g(x), & x \in \mathbb{R}^d, \end{cases} \quad (1.6)$$

where

$$\mathcal{L}^a u(t, x) = (b(x) + \varrho(x)a) \cdot D_x u(t, x) + \frac{1}{2} \text{tr}[\sigma \sigma^\top(x) D_x^2 u(t, x)]. \quad (1.7)$$

It is standard to relate equation (1.6), at least formally, to the following optimal stochastic control problem of a recursive type:

$$u(t, x) = \inf_{\alpha} \mathbb{E} \left[\int_t^T f(X_s^{t,x,\alpha}, \alpha_s, u(s, X_s^{t,x,\alpha})) ds + g(X_T^{t,x,\alpha}) \right], \quad (1.8)$$

where the infimum is taken over all \mathbb{R}^d -valued predictable processes α and the controlled diffusion $X^{t,x,\alpha}$ evolves according to the equation

$$X_s^{t,x,\alpha} = x + \int_t^s (b(X_r^{t,x,\alpha}) + \varrho(X_r^{t,x,\alpha}) \alpha_r) dr + \int_t^s \sigma(X_r^{t,x,\alpha}) dW_r, \quad t \leq s \leq T, \quad (1.9)$$

where W is a d -dimensional Brownian motion.

Rather than studying the control problem (1.8) directly, following [12], we introduce a *randomized control* formulation. For every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, we consider the following forward system of stochastic differential equations:

$$\begin{cases} X_s^{t,x,a} = x + \int_t^s (b(X_r^{t,x,a}) + \varrho(X_r^{t,x,a}) I_r^{t,a}) dr + \int_t^s \sigma(X_r^{t,x,a}) dW_r, \\ I_s^{t,a} = a + B_s - B_t, \end{cases} \quad (1.10)$$

where B is a d -dimensional Brownian motion, independent of W . In the next section, we shall check, under Assumption 1.1 (i) and (ii), that there exists a unique solution $(X^{t,x,a}, I^{t,a})$. System (1.10) is the randomized version of the controlled dynamics (1.9). More precisely, the randomization procedure is performed by introducing the independent Brownian motion B , which is the natural choice when the control process α takes values in the entire space \mathbb{R}^d as in the present case. On the contrary, if the control process α is A -valued, for some compact subset A of \mathbb{R}^d , then a natural randomization is carried out by means of an independent Poisson random measure μ on $[0, \infty) \times A$; see [12].

Now we introduce a stochastic representation for (1.6) and (1.1), via the following *BSDE with diffusion constraint*. Given $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, consider

$$Y_s = g(X_T^{t,x,a}) + \int_s^T f(X_r^{t,x,a}, I_r^{t,a}, Y_r) dr - (K_T - K_s) - \int_s^T Z_r dW_r - \int_s^T V_r dB_r, \quad (1.11)$$

for all $s \in [t, T]$, together with the constraint

$$V_s = 0, \quad ds \otimes d\mathbb{P}\text{-a.e.} \quad (1.12)$$

The presence of the constraint (1.12) forces the introduction of the nondecreasing process K . In Theorem 2.1 in Section 2 below, we construct and prove the existence of a unique maximal solution $(Y^{t,x,a}, Z^{t,x,a}, V^{t,x,a}, K^{t,x,a})$ to (1.11)-(1.12). This allows us to present the main result of this paper, whose proof is presented in Section 3.

Theorem 1.1 *For all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $a, a' \in \mathbb{R}^d$,*

$$Y_t^{t,x,a} = Y_t^{t,x,a'}.$$

Define

$$u(t, x) := Y_t^{t,x,a}, \quad \text{for all } (t, x) \in [0, T] \times \mathbb{R}^d. \quad (1.13)$$

Then u is a viscosity solution to equation (1.1) (or, equivalently, to (1.6)).

Remark 1.2 Theorem 1.1 establishes the existence of a solution to the viscous HJ equation (1.1) under Assumption 1.1, which covers the case of convexity and superquadratic growth in the gradient term, with possibly state dependent and unbounded coefficient ρ , and unbounded terminal data g . This solution is constructed and also represented by the maximal solution of the BSDE with diffusion constraint (1.10)-(1.11)-(1.12). This is in contrast to [7] where viscosity solution techniques are applied. Regarding the uniqueness of solutions, [7, Theorem 3.1] shows that, when ρ is bounded, the function u given by the nonlinear Feynman-Kac formula (1.13) is the *unique* viscosity solution to equation (1.1) in the class of functions satisfying a polynomial growth condition. \square

2 BSDE with diffusion constraint

Let us introduce our probabilistic setting. Consider a filtered probability space $(\Omega, \mathcal{F}_T, \mathbb{F} = (\mathcal{F}_s)_{0 \leq s \leq T}, \mathbb{P})$, where \mathbb{F} is the standard augmentation of the filtration generated by two d -dimensional independent Brownian motions $W = (W_s)_{0 \leq s \leq T}$ and $B = (B_s)_{0 \leq s \leq T}$. For $0 \leq$

$t \leq T$, we also consider the “shifted” version $(\Omega, \mathcal{F}_T^t, \mathbb{F}^t = (\mathcal{F}_s^t)_{t \leq s \leq T}, \mathbb{P})$, where $(W_s)_{0 \leq s \leq T}$ and $(B_s)_{0 \leq s \leq T}$ before are replaced by $(W_{s-t})_{t \leq s \leq T}$ and $(B_{s-t})_{t \leq s \leq T}$, respectively. We denote by \mathbb{E}_s^t the conditional expectation under \mathbb{P} given \mathcal{F}_s^t for $0 \leq t \leq s \leq T$, and see that \mathbb{E}_t^t coincides with the “ordinary” expectation \mathbb{E} . On this “shifted” probability space, we introduce the following spaces of stochastic processes.

- $\mathbb{S}^2(t, T)$: the set of real-valued càdlàg \mathbb{F}^t -adapted processes $Y = (Y_s)_{t \leq s \leq T}$ satisfying

$$\|Y\|_{\mathbb{S}^2(t, T)}^2 := \mathbb{E} \left[\sup_{t \leq s \leq T} |Y_s|^2 \right] < \infty.$$

- $\mathbb{H}^2(t, T)$: the set of \mathbb{R}^d -valued \mathbb{F}^t -predictable processes $Z = (Z_s)_{t \leq s \leq T}$ satisfying

$$\|Z\|_{\mathbb{H}^2(t, T)}^2 := \mathbb{E} \left[\int_t^T |Z_s|^2 ds \right] < \infty.$$

- $\mathbb{K}^2(t, T)$: the set of nondecreasing \mathbb{F}^t -predictable processes $K = (K_s)_{t \leq s \leq T} \in \mathbb{S}^2(t, T)$ with $K_t = 0$. We have

$$\|K\|_{\mathbb{S}^2(t, T)}^2 = \mathbb{E}[K_T^2].$$

This section focuses on the construction of a maximal solution to the forward backward SDE with diffusion constraint (1.10)-(1.11)-(1.12). We first check the existence of a unique solution to the randomized forward system (1.10), and show some useful moment estimates. Throughout this section, Assumption 1.1 is enforced.

Lemma 2.1 *For any $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, there exists a unique (up to indistinguishability) adapted and continuous process $\{(X_s^{t, x, a}, I_s^{t, a}), t \leq s \leq T\}$ solving (1.10). Moreover, for every $m \geq 1$, there exists a positive constant C , depending only on $m, T, L_{b, \sigma, \varrho}, M_{\varrho}$, and p_{ϱ} such that*

$$\mathbb{E}_s^t \left[\sup_{r \in [s, T]} |X_r^{t, x, a}|^m \right] \leq C \left(1 + |X_s^{t, x, a}|^m + \int_s^T \mathbb{E}_s^t [|I_r^{t, a}|^{\frac{m}{1-p_{\varrho}}}] dr \right), \quad \text{if } p_{\varrho} < 1, \quad (2.1)$$

and

$$\begin{aligned} \mathbb{E}_s^t \left[\sup_{r \in [s, T]} |X_r^{t, x, a}|^m \right] &\leq C \left(1 + |X_s^{t, x, a}|^m + \sqrt{\int_s^T \mathbb{E}_s^t [|I_r^{t, a}|^{2m}] dr} \right) \\ &\quad \cdot \left(1 + \sqrt{\int_s^T \mathbb{E}_s^t [(1 + |I_r^{t, a}|^2) e^{\int_r^T 2|I_u^{t, a}| du}] dr} \right), \quad \text{if } p_{\varrho} = 1, \end{aligned} \quad (2.2)$$

for all $0 \leq t \leq s \leq T$.

Proof. For every $(t, a) \in [0, T] \times \mathbb{R}^d$, there exists a unique process $I^{t, a} = (I_s^{t, a})_{t \leq s \leq T}$ satisfying the second equation in (1.10). Since the coefficients $b(x) + \varrho(x)a$ and $\sigma(x)$ of the diffusion system (1.10) for (X, I) are locally Lipschitz in (x, a) , it is well-known (see e.g. Exercise IX.2.10 in [20]) that for every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ there exists an adapted

process $X^{t,x,a} = (X_s^{t,x,a})_{t \leq s \leq T}$ such that, if $e := \inf\{s \geq t: |X_s^{t,x,a}| = \infty\}$ with $\inf \emptyset = \infty$, then $X^{t,x,a}$ is the unique (up to indistinguishability) adapted and continuous process on $[t, e) \cap [t, T]$ that satisfies the first equation in (1.10) on $[t, e) \cap [t, T]$. It then remains to prove that the explosion time e of $X^{t,x,a}$ satisfies: $\mathbb{P}(e = \infty) = 1$.

For simplicity of notation, we denote $I = I^{t,a}$ and $X = X^{t,x,a}$. Consider $T_n = \inf\{s \geq t: |X_s| > n\} \wedge T$ and apply Itô's formula to $|X|^m$, for $m \geq 1$. From the dynamics (1.10) of (X, I) and Young's inequality, we see, under Assumption 1.1 (i) and (ii), that there exists a constant C (which in the sequel may change from line to line) depending only on $m, T, L_{b,\sigma,\varrho}$, and M_ϱ , such that for all $t \leq s \leq T_n$,

$$\begin{aligned} \sup_{r \in [s, T_n]} |X_r|^m &\leq C \left(1 + |X_s|^m + \int_s^{T_n} (|X_r|^m + |I_r|^m + |X_r|^{m-1+p_\varrho} |I_r|) dr \right. \\ &\quad \left. + \sup_{r \in [s, T_n]} \left| \int_s^r |X_u|^{m-1} \sigma(X_u) dW_u \right| \right). \end{aligned} \quad (2.3)$$

Case $p_\varrho < 1$. By applying Young's inequality to $|X_r|^{m-1+p_\varrho} |I_r|$, taking conditional expectation on both sides of (2.3), and using the Burkholder-Davis-Gundy inequality, we get

$$\begin{aligned} \mathbb{E}_s^t \left[\sup_{r \in [s, T_n]} |X_r|^m \right] &\leq C \left(1 + |X_s|^m + \int_s^T \mathbb{E}_s^t [|I_r|^{\frac{m}{1-p_\varrho}} 1_{[s, T_n]}(r)] dr \right. \\ &\quad \left. + \int_s^T \mathbb{E}_s^t \left[\sup_{u \in [s, r]} (|X_u|^m 1_{[s, T_n]}(u)) \right] dr \right), \end{aligned}$$

which shows, by Gronwall's lemma,

$$\begin{aligned} \mathbb{E}_s^t \left[\sup_{r \in [s, T_n]} |X_r|^m \right] &\leq C \left(1 + |X_s|^m + \int_s^T \mathbb{E}_s^t [|I_r|^{\frac{m}{1-p_\varrho}} 1_{[s, T_n]}(r)] dr \right) \\ &\leq C \left(1 + |X_s|^m + \int_s^T \mathbb{E}_s^t [|I_r|^{\frac{m}{1-p_\varrho}}] dr \right). \end{aligned}$$

Since $T_n \nearrow e \wedge T$ as n goes to infinity, from Fatou's lemma we obtain

$$\mathbb{E}_s^t \left[\sup_{r \in [s, e \wedge T]} |X_r|^m \right] \leq C \left(1 + |X_s|^m + \int_s^T \mathbb{E}_s^t [|I_r|^{\frac{m}{1-p_\varrho}}] dr \right). \quad (2.4)$$

In particular, taking $s = t$, we have $\mathbb{P}(e = \infty) = 1$. Then, from (2.4) we deduce the required estimate (2.1).

Case $p_\varrho = 1$. Take the conditional expectation in (2.3) with respect to the σ -algebra $\mathcal{G}_s^t := \mathcal{F}_s^t \vee \sigma(I_s, t \leq s \leq T)$, and observe that $(W_s - W_t)_{t \leq s \leq T}$ is a Brownian motion with respect to $(\mathcal{G}_s^t)_{t \leq s \leq T}$ since W and I are independent. Therefore, we can still use the Burkholder-Davis-Gundy inequality, and obtain (we denote by $\mathbb{E}_s^{t, \mathcal{G}}$ the conditional expectation under \mathbb{P} given \mathcal{G}_s^t)

$$\mathbb{E}_s^{t, \mathcal{G}} \left[\sup_{r \in [s, T_n]} |X_r|^m \right] \leq C \left(1 + |X_s|^m + \int_s^T |I_r|^m \mathbb{E}_s^{t, \mathcal{G}} [1_{[s, T_n]}(r)] dr \right)$$

$$+ \int_s^T \mathbb{E}_s^{t,\mathcal{G}} \left[\sup_{u \in [s,r]} (|X_u|^m \mathbf{1}_{[s,T_n]}(u)) \right] (1 + |I_r|) dr \Big),$$

which gives, by Gronwall's lemma and noting that $T_n \leq T$,

$$\mathbb{E}_s^{t,\mathcal{G}} \left[\sup_{r \in [s,T_n]} |X_r|^m \right] \leq C \left(1 + |X_s|^m + \int_s^T |I_r|^m dr \right) \left(1 + \int_s^T (1 + |I_r|) e^{\int_r^T |I_u| du} dr \right).$$

Afterwards, by taking conditional expectation with respect to \mathcal{F}_s^t , using the Cauchy-Schwarz inequality, and the fact that $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ for any $a, b \in \mathbb{R}$, we get

$$\begin{aligned} \mathbb{E}_s^t \left[\sup_{r \in [s,T_n]} |X_r|^m \right] &\leq C \left(1 + |X_s|^m + \sqrt{\int_s^T \mathbb{E}_s^t [|I_r|^{2m}] dr} \right) \\ &\quad \cdot \left(1 + \sqrt{\int_s^T \mathbb{E}_s^t [(1 + |I_r|^2) e^{\int_r^T 2|I_u| du}] dr} \right). \end{aligned}$$

Recalling that $T_n \nearrow e \wedge T$ as n goes to infinity, from Fatou's lemma we obtain

$$\begin{aligned} \mathbb{E}_s^t \left[\sup_{r \in [s,e \wedge T]} |X_r|^m \right] &\leq C \left(1 + |X_s|^m + \sqrt{\int_s^T \mathbb{E}_s^t [|I_r|^{2m}] dr} \right) \\ &\quad \cdot \left(1 + \sqrt{\int_s^T \mathbb{E}_s^t [(1 + |I_r|^2) e^{\int_r^T 2|I_u| du}] dr} \right), \end{aligned} \tag{2.5}$$

where the second conditional expectation on the right-hand side is finite, since I is a Brownian motion. Therefore, (2.5) with $s = t$ implies $\mathbb{P}(e = \infty) = 1$. Finally, from (2.5) we get the required estimate (2.2). \square

Next, it is straight forward to check that Assumption 1.1 translate to the following properties on the generator f of the BSDE (1.11).

Lemma 2.2 *The map $a \mapsto f(x, y, a)$ is convex and satisfies*

$$|f(x, a, y) - f(x, a, y')| \leq L_F |y - y'|, \tag{2.6}$$

$$-M_F \left(1 + |x|^{q_F} - \frac{1}{q' M_F^{q'}} |a|^{q'} \right) \leq f(x, a, 0) \leq m_F \left(1 + |x|^{p_F} + \frac{1}{p' m_F^{p'}} |a|^{p'} \right), \tag{2.7}$$

for all $x \in \mathbb{R}^d$, $y, y' \in \mathbb{R}$, and $a \in \mathbb{R}^d$. Here $p' = p/(p-1)$ and $q' = q/(q-1)$ are the conjugate exponents of, respectively, p and q in Assumption 1.1.

We may then define the notion of maximal solution to the BSDE with diffusion constraint (1.11)-(1.12).

Definition 2.1 *For every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, we say that $(Y^{t,x,a}, Z^{t,x,a}, V^{t,x,a}, K^{t,x,a}) \in \mathbb{S}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{K}^2(t, T)$ is a maximal solution to (1.11)-(1.12) if it satisfies (1.11)-(1.12), and for any other solution $(\underline{Y}^{t,x,a}, \underline{Z}^{t,x,a}, \underline{V}^{t,x,a}, \underline{K}^{t,x,a}) \in \mathbb{S}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{K}^2(t, T)$ we have $Y_s^{t,x,a} \geq \underline{Y}_s^{t,x,a}$, for any $s \in [t, T]$.*

Such a maximal solution is constructed using a penalization approach as in [12, Theorem 2.1]. However, rather than employing an independent Poisson random measure as in [12], our randomization here is carried out by means of an independent Brownian motion in (1.10). Let us consider, for every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ and $n \in \mathbb{N}$, the following *penalized BSDE*:

$$\begin{aligned} Y_s^n &= g(X_T^{t,x,a}) + \int_s^T f(X_r^{t,x,a}, I_r^{t,a}, Y_r^n) dr - (K_T^n - K_s^n) - \int_s^T Z_r^n dW_r - \int_s^T V_r^n dB_r \\ &= g(X_T^{t,x,a}) + \int_s^T f_n(X_r^{t,x,a}, I_r^{t,a}, Y_r^n, V_r^n) dr - \int_s^T Z_r^n dW_r - \int_s^T V_r^n dB_r, \end{aligned} \quad (2.8)$$

for all $s \in [t, T]$, where

$$K_s^n = n \int_t^s |V_s^n| ds$$

and the generator $f_n(x, a, y, v) = f(x, a, y) - n|v|$. Notice by (2.6) that this generator f_n is Lipschitz in (y, v) , so that from standard result due to [17], we know that for every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ and $n \in \mathbb{N}$, there exists a unique solution $(Y^{n,t,x,a}, Z^{n,t,x,a}, V^{n,t,x,a}) \in \mathbb{S}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{H}^2(t, T)$ to BSDE (2.8).

Moreover, we have the following comparison results.

Lemma 2.3 *For every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, the following statements hold:*

- (i) *The sequence $(Y^{n,t,x,a})_n$ is nonincreasing, i.e., $Y^{n,t,x,a} \geq Y^{n+1,t,x,a}$, $n \in \mathbb{N}$.*
- (ii) *For any solution $(\bar{Y}^{t,x,a}, \bar{Z}^{t,x,a}, \bar{V}^{t,x,a}, \bar{K}^{t,x,a}) \in \mathbb{S}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{K}^2(t, T)$ to (1.11)-(1.12), we have $Y^{n,t,x,a} \geq \bar{Y}^{t,x,a}$, $n \in \mathbb{N}$.*

Proof. Since $f_n \geq f_{n+1}$, the first statement follows from a direct application of the comparison theorem for BSDEs. For the second statement, note that $f_n(X_r^{t,x,a}, I_r^{t,a}, \bar{Y}_r^{t,x,a}, \bar{V}_r^{t,x,a}) = f(X_r^{t,x,a}, I_r^{t,a}, \bar{Y}_r^{t,x,a})$, due to $\bar{V}_r^{t,x,a} = 0$, and $\bar{K}^{t,x,a}$ is a nondecreasing process. Then the second statement readily follows from [19, Theorem 1.3]. \square

The aim is to obtain a uniform bound on $(\|Y^{n,t,x,a}\|_{\mathbb{S}^2(t,T)})_n$, which, together with the monotonicity property stated in Lemma 2.3(i), allows to construct $Y^{t,x,a}$ as the limit of $(Y^{n,t,x,a})_n$. In contrast to [12, Lemma 3.1], where the compactness of the space of control actions A (which does not hold in the present case, since $A = \mathbb{R}^d$) is exploited to prove the \mathbb{S}^2 -bound, we utilize a dual (or randomized) representation of $Y^{n,t,x,a}$. To this end, we introduce some additional notations. Let \mathcal{D}_n denote the set of \mathcal{B}_n -valued predictable processes ν , where \mathcal{B}_n is the closed ball in \mathbb{R}^d with radius n and centered at the origin, and define $\mathcal{D} = \cup_n \mathcal{D}_n$. For $t \in [0, T]$ and $\nu \in \mathcal{D}$, define the probability measure \mathbb{P}^ν on $(\Omega, \mathcal{F}_T^t)$ via

$$\frac{d\mathbb{P}^\nu}{d\mathbb{P}} = \mathcal{E} \left(\int_t^\cdot \nu_s dB_s \right)_T = \exp \left(\int_t^T \nu_s dB_s - \frac{1}{2} \int_t^T |\nu_s|^2 ds \right),$$

where the Doléans-Dades stochastic exponential on the right-hand side is a martingale due to the definition of \mathcal{D} . In the sequel, we denote by $\mathbb{E}_s^{t,\nu}$ the condition expectation under \mathbb{P}^ν given \mathcal{F}_s^t , for any $s \in [t, T]$, and $\nu \in \mathcal{D}$.

Remark 2.1 Notice by Girsanov's theorem that W remains a Brownian motion under \mathbb{P}^ν for any $\nu \in \mathcal{D}$. Then, by the same argument as in the derivation of (2.1) in Lemma 2.1, we see that when $p_\varrho < 1$, for every $m \geq 1$, there exists a positive constant C , depending only on $m, T, L_{b,\sigma,\varrho}, M_\varrho$, and p_ϱ such that

$$\mathbb{E}_s^{t,\nu} \left[\sup_{r \in [s,T]} |X_r^{t,x,a}|^m \right] \leq C \left(1 + |X_s^{t,x,a}|^m + \int_s^T \mathbb{E}_s^{t,\nu} [|I_r^{t,a}|^{\frac{m}{1-p_\varrho}}] dr \right), \quad (2.9)$$

for all $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, $s \in [t, T]$, and $\nu \in \mathcal{D}$. \square

The next result provides a dual representation of the solution to the penalized BSDE.

Proposition 2.1 For every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ and $n \in \mathbb{N}$, it holds that

$$Y_s^{n,t,x,a} = \operatorname{ess\,inf}_{\nu \in \mathcal{D}_n} \mathbb{E}_s^{t,\nu} \left[\int_s^T e^{\int_s^r \gamma_u^n du} f(X_r^{t,x,a}, I_r^{t,a}, 0) dr + e^{\int_s^T \gamma_u^n du} g(X_T^{t,x,a}) \right], \quad (2.10)$$

for all $t \leq s \leq T$, where the predictable process $\gamma^n: \Omega \times [t, T] \rightarrow \mathbb{R}$ is given by

$$\gamma^n = \frac{f(X^{t,x,a}, I^{t,a}, Y^{n,t,x,a}) - f(X^{t,x,a}, I^{t,a}, 0)}{Y^{n,t,x,a}} \mathbf{1}_{\{Y^{n,t,x,a} \neq 0\}}.$$

In particular, $|\gamma^n|$ is bounded uniformly by the constant L_F in (2.6).

Proof. Applying Itô's formula to $e^{\int_s^r \gamma_u^n du} Y^{n,t,x,a}$ and integrating between s and T , we obtain

$$\begin{aligned} Y_s^{n,t,x,a} &= e^{\int_s^T \gamma_u^n du} g(X_T^{t,x,a}) + \int_s^T e^{\int_s^r \gamma_u^n du} f(X_r^{t,x,a}, I_r^{t,a}, 0) dr - n \int_s^T e^{\int_s^r \gamma_u^n du} |V_r^{n,t,x,a}| dr \\ &\quad - \int_s^T e^{\int_s^r \gamma_u^n du} Z_r^{n,t,x,a} dW_r - \int_s^T e^{\int_s^r \gamma_u^n du} V_r^{n,t,x,a} dB_r. \end{aligned}$$

Take $\nu \in \mathcal{D}_n$, then add and subtract the term $\int_s^T e^{\int_s^r \gamma_u^n du} V_r^{n,t,x,a} \nu_r dr$, so that we obtain

$$\begin{aligned} Y_s^{n,t,x,a} &= e^{\int_s^T \gamma_u^n du} g(X_T^{t,x,a}) + \int_s^T e^{\int_s^r \gamma_u^n du} f(X_r^{t,x,a}, I_r^{t,a}, 0) dr \\ &\quad - \int_s^T e^{\int_s^r \gamma_u^n du} (n |V_r^{n,t,x,a}| + V_r^{n,t,x,a} \nu_r) dr \\ &\quad - \int_s^T e^{\int_s^r \gamma_u^n du} Z_r^{n,t,x,a} dW_r - \int_s^T e^{\int_s^r \gamma_u^n du} V_r^{n,t,x,a} dB_r^\nu, \end{aligned} \quad (2.11)$$

where $B^\nu = -\int_t^\cdot \nu_s ds + B$ and W are both \mathbb{P}^ν -Brownian motions by Girsanov's theorem. Note that both local martingales in (2.11) are in fact \mathbb{P}^ν -martingales. Indeed, from Bayes formula and the Cauchy-Schwarz inequality, we have

$$\mathbb{E}_t^{t,\nu} \left[\sqrt{\left\langle \int_t^\cdot Z_r dW_r \right\rangle_T} \right] = \mathbb{E} \left[L_T^\nu \sqrt{\left\langle \int_t^\cdot Z_r dW_r \right\rangle_T} \right] \leq \|L_T^\nu\|_{\mathbb{L}^2(\Omega, \mathcal{F}_T^t, \mathbb{P})} \|Z\|_{\mathbb{H}^2(t, T)},$$

where $L_T^\nu = d\mathbb{P}^\nu/d\mathbb{P}$ is bounded in $\mathbb{L}^2(\Omega, \mathcal{F}_T^t, \mathbb{P})$ since ν is bounded. Combining the previous estimate with the Burkholder-Davis-Gundy inequality, we obtain that $\int_t^\cdot Z_r dW_r$ is of class

D under \mathbb{P}^ν , hence is a \mathbb{P}^ν -martingale. The same argument can be applied to the other local martingale. Projecting both sides of (2.11) on \mathcal{F}_s^t , and using the inequality $n|v| + v \cdot a \geq 0$, for any $v \in \mathbb{R}$ and $a \in B_n$, we obtain

$$Y_s^{n,t,x,a} \leq \operatorname{ess\,inf}_{\nu \in \mathcal{D}_n} \mathbb{E}_s^{t,\nu} \left[\int_s^T e^{\int_s^r \gamma_u^n du} f(X_r^{t,x,a}, I_r^{t,a}, 0) dr + e^{\int_s^T \gamma_u^n du} g(X_T^{t,x,a}) \right].$$

To prove the reverse inequality, denote $V^{n,i}$ the i -th component of $V^{n,t,x,a}$, for $i = 1, \dots, d$. Set $\nu_r^{*,i} = -nV_r^{n,i}/|V_r^n|1_{\{|V_r^n| \neq 0\}}$, for $r \in [t, T]$ and $i = 1, \dots, d$. Then $\nu^* \in \mathcal{D}_n$, and $n|V_r^{n,t,x,a}| + V_r^{n,t,x,a} \cdot \nu_r^* = 0$ for any $r \in [t, T]$. It then follows from (2.11) that

$$Y_s^{n,t,x,a} = \mathbb{E}_s^{t,\nu^*} \left[\int_s^T e^{\int_s^r \gamma_u^n du} f(X_r^{t,x,a}, I_r^{t,a}, 0) dr + e^{\int_s^T \gamma_u^n du} g(X_T^{t,x,a}) \right],$$

from which the claim follows. \square

Combining the above dual representation together with the moment estimate for X in Remark 2.1, we obtain the following uniform estimate.

Corollary 2.1 *There exists a positive constant C , independent of n , such that*

$$\begin{aligned} -C(1 + |X_s^{t,x,a}|^{q_F \vee p_g}) &\leq Y_s^{n,t,x,a} \\ &\leq \begin{cases} C(1 + |X_s^{t,x,a}|^{p_F \vee q_g} + |I_s^{t,a}|^{\frac{p_F \vee q_g}{1-p_\varrho} \vee p'}), & \text{if } p_\varrho < 1, \\ C\left(1 + |X_s^{t,x,a}|^{p_F \vee q_g} + |I_s^{t,a}|^{p_F \vee q_g \vee p'}\right) \left(1 + |I_s^{t,a}|^2 + e^{C|I_s^{t,a}|}\right) & \text{if } p_\varrho = 1, \end{cases} \end{aligned} \quad (2.12)$$

for all $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, $s \in [t, T]$, and $n \in \mathbb{N}$. In particular, $Y^{n,t,x,a} \in \mathbb{S}^2(t, T)$ and

$$\|Y^{n,t,x,a}\|_{\mathbb{S}^2(t,T)} \leq \begin{cases} \tilde{C}(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{\frac{p_F \vee q_g}{1-p_\varrho} \vee p'}), & \text{if } p_\varrho < 1, \\ \tilde{C}(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{p_F \vee q_g \vee p'}) \\ \quad \cdot (1 + |a|^2 + e^{\tilde{C}|a|}), & \text{if } p_\varrho = 1, \end{cases} \quad (2.13)$$

for some positive constant \tilde{C} , independent of n .

Proof. The upper bound in (2.12). From the dual representation formula (2.10), we have (recall that, when $\nu \equiv 0$, \mathbb{P}^ν coincides with \mathbb{P})

$$Y_s^{n,t,x,a} \leq \mathbb{E}_s^t \left[\int_s^T e^{\int_s^r \gamma_r^n dr} f(X_r^{t,x,a}, I_r^{t,a}, 0) dr + e^{\int_s^T \gamma_r^n dr} g(X_T^{t,x,a}) \right].$$

Then, recalling the fact that $|\gamma^n|$ is bounded by L_F , exploiting the polynomial growth condition of f in (2.7) and of g in Assumption 1.1 (v), we get

$$Y_s^{n,t,x,a} \leq C \left(1 + \mathbb{E}_s^t \left[\sup_{r \in [s, T]} |X_r^{t,x,a}|^{p_F \vee q_g} + \sup_{r \in [s, T]} |I_r^{t,a}|^{p'} \right] \right). \quad (2.14)$$

Together with the estimate (2.1), and the standard estimate $\mathbb{E}_s^t[\sup_{r \in [s, T]} |I_r^{t,a}|^m] \leq C(1 + |I_s^{t,a}|^m)$ for any $m \geq 1$, this shows the upper bound in (2.12) when $p_\varrho < 1$.

When $p_\varrho = 1$, by using the estimate (2.2), together with the fact that $ab \leq (a^2 + b^2)/2$ for any $a, b \in \mathbb{R}$, we obtain from (2.14):

$$Y_s^{n,t,x,a} \leq C \left(1 + |X_s^{t,x,a}|^{p_F \vee q_g} + |I_s^{t,a}|^{p_F \vee q_g \vee p'} \right) \left(1 + |I_s^{t,a}|^2 + \sqrt{\mathbb{E}_s^t [e^{4 \int_s^T |I_r^{t,a}| dr}]} \right).$$

For the expectation on the right-hand side, we have

$$\begin{aligned} \mathbb{E}_s^t [e^{4 \int_s^T |I_r^{t,a}| dr}] &\leq e^{4(T-s)|I_s^{t,a}|} \mathbb{E}_s^t [e^{4 \int_s^T |B_r - B_s| dr}] = e^{4(T-s)|I_s^{t,a}|} \mathbb{E} [e^{4 \int_0^{T-s} |B_t| dt}] \\ &\leq e^{4T|I_s^{t,a}|} \mathbb{E} [e^{4T \sup_{0 \leq t \leq T} |B_t|}] \leq e^{4T|I_s^{t,a}|} \sum_{n=0}^{\infty} e^{4T(n+1)} \mathbb{P} \left(n \leq \sup_{0 \leq t \leq T} |B_t| < n+1 \right) \\ &\leq e^{4T|I_s^{t,a}|} \sum_{n=0}^{\infty} e^{4T(n+1)} \mathbb{P} \left(\sup_{0 \leq t \leq T} |B_t| \geq n \right) \leq e^{4T|I_s^{t,a}|} \sum_{n=0}^{\infty} e^{4T(n+1)} 2 \mathbb{P} \left(\sup_{0 \leq t \leq T} B_t \geq n \right) \\ &\leq e^{4T|I_s^{t,a}|} \sum_{n=0}^{\infty} e^{4T(n+1)} 2 e^{-\frac{n^2}{2T}} \leq C e^{4T|I_s^{t,a}|}, \end{aligned} \quad (2.15)$$

where we used the exponential inequality $\mathbb{P}(\sup_{0 \leq t \leq T} B_t \geq n) \leq \exp(-n^2/(2T))$, see Proposition II.1.8 in [20]. Combining the previous two estimates, we confirm the upper bound in (2.12) for $p_\varrho = 1$ case.

The lower bound in (2.12). From formula (2.10), the polynomial growth condition of f and g , we find that $Y_s^{n,t,x,a}$ is greater than or equal to the following quantity

$$\operatorname{ess\,inf}_{\nu \in \mathcal{D}_n} \mathbb{E}_s^{t,\nu} \left[-M_F \int_s^T e^{\int_t^s \gamma_r^n dr} (1 + |X_r^{t,x,a}|^{q_F} - \frac{1}{q' M_F^{q'}} |I_r^{t,a}|^{q'}) dr - m_g e^{\int_t^T \gamma_r^n dr} (1 + |X_T^{t,x,a}|^{p_g}) \right].$$

In the case where $p_\varrho = 1$, we have $q_F = p_g = 0$, and since $|\gamma^n|$ is bounded by L_F , we then obtain the lower bound: $Y_s^{n,t,x,a} \geq -2(M_F T + m_g) e^{L_F T}$. When $p_\varrho < 1$, we use the estimate (2.9) and the fact that $|\gamma^n|$ is bounded by L_F , to deduce that there exists a positive constant \bar{C} , depending only on $q_F, p_g, T, m_f, q', L_b, \sigma_\varrho, L_F, M_\varrho$ and p_ϱ , such that

$$\begin{aligned} Y_s^{n,t,x,a} &\geq -\bar{C} (1 + |X_s^{t,x,a}|^{q_F} + |X_s^{t,x,a}|^{p_g}) \\ &\quad + \operatorname{ess\,inf}_{\nu \in \mathcal{D}_n} \mathbb{E}_s^{t,\nu} \left[\int_s^T \left(\frac{M_F^{1-q'}}{q'} e^{-L_F T} |I_r^{t,a}|^{q'} - \bar{C} |I_r^{t,a}|^{\frac{q_F}{1-p_\varrho}} - \bar{C} |I_r^{t,a}|^{\frac{p_g}{1-p_\varrho}} \right) dr \right]. \end{aligned}$$

Recalling that $q_F, p_g < (1 - p_\varrho)q'$, we see that the function $h(x) = \frac{M_F^{1-q'}}{q'} e^{-L_F T} |x|^{q'} - \bar{C} |x|^{\frac{q_F}{1-p_\varrho}} - \bar{C} |x|^{\frac{p_g}{1-p_\varrho}}$, $x \in \mathbb{R}^d$, is bounded from below on \mathbb{R}^d by a constant independent of n . Therefore, we obtain the lower bound.

Estimate (2.13). When $p_\varrho < 1$, it follows directly from bounds (2.12), Lemma 2.1 with $s = t$, the standard estimate $\mathbb{E}[\sup_{s \in [t, T]} |I_s^{t,a}|^m] \leq C(1 + |a|^m)$ for any $m \geq 1$. When $p_\varrho = 1$, (2.13) follows from the Cauchy-Schwarz inequality together with $\mathbb{E}[e^{C|I_s^{t,a}|}] \leq C e^{C|a|}$ (this latter inequality follows from $\mathbb{E}[e^{C|I_s^{t,a}|}] \leq e^{C|a|} \mathbb{E}[e^{C|B_s - B_t|}] = e^{C|a|} \mathbb{E}[e^{C|B_s - t|}] \leq e^{C|a|} \mathbb{E}[e^{C \sup_{0 \leq t \leq T} |B_t|}]$, afterwards we reason as in (2.15)). \square

The previous uniform norm estimate implies the following uniform norm estimate on $(Z^{n,t,x,a}, V^{n,t,x,a}, K^{n,t,x,a})_n$.

Corollary 2.2 *There exists a positive constant C , independent of n , such that*

$$\begin{aligned} & \|Z^{n,t,x,a}\|_{\mathbb{H}^2(t,T)}^2 + \|V^{n,t,x,a}\|_{\mathbb{H}^2(t,T)}^2 + \|K^{n,t,x,a}\|_{\mathbb{S}^2(t,T)}^2 \\ & \leq \begin{cases} C(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{\frac{p_F \vee q_g}{1-p_\varrho} \vee p'}), & \text{if } p_\varrho < 1 \\ C(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{p_F \vee q_g \vee p'}) \\ \quad \cdot (1 + |a|^2 + e^{C|a|}), & \text{if } p_\varrho = 1, \end{cases} \end{aligned} \quad (2.16)$$

for all $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, $n \in \mathbb{N}$.

Proof. By proceeding along the same arguments as in the proof of [12, Lemma 2.3], we have:

$$\begin{aligned} & \|Z^{n,t,x,a}\|_{\mathbb{H}^2(t,T)}^2 + \|V^{n,t,x,a}\|_{\mathbb{H}^2(t,T)}^2 + \|K^{n,t,x,a}\|_{\mathbb{S}^2(t,T)}^2 \\ & \leq C \left(\mathbb{E} \left[\int_t^T |f(X_s^{t,x,a}, I_s^{t,a}, 0)|^2 ds \right] + \sup_{n \in \mathbb{N}} \|Y^{n,t,x,a}\|_{\mathbb{S}^2(t,T)}^2 \right), \end{aligned}$$

Then, by exploiting the polynomial growth condition of f in (2.7) combined again with Lemma 2.1 for $s = t$, the standard estimate $\mathbb{E}[\sup_{r \in [t, T]} |I_r^{t,a}|^m] \leq C(1 + |a|^m)$ for any $m \geq 1$, and $\mathbb{E}[e^{C|I_s^{t,a}|}] \leq C e^{C|a|}$ when $p_\varrho = 1$, and using the estimate (2.13) for $Y^{n,t,x,a}$, we obtain the required uniform estimate. \square

Now the previous uniform norm estimates allow us to take limit as $n \rightarrow \infty$.

Theorem 2.1 *Let Assumption 1.1 hold. For every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, there exists a unique maximal solution $(Y^{t,x,a}, Z^{t,x,a}, V^{t,x,a}, K^{t,x,a}) \in \mathbb{S}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{K}^2(t, T)$ to (1.11)-(1.12), such that, as $n \rightarrow \infty$, $Y_s^{n,t,x,a} \searrow Y_s^{t,x,a}$; $K_s^{n,t,x,a}$ weakly converges to $K_s^{t,x,a}$ in $L^2(\Omega, \mathcal{F}_s, \mathbb{P})$, for all $s \in [t, T]$; and $(Z^{n,t,x,a}, V^{n,t,x,a})_n$ weakly converges to $(Z^{t,x,a}, V^{t,x,a})$ in $\mathbb{H}^2(t, T) \times \mathbb{H}^2(t, T)$. Moreover*

$$\|Y^{t,x,a}\|_{\mathbb{S}^2(t,T)} \leq \begin{cases} \tilde{C}(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{\frac{p_F \vee q_g}{1-p_\varrho} \vee p'}), & \text{if } p_\varrho < 1 \\ \tilde{C}(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{p_F \vee q_g \vee p'}) \\ \quad \cdot (1 + |a|^2 + e^{\tilde{C}|a|}), & \text{if } p_\varrho = 1 \end{cases} \quad (2.17)$$

where \tilde{C} is the same constant as in (2.13).

Proof. The proof follows the same passages of the proof for [12, Theorem 2.1], with some small modifications due to the fact that here we adopted a Brownian (rather than Poisson) randomization. For this reason, we just outline main steps of the proof. For every $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, from Lemma 2.3(i) and estimate (2.13), it follows that the sequence $(Y^{n,t,x,a})_n$ converges decreasingly to some \mathbb{F}^t -adapted process $Y^{t,x,a}$ satisfying the bound (2.17). Next, the monotonic limit theorem (see [19, Theorem 2.4]) implies the weak convergence stated in the theorem and that the limit $(Y^{t,x,a}, Z^{t,x,a}, V^{t,x,a}, K^{t,x,a}) \in \mathbb{S}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{H}^2(t, T) \times \mathbb{K}^2(t, T)$ satisfies BSDE (1.11). In order to prove the constraint (1.12), define the functional $F(V) = \mathbb{E} \int_t^T |V_s| ds$, for $V \in \mathbb{H}^2(t, T)$. Notice that $\mathbb{E} K_T^{n,t,x,a} = nF(V^{n,t,x,a})$. Hence estimate (2.16) implies $\sup_n \mathbb{E} |K_T^{n,t,x,a}|^2 < \infty$, therefore

$\lim_{n \rightarrow \infty} F(V^{n,t,x,a}) = 0$. Since F is lower semicontinuous with respect to the weak topology of $\mathbb{H}^2(t, T)$ (F is convex and strongly continuous), it then implies $F(V^{t,x,a}) = 0$ and (1.12) holds. Finally, the maximality of $(Y^{t,x,a}, Z^{t,x,a}, V^{t,x,a}, K^{t,x,a})$ follows from a direct application of Lemma 2.3(ii) and the uniqueness follows from [19, Proposition 1.6]. \square

3 Feynman-Kac representation formula

The present section is devoted to the proof of Theorem 1.1. Assumption 1.1 is enforced throughout this section. Let us first recall the definition of (discontinuous) viscosity solution to equation (1.6) (or, equivalently, to (1.1)). Define the *lower semicontinuous (lsc) envelope* u_* and *upper semicontinuous (usc) envelope* u^* of a locally bounded function $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ as follows:

$$u_*(t, x) = \liminf_{\substack{(t', x') \rightarrow (t, x) \\ t' < T}} u(t', x') \quad \text{and} \quad u^*(t, x) = \limsup_{\substack{(t', x') \rightarrow (t, x) \\ t' < T}} u(t', x')$$

for all $(t, x) \in [0, T] \times \mathbb{R}^d$.

Definition 3.1

- (i) We say that an usc (resp. lsc) function $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a viscosity subsolution (resp. supersolution) to (1.6) if

$$u(T, x) \leq (\text{resp. } \geq) g(x), \quad \forall x \in \mathbb{R}^d$$

and

$$-\frac{\partial \varphi}{\partial t}(t, x) - \inf_{a \in \mathbb{R}^d} [\mathcal{L}^a \varphi(t, x) + f(x, a, u(t, x))] \leq (\text{resp. } \geq) 0$$

for every $(t, x) \in [0, T] \times \mathbb{R}^d$ and every $\varphi \in C^{1,2}([0, T] \times \mathbb{R}^d)$ satisfying

$$(u - \varphi)(t, x) = \max_{[0, T] \times \mathbb{R}^d} (u - \varphi) \quad (\text{resp. } \min_{[0, T] \times \mathbb{R}^d} (u - \varphi)).$$

- (ii) We say that a locally bounded function $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a (discontinuous) viscosity solution to (1.6) whenever u^* is a viscosity subsolution and u_* is a viscosity supersolution to (1.6).

3.1 Penalized BSDE and corresponding semilinear parabolic PDE

Let us define, for every $n \in \mathbb{N}$, functions $v_n, v: [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ via

$$v_n(t, x, a) = Y_t^{n,t,x,a} \quad \text{and} \quad v(t, x, a) = Y_t^{t,x,a}, \quad \text{for } (t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d.$$

Notice that $v_n \searrow v$ pointwise as n goes to infinity. Moreover, it follows from (2.13) and (2.17) that

$$|v_n(t, x, a)| + |v(t, x, a)| \leq \begin{cases} \tilde{C}(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{\frac{p_F \vee q_g}{1-p_\rho} \vee p'}), & \text{if } p_\rho < 1 \\ \tilde{C}(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |a|^{p_F \vee q_g \vee p'}) \\ \cdot (1 + |a|^2 + e^{\tilde{C}|a|}), & \text{if } p_\rho = 1 \end{cases} \quad (3.1)$$

for all $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ and $n \in \mathbb{N}$. For each $n \in \mathbb{N}$, let us consider the following semilinear parabolic PDE

$$\begin{cases} -\frac{\partial v_n}{\partial t}(t, x, a) - \frac{1}{2}\Delta_a v_n(t, x, a) - \mathcal{L}^a v_n(t, x, a) \\ -f(x, a, v_n(t, x, a)) + n|D_a v_n(t, x, a)| = 0, & (t, x, a) \in [0, T) \times \mathbb{R}^d \times \mathbb{R}^d, \\ v_n(T, x, a) = g(x), & (x, a) \in \mathbb{R}^d \times \mathbb{R}^d, \end{cases} \quad (3.2)$$

where Δ_a is the Laplace operator with respect to a and \mathcal{L}^a is given in (1.7). Then, we have the following result.

Proposition 3.1 *The function v_n is a continuous viscosity solution to (3.2), i.e., $v_n = (v_n)_* = (v_n)^*$ and properties in Definition 3.1 are satisfied where the equation is replaced by (3.2).*

Proof. When $p_\varrho < 1$, so that v_n satisfies a polynomial growth condition, the result is well-known, see e.g. [18, Theorem 4.3]. When $p_\varrho = 1$, a change of variable is needed in order to deal with the exponential growth of v_n in a . More precisely, define the map $\beta: \mathbb{R} \rightarrow \mathbb{R}$ as follows

$$\beta(x) = \begin{cases} e^x - 1, & x \geq \log 2, \\ P(x), & -\log 2 \leq x < \log 2, \\ 1 - e^{-x}, & x < -\log 2, \end{cases}$$

where $P(x) = Ax^5 + Bx^3 + Cx$ (for some real constants A, B, C) is a polynomial of fifth degree which realizes a smooth paste of the two other branches of the map β , so that $\beta \in C^2(\mathbb{R})$ (since P is an odd function, in order to determine A, B, C it is enough to realize a C^2 -paste with the upper branch of β) and β is increasing, hence invertible. We denote by $\alpha: \mathbb{R} \rightarrow \mathbb{R}$ the inverse map of β , which is given by

$$\alpha(y) = \begin{cases} \log(1 + y), & y \geq 1, \\ P^{-1}(y), & -1 \leq y < 1, \\ -\log(1 - y), & y < -1. \end{cases} \quad (3.3)$$

For $a = (a_1, \dots, a_d), b = (b_1, \dots, b_d) \in \mathbb{R}^d$, we denote $\alpha(b) = (\alpha(b_1), \dots, \alpha(b_d)) \in \mathbb{R}^d$, $\beta(a) = (\beta(a_1), \dots, \beta(a_d)) \in \mathbb{R}^d$, and define

$$w_n(t, x, b) = v_n(t, x, \alpha(b)).$$

From (3.1), with $p_\varrho = 1$, we obtain that there exists a positive constant c such that

$$|w_n(t, x, b)| \leq c(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g} + |b|^{p_F \vee q_g \vee p'}) (1 + |b|^{2\vee \tilde{C}}), \quad (3.4)$$

for all $(t, x, b) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ and $n \in \mathbb{N}$. Notice that

$$D_a v_n(t, x, a) = \left(\frac{\partial w_n(t, x, \beta(a))}{\partial b_1} \beta'(a_1), \dots, \frac{\partial w_n(t, x, \beta(a))}{\partial b_d} \beta'(a_d) \right),$$

$$\Delta_a v_n(t, x, a) = \sum_{i=1}^d \frac{\partial^2 w_n(t, x, \beta(a))}{\partial b_i^2} (\beta'(a_i))^2 + \sum_{i=1}^d \frac{\partial w_n(t, x, \beta(a))}{\partial b_i} \beta''(a_i).$$

Then, v_n is a continuous viscosity solution to (3.2) if and only if w_n is a continuous viscosity solution to the following equation:

$$\begin{cases} -\frac{\partial w_n}{\partial t}(t, x, b) - \frac{1}{2} \Delta_b^\beta w_n(t, x, b) - \mathcal{L}^{\alpha(b)} w_n(t, x, b) \\ -f(x, \alpha(b), w_n(t, x, b)) + n |D_b^\beta w_n(t, x, b)| = 0, & (t, x, b) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d, \\ w_n(T, x, b) = g(x), & (x, b) \in \mathbb{R}^d \times \mathbb{R}^d, \end{cases} \quad (3.5)$$

where

$$\begin{aligned} D_b^\beta w_n(t, x, b) &= \left(\frac{\partial w_n(t, x, b)}{\partial b_1} \beta'(\alpha(b_1)), \dots, \frac{\partial w_n(t, x, b)}{\partial b_d} \beta'(\alpha(b_d)) \right), \\ \Delta_b^\beta w_n(t, x, b) &= \sum_{i=1}^d \frac{\partial^2 w_n(t, x, b)}{\partial b_i^2} (\beta'(\alpha(b_i)))^2 + \sum_{i=1}^d \frac{\partial w_n(t, x, b)}{\partial b_i} \beta''(\alpha(b_i)). \end{aligned}$$

Proceeding as in [18, Theorem 4.3] we can prove that w_n is a continuous viscosity solution to the previous equation, so the claim follows. \square

3.2 The invariance of the function v with respect to the variable a

We distinguish between the two cases $p_\varrho < 1$ and $p_\varrho = 1$. Let us consider the two following first-order PDEs:

$$|D_a v(t, x, a)| = 0, \quad (t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d, \quad (3.6)$$

$$|D_b w(t, x, b)| = 0, \quad (t, x, b) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d. \quad (3.7)$$

Here $w(t, x, b) = v(t, x, \alpha(b))$, where $\alpha(b)$ comes from (3.3).

Lemma 3.1 *When $p_\varrho < 1$ (resp. $p_\varrho = 1$) the function v (resp. w) is a (discontinuous) viscosity solution to (3.6) (resp. (3.7)).*

Proof. We firstly suppose that $p_\varrho < 1$, so that v_n and v satisfy a polynomial growth condition as reported in (3.1). Since the supersolution property clearly holds, let us focus on the subsolution property. To this end, since v is a pointwise limit of the decreasing sequence $(v_n)_n$, it is upper semi-continuous and $v = v^*$. Take $(t, x, a) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ and $\varphi \in C^{1,2}([0, T] \times \mathbb{R}^d \times \mathbb{R}^d)$ satisfying $(v - \varphi)(t, x, a) = \max_{[0, T] \times \mathbb{R}^d \times \mathbb{R}^d} (v - \varphi)$. From the polynomial growth property (3.1), we can assume without loss of generality (up to a polynomial perturbation of φ for large values of x and a) that the previous maximum is strict. Moreover, using standard techniques in viscosity solutions, we can deduce the existence of a bounded sequence $(t_n, x_n, a_n)_n \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ such that

$$\begin{aligned} (v_n - \varphi)(t_n, x_n, a_n) &= \max_{[0, T] \times \mathbb{R}^d \times \mathbb{R}^d} (v_n - \varphi) \quad \text{and} \\ (t_n, x_n, a_n, v_n(t_n, x_n, a_n)) &\longrightarrow (t, x, a, v(t, x, a)) \quad \text{as } n \rightarrow \infty. \end{aligned}$$

The viscosity subsolution property of v_n then implies that

$$|D_a \varphi(t_n, x_n, a_n)| \leq \frac{1}{n} \left(\frac{\partial \varphi}{\partial t}(t_n, x_n, a_n) + \frac{1}{2} \Delta_a \varphi(t_n, x_n, a_n) + \mathcal{L}^{a_n} \varphi(t_n, x_n, a_n) + f(x_n, a_n, v_n(t_n, x_n, a_n)) \right).$$

Sending n to infinity, and using the continuity of the coefficients b , σ , and f , we deduce the claim.

Suppose now that $p_\varrho = 1$. Reasoning as in the case $p_\varrho < 1$ and using the fact that $\beta' > 0$, we can prove that, for every $i = 1, \dots, d$, w is a viscosity solution to the first-order PDE:

$$\frac{\partial w(t, x, b)}{\partial b_i} = 0, \quad (t, x, b) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d.$$

As a consequence, we deduce that w is a viscosity solution to (3.7). \square

We can now state the following result on the independence of v with respect to a .

Proposition 3.2 *The function v satisfies*

$$v(t, x, a) = v(t, x, a'), \quad \text{for all } (t, x) \in [0, T] \times \mathbb{R}^d \text{ and } a, a' \in \mathbb{R}^d.$$

Proof. Let us firstly consider the case $p_\varrho < 1$. Then, this result can be proved proceeding as in [12, Lemma 3.4 and Proposition 3.2], with only two small modifications. First, in [12] the equation $-|D_a v(t, x, a)| = 0$ is considered rather than (3.6) (this is due to the fact that the function v therein is the *increasing* limit of $(v_n)_n$, due to the fact that a HJB equation with a “sup” operator, instead of “inf”, is studied there). However, the proofs of [12, Lemma 3.4 and Proposition 3.2] are not affected (apart from obvious modifications by the presence of the minus sign). Second, in [12] the variable a belongs to an open, bounded, and connected subset \mathring{A} of some Euclidean space \mathbb{R}^q , rather than to the entire space \mathbb{R}^d as here. To link to [12], let us consider the open ball $\mathring{\mathcal{B}}_r \subset \mathbb{R}^d$ for any $r > 0$. Lemma 3.1 implies that v is a viscosity solution to equation (3.6) on $[0, T] \times \mathbb{R}^d \times \mathring{\mathcal{B}}_r$, for every $r > 0$. At this point we can apply [12, Lemma 3.4 and Proposition 3.2] to deduce that v satisfies

$$v(t, x, a) = v(t, x, a'), \quad \text{for all } (t, x) \in [0, T] \times \mathbb{R}^d \text{ and } a, a' \in \mathcal{B}_r.$$

As a result, the claim follows since r is arbitrarily chosen.

Suppose now that $p_\varrho = 1$. Then, proceeding along the same lines as in the case $p_\varrho < 1$, we conclude that w does not depend on b . Since $v(t, x, a) = w(t, x, \beta(a))$, this implies that v does not depend on a . \square

3.3 From the BSDE with diffusion constraint to the viscous HJ equation

Following Proposition 3.2, we define the function u , as in (1.13), which satisfies (due to (3.1) with $a = 0$)

$$|u(t, x)| \leq \tilde{C}(1 + |x|^{p_F \vee q_F \vee p_g \vee q_g}), \quad \text{for all } (t, x) \in [0, T] \times \mathbb{R}^d.$$

Let us now prove that u is a viscosity solution to equation (1.6) (equivalently, to equation (1.1)), which, together with Proposition 3.2, concludes the proof of Theorem 1.1.

Proof of Theorem 1.1. In order to prove that u is a viscosity solution to equation (1.6), we begin by noticing that, as a direct consequence of Proposition 3.1, v_n is a viscosity solution to (3.2) on $[0, T] \times \mathbb{R}^d \times \mathcal{B}_r$, for any $r > 0$. Therefore, as a decreasing limit of $(v_n)_n$, v , hence u , is a viscosity solution to the following equation:

$$\begin{cases} -\frac{\partial u}{\partial t}(t, x) - \inf_{a \in \mathcal{B}_r} [\mathcal{L}^a u(t, x) + f(x, a, u(t, x))] = 0, & (t, x) \in [0, T] \times \mathbb{R}^d, \\ u(T, x) = g(x), & x \in \mathbb{R}^d. \end{cases} \quad (3.8)$$

The proof of this result can be done proceeding as in [12, Section 3.4], with some small modifications, due to the Poisson, rather than Brownian, randomization used in [12], which affects in part the form of the penalized PDE (3.2). From the definition of viscosity solution to (3.8) we have, for any $(t, x) \in [0, T] \times \mathbb{R}^d$, $r > 0$, and any $\varphi \in C^{1,2}([0, T] \times \mathbb{R}^d)$,

$$-\frac{\partial \varphi}{\partial t}(t, x) - \inf_{a \in \mathcal{B}_r} [\mathcal{L}^a \varphi(t, x) + f(x, a, u(t, x))] \leq (\text{resp. } \geq) 0$$

whenever $(u - \varphi)(t, x) = \max_{[0, T] \times \mathbb{R}^d}(u - \varphi)$ (resp. $\min_{[0, T] \times \mathbb{R}^d}(u - \varphi)$). Noticing that the choice of the test function φ is independent of r , we deduce

$$-\frac{\partial \varphi}{\partial t}(t, x) - \inf_{a \in \mathbb{R}^d} [\mathcal{L}^a \varphi(t, x) + f(x, a, u(t, x))] \leq (\text{resp. } \geq) 0,$$

which implies that u is a viscosity solution to equation (1.6). Finally, the terminal condition is also satisfied thanks to the terminal condition in (3.2) and the fact that v is a monotone limit of $(v_n)_n$. \square

References

- [1] K. Bahlali, M. Eddahbi, and Y. Ouknine. Quadratic BSDEs with \mathbb{L}^2 -terminal data: Existence results, Krylov's estimate and Itô-Krylov's formula. Preprint arXiv:1402.6596, 2014.
- [2] P. Barrieu and N. El Karoui. Monotone stability of quadratic semimartingales with applications to unbounded general quadratic BSDEs. *Ann. Probab.*, 41(3B):1831–1863, 2013.
- [3] M. Ben-Artzi, P. Souplet, and F. B. Weissler. The local theory for viscous Hamilton-Jacobi equations in Lebesgue spaces. *J. Math. Pures. Appl.*, 81:343–378, 2002.
- [4] P. Briand and Y. Hu. BSDE with quadratic growth and unbounded terminal value. *Probab. Theory Related Fields*, 136(4):604–618, 2006.
- [5] P. Briand and Y. Hu. Quadratic BSDEs with convex generators and unbounded terminal conditions. *Probab. Theory Related Fields*, 141(3-4):543–567, 2008.
- [6] P. Cheridito and K. Nam. BSDEs with terminal conditions that have bounded Malliavin derivative. *J. Funct. Anal.*, 266(3):1257–1285, 2014.
- [7] F. Da Lio and O. Ley. Uniqueness results for convex Hamilton-Jacobi equations under $p > 1$ growth conditions on data. *Appl. Math. Optim.*, 63(3):309–339, 2011.

- [8] F. Delbaen, Y. Hu, and X. Bao. Backward SDEs with superquadratic growth. *Probab. Theory Related Fields*, 150(1-2):145–192, 2011.
- [9] F. Delbaen, Y. Hu, and A. Richou. On the uniqueness of solutions to quadratic BSDEs with convex generators and unbounded terminal conditions. *Ann. Inst. Henri Poincaré Probab. Stat.*, 47(2):559–574, 2011.
- [10] B. H. Gilding, M. Guedda, and R. Kersner. The Cauchy problem for $u_t = \Delta u + |\nabla u|^q$. *J. Math. Anal. Appl.*, 284(2):733–755, 2003.
- [11] A. Gladkov, M. Guedda, and R. Kersner. A KPZ growth model with possibly unbounded data: correctness and blow-up. *Nonlinear Anal.*, 68(7):2079–2091, 2008.
- [12] I. Kharroubi and H. Pham. Feynman-Kac representation for Hamilton-Jacobi-Bellman IPDE. Preprint arXiv:1212.2000v2, to appear on *The Annals of Probability*, 2015.
- [13] M. Kobylanski. Backward stochastic differential equations and partial differential equations with quadratic growth. *Ann. Probab.*, 28(2):558–602, 2000.
- [14] F. Masiero and A. Richou. A note on the existence of solutions to Markovian superquadratic BSDEs with an unbounded terminal condition. *Electron. J. Probab.*, 18:no. 50, 15, 2013.
- [15] M. Musiela and T. Zariphopoulou. An example of indifference prices under exponential preferences. *Finance Stoch.*, 8(229-239), 2004.
- [16] H. Nagai. Optimal strategies for risk-sensitive portfolio optimization problems for general factor models. *SIAM J. Control. Optim.*, 41(6):1779–1800, 2006.
- [17] É. Pardoux and S. Peng. Adapted solution of a backward stochastic differential equation. *Systems Control Lett.*, 14(1):55–61, 1990.
- [18] É. Pardoux and S. Peng. Backward stochastic differential equations and quasilinear parabolic partial differential equations. In *Stochastic partial differential equations and their applications (Charlotte, NC, 1991)*, volume 176 of *Lecture Notes in Control and Inform. Sci.*, pages 200–217. Springer, Berlin, 1992.
- [19] S. Peng. Monotonic limit theorem of BSDE and nonlinear decomposition theorem of Doob-Meyer’s type. *Probab. Theory Related Fields*, 113(4):473–499, 1999.
- [20] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 1999.
- [21] A. Richou. Markovian quadratic and superquadratic BSDEs with an unbounded terminal condition. *Stochastic Process. Appl.*, 122(9):3173–3208, 2012.