



HAL
open science

Coupling local and non-local damage evolutions with thick level set model.

Nicolas Moes, Claude Stolz, Nicolas Chevaugnon

► **To cite this version:**

Nicolas Moes, Claude Stolz, Nicolas Chevaugnon. Coupling local and non-local damage evolutions with thick level set model.. IACM-2014, Jul 2014, Barcelone, Spain. hal-01154938

HAL Id: hal-01154938

<https://hal.science/hal-01154938>

Submitted on 17 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Coupling local and non-local damage evolutions with the Thick Level Set Model

Moës, N.¹, Stolz, C.^{1,2} and Chevaugeon, N.¹

¹ Ecole Centrale de Nantes,
GeM, UMR CNRS 6183 1 Rue de la Noe, 44321 Nantes, France,
{nicolas.moes,claud.stolz,nicolas.chevaugeon}@ec-nantes.fr

² LAMSID, EdF-CEA-CNRS UMR 2832, 1 Avenue du Général de Gaulle, 92141 Clamart, France

Key Words: , *Damage, Fracture, non-local, Thick Level Set*

It is well known that damage evolution models requires a length to avoid spurious localization. We gather these type of models under the name “non-local damage models”. In fact, non-locality is strictly needed only when the stress-strain curve starts to have a negative slope. Before this point, local models may be used. For instance, when damage exhibits some hardening, the stress-strain curve may be growing for quite some time before a peak point is reached.

Most if not all current non-local models are not able to distinguish between local and non-local zones, thus enforcing non-locality at every point and every instant. This leads to important computing times.

The Thick Level Set (TLS) model is a new non-local damage model [1,2,3] that allows to combine nicely zones in the domain where local model is used and zones where non-local model is used. This allows to reduce dramatically the computing time. Moreover, the TLS model is able to locate automatically zones fully damaged across displacement are allowed to be discontinuous.

The theoretical basis of the TLS model will be described. In short, it is a model in which the damage gradient is bounded (and not the Laplacian as in other modelling). The equation at stake is of Eikonal type (this explains the words level set is the name of the model). A 1D axisymmetric case will be fully developed both analytically and numerically to show the concurrent development of local and non-local damage zones.

REFERENCES

- [1] N. Moës, C. Stolz, P.-E. Bernard, N. Chevaugeon, A level set based model for damage growth : the thick level set approach. *International Journal For Numerical Methods in Engineering*, **86**, 358–380, 2011.
- [2] C. Stolz, N. Moës, A new model of damage : a moving thick layer approach. *International Journal of Fracture*, **174**, 49–60, 2012.

-
- [3] P.-E. Bernard, N. Moës, N. Chevaugeon, Damage growth modeling using the Thick Level Set (TLS) approach: Efficient discretization for quasi-static loadings. *Computer Methods in Applied Mechanics and Engineering*, **233-236**, 11–27, 2012.
- [4] Pradeilles-Duval, R. M., & Stolz, C. (1995). Mechanical transformations and discontinuities along a moving surface. *Journal of the Mechanics and Physics of Solids*, *43*(1), 91–121. Retrieved from <http://linkinghub.elsevier.com/retrieve/pii/0022509694000619>