



HAL
open science

DISTRIBUTION OF POSTCRITICALLY FINITE POLYNOMIALS II: SPEED OF CONVERGENCE

Thomas Gauthier, Gabriel Vigny

► **To cite this version:**

Thomas Gauthier, Gabriel Vigny. DISTRIBUTION OF POSTCRITICALLY FINITE POLYNOMIALS II: SPEED OF CONVERGENCE. 2015. hal-01155711v1

HAL Id: hal-01155711

<https://hal.science/hal-01155711v1>

Preprint submitted on 27 May 2015 (v1), last revised 1 Dec 2016 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DISTRIBUTION OF POSTCRITICALLY FINITE POLYNOMIALS II: SPEED OF CONVERGENCE

by

Thomas Gauthier & Gabriel Vigny

Abstract. — In the moduli space of degree d polynomials, we prove the equidistribution of postcritically finite polynomials toward the bifurcation measure. More precisely, using complex analytic arguments and pluripotential theory, we prove the exponential speed of convergence for \mathcal{C}^2 -observables. This improves results obtained with arithmetic methods by Favre and Rivera-Letellier in the unicritical family and Favre and the first author in the space of degree d polynomials.

We deduce from that the equidistribution of hyperbolic parameters with $(d - 1)$ distinct attracting cycles of given multipliers toward the bifurcation measure with exponential speed for \mathcal{C}^1 -observables. As an application, we prove the equidistribution (up to an explicit extraction) of parameters with $(d - 1)$ distinct cycles with prescribed multiplier toward the bifurcation measure for any $(d - 1)$ multipliers outside a pluripolar set.

Contents

1. Introduction.....	1
2. Background material.....	6
3. Preliminary results.....	8
4. Speed of convergence in the unicritical family.....	11
5. Initial estimates in the moduli space of polynomials.....	16
6. Distribution of postcritically finite hyperbolic polynomials.....	22
7. Distribution of polynomials with prescribed multipliers.....	27
8. Distribution of parametric preimages.....	32
References.....	34

1. Introduction

In a holomorphic family $(f_\lambda)_{\lambda \in \Lambda}$ of degree $d \geq 2$ rational maps, Mañé, Sad and Sullivan [MSS] studied quite precisely the notion of structural stability. In particular, they related the instability of critical orbits under small perturbations of the dynamics and instability of periodic cycles (see also [Ly]).

This description has been enriched by DeMarco [DeM1] who introduced a current T_{bif} which is supported exactly on the bifurcation locus. This current and its self-intersections reveal to be an appropriate tool to study bifurcations from a measurable viewpoint. Now, consider the particular case of the moduli space \mathcal{P}_d of critically marked complex polynomials of degree d modulo conjugacy by affine transformations. In that space, the maximal self-intersection of the bifurcation current induces a *bifurcation measure* μ_{bif} , introduced by Bassanelli and Berteloot [BB1], which may be considered as the analogue of the harmonic measure of the Mandelbrot set when $d \geq 3$. The support of this measure is where maximal bifurcation phenomena occur. Recall that a polynomial is *postcritically finite* if

all its critical points have finite orbit. Using the bifurcation measure, it is proved in [BB1] that its support is accumulated by postcritically finite hyperbolic parameters (which are in a certain way the most stable parameters) and that it coincides with the closure of parameters having a maximal number of neutral cycles. Still using the bifurcation measure, the first author also showed in [G1] that its support has maximal Hausdorff dimension.

As the bifurcation locus of the moduli space \mathcal{P}_d is a complicated fractal set, a natural approach is to study it on some dynamical slices. In particular, one can study the maps having a superattracting orbit of fixed period n . The study of such a set is difficult and involves naturally arithmetic, combinatorics, topology and complex analysis (see e.g. [M2]). Furthermore, to understand the global geography of the moduli space \mathcal{P}_d , it is useful to approximate the bifurcation current (resp. the bifurcation measure) by dynamically defined hypersurfaces (resp. finite sets). Following [MSS], one can try to approximate the bifurcation currents by different types of phenomena: existence of critical orbit relations or periodic cycles of given nature.

Let us focus first on the simplest case of the unicritical family, i.e. the family defined by $p_c(z) := z^d + c$, $c \in \mathbb{C}$. Consider the set $\text{Per}^*(n) := \{c \in \mathbb{C}; p_c^n(0) = 0\}$ of parameters that admit a superattractive periodic point of period dividing n . Recall that the *Multibrot set* \mathbf{M}_d is defined by $\mathbf{M}_d := \{c \in \mathbb{C}; \mathcal{J}_c \text{ is connected}\}$. Finally, let $\mu_{\mathbf{M}_d}$ be the harmonic measure of \mathbf{M}_d . Then, the first result in this direction goes back to Levin [Le]. In the quadratic family ($d = 2$), he proved that the measure equidistributed on the set $\text{Per}^*(n)$ converges to the harmonic measure of the Mandelbrot set, as $n \rightarrow \infty$. Favre and Rivera-Letelier [FRL] gave a *quantitative* version of Levin's in the unicritical family, using arithmetic methods. Namely, they proved

$$\left| \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} \right| \leq C \left(\frac{n}{d^n} \right)^{\frac{1}{2}} \|\varphi\|_{C^1},$$

for any $\varphi \in C_c^1(\mathbb{C})$ and any $n \geq 1$. Using potential theory, we prove here the following.

Theorem A. — *Let $d \geq 2$, $\mu_{\mathbf{M}_d}$ be the harmonic measure of the Multibrot set \mathbf{M}_d and consider the set $\text{Per}^*(n) := \{c \in \mathbb{C}; p_c^n(0) = 0\}$. There exists a constant $C > 0$ depending only on d such that for any $\varphi \in C_c^2(\mathbb{C})$ and any $n \geq 1$,*

$$\left| \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} \right| \leq C \frac{n}{d^n} \|\varphi\|_{C^2}.$$

A classical interpolation argument immediately gives Favre and Rivera-Letelier's above result. We shall also give a similar estimate for non-postcritically finite hyperbolic parameters in Section 4.

More recently, this subject has been intensively studied in the moduli space \mathcal{P}_d for $d \geq 3$. As the critical points are marked, i.e. can be followed holomorphically in the whole moduli space \mathcal{P}_d , the bifurcation current decomposes as $T_{\text{bif}} = \sum_{P'(c)=0} T_c$. For $n > k \geq 0$, let

$$\text{Per}_c(n, k) := \{[P] \in \mathcal{P}_d; P^n(c) = P^k(c)\}.$$

Refining Levin's approach, Dujardin and Favre prove in [DF] that for any sequence $k(n)$ satisfying $0 \leq k(n) < n$, the sequence $d^{-n}[\text{Per}_c(n, k(n))]$ converges to T_c in the weak sense

of currents on \mathcal{P}_d . Recently, Okuyama [Oku1] gave a simplified proof of their result in the case $k(n) = 0$.

Compared to the unicritical family, a significant difficulty comes from the fact that two distinct critical points can very well be, at some parameter, in the immediate Fatou component of the same attractive periodic point. To overcome that, for $n \geq 1$ and $w \in \mathbb{C}$, we let

$$\text{Per}(n, w) := \{[P] \in \mathcal{P}_d; P \text{ has a cycle of multiplier } w \text{ and exact period } n\} .$$

The set $\text{Per}(n, w)$ happens to be a complex hypersurface of degree $(d-1)d_n \sim (d-1)d^n$ and it is known that, for a fixed $w \in \mathbb{C}$, the sequence $d^{-n}[\text{Per}(n, w)]$ converges weakly to T_{bif} on the moduli space \mathcal{P}_d . The case $|w| \leq 1$ has been established by Bassanelli and Berteloot [BB3] using an approximation formula for the Lyapunov exponent (see also [BB2]). The more delicate case $|w| > 1$ has been proved recently by the first author in [G2] (see also [BG2] for the case of quadratic polynomials with changing multipliers).

Building on arithmetic methods, Favre and the first author [FG] proved that postcritically finite hyperbolic parameters with $(d-1)$ distinct super-attracting cycles (resp. strictly postcritically finite parameters with given combinatorics) equidistribute towards the bifurcation measure. The proof developed in that work is only qualitative, since there exists no effective version of Yuan’s arithmetic equidistribution Theorem. It also raises the question to know whether the result is of purely arithmetic nature or not. We consider the present work as a continuation of [FG].

Statement of the main results. — Our main goal here is twofold. First, we want to establish a *quantitative* equidistribution theorem for postcritically finite hyperbolic parameters. Second, we aim at giving a simpler proof than the one of [FG], relying only on pluripotential theoretic and complex analytic arguments. To our purposes, as in the recent works [DF, BB3, G2, FG], we shall use the following “orbifold” parametrization of the moduli space \mathcal{P}_d . For $(c, a) = (c_1, \dots, c_{d-2}, a) \in \mathbb{C}^{d-1}$, we let

$$P_{c,a}(z) := \frac{1}{d}z^d + \sum_{j=2}^{d-1} (-1)^{d-j} \sigma_{d-j}(c) \frac{z^j}{j} + a^d, \quad z \in \mathbb{C},$$

where $\sigma_\ell(c)$ is the monic symmetric polynomial of degree ℓ in (c_1, \dots, c_{d-2}) . Observe that the canonical projection $(c, a) \in \mathbb{C}^{d-1} \mapsto \{P_{c,a}\} \in \mathcal{P}_d$ is a finite branched cover of degree $d(d-1)$ and that the critical points of $P_{c,a}$ are exactly c_0, c_1, \dots, c_{d-2} with the convention that $c_0 := 0$ (see Section 5.1 for details). For an integer $n \in \mathbb{N}^*$, we let $\sigma(n)$ be the sum of its divisors $\sigma(n) := \sum_{k|n} k$. The function σ is known to be bounded by $Cn \log \log n$ for some constant C independent of n .

Our main result may be stated as follows.

Theorem B. — *Let $\underline{n} = (n_0, \dots, n_{d-2})$ be a $(d-1)$ -tuple of pairwise distinct positive integers with $n_0 \geq 2$ and let $\mu_{\underline{n}}$ be the probability measure equidistributed on the set of parameters in \mathbb{C}^{d-1} for which the critical point c_i is periodic of exact period n_i for all i . There exists a constant $C > 0$ depending only on d such that*

$$\left| \int \varphi \mu_{\underline{n}} - \int \varphi \mu_{\text{bif}} \right| \leq C \max_{1 \leq j \leq d-1} \left(\frac{\sigma(n_j)}{d^{n_j}} \right) \|\varphi\|_{\mathcal{C}^2},$$

for any test function $\varphi \in \mathcal{C}_c^2(\mathbb{C}^{d-1})$.

The first ingredient of the proof is a very general dynamical property established by Przytycki: if a critical point c of a rational map f does not lie in an attracting basin f , the points c and $f^n(c)$ can not be too close (see Lemma 3.1). The idea to use Przytycki estimate in this context has been introduced by Okuyama [Oku1] (see also [Oku2]) and constitutes the starting point of our work. Combined with known global properties of the moduli space \mathcal{P}_d , this allows us to have precise pointwise estimates outside some specific “bad” hyperbolic components. The second important tool is a transversality result for critical periodic orbit relations proved in [FG] and relying on Epstein’s transversality theory (see [E]). The last important ingredient we use is a L^1 estimate for specific solutions of the Laplacian in a bounded topological disk of an affine curve of \mathcal{P}_d , which proof crucially relies on length-area estimates (see Theorem 3.2). This allows us to replace an estimate involving the *diameter* of hyperbolic components with their *volume*. This actually is a key step, since even in the quadratic family, estimating the diameter of hyperbolic components is a very delicate problem related to the famous hyperbolicity conjecture.

Nevertheless, notice that, in the context of the unicritical family, the equation $p_c^n(0) = 0$ is known to have simple roots so we do not need to exclude parameters with a periodic critical point of period *dividing* n . As a consequence, we do not use transversality statements à la Epstein. Hence, we will start by the proof of Theorem A which is simpler than in the general case and more efficient. We may regard this as a model for the general case.

Following the strategy of the proof of [FG, Theorem 3], we can deduce from Theorem B a speed of convergence for the measure equidistributed on the (finite) set of parameters admitting $(d-1)$ distinct attracting cycles of given respective multipliers $w_0, \dots, w_{d-2} \in \mathbb{D}$ towards μ_{bif} . Let us be more precise and pick a $(d-1)$ -tuple $\underline{n} := (n_0, \dots, n_{d-2})$ of positive integers and $\underline{w} := (w_1, \dots, w_{d-1}) \in \mathbb{C}^{d-1}$. When the set $\bigcap_{i=1}^{d-1} \text{Per}(n_i, w_i)$ is finite, let

$$\mu_{\underline{n}, \underline{w}} := \frac{1}{(d-1)! \prod_j d_{n_j}} \bigwedge_{j=1}^{d-1} [\text{Per}(n_j, w_j)] .$$

Notice that $\mu_{\underline{n}, \underline{w}}$ is a probability measure and that, when $\underline{w} \in \mathbb{D}^{d-1}$, the measure $\mu_{\underline{n}, \underline{w}}$ is exactly the measure equidistributed on the set of parameters in \mathbb{C}^{d-1} having $(d-1)$ cycles of respective exact period n_0, \dots, n_{d-2} and multipliers w_0, \dots, w_{d-2} .

The precise result we prove may be stated as follows.

Theorem C. — *Pick $\underline{w} = (w_0, \dots, w_{d-2}) \in \mathbb{D}^{d-1}$, let $\underline{n} = (n_0, \dots, n_{d-2})$ be a $(d-1)$ -tuple of pairwise distinct positive integers with $n_0 \geq 2$ and let $\mu_{\underline{n}, \underline{w}}$ be as above. There exists a constant $C > 0$ depending only on d such that for any test function $\varphi \in \mathcal{C}_c^1(\mathbb{C}^{d-1})$,*

$$\left| \int \varphi \mu_{\underline{n}, \underline{w}} - \int \varphi \mu_{\text{bif}} \right| \leq C \left(\max_{0 \leq j \leq d-2} \left\{ \frac{\sigma(n_j)}{d^{n_j}}, \frac{-1}{d^{n_j} \log |w_j|} \right\} \right)^{\frac{1}{2}} \|\varphi\|_{\mathcal{C}^1} .$$

Combining Theorem C with techniques from pluripotential theory (see e.g. [DTV] and [DS3]), we can actually prove that for all $(d-1)$ -tuples of multipliers (w_0, \dots, w_{d-2}) lying outside a pluripolar set of \mathbb{C}^{d-1} , the measure $\mu_{\underline{n}, \underline{w}}$ equidistributed on parameters having $(d-1)$ cycles of respective multipliers w_0, \dots, w_{d-2} converges towards the bifurcation measure, as soon as the periods of the given cycles grow fast enough:

Theorem D. — *Pick any sequence $\underline{n}_k = (n_{1,k}, \dots, n_{d-1,k})$ of $(d-1)$ -tuples of pairwise distinct positive integers such that the series $\sum_k \max_j \{n_{j,k}^{-1}\}$ converges. Then there exists*

a pluripolar set of $\mathcal{E} \subset \mathbb{C}^{d-1}$ such that for any $w \notin \mathcal{E}$, the set $\bigcap_{i=1}^{d-1} \text{Per}(n_{i,k}, w_i)$ is finite for any k and the sequence $(\mu_{\underline{n}_k, \underline{w}})_k$ converges to μ_{bif} in the sense of measures.

As an obvious corollary, we can deduce that if we only assume $\min_j \{n_{j,k}\} \rightarrow \infty$, then, for \underline{w} outside a pluripolar set, up to extraction $(\mu_{\underline{n}_k, \underline{w}})_k$ converges to μ_{bif} in the sense of measures. Here is another immediate and interesting consequence of Theorem D.

Corollary E. — *Pick any sequence $\underline{n}_k = (n_{1,k}, \dots, n_{d-1,k})$ of $(d-1)$ -tuples of pairwise distinct positive integers such that the series $\sum_k \max_j (n_{j,k}^{-1})$ converges. Then, for almost any $\Theta = (\theta_1, \dots, \theta_{d-1}) \in]0, 1[^{d-1}$, if $\underline{w}(\Theta) = (e^{2i\pi\theta_1}, \dots, e^{2i\pi\theta_{d-1}})$, the sequence $(\mu_{\underline{n}_k, \underline{w}(\Theta)})_k$ of measures equidistributed on parameters having cycles of respective exact period $n_{j,k}$ and multipliers $e^{2i\pi\theta_j}$ converges to μ_{bif} in the sense of measures.*

Notice that Bassanelli and Berteloot [BB2] proved a weaker version of Corollary E: they prove that the average measures $\int_{]0, 1[^{d-1}} \mu_{\underline{n}_k, \underline{w}(\Theta)} dm(\Theta)$ converge weakly to the bifurcation measure. Their proof also works in any codimension.

We view these results as parametric analogues of important dynamical phenomena. Indeed, Theorem B is an analogue of the equidistribution of repelling periodic points of a holomorphic endomorphism F of \mathbb{P}^k towards its maximal entropy measure μ_F , and Theorem D is an analogue of the equidistribution of preimages of a generic points, again towards the measure μ_F (see [Lj, BD]).

Perspectives. — The questions we discuss here may be addressed in a more general setting. A first natural generalization is the case when critical points can have the same period. In that case, the transversality theory à la Epstein fails at parameters admitting multiple critical points and we a priori have no control of the multiplicity of intersection at those parameters.

A second natural question is concerned with the case of the moduli space of degree d rational maps. Even in the case of quadratic rational maps which is much better understood than the general case, important difficulties occur. The main problem comes from the fact that, contrary to the case of polynomials, the support of the bifurcation measure is *not* compact in the moduli space of quadratic rational maps and that the collection of relatively compact hyperbolic components cluster at infinity (see [BG1]).

We shall study both cases in future works.

Structure of the article. — Section 2 is devoted to needed material. In Section 3, we establish two preliminary results: Przytycki distance estimate and the L^1 estimate for solutions of the Laplacian. We then present the proof of Theorem A and its corollaries in Section 4. The initial estimates relying on Przytycki Lemma are established in Section 5. Section 6 is dedicated to the proof of the main Theorem B and Section 7 to proving Theorems C and D. Finally, in Section 8, we investigate other approximation phenomena. We try here to understand the distribution of maps for which the critical points are sent to some prescribed target (and not necessarily themselves). We are especially interested in a theorem of Dujardin [Du] that proves the convergence outside some pluripolar set. We give here some convergence estimates and show that in some cases, the pluripolar set can be described explicitly.

Acknowledgment. — The research of both authors is partially supported by the ANR project Lambda ANR-13-BS01-0002. We would like to thank Vincent Guedj for very helpful discussions concerning L^1 estimates for solutions of the Laplacian and François Berteloot and Charles Favre for useful comments on preliminary versions.

2. Background material

In this section, we want to recall briefly background material on bifurcation currents and on classical complex analytic tools we will rely on in the whole paper.

2.1. Holomorphic families with marked critical points

Let us recall classic facts concerning holomorphic families of rational maps.

A *holomorphic family* $(f_\lambda)_{\lambda \in \Lambda}$ of degree $d \geq 2$ rational maps parametrized by Λ is a holomorphic map

$$f : \Lambda \times \mathbb{P}^1 \longrightarrow \mathbb{P}^1$$

such that the map $f_\lambda := f(\lambda, \cdot) : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is a degree d rational map, or equivalently if the map $f : \lambda \in \Lambda \mapsto f_\lambda \in \text{Rat}_d$ is holomorphic. We say that a holomorphic family $(f_\lambda)_{\lambda \in \Lambda}$ is *with a marked critical point* if there exists a holomorphic map $c : \Lambda \rightarrow \mathbb{P}^1$ such that $f'_\lambda(c(\lambda)) = 0$ for all $\lambda \in \Lambda$.

We say that a marked critical point c is *passive* at $\lambda_0 \in \Lambda$ if there exists a neighborhood $U \subset \Lambda$ of λ_0 such that the sequence F_n of holomorphic maps defined by $F_n(\lambda) := f_\lambda^n(c(\lambda))$ is a normal family on U . We say that c is *active* at λ_0 if it is not passive at λ_0 . The *activity locus* of c is the set of parameters $\lambda_0 \in \Lambda$ such that c is active at λ_0 .

Theorem 2.1 (Dujardin-Favre). — *The sequence $d^{-n}(F_n)^*\omega_{\mathbb{P}^1}$ converges in the weak sense of currents to a closed positive $(1, 1)$ -current T_c which is supported by the activity locus of c .*

More precisely, there exists a locally uniformly bounded sequence of continuous function $u_n : \Lambda \times \mathbb{P}^1 \rightarrow \mathbb{R}$ such that $\frac{1}{d^n}(F_n)^*\omega_{\mathbb{P}^1} = T_c + \frac{1}{d^n}dd^c u_n$, see e.g. [DF, Proposition-Definition 3.1]. It is also known that $T_c \wedge T_c = 0$ (see [G1, Theorem 6.1]).

Let us also recall that, when a holomorphic family is with $2d - 2$ marked critical points c_1, \dots, c_{2d-2} , the current $T_{\text{bif}} := \sum_{i=1}^{2d-2} T_{c_i}$ is supported by the bifurcation locus in the sense of Mañé, Sad and Sullivan (see [DeM2]).

We define the *bifurcation measure* of a family Λ as

$$\mu_{\text{bif}} := \frac{1}{m!} T_{\text{bif}}^m, \quad m = \dim \Lambda.$$

This measure detects, in a certain sense, the strongest bifurcations which occur in Λ . Finally, when $(f_\lambda)_{\lambda \in \Lambda}$ is a holomorphic family of polynomials with $d - 1$ marked critical point $c_1, \dots, c_{d-1} : \Lambda \rightarrow \mathbb{C}$, we let

$$g_\lambda(z) := \lim_{n \rightarrow \infty} d^{-n} \log^+ |f_\lambda^n(z)|,$$

for $\lambda \in \Lambda$ and $z \in \mathbb{C}$. The current T_i is then given by $T_{c_i} = dd^c g_\lambda(c_i(\lambda))$ (see e.g. [DF]). We also can remark that it actually can be considered as equipped with $2d - 2$ marked critical points letting $c_d \equiv \dots \equiv c_{2d-2} \equiv \infty$ and that $T_i = 0$ for all $d \leq i \leq 2d - 2$.

2.2. A specific family

Recall that the *moduli space* \mathcal{P}_d of degree d polynomials is the space of affine conjugacy classes of degree d polynomials with $d - 1$ marked critical points. We define a finite branched cover of $\mathbb{C}^{d-1} \rightarrow \mathcal{P}_d$ as follows. For $c = (c_1, \dots, c_{d-2}) \in \mathbb{C}^{d-2}$ and $a \in \mathbb{C}$, let

$$P_{c,a}(z) := \frac{1}{d}z^d + \sum_{j=2}^{d-1} (-1)^{d-j} \frac{\sigma_{d-j}(c)}{j} z^j + a^d, \quad z \in \mathbb{C},$$

where $\sigma_k(c)$ is the monic homogeneous degree k symmetric function in the c_i 's. This family is known to be a finite branched cover (see e.g. [DF, §5]). Remark also that the critical points of $P_{c,a}$ are exactly c_0, \dots, c_{d-2} , where we set $c_0 := 0$, and that they depend algebraically on $(c, a) \in \mathbb{C}^{d-1}$.

We define a continuous psh function $G : \mathbb{C}^{d-1} \rightarrow \mathbb{R}_+$ by setting

$$G(c, a) := \max_{0 \leq j \leq d-2} g_{c,a}(c_j), \quad (c, a) \in \mathbb{C}^{d-1}.$$

It is known that the *connectedness locus*

$$\mathcal{C}_d := \{(c, a) \in \mathbb{C}^{d-1}; \mathcal{J}_{c,a} \text{ is connected}\}$$

is compact and satisfies $\mathcal{C}_d = \{G = 0\}$ (see [BH]). Moreover, the bifurcation measure, in this actual family, coincides with the Monge-Ampère mass of G , i.e.

$$\mu_{\text{bif}} = (dd^c G)^{d-1}$$

as probability measures on \mathbb{C}^{d-1} . It is also known that

$$G(c, a) = \log^+ \max\{|c|, |a|\} + O(1),$$

where we set $|c| := \max_{1 \leq j \leq d-2} |c_j|$, and that the function G is the pluricomplex Green function of \mathcal{C}_d . In particular, $\mu_{\text{bif}} = (dd^c G)^{d-1}$ is supported by the Shilov boundary $\partial_S \mathcal{C}_d$ of \mathcal{C}_d (see [DF, §6]). In particular, the estimates of [DF] give

Lemma 2.2. — *There exists a constant $C > 0$ independent of (c, a) such that for any $(c, a) \in \mathbb{C}^{d-1}$, any $z \in \mathbb{C}$ and any n :*

$$\left| \frac{1}{d^n} \log^+ |P_{c,a}^n(z)| - g_{c,a}(z) \right| \leq \frac{1}{d^n} (\log^+ \max\{|c|, |a|\} + C').$$

Proof. — This actually follows from [DF, Proposition-Definition 3.1] and the precise definition of $P_{c,a}$. Indeed, in the proof of that proposition, observe that the function \hat{g} satisfies the estimate $|\hat{g}((c, a), z)| \leq \log^+ \max\{|c|, |a|\} + C'$ for some C' that does not depend on (c, a) nor z . The result then follows by summing. \square

2.3. Complex analytic tools

We will denote in what follows $d_{\mathbb{P}^1}$ the classical spherical distance on \mathbb{P}^1 . For a C^1 map f , we denote $f^\#$ the spherical derivative of f :

$$\forall z \in \mathbb{P}^1, \quad f^\#(z) := \lim_{d_{\mathbb{P}^1}(z,y) \rightarrow 0} \frac{d_{\mathbb{P}^1}(f(z), f(y))}{d_{\mathbb{P}^1}(z, y)}.$$

We also shall denote by $\|D^2 f(z)\|$ the norm of the bilinear map $D^2 f(z)$ induced by the spherical metric.

Recall that if A is an annulus and if A is conformally equivalent to $A' = \{z \in \mathbb{C}; r < |z| < R\}$ with $0 < r < R < +\infty$, the *modulus* of A is the same as the modulus of A' :

$$\text{mod}(A) = \text{mod}(A') = \frac{1}{2\pi} \log \left(\frac{R}{r} \right) .$$

We will rely on the following classical estimate (see [BD, Appendix]).

Lemma 2.3 (Briend-Duval). — *There exists a constant $\tau > 0$ depending only on $k \geq 1$ such that for any holomorphic disks $D_1 \Subset D_2 \Subset \mathbb{C}^k$, and any hermitian metric α on \mathbb{C}^k ,*

$$(\text{diam}_\alpha(D_1))^2 \leq \tau \cdot \frac{\text{Area}_\alpha(D_2)}{\min(1, \text{mod}(A))} ,$$

where A is the annulus $D_2 \setminus \overline{D_1}$ and distances are computed with respect to α .

We also rely on the following classical integration by part formula, which can be stated as follows (see [Dem, Formula 3.1 page 144]).

Lemma 2.4. — *Let $\Omega \Subset \Omega' \subset \mathbb{C}^k$ be bounded open sets. Assume that Ω has smooth boundary. Let u, v be psh functions on Ω' and let T be a closed positive $(k-1, k-1)$ -current on Ω' such that $dd^c u \wedge T$ and $dd^c v \wedge T$ are well-defined. Then*

$$\int_{\Omega} (v dd^c u - u dd^c v) \wedge T = \int_{\partial\Omega} (v d^c u - u d^c v) \wedge T .$$

3. Preliminary results

In this section, we establish the two main technical estimates we will rely on. The first one is of dynamical nature and follows very closely a classical result of Przytycki. The second is of more geometric nature and might be of independent interest.

3.1. Local dynamical estimates

We shall use the following estimate in a crucial way. The proof follows very closely that of [P, Lemma 1]. The idea to use this result for proving equidistribution phenomena in parameters spaces first appeared in the recent work [Oku1] of Okuyama.

Lemma 3.1. — *Let $(f_\lambda)_{\lambda \in \Lambda}$ be a holomorphic family of degree d rational maps and let $c : \Lambda \rightarrow \mathbb{P}^1$ be a marked critical point of $(f_\lambda)_{\lambda \in \Lambda}$. Assume that $c(\lambda)$ does not lie persistently in a parabolic basin of f_λ . There exists a universal constant $\kappa > 0$ and a continuous function $M : \Lambda \rightarrow]1, +\infty[$ depending only on the family $(f_\lambda)_{\lambda \in \Lambda}$ such that, for any $n \geq 1$ and any $\lambda \in \Lambda$ with $f_\lambda^n(c(\lambda)) \neq c(\lambda)$,*

- either $d_{\mathbb{P}^1}(f_\lambda^n(c(\lambda)), c(\lambda)) \geq \kappa \cdot M(\lambda)^{-n}$,
- or $c(\lambda)$ lies in the immediate basin of an attracting cycle of period p dividing n and $d_{\mathbb{P}^1}(c(\lambda), \mathcal{J}_\lambda) \geq \kappa M(\lambda)^{-n}$. In that case, $d_{\mathbb{P}^1}(f_\lambda^n(c(\lambda)), c(\lambda)) \geq \kappa \cdot |\rho(\lambda)|^{n/p} \cdot M(\lambda)^{-3n}$, where $\rho(\lambda) \in \mathbb{D}$ is the multiplier of the attracting cycle.

Proof. — Notice that the function $M : \Lambda \rightarrow]1, +\infty[$ defined by

$$M(\lambda) := \max \left\{ \sup_{z \in \mathbb{P}^1} f_\lambda^\#(z), \sup_{z \in \mathbb{P}^1} \|D^2 f_\lambda(z)\| \right\} \in]1, +\infty[$$

is continuous on Λ . Notice that, if $\|\cdot\|$ is the metric $T^*\mathbb{P}^1$ induced by the spherical metric, as $Df_\lambda : \mathbb{P}^1 \rightarrow T^*\mathbb{P}^1$ is continuous, it has compact image $\mathcal{K} := Df_\lambda(\mathbb{P}^1)$. This guarantees that $M(\lambda)$ is finite. Moreover, the map $f_\lambda : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is Lipschitz with constant $M(\lambda)$ with respect to $d_{\mathbb{P}^1}$, i.e.

$$d_{\mathbb{P}^1}(f_\lambda(z), f_\lambda(w)) \leq M(\lambda)d_{\mathbb{P}^1}(z, w) ,$$

for any $z, w \in \mathbb{P}^1$ and any $\lambda \in \Lambda$. Let $\kappa := 1/16$.

We follow closely the proof of [**P**, Lemma 1]. Assume that $d_{\mathbb{P}^1}(c(\lambda), \mathcal{J}_\lambda) < \kappa M(\lambda)^{-n}$ and $d_{\mathbb{P}^1}(f_\lambda^n(c(\lambda)), c(\lambda)) < \kappa M(\lambda)^{-n}$. Let

$$\epsilon := \max\{d_{\mathbb{P}^1}(c(\lambda), \mathcal{J}_\lambda), d_{\mathbb{P}^1}(c(\lambda), f_\lambda^n(c(\lambda)))\} > 0 .$$

Then $\overline{\mathbb{B}}(c(\lambda), \epsilon) \cap \mathcal{J}_\lambda \neq \emptyset$. Then, since $c(\lambda) \in C(f_\lambda)$, by the Mean Value Theorem,

$$\text{diam } f_\lambda(\mathbb{B}(c(\lambda), 2\epsilon)) \leq 16M(\lambda)\epsilon^2$$

so

$$\text{diam } f_\lambda^n(\mathbb{B}(c(\lambda), 2\epsilon)) \leq 16M(\lambda)^n\epsilon^2 < \epsilon .$$

In particular, $f_\lambda^n(\mathbb{B}(c(\lambda), 2\epsilon)) \subset \mathbb{B}(c(\lambda), 2\epsilon)$ and $f_\lambda^{kn}(\mathbb{B}(c(\lambda), 2\epsilon)) \subset \mathbb{B}(c(\lambda), 2\epsilon)$ follows for any $k \geq 1$ by an immediate induction. By Montel's Theorem, the sequence $(f_\lambda^{kn})_{k \geq 1}$ is a normal family on $\mathbb{B}(c(\lambda), 2\epsilon)$, which is a contradiction, since $\mathcal{J}_\lambda \cap \mathbb{B}(c(\lambda), 2\epsilon) \neq \emptyset$. That implies in particular the lemma in the case where $c(\lambda) \in \mathcal{J}_\lambda$.

Assume from now on that $c(\lambda) \notin \mathcal{J}_\lambda$. Consider first the case when $c(\lambda)$ is either attracted by an attracting cycle of period k that does not divide n , or not attracted by an attracting cycle or a parabolic cycle. Then $f_\lambda^n(c(\lambda))$ lies in a different Fatou component of f_λ than $c(\lambda)$. Then we either have $d_{\mathbb{P}^1}(c(\lambda), \mathcal{J}_\lambda) < \kappa \cdot M(\lambda)^{-n}$, which has just been proven to imply the wanted result, or $d_{\mathbb{P}^1}(c(\lambda), \mathcal{J}_\lambda) \geq \kappa \cdot M(\lambda)^{-n}$ and we have

$$d_{\mathbb{P}^1}(f_\lambda^n(c(\lambda)), c(\lambda)) \geq d_{\mathbb{P}^1}(c(\lambda), \mathcal{J}_\lambda) \geq \kappa \cdot M(\lambda)^{-n} .$$

Second, if $c(\lambda)$ lies in a parabolic basin of f_λ , by assumption, the critical point is active at λ , hence there exists $\lambda_n \rightarrow \lambda$ with $c(\lambda_n) \in \mathcal{J}_{\lambda_n}$, and the conclusion follows by continuity of the function $d_{\mathbb{P}^1}(f_\lambda^n(c(\lambda)), c(\lambda)) - \kappa \cdot M(\lambda)^{-n}$.

Finally, if $c(\lambda)$ is attracted by an attracting cycle of period p dividing n and multiplier $\rho(\lambda) \in \mathbb{D}$, and if $\epsilon = d_{\mathbb{P}^1}(c(\lambda), f_\lambda^n(c(\lambda))) < \kappa M(\lambda)^{-n}$ then

$$f_\lambda^n(\mathbb{B}(c(\lambda), \epsilon)) \subset \mathbb{B}(c(\lambda), \epsilon) .$$

By Brouwer fixed point Theorem, f_λ^n has a fixed point in $\overline{\mathbb{B}}(c(\lambda), \epsilon)$ and thus $p_\lambda \in \overline{\mathbb{B}}(c(\lambda), \epsilon)$. In particular, reducing slightly ϵ , we may assume $\overline{\mathbb{B}}(c(\lambda), \epsilon) \cap \mathcal{J}_\lambda = \emptyset$, i.e. $\overline{\mathbb{B}}(c(\lambda), \epsilon)$ is contained in the attracting basin of the attracting point p_λ attracting $c(\lambda)$. This can be rewritten as

$$d_{\mathbb{P}^1}(p_\lambda, c(\lambda)) \leq 2d_{\mathbb{P}^1}(c(\lambda), f_\lambda^n(c(\lambda))) .$$

By the Mean Value Theorem, one has

$$|\rho(\lambda)|^{n/p} = \|D(f_\lambda^n)(p_\lambda) - D(f_\lambda^n)(c(\lambda))\| \leq nM(\lambda)^{2n}d_{\mathbb{P}^1}(p_\lambda, c(\lambda)) .$$

This concludes the proof since we have $2 < \kappa^{-1}$ by the above. \square

3.2. L^1 -estimate for local solutions of the Laplacian on affine curves

Our aim, in the present, is to give the following L^1 -estimate for solutions of the Laplacian in disks of algebraic curves. We let β be the standard hermitian metric on \mathbb{C}^k . Precisely, we want to prove the following.

Theorem 3.2. — *Let $k \geq 1$, let $S \subset \mathbb{C}^k$ be an affine algebraic curve and let Ω be a simply connected bounded open set of S and $z_0 \in \Omega$. Let also $f \in \mathcal{C}(\overline{\Omega}, \mathbb{C})$ be holomorphic on Ω . Assume that $\Omega \cap S_{\text{sing}} = \emptyset$ and that $dd^c \log |f| = \delta_{z_0}$. Then, there exists a constant $C > 0$ depending neither on Ω , f , z_0 and S , such that*

$$\|\log |f|\|_{L^1(\Omega)} \leq C \max \left\{ \|\log |f|\|_{L^\infty(\partial\Omega)}, 1 \right\} \text{Area}_\beta(\Omega) .$$

Proof. — Set $u := \log |f|$. First, if f vanishes on $\partial\Omega$, then the estimate is trivial since $\|u\|_{L^\infty(\partial\Omega)} = +\infty$ then. We thus assume $f(\zeta) \neq 0$ for all $\zeta \in \partial\Omega$.

Let $h : \Omega \rightarrow \mathbb{D}$ be a biholomorphic map with $h(z_0) = 0$. It is clear that $|h|$ extends continuously to $\partial\Omega$ with $|h| = 1$ on $\partial\Omega$. We set $\chi := \log |h|$ and $K := \max \left\{ \|u\|_{L^\infty(\partial\Omega)}, 1 \right\}$. The function χ is subharmonic on Ω , satisfies $dd^c \chi = \delta_{z_0}$ and $\chi \leq 0$. The functions h/f and f/h are holomorphic on Ω and satisfy

$$\left| \frac{f}{h} \right| \leq \exp(K) \quad \text{and} \quad \left| \frac{h}{f} \right| \leq \exp(K) \quad \text{on } \partial\Omega .$$

Using twice the maximum principle, we get $\chi - K \leq u \leq K + \chi \leq K - \chi$ on Ω . Define

$$\Omega' := \{z \in \Omega; \chi(z) < -3K\} \quad \text{and} \quad \Omega'' := \left\{ z \in \Omega; \chi(z) < -\frac{5}{2}K \right\} .$$

We can decompose $\|u\|_{L^1(\Omega)}$ as follows

$$\begin{aligned} \|u\|_{L^1(\Omega)} &= \int_\Omega |u| \beta \leq \int_\Omega (K - \chi) \beta \leq K \text{Area}_\beta(\Omega) - \int_\Omega \chi \beta \\ &\leq K \text{Area}_\beta(\Omega) - \int_{\Omega \setminus \Omega'} \chi \beta - \int_{\Omega'} \chi \beta \\ &\leq 4K \text{Area}_\beta(\Omega) - \int_{\Omega'} \chi \beta . \end{aligned}$$

We are thus left with estimating $\int_{\Omega'} \chi \beta$. Writing $v(z) := \|z - z_0\|^2$, where $\|\cdot\|$ is the euclidean norm of \mathbb{C}^k , we have $\beta = dd^c v$ and by Stokes (see Lemma 2.4):

$$\int_{\Omega'} \chi \beta = \int_{\Omega'} v dd^c \log |h| - \int_{\partial\Omega'} v d^c \log |h| + \int_{\partial\Omega'} \log |h| d^c v .$$

As $dd^c \log |h| = \delta_{z_0}$, $d^c \log |h|^2 = d^c |h|^2 / |h|^2$ and by definition of Ω' , we find

$$\begin{aligned} \int_{\Omega'} \chi \beta &= -3K \int_{\partial\Omega'} d^c v - \int_{\partial\Omega'} v d^c \log |h| \\ &= -3K \text{Area}_\beta(\Omega') - \frac{1}{2} \exp(6K) \int_{\partial\Omega'} v d^c |h|^2 . \end{aligned}$$

Applying again Stokes yields

$$\begin{aligned} \int_{\partial\Omega'} v d^c |h|^2 &= \int_{\Omega'} v dd^c |h|^2 - \int_{\Omega'} |h|^2 \beta + \int_{\partial\Omega'} |h|^2 d^c v \\ &\leq \int_{\Omega'} v dd^c |h|^2 + \exp(-6K) \text{Area}_\beta(\Omega') \end{aligned}$$

since $|h| \leq 1$ and $v \geq 0$.

All these estimates summarize as follows

$$(1) \quad 0 \geq \int_{\Omega'} \chi \beta \geq -4K \cdot \text{Area}_\beta(\Omega') - \frac{1}{2} \exp(6K) \int_{\Omega'} v dd^c |h|^2 ,$$

since $K \geq 1$ and $|h| \geq 0$. Since $v(z) = \|z - z_0\|^2$, we can bound it in Ω' by $(\text{diam}_\beta(\Omega'))^2$. We now apply Lemma 2.3 to $\Omega' \Subset \Omega''$ and find

$$\begin{aligned} \int_{\Omega'} v dd^c |h|^2 &\leq (\text{diam}_\beta(\Omega'))^2 \cdot \int_{\Omega'} dd^c |h|^2 = (\text{diam}_\beta(\Omega'))^2 \cdot \int_{\mathbb{D}(0, e^{-3K})} \beta \\ &\leq (\text{diam}_\beta(\Omega'))^2 \cdot \pi \exp(-6K) \\ &\leq \pi \exp(-6K) \tau \cdot \frac{\text{Area}_\beta(\Omega'')}{\min\{1, \text{mod}(\Omega'' \setminus \overline{\Omega'})\}} . \end{aligned}$$

Since h is a biholomorphism and since $K \geq 1$, we have

$$\text{mod}(\Omega'' \setminus \overline{\Omega'}) = \text{mod}(\mathbb{D}(0, e^{-5K/2}) \setminus \overline{\mathbb{D}(0, e^{-3K})}) = \frac{K}{2\pi} \geq \frac{1}{2\pi} .$$

Taking $C := 8 + \pi^2 \tau$ ($K \geq 1$) ends the proof. \square

4. Speed of convergence in the unicritical family

This section is devoted to the proof of Theorem A. The method we use can be seen as a toy-model for the proof of Theorem B. Our idea consists in giving estimates in L_{loc}^1 for the sequence $c \mapsto d^{-n+1} \log |p_c^n(0)| - g_{\mathbf{M}_d}$ of DSH functions (difference of subharmonic functions) and then to deduce Theorem A from these estimates.

4.1. Preliminaries

Recall that the *Multibrot set* \mathbf{M}_d is defined by

$$\mathbf{M}_d := \{c \in \mathbb{C} ; \mathcal{J}_c \text{ is connected}\} .$$

It coincides with the set $\{c \in \mathbb{C} ; g_c(0) = 0\}$. The bifurcation locus of the family $(p_c)_{c \in \mathbb{C}}$ is known to coincide with the boundary $\partial \mathbf{M}_d$ of the Multibrot set. In this family, the bifurcation measure is given by

$$dd^c g_c(0) = \frac{1}{d} dd^c g_c(c) = \frac{1}{d} dd^c g_{\mathbf{M}_d}$$

where $g_{\mathbf{M}_d}$ is the Green function of the compact set \mathbf{M}_d (see e.g. [St]). In particular, $dd^c g_{\mathbf{M}_d} = \mu_{\mathbf{M}_d}$ is the equilibrium measure (and the harmonic measure) of \mathbf{M}_d .

Let us first prove the following.

Lemma 4.1. — *There exists $C_1 > 0$ depending only on d such that*

$$|d^{-n+1} \log |p_c^n(0)| - g_{\mathbf{M}_d}(c)| \leq \frac{n}{d^{n-1}} C_1 ,$$

for any $c \in \mathbb{C} \setminus \mathbf{M}_d$ and any $n \geq 1$.

Proof. — Let $u_n(c) := d^{-n+1} \log |p_c^n(0)|$, $c \in \mathbb{C}$. Recall that, for any $c \in \mathbf{M}_d$, one has $|p_c^n(0)| \leq 2$. Let $M := \sup_{c \in \partial \mathbf{M}_d} M(c)$, where $M(c)$ is the constant given by Lemma 3.1. Then, for all $c \in \partial \mathbf{M}_d$,

$$-\frac{1}{d^{n-1}} (n \log M_1 - \log \kappa) \leq u_n(c) - g_{\mathbf{M}_d}(c) \leq \frac{1}{d^{n-1}} \log 2 ,$$

since $g_{\mathbf{M}_d}(c) = 0$. As $\kappa = 1/16$, we get for any $c \in \partial \mathbf{M}_d$

$$(2) \quad -\frac{n}{d^{n-1}} \log (\max (16M_1, 2)) \leq u_n(c) - g_{\mathbf{M}_d}(c) \leq \frac{n}{d^{n-1}} \log (\max (16M_1, 2)) .$$

Hence we let $C_1 := \max (16M_1, 2)$. Now, as $c \rightarrow \infty$, by definition of p_c , one has

$$\lim_{c \rightarrow \infty} (u_n(c) - \log |c|) = \lim_{c \rightarrow \infty} \left(d^{-(n-1)} \log \left| c^{d^{n-1}} \right| - \log |c| \right) = 0 .$$

As $g_{\mathbf{M}_d}(c) = \log |c| + o(1)$ as $c \rightarrow \infty$ (see e.g. [CG, Chapter VIII]), the function $u_n - g_{\mathbf{M}_d}$ extends continuously on $\mathbb{P}^1 \setminus \mathbf{M}_d$. Moreover, $h := u_n - g_{\mathbf{M}_d}$ is harmonic on $\mathbb{C} \setminus \mathbf{M}_d$.

The function h is thus subharmonic and continuous on $\mathbb{P}^1 \setminus \mathbf{M}_d$. Moreover, since h is harmonic on $\mathbb{C} \setminus \mathbf{M}_d$, one has $\text{supp}(\Delta h) \subset \{\infty\}$. As h is continuous at ∞ , it can not have a Dirac mass here, hence $\Delta h = 0$ on $\mathbb{P}^1 \setminus \mathbf{M}_d$, i.e. h is harmonic on $\mathbb{P}^1 \setminus \mathbf{M}_d$. By the maximum principle, applied successively to h and $-h$, (2) gives the wanted estimate. \square

4.2. In the Multibrot set

We denote $v_n := \log |p_c^n(0)|$. We will prove that:

Theorem 4.2. — *There exists $C > 0$ depending only on d such that for any $n \geq 1$,*

$$\|v_n\|_{L^1(\mathbf{M}_d)} \leq C \cdot n \cdot \text{Area}(\mathbf{M}_d) .$$

The next lemma will be use in the proof of Theorem 4.2. Though it is classical, we include a proof for the sake of completeness.

Lemma 4.3. — *The polynomial $p_c^n(0) \in \mathbb{C}[c]$ has simple roots.*

Proof. — Let $Q_n(c) := p_c^n(0) \in \mathbb{Z}[c]$. By definition, $Q_n(c) = Q_{n-1}(c)^d + c$, hence $Q'_n = dQ'_{n-1}Q_{n-1}^{d-1} + 1$. In particular, the discriminant $\Delta(Q_n) = \text{Res}(Q_n, Q'_n)$ is the determinant of a matrix with entries divisible by d below the diagonal and equal to 1 on the diagonal. Hence reducing modulo d , it is upper triangular with 1 on the diagonal, i.e. $\Delta(Q_n) \equiv 1 \pmod{d}$. In particular, if $Q_n(c_0) = 0$, then $Q'_n(c_0) \neq 0$ and the point c_0 is a simple root of Q_n . \square

Let $n \geq 1$ and \mathcal{H}_n be the union of connected components Ω of the interior of \mathbf{M}_d such that $\text{Per}^*(n) \cap \Omega \neq \emptyset$. For Ω such a connected component, recall that Ω is simply connected and there exists only one $c_c \in \Omega$ for which $p_{c_c}^n(0) = 0$ ([DH], this can be seen using the multiplier map $\rho : \Omega \rightarrow \mathbb{D}$ which is a branched cover of degree $d - 1$ which is totally ramified at $\{c_c\} := \rho^{-1}\{0\} = \text{Per}^*(n) \cap \Omega$). The proof of Theorem 4.2 is an application of Theorem 3.2.

Proof. — We decompose $\|v_n\|_{L^1(\mathbf{M}_d)}$ as follows

$$\|v_n\|_{L^1(\mathbf{M}_d)} = \int_{\mathbf{M}_d} |v_n| \beta = \int_{\mathbf{M}_d \setminus \mathcal{H}_n} |v_n| \beta + \int_{\mathcal{H}_n} |v_n| \beta .$$

Let U be a connected component of $\mathbf{M}_d \setminus \mathcal{H}_n$. By Lemma 4.1 (recall that $g_{\mathbf{M}_d}(c) = 0$ in \mathbf{M}_d), we have that

$$|v_n| \leq nC_1 \text{ on } \partial U .$$

As v_n is harmonic on U , by the maximum principle, the estimate extends to U :

$$|v_n| \leq nC_1 \text{ on } U .$$

Hence:

$$\int_{\mathbf{M}_d \setminus \mathcal{H}_n} |v_n| \beta \leq nC_1 \text{Area}_\beta(\mathbf{M}_d \setminus \mathcal{H}_n) .$$

Now, let Ω be a connected component of \mathcal{H}_n . Then, by Lemma 4.3, the function v_n satisfies $dd^c v_n = \delta_{c_c}$ where c_c is the center of the component Ω . Hence using Theorem 3.2 and Lemma 4.1 gives (take $\log |f| = v_n$):

$$\|v_n\|_{L^1(\Omega)} \leq C \max \left\{ \|v_n\|_{L^\infty(\partial\Omega)}, 1 \right\} \text{Area}_\beta(\Omega) \leq CnC_1 \text{Area}_\beta(\Omega) ,$$

where C is a universal constant that depends only on d . Summing on all the connected components Ω of \mathcal{H}_n gives:

$$\int_{\mathcal{H}_n} |v_n| \beta \leq CnC_1 \text{Area}_\beta(\mathcal{H}_n) .$$

Summing over \mathcal{H}_n and $\mathbf{M}_d \setminus \mathcal{H}_n$ ends the proof. \square

4.3. Proof of Theorem A

Let us now explain how to deduce Theorem A from Theorem 4.2.

Proof of Theorem A. — Recall that $u_n(c) = d^{-n+1} \log |p_c^n(0)| = d^{-n+1} v_n$. Let $\varphi \in \mathcal{C}_c^2(\mathbb{C})$. Then, for any $n \geq 1$, by Stokes formula,

$$\frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} = \int \varphi dd^c u_n - \int \varphi dd^c g_{\mathbf{M}_d} = \int (u_n - g_{\mathbf{M}_d}) dd^c \varphi .$$

We cut the integral into two parts

$$\frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} = \int_{\mathbf{M}_d} (u_n - g_{\mathbf{M}_d}) dd^c \varphi + \int_{\mathbb{C} \setminus \mathbf{M}_d} (u_n - g_{\mathbf{M}_d}) dd^c \varphi .$$

Now, as φ is \mathcal{C}^2 , we have (up to a constant that depends on the choice of the \mathcal{C}^2 -norm) that $\|\varphi\|_{\mathcal{C}^2} \omega \pm dd^c \varphi \geq 0$, where ω is the Fubini-Study form on \mathbb{P}^1 , hence we can write

$$\left| \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} \right| \leq \|\varphi\|_{\mathcal{C}^2} \left(\int_{\mathbf{M}_d} |u_n - g_{\mathbf{M}_d}| \omega + \int_{\mathbb{C} \setminus \mathbf{M}_d} |u_n - g_{\mathbf{M}_d}| \omega \right) .$$

As $\omega \leq \beta$, where β is the standard hermitian metric on \mathbb{C} ,

$$\left| \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} \right| \leq \|\varphi\|_{\mathcal{C}^2} \left(\int_{\mathbf{M}_d} |u_n - g_{\mathbf{M}_d}| \beta + \int_{\mathbb{C} \setminus \mathbf{M}_d} |u_n - g_{\mathbf{M}_d}| \omega \right) .$$

By Lemma 4.1, we have the bound:

$$\int_{\mathbb{C} \setminus \mathbf{M}_d} |u_n - g_{\mathbf{M}_d}| \omega \leq \frac{n}{d^{n-1}} C_1 \int \omega \leq \frac{n}{d^{n-1}} C_1 .$$

Now, by Theorem 4.2, there exists $C > 0$ depending only on d such that for any $n \geq 1$,

$$\|v_n\|_{L^1(\mathbf{M}_d)} \leq C \cdot n \cdot \text{Area}(\mathbf{M}_d) .$$

Hence

$$\begin{aligned} \int_{\mathbf{M}_d} |u_n - g_{\mathbf{M}_d}|^\beta &= d^{-n+1} \int_{\mathbf{M}_d} |v_n|^\beta \\ &\leq \text{Area}(\mathbf{M}_d) \cdot \frac{C'n}{d^n}, \end{aligned}$$

where C' depends only on d . Combining the estimates gives that there exists a constant C that depends only on d such that:

$$\left| \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} \right| \leq \frac{Cn}{d^n} \|\varphi\|_{\mathcal{C}^2} .$$

This ends the proof. \square

Remark. — 1. Observe that in fact, one can replace in Theorem A the norm $\|\varphi\|_{\mathcal{C}^2}$ by the L^∞ norm of $dd^c\varphi$.
2. On the other hand, as the measures

$$\frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \delta_c \text{ and } \mu_{\mathbf{M}_d}$$

all have supports in \mathbf{M}_d , if θ denotes a cut-off function equal to 1 in a neighborhood of \mathbf{M}_d then for all φ :

$$\frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} = \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \theta(c)\varphi(c) - \int \theta\varphi \mu_{\mathbf{M}_d}.$$

Then, one easily gets that $\|\theta\varphi\|_{\mathcal{C}^2} \leq A\|\varphi\|_{\mathcal{C}^2(K)}$ where $K = \text{supp}(\theta)$ and A is a constant that depends only on θ . Then we have the estimate, for all $\varphi \in \mathcal{C}^2(\mathbb{C})$:

$$\left| \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n)} \varphi(c) - \int \varphi \mu_{\mathbf{M}_d} \right| \leq CA \frac{n}{d^n} \|\varphi\|_{\mathcal{C}^2(K)} .$$

4.4. An application in the spirit of Theorem C

We here want to extend the \mathcal{C}^1 -estimate to non-postcritically finite hyperbolic parameters, i.e. parameters $c \in \mathbf{M}_d$ for which p_c has a cycle of period $k|n$ and multiplier $w^{k/n}$ where $w \in \mathbb{D}$ has been fixed. The precise statement is the following.

Theorem 4.4. — *For any $n \geq 1$ and any $w \in \mathbb{D}$, let*

$$\text{Per}^*(n, w) := \{c \in \mathbb{C} ; \exists z \in \mathbb{C}, p_c^n(z) = z \text{ and } (p_c^n)'(z) = w\} .$$

The set $\text{Per}^(n, w)$ is a finite set of cardinal d^{n-1} (resp. $(d-1)d^{n-1}$) if $w = 0$ (resp. $w \in \mathbb{D}^*$) and there exists a constant $C > 0$ depending only on d such that*

$$\left| \frac{1}{d^{n-1}} \sum_{c \in \text{Per}^*(n, w)} \frac{\varphi(c)}{(d-1)} - \int \varphi \mu_{\mathbf{M}_d} \right| \leq C \left(\max \left\{ \frac{n}{d^n}, \frac{-1}{d^n \log |w|} \right\} \right)^{\frac{1}{2}} \|\varphi\|_{\mathcal{C}^1},$$

for any $w \in \mathbb{D}^$, any $\varphi \in \mathcal{C}_c^1(\mathbb{C})$ and any $n \geq 1$.*

The proof of Theorem 4.4 uses the length-area estimates of Briend and Duval [BD] in a crucial way. Notice that we may as well have used Koebe Distortion estimates, but we again want to present the proof of Theorem 4.4 as a toy model for that of Theorem C.

Notice that, though this estimate looks weaker than the one obtained in Theorem C, it is actually more general. Indeed, the set $\text{Per}^*(n, w)$ as we defined it here is the set of parameters c for which there exists a cycle of period *dividing* n and multiplier *a root of* w . Hence, this is a much bigger set than the one involved in Theorem C.

Proof. — Let us first prove that $\text{Per}^*(n, w)$ is finite and determine its cardinal. If $c \in \text{Per}^*(n, w)$, then p_c has an attracting cycle of exact period $k|n$ and multiplier $w^{k/n} \in \mathbb{D}$. But the set of parameters c admitting a k -cycle of multiplier $t \in \mathbb{D}$ is finite and its cardinal is d_{k-1} if $t = 0$ and $(d-1)d_{k-1}$ otherwise. In particular, $\text{Per}^*(n, w)$ is finite and has cardinal $\sum_{k|n} d_{k-1} = d^{n-1}$ if $w = 0$ and $\sum_{k|n} (d-1)d_{k-1} = (d-1)d^{n-1}$ otherwise (recall that d_n was defined in the introduction).

Pick $\varphi \in \mathcal{C}_c^1(\mathbb{C})$ and let

$$\mu_n := \frac{1}{d^n} \sum_{c \in \text{Per}^*(n)} \delta_c \quad \text{and} \quad \mu_{n,w} := \frac{1}{(d-1)d^n} \sum_{c \in \text{Per}^*(n,w)} \delta_c .$$

Recall that we want to estimate $|\langle \mu_n, \varphi \rangle - \langle \mu_{\mathbf{M}_d}, \varphi \rangle|$. Let us remark that, by a classical interpolation argument, Theorem A gives

$$|\langle \mu_n, \varphi \rangle - \langle \mu_{\mathbf{M}_d}, \varphi \rangle| \leq C_1 \left(\frac{n}{d^n} \right)^{1/2} \|\varphi\|_{\mathcal{C}^1} ,$$

where C_1 depends only on d . Pick $w \in \mathbb{D}^*$ and $n \geq 1$. We are left with estimating $|\langle \mu_n, \varphi \rangle - \langle \mu_{n,w}, \varphi \rangle|$. For any $c_0 \in \text{Per}^*(n)$, let us denote by Ω_{c_0} the hyperbolic component containing c_0 and let c_w^1, \dots, c_w^{d-1} be the $d-1$ parameters $c_w^j \in \Omega_{c_0} \cap \text{Per}^*(n, w)$. Let also Ω'_{c_0} be the open set

$$\Omega'_{c_0} := \bigcap_{|t| < |w|} \text{Per}^*(n, t) \cap \Omega_{c_0} \in \Omega_{c_0} .$$

By Cauchy-Schwarz inequality,

$$\begin{aligned} |\langle \mu_n, \varphi \rangle - \langle \mu_{n,w}, \varphi \rangle| &\leq \frac{1}{(d-1)d^n} \sum_{c_0 \in \text{Per}^*(n)} \sum_{j=1}^{d-1} |\varphi(c_0) - \varphi(c_w^j)| \\ &\leq \frac{1}{d^n} \sum_{c_0 \in \text{Per}^*(n)} \|\varphi\|_{\mathcal{C}^1} \text{Diam}_\beta(\Omega'_{c_0}) \\ &\leq \frac{1}{d^{n/2}} \|\varphi\|_{\mathcal{C}^1} \left(\sum_{c_0 \in \text{Per}^*(n)} (\text{Diam}_\beta(\Omega'_{c_0}))^2 \right)^{1/2} \end{aligned}$$

Let us write $\text{Per}^*(n) = \bigcup_{k|n} \text{Per}(k)$, where $\text{Per}(k)$ denote the set of parameters admitting a super-attracting cycle of exact period k . Moreover, if $k|n$ and $c_0 \in \text{Per}^*(k)$,

$$\Omega'_{c_0} = \rho^{-1} \left(\mathbb{D}(0, |w|^{k/n}) \right) ,$$

where $\rho : \Omega_{c_0} \rightarrow \mathbb{D}$ is the map which, to c , associates the multiplier of the attracting cycle of p_c . Recall that ρ is a $(d-1)$ -branched cover ramifying exactly at c_0 . In particular,

$$\text{mod} \left(\mathbb{D} \setminus \overline{\mathbb{D}(0, |w|^{k/n})} \right) = (d-1) \cdot \text{mod}(\Omega_{c_0} \setminus \overline{\Omega'_{c_0}}) ,$$

whence $\text{mod}(\Omega_{c_0} \setminus \overline{\Omega'_{c_0}}) = \frac{-k}{2\pi n(d-1)} \log |w| \geq \frac{-1}{2\pi n(d-1)} \log |w|$. By Lemma 2.3, we deduce

$$|\langle \mu_n, \varphi \rangle - \langle \mu_{n,w}, \varphi \rangle| \leq \frac{1}{d^{n/2}} \|\varphi\|_{C^1} \left(\sum_{c_0 \in \text{Per}^*(n)} \frac{\tau}{\min(1, \frac{-1}{2\pi n(d-1)} \log |w|)} \text{Area}_\beta(\Omega_{c_0}) \right)^{1/2}.$$

On the other hand, since $\Omega_{c_0} \cap \Omega_{c'_0} = \emptyset$ for $c_0 \neq c'_0 \in \text{Per}^*(n)$ and $\Omega_{c_0} \subset \mathbf{M}_d \subset \overline{D(0, 2)}$,

$$\sum_{\text{Per}^*(n)} \text{Area}_\beta(\Omega_{c_0}) \leq \text{Area}_\beta(\mathbb{D}(0, 2)) = 4\pi.$$

Combined with the above, this gives

$$|\langle \mu_n, \varphi \rangle - \langle \mu_{n,w}, \varphi \rangle| \leq \frac{2\sqrt{\pi\tau}}{d^{n/2}} \|\varphi\|_{C^1} \max \left\{ n, \frac{-2\pi(d-1)}{\log |w|} \right\}^{1/2},$$

which, letting $C := C_1 + 2\pi\sqrt{2(d-1)\tau}$, finally gives the wanted result. \square

5. Initial estimates in the moduli space of polynomials

We aim, here, at giving estimates in the spirit of the one provided by Lemma 4.1. We begin the section with preliminaries on the Moebius function and dynatomic polynomials $P_{m,j}$. We then give estimates using Lemma 3.1 for the potentials of the currents $dd^c \log |P_{m,j}|$ (renormalized by their mass) at suitable parameters.

5.1. Basics

For $0 \leq j \leq d-2$ and $m \geq 1$, following [Si, §4.1], we define the *dynatomic* polynomials $P_{m,j}$ of period m for the critical point c_j by

$$P_{m,j}(c, a) := \prod_{k|m} \left(P_{c,a}^k(c_j) - c_j \right)^{\mu(m/k)},$$

where μ stands for the Moebius function. The degree of $P_{m,j}$ is equivalent to d^m . It does not depend on j so we denote it by d_m as we have the formula:

$$(3) \quad d_m = \sum_{k|m} \mu(m/k) d^k = d^m + \sum_{k|m, k \neq m} \mu(m/k) d^k.$$

Recall that the function $\sigma : \mathbb{N}^* \rightarrow \mathbb{N}^*$ is the *sum of divisors*:

$$\sigma(n) := \sum_{k|n} k.$$

The following is classical (see e.g. [Si, §4.1]).

Lemma 5.1. — *Let $m \geq 1$, $0 \leq j \leq d-2$ and $(c, a) \in \mathbb{C}^{d-1}$. Then $P_{m,j}(c, a) = 0$ if and only if c_j is periodic under iteration of $P_{c,a}$ with period m .*

We also let $\text{Per}_j(m)$ be the algebraic variety

$$\text{Per}_j(m) := \{(c, a) \in \mathbb{C}^{d-1}; P_{m,j}(c, a) = 0\}$$

defined by $(c, a) \in \text{Per}_j(m)$ if and only if the critical point c_j is periodic under iteration of $P_{c,a}$ with exact period m . We shall use in the sequel the following result (see [FG, Theorem 6.1]).

Theorem 5.2 (Favre-Gauthier). — *Let $\underline{m} = (m_0, \dots, m_{d-2})$ be a $(d-1)$ -tuple of pairwise distinct positive integers such that $m_0 \geq 2$. If the hypersurfaces $\{\text{Per}_j(m_j)\}_{0 \leq j \leq d-2}$ intersect at $(c, a) \in \mathbb{C}^{d-1}$, their intersection at (c, a) is smooth and transverse.*

Let $\mathcal{H}_{\underline{n}}$ be the open set of hyperbolic parameters (c, a) for which, for all j , the critical point c_j is in the immediate basin of an attracting cycle of exact period n_j . For $(c_c, a_c) \in \bigcap_j \text{Per}_j(n_j)$, we let Ω_{c_c, a_c} be the connected component of \mathcal{C}_d that contains (c_c, a_c) which is the center of the component Ω_{c_c, a_c} . In the case where all n_j are distinct, it is known that the set Ω_{c_c, a_c} is simply connected and contains only one postcritically finite parameter, the parameter (c_c, a_c) . Notice that, according to Theorem 5.2 and to Bézout Theorem, the set $\mathcal{H}_{\underline{n}}$ has $\text{Card}(\bigcap_j \text{Per}_j(n_j)) = \prod_j d_{n_j}$ distinct hyperbolic components.

5.2. Przytycki's estimates in the space of polynomials

We first give estimates for the functions $d^{-n} \log |P_{c,a}^{n_j}(c_j) - c_j|$ that will enable us to deal with the dynamomic polynomials. The two next lemmas are consequences of [DF, §6] and of Lemma 3.1.

Lemma 5.3. — *There exists a constant $C_1 > 0$ depending only on d such that for any $(c, a) \in \mathbb{C}^{d-1}$ and any j such that $g_{c,a}(c_j) = G(c, a)$, for any $n \geq 1$, then*

$$\frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| - g_{c,a}(c_j) \leq C_1 \frac{1}{d^n}.$$

Proof. — Assume first that $(c, a) \in \mathcal{C}_d$. Then for any j , $g_{c,a}(c_j) = 0 = G(c, a)$. As $|P_{c,a}(z)| \geq \frac{1}{2}|z|^2$ as soon as $|z| \geq 2d^2 \max\{|c|, |a|\}$ (see e.g. [I]), we deduce that for all n :

$$|P_{c,a}^n(c_j) - c_j| \leq |P_{c,a}^n(c_j)| + |c_j| \leq 4d^2 \max_{(c,a) \in \partial \mathcal{C}_d} \{|c|, |a|, 2\}.$$

In particular, there exists a constant C , independent of n such that for all $(c, a) \in \mathcal{C}_d$:

$$\frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| \leq \frac{C}{d^n}.$$

Since $g_{c,a}(c_j) = 0$ in \mathcal{C}_d , the lemma follows in the case where $(c, a) \in \mathcal{C}_d$.

We now consider the case where $(c, a) \in \mathbb{C}^{d-1}$ satisfies $g_{c,a}(c_j) = G(c, a) > 0$ and pick $(c, a) \in \{G = R\}$ with $g_{c,a}(c_j) = G(c, a) > 0$. According to Lemma 2.2, taking R large enough, say $R \geq R_0$, we can assume that for all $n \geq 1$:

$$|P_{c,a}^n(c_j)| \geq 2 \max\{|c|, |a|\} \geq 2|c_j| \geq 1.$$

Hence:

$$\frac{1}{2}|P_{c,a}^n(c_j)| \leq |P_{c,a}^n(c_j) - c_j| \leq \frac{3}{2}|P_{c,a}^n(c_j)|$$

and if $\delta = (d-1)^{-1} \sum_l c_l$, then

$$\frac{1}{2}|P_{c,a}^n(c_j)| \leq |P_{c,a}^n(c_j) - \delta|.$$

We take the logarithm, divide by d^n , and use that $\log |P_{c,a}^n(c_j)| = \log^+ |P_{c,a}^n(c_j)|$:

$$\frac{1}{d^n} \log^+ |P_{c,a}^n(c_j)| - \frac{\log 2}{d^n} \leq \frac{1}{d^n} \log^+ |P_{c,a}^n(c_j) - c_j| \leq \frac{1}{d^n} \log^+ |P_{c,a}^n(c_j)| + \frac{\log \frac{3}{2}}{d^n}.$$

By invariance, $g_{c,a}(c_j) = d^{-n}g_{c,a}(P_{c,a}^n(c_j)) = G(c, a)$ so that $\max\{g_{c,a}(P_{c,a}^n(c_j)), G(c, a)\} = g_{c,a}(P_{c,a}^n(c_j))$. Hence, by [DF, Lemma 6.5], this gives a constant $C'' > 0$ depending only on d and R_0 such that

$$\begin{aligned} g_{c,a}(c_j) &= d^{-n}g_{c,a}(P_{c,a}^n(c_j)) \geq d^{-n}(\log |P_{c,a}^n(c_j) - \delta| - \log 4) \\ &\geq d^{-n}(\log |P_{c,a}^n(c_j) - c_j| - C''). \end{aligned}$$

We now may assume that $0 < G(c, a) = g_{c,a}(c_j) \leq R_0$. We treat two cases separately. Assume first that $|P_{c,a}^n(c_j)| \leq 1/2|c_j|$ so that $|P_{c,a}^n(c_j) - c_j| \leq 3/2|c_j|$. In particular, taking the logarithm, dividing by d^n and using that $\log \leq \log^+$ imply:

$$\frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| \leq \frac{1}{d^n} \log |c_j| + \frac{\log \frac{3}{2}}{d^n} \leq \frac{1}{d^n} G(c, a) + \frac{C''}{d^n}$$

where C'' is a constant that does not depend on (c, a) nor n and where we used that $G(c, a) = \log^+ \max\{|c|, |a|\} + O(1)$. In particular, since $G \geq 0$ and $n \geq 0$,

$$\frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| \leq g_{c,a}(c_j) + \frac{C''}{d^n}.$$

Assume now that $|P_{c,a}^n(c_j)| \geq 1/2|c_j|$ so that $|P_{c,a}^n(c_j) - c_j| \leq 3/2|P_{c,a}^n(c_j)|$. Proceeding as above gives:

$$\frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| \leq \frac{1}{d^n} \log^+ |P_{c,a}^n(c_j)| + \frac{\log \frac{3}{2}}{d^n}.$$

Using Lemma 2.2 implies that there is a constant that depends only on d and R_0 , but neither on (c, a) nor on n that we still denote C'' such that:

$$\frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| \leq g_{c,a}(c_j) + \frac{C''}{d^n}.$$

This concludes the proof in the case where $0 < g_{c,a}(c_j) = G(c, a) \leq R_0$. \square

The second lemma we will need is the following.

Lemma 5.4. — *There exists a constant $C_2 \geq C_1 > 0$ depending only on d such that for any $(c, a) \in \mathbb{C}^{d-1}$ such that either $g_{c,a}(c_j) = G(c, a) > 0$ or $(c, a) \in \mathcal{C}_d \cap \text{supp}(T_j)$ and any $n \geq 1$, then:*

$$\frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| - g_{c,a}(c_j) \geq -C_1 \frac{n}{d^n}.$$

Proof. — Consider first the case where $(c, a) \in \mathcal{C}_d \cap \text{supp}(T_j)$. Hence c_j is active at (c, a) . By Lemma 3.1, there exist a universal constant $0 < \kappa < 1$ and a continuous function $C(c, a) > 1$ such that:

$$|P_{c,a}^n(c_j) - c_j| \geq \frac{|P_{c,a}^n(c_j) - c_j|}{\sqrt{1 + |P_{c,a}^n(c_j)|^2} \cdot \sqrt{1 + |c_j|^2}} = d_{\mathbb{P}^1}(P_{c,a}^n(c_j), c_j) \geq \frac{\kappa}{C(c, a)^n}.$$

Hence, if $C := \max_{(c,a) \in \partial \mathcal{C}_d} C(c, a) > 1$, then

$$\begin{aligned} \frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| &\geq -\frac{n}{d^n} \log C + \frac{1}{d^n} \log \kappa \\ (4) \quad \frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j| &\geq -\frac{n}{d^n} \log \left(\frac{C}{\kappa} \right) + g_{c,a}(c_j) \end{aligned}$$

for any $n \in \mathbb{N}^*$, since $g_{c,a}(c_j) = 0$ on $\partial \mathcal{C}_d$. This is the expected result in the case where $(c, a) \in \mathcal{C}_d \cap \text{supp}(T_j)$.

Assume now that (c, a) satisfies $G(c, a) = g_{c,a}(c_j) = R > 0$. Consider the compact domain $\Omega_R := \{G \leq R\}$. As above, since $G(c, a) = \log^+ \max\{|c|, |a|\} + O(1)$, according to Lemma 2.2, taking $R \geq R_0$ large enough, we can assume that for all $n \geq 1$:

$$|P_{c,a}^n(c_j)| \geq 2 \max\{|c|, |a|\} \geq 2|c_j| \geq 1.$$

Hence:

$$|P_{c,a}^n(c_j) - c_j| \geq \frac{1}{2}|P_{c,a}^n(c_j)|.$$

Up to increasing R_0 , combining $G(c, a) = \log^+ \max\{|c|, |a|\} + O(1)$ with [DF, Lemma 6.4] gives for (c, a) , providing that $R \geq R_0$, for any $n \geq 1$:

$$(5) \quad d^{-n} \log |P_{c,a}^n(c_j)| = d^{-n} \log^+ |P_{c,a}^n(c_j)| \geq g_{c,a}(c_j) + O(d^{-n}),$$

where the O is independent of n and (c, a) . This gives a constant C'' (independent of $R \geq R_0$) such that for all $n \geq 1$:

$$g_{c,a}(c_j) - C''(d^{-n}) \leq \frac{1}{d^n} \log |P_{c,a}^n(c_j) - c_j|.$$

The only case that remains to consider is when $0 < G(c, a) = g_{c,a}(c_j) < R_0$. Pick such a parameter (c, a) . Notice that it belongs to the compact set Ω_{R_0} .

Claim. — For all $(c, a) \in \Omega_{R_0}$, all $n \geq 1$ and all $0 \leq j \leq d-2$, we have

$$d_{\mathbb{P}^1}(P_{c,a}^{n_j}(c_j), c_j) \geq \frac{\kappa}{C^n},$$

where $C > 1$ depends only on d and R_0 .

Let us first finish the proof of Lemma 5.4. As a consequence,

$$\frac{|P_{c,a}^n(c_j) - c_j|}{\sqrt{1 + |P_{c,a}^n(c_j)|^2} \cdot \sqrt{1 + |c_j|^2}} = d_{\mathbb{P}^1}(P_{c,a}^{n_j}(c_j), c_j) \geq \frac{\kappa}{C^n}.$$

Taking the logarithm, dividing by d^n and using Lemma 2.2 uniformly on Ω_{R_0} imply:

$$\begin{aligned} -\frac{n}{d^n} \log \frac{C}{\kappa} &\leq \frac{1}{d^n} |P_{c,a}^n(c_j) - c_j| - \frac{1}{2d^n} \log \sqrt{1 + |P_{c,a}^n(c_j)|^2} - \frac{1}{2d^n} \log (1 + |c_j|^2) \\ &\leq \frac{1}{d^n} |P_{c,a}^n(c_j) - c_j| - g_{c,a}(c_j) + C' \frac{1}{2d^n}, \end{aligned}$$

where C' is yet another constant that depends on d and R_0 but neither on (c, a) nor on n . This concludes the proof. \square

Proof of the Claim. — Let us first set

$$C := \sup_{(c,a) \in \Omega_{R_0}} \sup_{z \in \mathbb{P}^1} P_{c,a}^\#(z) > 1.$$

We proceed by contradiction, assuming that for some $n \geq 2$, $d_{\mathbb{P}^1}(P_{c,a}^{n_j}(c_j), c_j) < \frac{\kappa}{C^n}$. As seen in the proof of Lemma 3.1, since c_j lies in the attracting basin of ∞ of $P_{c,a}$, we have

$$\frac{1}{\sqrt{1 + |c_j|^2}} \leq d_{\mathbb{P}^1}(\infty, c_j) \leq 2d_{\mathbb{P}^1}(c_j, P_{c,a}^n(c_j)) < \frac{2\kappa}{C^n}.$$

This may be rephrased as $C^2 < C^{2n} < 4\kappa^2 (1 + C(R_0)^2)$ where $C(R_0) = \max_{(c,a) \in \Omega_{R_0}} |c_j|$. Up to increasing C , we may assume $C^2 \geq 2\kappa^2 (1 + C(R_0)^2)$, which is a contradiction. \square

5.3. Estimates for the dynatomic polynomials

In the sequel, we shall use the notation

$$g_{n,j}(c, a) := \frac{1}{d_n} \log |P_{n,j}(c, a)|, \quad (c, a) \in \mathbb{C}^{d-1}, \quad n \geq 2.$$

The following is a key to the proof of Theorem B.

Proposition 5.5. — *There exists a constant $C \geq 1$ depending only on d such that for all $n \geq 1$, $(c, a) \in \mathbb{C}^{d-1}$ and any $0 \leq j \leq d-2$ such that $g_{c,a}(c_j) = G(c, a)$. Then:*

$$g_{n,j}(c, a) - g_{c,a}(c_j) \leq C \frac{\sigma(n)}{d_n}.$$

In particular, for any $0 \leq j \leq d-2$ if $(c, a) \in \mathcal{C}_d$, we have:

$$g_{n,j}(c, a) \leq C \frac{\sigma(n)}{d_n}.$$

Proof. — By definition of $P_{n,j}$, we have:

$$\log |P_{n,j}(c, a)| = \sum_{k|n} \mu(n/k) \log \left| P_{c,a}^k(c_j) - c_j \right|,$$

Hence, dividing by d_n and using (3), we deduce:

$$\frac{1}{d_n} \log |P_{n,j}(c, a)| - g_{c,a}(c_j) = \sum_{k|n} \frac{d^k}{d_n} \mu(n/k) \left(\frac{1}{d^k} \log \left| P_{c,a}^k(c_j) - c_j \right| - g_{c,a}(c_j) \right).$$

Let $(c, a) \in \mathbb{C}^{d-1}$ be such that either $g_{c,a}(c_j) = G(c, a) > 0$ or $(c, a) \in \mathcal{C}_d \cap \text{supp}(T_j)$. Then Lemmas 5.3 and 5.4 imply that:

$$\left| \frac{1}{d^k} \log \left| P_{c,a}^k(c_j) - c_j \right| - g_{c,a}(c_j) \right| \leq C \frac{k}{d^k},$$

for all k . Hence, since $\mu(n_j/k) = \pm 1$:

$$\left| g_{c,a}(c_j) - \frac{1}{d_n} \log |P_{n,j}(c, a)| \right| \leq C \frac{1}{d_n} \sum_{k|n} k.$$

Hence by the definition of the σ function, we have:

$$(6) \quad \left| g_{c,a}(c_j) - \frac{1}{d_n} \log |P_{n,j}(c, a)| \right| \leq C \frac{\sigma(n)}{d_n}.$$

This ends the proof in the case where $g_{c,a}(c_j) = G(c, a) > 0$ or $(c, a) \in \mathcal{C}_d \cap \text{supp}(T_j)$.

Finally, as the Shilov boundary $\partial_S \mathcal{C}_d$ of \mathcal{C}_d is contained in $\cap_j \text{supp} T_j$, for all $(c, a) \in \partial_S \mathcal{C}_d$, all $n \geq 2$ and all $0 \leq j \leq d-2$, we have

$$\frac{1}{d_n} \log |P_{n,j}(c, a)| \leq C \frac{\sigma(n)}{d_n}.$$

in other words, for all $(c, a) \in \partial_S \mathcal{C}_d$, all $n \geq 2$ and all $0 \leq j \leq d-2$,

$$|P_{n,j}(c, a)| \leq \exp(C\sigma(n)).$$

By the maximum principle, the bound extends to \mathcal{C}_d , which gives

$$|P_{n,j}(c, a)| \leq \exp(C\sigma(n)),$$

for any $(c, a) \in \mathcal{C}_d$, $n \geq 2$ and $0 \leq j \leq d-2$. Taking the logarithm and dividing by d_n finishes the proof. \square

Observe that in the above proof, we proved the following crucial estimate (see (6)):

Proposition 5.6. — *Let $C \geq 1$ be the constant given by Proposition 5.5 and let $n \geq 1$. Assume that either $g_{c,a}(c_j) = G(c, a) > 0$, or $(c, a) \in \mathcal{C}_d \cap \text{supp}(T_j)$. Then*

$$g_{n,j}(c, a) - g_{c,a}(c_j) \geq -C \frac{\sigma(n)}{d_n} .$$

5.4. Locating “bad” parameters

We shall now give two basic consequences of the above estimates which will be crucial to our aim. Namely, the following is a consequence of Propositions 5.5 and 5.6.

Corollary 5.7. — *Let $C \geq 1$ be the constant given by Proposition 5.6. Then, for any $(d-1)$ -tuple $\underline{n} = (n_0, \dots, n_{d-2})$ of positive integers,*

$$\mathcal{B}_{\underline{n}} := \bigcap_{j=0}^{d-2} \left\{ |g_{n_j,j}(c, a) - g_{c,a}(c_j)| > C \frac{\sigma(n_j)}{d_{n_j}} \right\} = \bigcap_{j=0}^{d-2} \left\{ |P_{n_j,j}| < e^{-C\sigma(n_j)} \right\} \subset \mathcal{H}_{\underline{n}} \subset \mathcal{C}_d .$$

Proof. — Let us first prove that the set

$$\mathcal{B}_{\underline{n}} = \bigcap_{j=0}^{d-2} \left\{ |g_{n_j,j}(c, a) - g_{c,a}(c_j)| > C \frac{\sigma(n_j)}{d_{n_j}} \right\}$$

is contained in the connectedness locus \mathcal{C}_d . If not, there exists $(c, a) \in \mathcal{B}_{\underline{n}} \setminus \mathcal{C}_d$. Then there exists $0 \leq j \leq d-2$ such that $G(c, a) = g_{c,a}(c_j) > 0$. By Propositions 5.5 and 5.6, we have

$$-C \frac{\sigma(n_j)}{d_{n_j}} \leq g_{n_j,j}(c, a) - g_{c,a}(c_j) \leq C \frac{\sigma(n_j)}{d_{n_j}} ,$$

which is a contradiction. The same argument implies in fact that $\mathcal{B}_{\underline{n}}$ is contained in the interior of \mathcal{C}_d (use the case $(c, a) \in \mathcal{C}_d \cap \text{supp}(T_j)$ in Proposition 5.6). Now from Proposition 5.5, we have that for all $(c, a) \in \mathcal{C}_d$ and all $0 \leq j \leq d-2$:

$$g_{n_j,j}(c, a) \leq C \frac{\sigma(n_j)}{d_{n_j}} .$$

As $g_{c,a}(c_j) = 0$ in \mathcal{C}_d , this implies:

$$\mathcal{B}_{\underline{n}} = \bigcap_{j=0}^{d-2} \left\{ g_{n_j,j} < -C \frac{\sigma(n_j)}{d_{n_j}} \right\} \cap \mathcal{C}_d .$$

This can be rewritten as:

$$\bigcap_{j=0}^{d-2} \left\{ |P_{n_j,j}| < e^{-C\sigma(n_j)} \right\} \cap \mathcal{C}_d = \bigcap_{j=0}^{d-2} \left\{ |g_{n_j,j}(c, a) - g_{c,a}(c_j)| > C \frac{\sigma(n_j)}{d_{n_j}} \right\} .$$

Arguing as above, we see that the set $\bigcap_{j=0}^{d-2} \{|P_{n_j,j}| < e^{-C\sigma(n_j)}\}$ is contained in \mathcal{C}_d : if not, for some (c, a) in that set such that $g_{c,a}(c_j) = G(c, a) > 0$, we have

$$g_{n_j,j}(c, a) - g_{c,a}(c_j) < g_{n_j,j}(c, a) < -C \frac{\sigma(n_j)}{d_{n_j}} ,$$

which contradicts Proposition 5.6.

It remains to prove that $\mathcal{B}_{\underline{n}} \subset \mathcal{H}_{\underline{n}}$. Take a stable component $U \subset \mathcal{C}_d$ which is not contained in $\mathcal{H}_{\underline{n}}$. Consider the plurisubharmonic function:

$$\phi_{\underline{n}} := \max_j (\log |P_{n_j, j}| + C\sigma(n_j)).$$

Then its Monge-Ampère $(dd^c \phi_{\underline{n}})^{d-1}$ is 0 on U by hypothesis (see e.g. [Dem]). On the other hand, by Proposition 5.6, it is non negative in ∂U ($\partial U \subset \partial \mathcal{C}_d \subset \cup_j \mathcal{C}_d \cap \text{supp}(T_j)$). The comparison principle of Bedford and Taylor [BT] implies that it is non negative on U . Hence $U \cap \mathcal{B}_{\underline{n}} = \emptyset$. \square

Here is a consequence of Corollary 5.7.

Corollary 5.8. — *For any $(d-1)$ -tuple $\underline{n} = (n_0, \dots, n_{d-2})$ of pairwise distinct positive integers with $n_0 \geq 2$ and any connected component U of $\mathcal{B}_{\underline{n}}$, the map*

$$P_{\underline{n}} := (P_{n_0, 0}, \dots, P_{n_{d-2}, d-2}) : \mathbb{C}^{d-1} \longrightarrow \mathbb{C}^{d-1}$$

is a biholomorphism from U to the polydisk $\mathbb{D}_{\underline{n}}^{d-1} := \prod_j \mathbb{D}(0, e^{-C\sigma(n_j)})$.

Proof. — By Corollary 5.7 and by the maximum principle, $P_{\underline{n}}$ maps U to $\mathbb{D}_{\underline{n}}^{d-1}$ surjectively. Moreover, by definition of U , the map $P_{\underline{n}}$ is also proper on U , hence is a finite branched cover from U to $\mathbb{D}_{\underline{n}}^{d-1}$. Let (c_c, a_c) be the unique postcritically finite parameter of U . By Theorem 5.2, the map $P_{\underline{n}}$ is a local biholomorphism at (c_c, a_c) and, since $P_{\underline{n}}^{-1}\{0\} = \{(c_c, a_c)\}$, the degree of $P_{\underline{n}}$ is 1. \square

6. Distribution of postcritically finite hyperbolic polynomials

In the present section, we prove Theorem B. We adapt the strategy we used in the unicritical family $(p_c)_{c \in \mathbb{C}}$ in Section 4 to the present situation. We use all along the present section the notations introduced in Sections 5.1 & 5.2. We assume from now on that $\underline{n} = (n_0, \dots, n_{d-2})$ is a $(d-1)$ -tuple of pairwise distinct integers with $n_0 \geq 2$. We also let $D_{\underline{n}} > 0$ be the integer

$$D_{\underline{n}} := \prod_j d_{n_j}.$$

6.1. Estimates along specific curves

Recall that we denoted by $\mathcal{H}_{\underline{n}}$ the union of hyperbolic components of \mathcal{C}_d intersecting $\bigcap_j \text{Per}_j(n_j)$. Let us first introduce some notations. For $\delta \in \mathbb{C}$, $n \geq 1$ and $0 \leq j \leq d-2$, we let

$$g_{n_j, j}^\delta := \frac{1}{d_{n_j}} \log |P_{n_j, j} - \delta|, \text{ and } T_{n_j, j}^\delta := dd^c g_{n_j, j}^\delta.$$

For the rest of the section, we also shall write for $0 \leq j \leq d-2$,

$$g_j(c, a) := g_{c, a}(c_j), \quad T_j := dd^c g_j, \quad g_{n_j, j} := \frac{1}{d_{n_j}} \log |P_{n_j, j}| \text{ and } T_{n_j, j} := dd^c g_{n_j, j}.$$

The key step of our proof can be summarized in the following proposition.

Theorem 6.1. — *Let $C \geq 1$ be the constant given by Proposition 5.6. Let $0 \leq j \leq d-2$ and, for all $0 \leq \ell \leq j-1$, pick $\delta_\ell \in \mathbb{D}(0, \exp(-C\sigma(n_j)))$. There exists a constant $C' \geq 1$ which depends only on d such that for any $\varphi \in \mathcal{C}_c^2(\mathbb{C}^{d-1})$, one has*

$$\left| \left\langle \bigwedge_{\ell=0}^{j-1} T_{n_\ell, \ell}^{\delta_\ell} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k}, \varphi \right\rangle \right| \leq C' \|\varphi\|_{\mathcal{C}^2} \frac{\sigma(n_j)}{d_{n_j}}.$$

Proof. — First, the support of $\bigwedge_{\ell < j} T_{n_\ell, \ell}^{\delta_\ell} \wedge \bigwedge_{k > j} T_{n_k, k}$ is the algebraic curve

$$S := \bigcap_{\ell=0}^{j-1} \{P_{n_\ell, \ell} = \delta_\ell\} \cap \bigcap_{k=j+1}^{d-2} \{P_{n_k, k} = 0\}.$$

Pick $\varphi \in \mathcal{C}_c^2(\mathbb{C}^{d-1})$. Then

$$\begin{aligned} I(\varphi) &:= \left\langle \bigwedge_{\ell=0}^{j-1} T_{n_\ell, \ell}^{\delta_\ell} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k}, \varphi \right\rangle \\ &= \int (g_{n_j, j} - g_j) dd^c \varphi \wedge \bigwedge_{\ell=0}^{j-1} T_{n_\ell, \ell}^{\delta_\ell} \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k} \\ &= \frac{d_{n_j}}{D_{\underline{n}}} \int_S (g_{n_j, j} - g_j) dd^c \varphi. \end{aligned}$$

As φ is \mathcal{C}^2 , we have (up to a constant that depends on the choice of the \mathcal{C}^2 -norm) that $\|\varphi\|_{\mathcal{C}^2} \omega \pm dd^c \varphi \geq 0$, where ω is the Fubini-Study form on \mathbb{P}^{d-1} , hence we can write

$$|I(\varphi)| \leq \|\varphi\|_{\mathcal{C}^2} \frac{d_{n_j}}{D_{\underline{n}}} \int_S |g_{n_j, j} - g_j| \omega.$$

According to Corollary 5.7, the set $E := S \cap \{|g_{n_j, j} - g_j| > C\sigma(n_j)/d_{n_j}\}$ is contained in $S \cap \mathcal{H}_{\underline{n}}$ and coincides with $\{(c, a) \in S; |P_{n_j, j}| < e^{-C\sigma(n_j)}\}$. Hence

$$\int_S |g_{n_j, j} - g_j| \omega \leq \int_E |g_{n_j, j}| \omega + C \frac{\sigma(n_j)}{d_{n_j}} \int_S \omega.$$

By Corollary 5.8, the curve S is smooth in the open set $\mathcal{B}_{\underline{n}}$. By the maximum principle, E is a finite union of $D_{\underline{n}}$ topological disks (see § 5.1), which are bounded since contained in \mathcal{C}_d . We now let $u := d_{n_j} g_{n_j, j} = \log |P_{n_j, j}|$. Then $dd^c(u|_S) = \sum \alpha_{z_0} \delta_{z_0}$, for some collection $\{\alpha_{z_0}\}$ of positive integers, where the sum ranges over $\text{Per}_j(n_j) \cap S$. Moreover, by Bézout, we see that the finite measure $dd^c u \wedge S$ has mass $D_{\underline{n}}$. In particular, $dd^c(u|_S) = \sum \delta_{z_0}$.

Let now Ω be a connected component of $E = S \cap \{g_{n_j, j} < -C\sigma(n_j)/d_{n_j}\}$. According to Theorem 3.2, there exists $C_1 > 0$ universal such that

$$\int_\Omega |u| \omega \leq \int_\Omega |u| \beta \leq C_1 \max\{\|u\|_{L^\infty(\partial\Omega)}, 1\} \text{Area}_\beta(\Omega).$$

Now, since $g_{n_j, j} = -C \frac{\sigma(n_j)}{d_{n_j}}$ on $\partial\Omega$, we have $\|u\|_{L^\infty(\partial\Omega)} = C\sigma(n_j) \geq 1$ and we find

$$\int_\Omega |g_{n_j, j}| \omega \leq C_1 C \frac{\sigma(n_j)}{d_{n_j}} \text{Area}_\beta(\Omega).$$

Since $\mathcal{C}_d \in \mathbb{B}(0, 16\sqrt{d-1})$, see e.g. [DF, Proof of Proposition 6.3], we can find $C_2 > 0$ depending only on d such that $\beta \leq C_2\omega$ on \mathcal{C}_d . In particular, $\text{Area}_\beta(\Omega) \leq C_2\text{Area}_\omega(\Omega)$. Summing up on the components Ω of E , this gives

$$\int_E |g_{n_j,j}| \omega \leq C_2 C_1 C \frac{\sigma(n_j)}{d_{n_j}} \int_S \omega .$$

Finally, by Bézout, the integral $\int_S \omega$ is equal the degree of the curve, i.e. $D_{\underline{n}}/d_{n_j}$ (see e.g. [C]). Summarizing what we did to now, we find

$$|I(\varphi)| \leq \|\varphi\|_{\mathcal{C}^2} (1 + C_2 C_1) C \frac{\sigma(n_j)}{d_{n_j}} ,$$

which gives the wanted result since $C' := (1 + C_2 C_1) C$ depends only on d . \square

6.2. Intermediate estimates

We introduce some notations. For all $0 \leq j \leq d-2$, we let

$$\tilde{g}_{n_j,j} := \max \left\{ g_{n_j,j}, -2C \frac{\sigma(n_j)}{d_{n_j}} \right\} , \quad \text{and} \quad \tilde{T}_{n_j,j} := dd^c \tilde{g}_{n_j,j} .$$

Let us set $\delta_j(\theta) := e^{-2C\sigma(n_j)+i\theta}$, $\theta \in \mathbb{R}$. It is classical that

$$\tilde{g}_{n_j,j} = \frac{1}{2\pi d_{n_j}} \int_0^{2\pi} \log |P_{n_j,j} - \delta_j(\theta)| d\theta , \quad \text{and} \quad \tilde{T}_{n_j,j} = \frac{1}{2\pi} \int_0^{2\pi} T_{n_j,j}^{\delta_j(\theta)} d\theta .$$

We now want to prove the following technical step of the proof of Theorem B.

Lemma 6.2. — *Let C be the constant given by Proposition 5.6. Then, for any $j \geq 1$, any $0 \leq r \leq j-1$ and any $\varphi \in \mathcal{C}_c^2(\mathbb{C}^{d-1})$, we have*

$$\left| \left\langle \bigwedge_{\ell=0}^{r-1} T_\ell \wedge (T_r - \tilde{T}_{n_r,r}) \wedge \bigwedge_{q=r+1}^{j-1} \tilde{T}_{n_q,q} \wedge (T_{n_j,j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k,k}, \varphi \right\rangle \right| \leq 4C \frac{\sigma(n_r)}{d_{n_r}} \|\varphi\|_{\mathcal{C}^2} .$$

Proof. — Pick $\varphi \in \mathcal{C}_c^2(\mathbb{C}^{d-1})$ and let

$$\begin{aligned} L(\varphi) &:= \left\langle \bigwedge_{\ell=0}^{r-1} T_\ell \wedge \bigwedge_{q=r}^{j-1} (T_q - \tilde{T}_{n_q,q}) \wedge (T_{n_j,j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k,k}, \varphi \right\rangle \\ &= \int (g_r - \tilde{g}_{n_r,r}) dd^c \varphi \wedge \bigwedge_{\ell=0}^{r-1} T_\ell \wedge \bigwedge_{q=r+1}^{j-1} \tilde{T}_{n_q,q} \wedge (T_{n_j,j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k,k} . \end{aligned}$$

As φ is \mathcal{C}^2 , we have (up to a constant that depends on the choice of the \mathcal{C}^2 -norm) that $\|\varphi\|_{\mathcal{C}^2} \omega \pm dd^c \varphi \geq 0$, where ω is the Fubini-Study form on \mathbb{P}^{d-1} , hence we can write

$$|L(\varphi)| \leq \|\varphi\|_{\mathcal{C}^2} \int |g_r - \tilde{g}_{n_r,r}| \omega \wedge \bigwedge_{\ell=0}^{r-1} T_\ell \wedge \bigwedge_{q=r+1}^{j-1} \tilde{T}_{n_q,q} \wedge (T_{n_j,j} + T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k,k} .$$

Each of the $(1,1)$ -currents T_ℓ , $T_{n_\ell,\ell}$ and $\tilde{T}_{n_\ell,\ell}$ having projective mass 1, by Bézout, we find

$$(7) \quad |L(\varphi)| \leq 2\|\varphi\|_{\mathcal{C}^2} \sup |g_r - \tilde{g}_{n_r,r}| ,$$

where the supremum is taken over the support of the current

$$W := \bigwedge_{\ell=0}^{r-1} T_\ell \wedge \bigwedge_{q=r+1}^{j-1} \tilde{T}_{n_q, q} \wedge (T_{n_j, j} + T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k}.$$

Owing to (7), the next lemma ends the proof. \square

Lemma 6.3. — *Let $C > 0$ be as above. For any $(c, a) \in \text{supp}(W)$, we have*

$$|\tilde{g}_{n_r, r}(c, a) - g_r(c, a)| \leq 2C \frac{\sigma(n_r)}{d_{n_r}}.$$

Proof. — On $\text{supp}(W)$, we have $g_\ell = 0$ for all $0 \leq \ell \leq r-1$ and all $j \leq \ell \leq d-2$. Moreover, for any $r+1 \leq \ell \leq j-1$, we have $|P_{n_\ell, \ell}| = e^{-2C\sigma(n_\ell)}$. Let $(c, a) \in W$ such that $g_r(c, a)$ is not maximal. This implies the existence of $r+1 \leq \ell \leq j-1$ such that $G(c, a) = g_\ell(c, a) > g_r(c, a) \geq 0$. Then, by Proposition 5.6:

$$-2C \frac{\sigma(n_\ell)}{d_{n_\ell}} - g_\ell(c, a) = g_{n_\ell, \ell}(c, a) - g_\ell(c, a) \geq -C \frac{\sigma(n_\ell)}{d_{n_\ell}},$$

which implies $g_\ell < 0$. This is a contradiction so we know that $g_r(c, a)$ is maximal.

Assume first that $(c, a) \in \text{supp}(W)$ satisfies $g_r(c, a) = 0$. By Proposition 5.5:

$$g_{n_r, r}(c, a) \leq C \frac{\sigma(n_r)}{d_{n_r}} \quad \text{so} \quad \tilde{g}_{n_r, r}(c, a) \leq C \frac{\sigma(n_r)}{d_{n_r}}.$$

Hence:

$$-2C \frac{\sigma(n_r)}{d_{n_r}} \leq \tilde{g}_{n_r, r}(c, a) = \tilde{g}_{n_r, r}(c, a) - g_r(c, a) \leq C \frac{\sigma(n_r)}{d_{n_r}}.$$

We now consider $(c, a) \in \text{supp}(W)$ with $G(c, a) = g_r(c, a) > 0$. By Propositions 5.5 and 5.6, we find

$$|g_{n_r, r}(c, a) - g_r(c, a)| \leq C \frac{\sigma(n_r)}{d_{n_r}}.$$

In particular, $g_{n_r, r}(c, a) = \tilde{g}_{n_r, r}(c, a)$ and the estimate follows. \square

6.3. Speed of convergence: proof of Theorem B

We are now in position to prove Theorem B. Pick $\varphi \in \mathcal{C}_c^2(\mathbb{C}^{d-1})$. We want to estimate

$$\Lambda(\varphi) := \langle \mu_{\underline{n}} - \mu_{\text{bif}}, \varphi \rangle = \left\langle \bigwedge_{j=0}^{d-2} T_{n_j, j} - \bigwedge_{j=0}^{d-2} T_j, \varphi \right\rangle.$$

We shall decompose $\mu_{\underline{n}} - \mu_{\text{bif}}$ into several pieces:

$$(8) \quad \mu_{\underline{n}} - \mu_{\text{bif}} = \sum_{j=0}^{d-2} \left(\bigwedge_{\ell=0}^{j-1} T_\ell \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k} \right) = \sum_{j=0}^{d-2} S_j.$$

For $j \geq 1$, using $T_j = \tilde{T}_{n_j, j} + (T_j - \tilde{T}_{n_j, j})$, we rewrite the j -th term S_j of (8) and find

$$S_j = \bigwedge_{\ell=0}^{j-1} \tilde{T}_{n_\ell, \ell} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k} \\ + \sum_{r=0}^{j-2} \left(\bigwedge_{\ell=0}^{r-1} T_\ell \wedge (T_r - \tilde{T}_{n_r, r}) \wedge \bigwedge_{q=r+1}^{j-1} \tilde{T}_{n_q, q} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k} \right).$$

According to Lemma 6.2, for $j \geq 1$, we find

$$|\langle S_j, \varphi \rangle| \leq \left| \left\langle \bigwedge_{\ell=0}^{j-1} \tilde{T}_{n_\ell, \ell} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k}, \varphi \right\rangle \right| + 4C \sum_{r=0}^{j-1} \frac{\sigma(n_r)}{d_{n_r}} \|\varphi\|_{\mathcal{C}^2}.$$

We now want to give an estimate for

$$J_j := \left\langle \bigwedge_{\ell=0}^{j-1} \tilde{T}_{n_\ell, \ell} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k}, \varphi \right\rangle.$$

To do so, we shall use the decomposition $\tilde{T}_{n_\ell, \ell} = (2\pi)^{-1} \int_{[0, 2\pi]} T_{n_\ell, \ell}^{\delta_\ell(\theta)} d\theta$ and Fubini:

$$J_j = \frac{1}{(2\pi)^{j-1}} \int_{[0, \pi]^{j-1}} \left\langle \bigwedge_{\ell=0}^{j-1} T_{n_\ell, \ell}^{\delta_\ell(\theta_\ell)} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k}, \varphi \right\rangle d\theta_0 \cdots d\theta_{j-1}.$$

By Theorem 6.1, there exists a constant $C' > 0$ depending only on d such that

$$|J_j| \leq \frac{1}{(2\pi)^{j-1}} \int_{[0, \pi]^{j-1}} \left| \left\langle \bigwedge_{\ell=0}^{j-1} T_{n_\ell, \ell}^{\delta_\ell(\theta_\ell)} \wedge (T_{n_j, j} - T_j) \wedge \bigwedge_{k=j+1}^{d-2} T_{n_k, k}, \varphi \right\rangle \right| d\theta_0 \cdots d\theta_{j-1} \\ \leq \frac{1}{(2\pi)^{j-1}} \int_{[0, \pi]^{j-1}} C' \frac{\sigma(n_j)}{d_{n_j}} \|\varphi\|_{\mathcal{C}^2} d\theta_0 \cdots d\theta_{j-1} = C' \frac{\sigma(n_j)}{d_{n_j}} \|\varphi\|_{\mathcal{C}^2}.$$

All we have done so far summarizes as follows

$$\sum_{j=1}^{d-2} |\langle S_j, \varphi \rangle| \leq \sum_{j=1}^{d-2} \left(C' \frac{\sigma(n_j)}{d_{n_j}} + 4C \sum_{r=0}^{j-1} \frac{\sigma(n_r)}{d_{n_r}} \right) \|\varphi\|_{\mathcal{C}^2}.$$

All which is left to do is to estimate $\langle S_0, \varphi \rangle$. By Theorem 6.1, we have

$$|\langle S_0, \varphi \rangle| \leq C' \frac{\sigma(n_0)}{d_{n_0}} \|\varphi\|_{\mathcal{C}^2},$$

and we have finally proven

$$|\Lambda(\varphi)| \leq \sum_{j=0}^{d-2} |\langle S_j, \varphi \rangle| \leq \left((d-1)C' + 4C \sum_{j=1}^{d-2} (j-1) \right) \max_{0 \leq j \leq d-2} \left(\frac{\sigma(n_j)}{d_{n_j}} \right) \|\varphi\|_{\mathcal{C}^2},$$

which ends the proof, since $(d-1)C' + 4C \sum_{j=1}^{d-2} (j-1) = (d-1)C' + 2C(d-1)(d-2)$ depends only on d .

7. Distribution of polynomials with prescribed multipliers

The aim of this section is to derive Theorem C and Theorem D from Theorem B. We begin with the proof of Theorem C which is based on the same idea as that of [FG, Theorem 3]. For our purpose, we have to refine the techniques used in [FG]. We then prove Theorem D using DSH techniques (see e.g. [DTV]).

7.1. Distribution of polynomials with $(d - 1)$ attracting cycles

Pick any $(d - 1)$ -tuple $\underline{n} = (n_0, \dots, n_{d-2})$ of pairwise distinct positive integers with $n_0 \geq 2$.

Let Ω be hyperbolic component of \mathcal{C}_d . Assume that for any $(c, a) \in \Omega$, the polynomial $P_{c,a}$ admits $(d - 1)$ distinct attracting cycles C_0, \dots, C_{d-2} of respective exact periods n_0, \dots, n_{d-2} and let $\rho_i(c, a) \in \mathbb{D}$ be the multiplier of the attracting cycle C_i . We call the map $\rho := (\rho_0, \dots, \rho_{d-2}) : \Omega \rightarrow \mathbb{D}^{d-1}$ the *multipliers map* of the component Ω where \mathbb{D}^{d-1} is the unit polydisk in \mathbb{C}^{d-1} . It is known that it is a biholomorphism (see § 5.1).

Recall that we denoted $D_{\underline{n}} := \prod_j d_{n_j}$ and that there exists polynomials

$$p_n : \mathbb{C}^{d-1} \times \mathbb{C} \longrightarrow \mathbb{C}$$

such that $p_n(c, a, w) = 0$ if and only if one of the following occurs (see e.g. [FG, §6]):

- if $w \neq 1$, $P_{c,a}$ has a cycle of exact period n and multiplier w ,
- if $w = 1$, there exists $k|n$ such that $P_{c,a}$ has a cycle of exact period k and multiplier ρ a primitive n/k -root of unity.

Moreover, we have

- $\deg_{c,a} p_n(\cdot, w) = (d - 1)d_n$ for any $w \in \mathbb{C}$. Indeed, $\deg_{c,a} p_n(\cdot, w)$ is independent of w , we can write $[p_n(\cdot, 0) = 0] = \sum_j [\text{Per}_j(n)]$ and $\text{Per}_j(n)$ has degree d_n ,
- $\deg_w p_n(c, a, \cdot) = d_n/n$ for any $(c, a) \in \mathbb{C}^{d-1}$ (see [BB3, §2.1]).

Let $\text{Per}(n, w) = \{(c, a) \in \mathbb{C}^{d-1}; p_n(c, a, w) = 0\}$. We have the following.

Proposition 7.1. — *Let $w_0, \dots, w_{d-2} \in \mathbb{D}$ and let n_1, \dots, n_{d-2} be pairwise distinct positive integers with $n_0 \geq 2$. Then for any $(c, a) \in \bigcap_j \text{Per}(n_j, w_j)$, the intersection is smooth and transverse at (c, a) . Moreover, $\text{Card}(\bigcap_j \text{Per}(n_j, w_j)) = (d - 1)!D_{\underline{n}}$.*

Proof. — According to [FG, §6], for any $(c, a) \in \bigcap_j \text{Per}(n_j, w_j)$, the intersection is smooth and transverse at (c, a) and $\text{Card}(\bigcap_j \text{Per}(n_j, w_j))$ is independent of (w_0, \dots, w_{d-2}) . It is thus sufficient to estimate this cardinal for $w_0 = \dots = w_{d-2} = 0$. For $n \geq 1$, it is easy to see that $\text{Per}(n, 0) = \bigcup_{j=0}^{d-2} \text{Per}_j(n)$. In particular,

$$\text{Card} \left(\bigcap_{j=0}^{d-2} \text{Per}(n_j, 0) \right) = \sum_{s \in \mathfrak{S}_{d-1}} \text{Card} \left(\bigcap_{j=0}^{d-2} \text{Per}_{s(j)}(n_j) \right),$$

which ends the proof, since $\text{Card}(\bigcap_{j=0}^{d-2} \text{Per}_{s(j)}(n_j)) = D_{\underline{n}}$ for any $s \in \mathfrak{S}_{d-1}$. \square

We are now in position to prove Theorem C. We deduce Theorem C from Theorem B. Our strategy mixes arguments from the proof of Corollary 4.4 and from the proof of [FG, Theorem 3].

Proof of Theorem C. — Let us first prove our result for $\underline{w} = (0, \dots, 0)$. The set of parameters $(c, a) \in \mathbb{C}^{d-1}$ such that $P_{c,a}$ has $(d-1)$ distinct super-attracting cycles of respective exact periods n_0, \dots, n_{d-2} coincides with

$$\bigcup_{\sigma \in \mathfrak{S}_{d-1}} \bigcap_{j=0}^{d-2} \text{Per}_{\sigma(j)}(n_j) .$$

As the intersection $\bigcap_{j=0}^{d-2} \text{Per}_{\sigma(j)}(n_j)$ is smooth and transverse for all $\sigma \in \mathfrak{S}_{d-1}$, the measure $\mu_{\underline{0}}^{\underline{n}}$ is the probability measure which is proportional to

$$\sum_{\sigma \in \mathfrak{S}_{d-1}} \frac{1}{D_{\underline{n}}} \bigwedge_{j=0}^{d-2} [\text{Per}_{\sigma(j)}(n_j)]$$

which has mass $(d-1)!$. As a consequence, we have

$$\mu_{\underline{0}}^{\underline{n}} = \sum_{\sigma \in \mathfrak{S}_{d-1}} \frac{1}{(d-1)! D_{\underline{n}}} \bigwedge_{j=0}^{d-2} [\text{Per}_{\sigma(j)}(n_j)] .$$

Fix now $\varphi \in \mathcal{C}_c^1(\mathbb{C}^{d-1})$. By a classical interpolation argument, Theorem B gives a constant C_1 depending only on d , such that

$$|\langle \mu_{\underline{0}}^{\underline{n}}, \varphi \rangle - \langle \mu_{\text{bif}}, \varphi \rangle| \leq C_1 \left(\max_j \left(\frac{\sigma(n_j)}{d_{n_j}} \right) \right)^{1/2} \|\varphi\|_{\mathcal{C}^1} .$$

This is the wanted result when $\underline{w} = \underline{0}$.

Pick now $\underline{w} \in \mathbb{D}^{d-1} \setminus \{\underline{0}\}$ and $\varphi \in \mathcal{C}_c^1(\mathbb{C}^{d-1})$. We write $\underline{w}[0] := (0, \dots, 0)$ and for any $1 \leq j \leq d-1$, we set $\underline{w}[j] := (w_0, \dots, w_{j-1}, 0, \dots, 0)$ and $\mu_j := \mu_{\underline{w}[j]}^{\underline{n}}$, so that $\underline{w}[d-1] = \underline{w}$, $\mu_0 = \mu_{\underline{0}}^{\underline{n}}$ and $\mu_{d-1} = \mu_{\underline{w}}^{\underline{n}}$. To conclude, it is sufficient to prove that

$$|\langle \mu_j, \varphi \rangle - \langle \mu_{j+1}, \varphi \rangle| \leq C' \left(\max_{0 \leq i \leq d-2} \left\{ \frac{1}{d_{n_j}}, \frac{-1}{d_{n_j} \log |w_i|} \right\} \right)^{1/2} \|\varphi\|_{\mathcal{C}^1},$$

for some constant $C' > 0$ depending only on d . If $w_j = 0$, there is nothing to prove, since in that case $\underline{w}[j] = \underline{w}[j+1]$. We thus may assume that $w_j \neq 0$. We consider the algebraic subvariety

$$C_j := \bigcap_{h < j} \text{Per}(n_h, w_h) \cap \bigcap_{l > j} \text{Per}(n_l, 0) .$$

Observe that $C_j \cap \text{Per}(n_j, 0) = \bigcap_k \text{Per}(n_k, \underline{w}[j]_k)$ is finite, hence C_j is an algebraic curve and that the intersections are smooth and transverse, by Proposition 7.1.

Let $X_j := \bigcap_k \text{Per}(n_k, \underline{w}[j]_k)$. Pick any point $(c, a) \in X_j$, and let $\Omega_{c,a}$ be the hyperbolic component containing (c, a) . Using [FG, Theorem 6.8], we define $\phi_{c,a} : \mathbb{D}(0, |w_j|^{-1/2}) \rightarrow \Omega_{c,a}$ by setting

$$\phi_{c,a}(t) := \rho^{-1}(w_0, \dots, w_{j-1}, tw_j, 0, \dots, 0) .$$

By construction, the disks

$$\mathbb{D}_{c,a} := \phi_{c,a}(\mathbb{D}(0, |w_j|^{-1/2})) \text{ and } \mathbb{D}'_{c,a} := \phi_{c,a}(\mathbb{D}(0, 1)) \in \mathbb{D}_{c,a}$$

are included in $\Omega_{c,a} \cap C_j$, $\phi_{c,a}(0) = (c, a)$ and $\phi_{c,a}(1)$ belongs to X_{j+1} . Any hyperbolic component contains a unique point in X_j , hence the collection of disks $\mathbb{D}_{c,a}$ is disjoint.

Note also that any point in X_{j+1} belongs to a hyperbolic component, and thus is equal to $\phi_{c,a}(1)$ for a unique $(c, a) \in X_j$.

By Cauchy-Schwarz inequality,

$$\begin{aligned} |\langle \mu_j, \varphi \rangle - \langle \mu_{j+1}, \varphi \rangle| &\leq \frac{1}{D_{\underline{n}}} \sum_{X_j} |\varphi(\phi_{c,a}(0)) - \varphi(\phi_{c,a}(1))| \\ &\leq \frac{1}{D_{\underline{n}}} \|\varphi\|_{C^1} \sum_{X_j} \text{Diam}_{\beta}(\mathbb{D}'_{c,a}) \\ &\leq \frac{1}{\sqrt{D_{\underline{n}}}} \|\varphi\|_{C^1} \left(\sum_{X_j} (\text{Diam}_{\beta}(\mathbb{D}'_{c,a}))^2 \right)^{1/2}. \end{aligned}$$

Recall that ρ is a biholomorphism. In particular,

$$\text{mod}(\mathbb{D}_{c,a} \setminus \overline{\mathbb{D}'_{c,a}}) = \text{mod}(\mathbb{D}(0, |w_j|^{-1/2}) \setminus \overline{\mathbb{D}}) = \frac{-1}{4\pi} \log |w_j|.$$

By Lemma 2.3, we deduce

$$|\langle \mu_j, \varphi \rangle - \langle \mu_{j+1}, \varphi \rangle| \leq \frac{1}{\sqrt{D_{\underline{n}}}} \|\varphi\|_{C^1} \left(\sum_{(c,a) \in X_j} \frac{\tau}{\min(1, \frac{1}{4\pi} \log |w|)} \text{Area}_{\beta}(\mathbb{D}_{c,a}) \right)^{1/2}.$$

On the other hand, by Bézout, we have

$$\deg(C_j) = \sum_{i=0}^{d-2} \sum_{\substack{\sigma \in \mathfrak{S}_{d-1}, \\ \sigma(j)=i}} \deg \left(\bigcap_{k \neq j} \text{Per}_{\sigma(k)}(n_k) \right) = (d-1)! \prod_{k \neq j} d_{n_k} = (d-1)! \frac{D_{\underline{n}}}{d_{n_j}}.$$

From which, since $\Omega_{c,a} \cap \Omega_{c',a'} = \emptyset$ for $(c, a) \neq (c', a') \in X_j$ and $\mathbb{D}_{c,a} \subset \Omega_{c,a} \cap C_j$, we deduce

$$\sum_{(c,a) \in X_j} \text{Area}_{\omega}(\mathbb{D}_{c,a}) \leq \text{Area}_{\omega}(C_j) \leq \deg(C_j) = (d-1)! \frac{D_{\underline{n}}}{d_{n_j}}.$$

Now, since $\Omega_{c,a} \subset \mathcal{C}_d \subset \mathbb{B}(0, 16\sqrt{d-1})$ (see the proof of Theorem B), there exists a constant $C_2 > 0$ such that $\beta \leq C_2\omega$ on $\Omega_{c,a}$, for all $(c, a) \in X_j$. Hence $\text{Area}_{\beta}(\mathbb{D}_{c,a}) \leq C_2 \text{Area}_{\omega}(\mathbb{D}_{c,a})$. Combined with the above, this gives

$$\begin{aligned} |\langle \mu_j, \varphi \rangle - \langle \mu_{j+1}, \varphi \rangle| &\leq \frac{16\sqrt{\pi\tau}(d-1)!(d-1)C_2}{\sqrt{D_{\underline{n}}}} \|\varphi\|_{C^1} \left(\max \left\{ 1, \sqrt{\frac{-4\pi}{\log |w_j|}} \right\} \right) \sqrt{\frac{D_{\underline{n}}}{d_{n_j}}}, \\ &\leq C_3 \|\varphi\|_{C^1} \left(\max_{0 \leq \ell \leq d-2} \left\{ \frac{1}{d_{n_{\ell}}}, \frac{-1}{d_{n_{\ell}} \log |w_{\ell}|} \right\} \right)^{1/2}, \end{aligned}$$

where $C_3 = 32\pi\sqrt{\tau(d-1)!(d-1)C_2}$. □

7.2. Convergence for multipliers outside a pluripolar set: Theorem D

For the content of the present, we are inspired by the techniques of [DTV] (see also [DS3]). Let ω be the Fubini-Study form on \mathbb{P}^1 and let $\pi_j : (\mathbb{P}^1)^k \rightarrow \mathbb{P}^1$ be the canonical projection to the j -th factor. We denote $\omega_j := \pi_j^*(\omega)$. We define a smooth probability measure Ω on $(\mathbb{P}^1)^k$ by $\Omega := \omega_1 \wedge \cdots \wedge \omega_k$.

First, recall some facts on DSH functions. Let $k \geq 1$. Recall that a probability measure ν in $(\mathbb{P}^1)^k$ has *bounded quasi-potentials* (or is PB) if ν admits a negative quasi-potential U ($dd^c U + \Omega = \nu$) such that $|\langle U, S \rangle| \leq C$ for any positive smooth form S of bidegree $(1, 1)$ and mass 1 ([DS1]). Such notion of quasi-potential can be extended to any positive closed current of bidegree $(1, 1)$ and mass 1 with the same bound $|\langle U, S \rangle| \leq C$ (using structural varieties in the space of currents). An interesting example of PB measure is the tensor product of measures in \mathbb{P}^1 with bounded quasi-potentials.

We say that a function φ on $(\mathbb{P}^1)^{d-1}$ is DSH if, outside a pluripolar set, it can be written as a difference of qpsH functions (for example, $\varphi \in \mathcal{C}^2$). Let $DSH\left((\mathbb{P}^1)^{d-1}\right)$ be the space of such functions. We write $dd^c \varphi = T^+ - T^-$ where T^\pm are positive closed currents of bidegree $(1, 1)$. Let ν be a PB measure on $(\mathbb{P}^1)^{d-1}$. Then, the following defines a norm on the space $DSH\left((\mathbb{P}^1)^{d-1}\right)$:

$$\|\varphi\|_\nu := \|\varphi\|_{L^1(\nu)} + \inf \|T^\pm\|$$

where the infimum is on all the decompositions $dd^c \varphi = T^+ - T^-$. It turns out that taking another PB measure ν' gives an equivalent norm on $DSH\left((\mathbb{P}^1)^{d-1}\right)$ (see e.g. [DS3, p. 283]).

Let φ be a smooth \mathcal{C}^2 function on $(\mathbb{P}^1)^{d-1}$, in particular φ is DSH and let \underline{n} be a $(d-1)$ -uple of pairwise distinct positive integers with $n_0 \geq 2$. Consider the function:

$$\Phi_{\underline{n}} := \underline{w} \in \mathbb{C}^{d-1} \mapsto \langle \varphi, \mu_{\underline{n}, \underline{w}} \rangle$$

where

$$\mu_{\underline{n}, \underline{w}} := \frac{1}{(d-1)! \prod_j d_{n_j}} \bigwedge_{j=1}^{d-1} [\text{Per}(n_j, w_j)].$$

This intersection is well defined outside an analytic (hence pluripolar) set $\mathcal{E}_{\underline{n}}$. So, if we consider the pluripolar set $\mathcal{E} := \cup_{\underline{n}} \mathcal{E}_{\underline{n}}$, the map $\Phi_{\underline{n}}$ is well defined outside \mathcal{E} for all \underline{n} . Adding a pluripolar set if necessary, we can assume that for $w \notin \mathcal{E}$ and all j then $w_j \neq 1$.

For the rest of the subsection, we let ν be a smooth probability measure with support in $\mathbb{D}(0, 1/2)^{d-1}$. In particular, ν is PB. The main result we need here is the following.

Lemma 7.2. — *There exists a constant C_ν depending only on ν and on d such that for all \underline{n} , the function $\Phi_{\underline{n}}$ is DSH with:*

$$\|\Phi_{\underline{n}} - \langle \mu_{\text{bif}}, \varphi \rangle\|_\nu \leq C_\nu \max_j \left(\frac{1}{n_j} \right).$$

Proof. — Let Π_1 (resp. Π_2) denote the canonical projection from $\mathbb{P}^{d-1} \times (\mathbb{P}^1)^{d-1}$ to \mathbb{P}^{d-1} (resp. $(\mathbb{P}^1)^{d-1}$). We let also $\tilde{\pi}_j : (\mathbb{P}^1)^{d-1} \rightarrow \mathbb{P}^1$ be the canonical projection to the j -th factor. Let ω_1 (resp. ω_{d-1}) denote the Fubini-Study form on \mathbb{P}^1 (resp. \mathbb{P}^{d-1}). Consider in $\mathbb{P}^{d-1} \times (\mathbb{P}^1)^{d-1}$ the (trivial extension of the) analytic set:

$$\mathcal{P}(n, j) := \{(c, a, \underline{w}), (c, a) \in \text{Per}(n, w_j)\}.$$

Then $(c, a, \underline{w}) \in \mathcal{P}(n, j)$ if and only if $p_n(c, a, w_j) = 0$ where the polynomials p_{n_j} where defined in the previous section. Recall that $\deg_w p_n = d_n/n$ and $\deg_{(c,a)} p_n = (d-1)d_n$ (see Section 7.1).

In particular, the current of integration $[\mathcal{P}(n_j, j, w_j)]$ is cohomologous to

$$(d-1)d_n \Pi_1^*(\omega_{d-1}) + \frac{d_n}{n} (\tilde{\pi}_j \circ \Pi_2)^*(\omega_1) .$$

Observe that the function $\Phi_{\underline{n}}$ can be defined in $\mathbb{C}^{d-1} \setminus \mathcal{E}$ as the following slice (we refer to [DS3, p. 280] for slicing on currents):

$$\Phi_{\underline{n}}(\underline{w}) = \left\langle \Pi_1^*(\varphi) \frac{1}{(d-1)! \prod_j d_{n_j}} \bigwedge_{i=1}^{d-1} [\mathcal{P}(n_j, j)], \Pi_2, \underline{w} \right\rangle .$$

Since the slicing commutes with the operator dd^c , we have

$$dd^c \Phi_{\underline{n}} = \left\langle \Pi_1^*(dd^c \varphi) \wedge \frac{1}{(d-1)! \prod_j d_{n_j}} \bigwedge_{i=1}^{d-1} [\mathcal{P}(n_j, j)], \Pi_2, \underline{w} \right\rangle .$$

Write $dd^c \varphi = T^+ - T^-$. As $\Pi_1^*(T^\pm)$ is cohomologous to $\|T^\pm\| \cdot \Pi_1^*(\omega_{d-1})$, the mass of

$$\mathcal{T}_{\underline{n}}^\pm = \left\langle \Pi_1^*(T^\pm) \wedge \frac{1}{(d-1)! \prod_j d_{n_j}} \bigwedge_{i=1}^{d-1} [\mathcal{P}(n_j, j)], \Pi_2, \underline{w} \right\rangle$$

can be computed in cohomology. In particular, it is bounded from above:

$$\|\mathcal{T}_{\underline{n}}^\pm\| \leq C_d \cdot \|T^+\| \sum_j \frac{1}{n_j} ,$$

for some constant C_d that depends only d ($C_d = (d-1)^{d-1}/(d-1)!$ works).

On the other hand, according to Theorem C above, for all $\underline{w} \in \text{supp}(\nu)$, we have

$$|\Phi_{\underline{n}} - \langle \mu_{\text{bif}}, \varphi \rangle| \leq C_\varphi \sum_j \frac{1}{n_j}$$

where C_φ is a constant that depends only on φ . The result follows. \square

We finish the proof of Theorem D.

Proof of Theorem D. — We assume that the series

$$\sum_k \max_j \left(\frac{1}{n_{j,k}} \right)$$

converges. Pick $\varphi \in \mathcal{C}_c^2(\mathbb{C}^{d-1})$. By Lemma 7.2, $\Phi_{\underline{n}_k} - \langle \mu_{\text{bif}}, \varphi \rangle \in L^1(\nu)$ and

$$\|\Phi_{\underline{n}_k} - \langle \mu_{\text{bif}}, \varphi \rangle\|_{L^1(\nu)} \leq C_\nu \max_j \left(\frac{1}{n_{j,k}} \right) .$$

Hence, it converges ν -almost everywhere to 0. As all DSH-norms are equivalent, we have that for another PB measure ν' ,

$$\|\Phi_{\underline{n}_k} - \langle \mu_{\text{bif}}, \varphi \rangle\|_{L^1(\nu')} \leq C_{\nu'} \max_j \left(\frac{1}{n_{j,k}} \right)$$

and we can apply the same argument: it converges to 0, ν' -a.e.

Finally, when a set E is of ν' -measure 0 for all PB measure ν' , it is pluripolar. Indeed, its logarithmic capacity has to vanish. We deduce, in particular, that the sequence $\Phi_{\underline{n}_k} - \langle \mu_{\text{bif}}, \varphi \rangle$ converges to 0 outside a pluripolar set.

By separability, we apply the same argument to a dense countable family in $\mathcal{C}_c(\mathbb{C}^{d-1})$ of $\varphi \in \mathcal{C}^2(\mathbb{P}^1)^{d-1}$. As a countable union of pluripolar sets is again pluripolar, we deduce

that outside a pluripolar set the measure $\mu_{\underline{n}, \underline{w}}$ converges to μ_{bif} . This ends the first part of the proof of Theorem D. \square

Remark. —

1. In the moduli space of quadratic *rational maps* (which is biholomorphic to \mathbb{C}^2), the curves $\text{Per}(3, 1)$ and $\text{Per}(2, -3)$ have a common irreducible component (see [M1]). Such a behavior is expected to be impossible in the moduli space of degree d *polynomials* \mathcal{P}_d . We even expect the exceptional set \mathcal{E} appearing in Theorem D to be empty.
2. We also expect the convergence of Theorem D to hold for *any* sequence \underline{n}_k of $(d-1)$ -tuple, though we don't know how to prove it.

We now give a quick proof of Corollary E.

Proof of Corollary E. — The Shilov boundary of \mathbb{D}^{d-1} is exactly $(\mathbb{S}^1)^{d-1}$ and the pluricomplex Green function of \mathbb{D}^{d-1} , which is given by $g = \log^+ \max_i \{|z_i|\}$, is continuous and $\nu := (dd^c g)^{d-1}$ coincides with the Lebesgue measure of $(\mathbb{S}^1)^{d-1}$, hence does not give mass to pluripolar sets. By Theorem D, for ν -a.e. \underline{w} , the sequence $(\mu_{\underline{n}_k, \underline{w}})_k$ converges to μ_{bif} . \square

8. Distribution of parametric preimages

Let $(f_\lambda)_{\lambda \in \Lambda}$ be a holomorphic family of rational maps of degree $d \geq 2$ on \mathbb{P}^1 with $\dim(\Lambda) = m$. Let ω_Λ be a Kähler form on Λ . Assume that c_1, \dots, c_k are marked critical points and let T_1, \dots, T_k be their respective bifurcation currents (see Section 2.1).

For $1 \leq j \leq k$, we let $v_n^j : \Lambda \rightarrow \mathbb{P}^1$ be the map defined by $v_n^j(\lambda) = f_\lambda^n(c_j(\lambda))$ and set $v_n := (v_n^1, \dots, v_n^k)$. Let ω_j and Ω be defined as in Section 7. Then [Du, Theorem 0.3] asserts that there exists a pluripolar set $\mathcal{E} \subset (\mathbb{P}^1)^k$ such that if $(z_1, \dots, z_k) \in (\mathbb{P}^1)^k \setminus \mathcal{E}$, the following equidistribution statement holds:

$$\frac{1}{d^{nk}} \{f_\lambda^n(c_1(\lambda)) = z_1\} \cap \dots \cap \{f_\lambda^n(c_k(\lambda)) = z_k\} \rightarrow T_1 \wedge \dots \wedge T_k.$$

Observe that the equidistribution statement can be rewritten as:

$$\frac{1}{d^{nk}} (v_n^1)^*(\delta_{z_1}) \wedge \dots \wedge (v_n^k)^*(\delta_{z_k}) = \frac{1}{d^{nk}} (v_n)^*(\delta_{(z_1, \dots, z_k)}) \rightarrow T_1 \wedge \dots \wedge T_k.$$

We want to give here some improved versions of that theorem with speed estimates.

The proof is quite elementary. Recall that PB measures were defined in the previous section. Our precise result may be formulated as follows.

Theorem 8.1. — *Assume that the family $(f_\lambda)_{\lambda \in \Lambda}$ is algebraic, i.e. that Λ is a quasi-projective variety and let $m := \dim(\Lambda)$. Let K be a compact subset $K \Subset \Lambda$, then with the above notations and hypotheses we have:*

1. *Let ν be a probability measure on $(\mathbb{P}^1)^k$ with bounded potentials, then there exists $C(K) > 0$ such that for any \mathcal{C}^2 -form φ of bidegree $(m-k, m-k)$ and support in K and any $n \geq 1$:*

$$\left| \left\langle \frac{1}{d^{nk}} (v_n)^*(\nu) - T_1 \wedge \dots \wedge T_k, \varphi \right\rangle \right| \leq C(K) \frac{1}{d^n} \|\varphi\|_{\mathcal{C}^2(K)}.$$

2. Let (a_n) be a sequence of positive number such that $\sum a_n < \infty$, then there exists a pluripolar set $\mathcal{E} \subset (\mathbb{P}^1)^k$ such that if $(z_1, \dots, z_k) \in (\mathbb{P}^1)^k \setminus \mathcal{E}$, then there exists $C(K) > 0$ such that for any \mathcal{C}^2 -form φ of bidegree $(m-k, m-k)$ and support in K and any $n \geq 1$:

$$\left| \left\langle \frac{1}{d^{nk}} (v_n)^* (\delta_{(z_1, \dots, z_k)}) - T_1 \wedge \dots \wedge T_k, \varphi \right\rangle \right| \leq C(K) \frac{1}{a_n d^n} \|\varphi\|_{\mathcal{C}^2(K)} .$$

Proof. — Let us first prove the first point. Recall that if we let $T_n^j = d^{-n} (v_n^j)^* (\omega)$, then $T_n^j - T_j = dd^c V_n^j$ where $V_n^j = O(d^{-n})$ on K (see e.g. [Du]) hence using the same arguments than above, it is enough to prove the bound:

$$\left| \left\langle \frac{1}{d^{nk}} (v_n)^* (\nu) - T_n^1 \wedge \dots \wedge T_n^k, \varphi \right\rangle \right| \leq C(K) \frac{1}{d^n} \|\varphi\|_{\mathcal{C}^2(K)}$$

for any \mathcal{C}^2 -form φ of bidegree $(m-k, m-k)$ and support in K .

Writing $T_n^1 \wedge \dots \wedge T_n^k = d^{-kn} (v_n)^* (\Omega)$ and applying Stokes formula (Lemma 2.4), we find

$$\begin{aligned} \left\langle \frac{1}{d^{nk}} (v_n)^* (\nu) - T_n^1 \wedge \dots \wedge T_n^k, \varphi \right\rangle &= \left\langle \frac{1}{d^{nk}} (v_n)^* (U), dd^c \varphi \right\rangle \\ &= \left\langle U, \frac{1}{d^{nk}} (v_n)_* (dd^c \varphi) \right\rangle. \end{aligned}$$

Now, as φ is \mathcal{C}^2 , we have $\|\varphi\|_{\mathcal{C}^2} \omega_\Lambda^{m-k+1} \pm dd^c \varphi \geq 0$. As we assumed that Λ is a quasi-projective variety, the mass of the current $d^{-n(k-1)} (v_n)_* (\omega_\Lambda^{m-k+1})$ is uniformly bounded. The fact that ν is PB gives the wanted result.

We now prove the second point. Let (a_n) be a sequence of positive numbers such that $\sum a_n < \infty$. Observe that the positive closed current $S = \sum_n a_n d^{-n(k-1)} (v_n)_* (\omega_\Lambda^{m-k+1})$ is well defined and has finite mass. For $z \in (\mathbb{P}^1)^k$, we let U_z be the Green quasi-potential of δ_z (see [DS2]). The complementary of the set of points $z \in (\mathbb{P}^1)^k$ such that $\langle U_z, S \rangle > -\infty$ is a pluripolar set we shall denote by \mathcal{E} . For any $z \notin \mathcal{E}$, following the above proof, we find

$$0 \geq \langle U_z, \frac{1}{d^{nk}} (v_n)_* (dd^c \varphi) \rangle \geq \langle U_z, \frac{1}{d^{nk}} \|\varphi\|_{\mathcal{C}^2} (v_n)_* (\omega_\Lambda^{m-k+1}) \rangle \geq \frac{1}{a_n d^n} \|\varphi\|_{\mathcal{C}^2} \langle U_z, S \rangle.$$

This ends the proof. \square

Removing the assumption on the algebraicity of the family, one can using the same techniques, prove the following.

Proposition 8.2. — *Let K be a compact subset $K \Subset \Lambda$. For $j \leq k$ let ν_j be a measure in \mathbb{P}^1 with bounded potentials and let $\nu := \nu_1 \otimes \dots \otimes \nu_k$. Then there exists $C(K) > 0$ such that for any \mathcal{C}^2 -form φ of bidegree $(m-k, m-k)$ and support in K and any $n \geq 1$:*

$$\left| \left\langle \frac{1}{d^{nk}} (v_n)^* (\nu) - T_1 \wedge \dots \wedge T_k, \varphi \right\rangle \right| \leq C(K) \frac{1}{d^n} \|\varphi\|_{\mathcal{C}^2(K)} .$$

The proof is similar to the above one although we do not need to push forward by v_n here and use instead that the sequence $(d^{-n} (v_n^j)^* (\nu_j))$ is locally uniformly bounded in mass.

As observed in [Du], the question is parallel to the equidistribution of preimages of points under a rational map or more generally under a holomorphic endomorphism of \mathbb{P}^k

(see e.g. [Lj, BD]). In the case of such a single map, it is known that the exceptional set, i.e. the set of points for which one does not have equidistribution, is algebraic.

In the particular case of the unicritical family, let $v_n(c) := p_c^n(0)$, $c \in \mathbb{C}$. Using the classical proof of the equidistribution of centers of hyperbolic components, we can prove that the exceptional set is reduced to the point ∞ which is exceptional for any p_c :

Theorem 8.3. — *The sequence $(\frac{1}{d^{n-1}}(v_n)^*(\delta_z))_{n \geq 1}$ converges to $\mu_{\mathbf{M}_d}$ in the weak sense of currents if and only if $z \in \mathbb{C}$, i.e. the exceptional set is $\mathcal{E} = \{\infty\}$.*

Proof. — We follow closely Levin [Le]. Indeed, take $u_n(c) = \frac{1}{d^{n-1}} \log |v_n(c) - z|$ so that $\Delta u_n = \frac{1}{d^{n-1}}(v_n)^*(\delta_z)$. Then u_n is a sequence of subharmonic functions that are bounded in L^1_{loc} . Extracting a converging subsequence, we have that the limit u is equal to $g_{\mathbf{M}_d}$ outside the Mandelbrot set since $|v_n(c) - z| \simeq |v_n(c)|$ outside the connectedness locus. Similarly, $u \leq g_{\mathbf{M}_d}$ elsewhere. As $g_{\mathbf{M}_d}$ is continuous and $\mu_{\mathbf{M}_d}$ gives no mass to the boundary of the component of the interior of \mathbf{M}_d , it implies that $u = g_{\mathbf{M}_d}$.

Finally, remark that the polynomial v_n extends at infinity of the parameter space by setting $v_n(\infty) = \infty$. We get this way a polynomial map $v_n : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ of degree d^{n-1} . Hence $(v_n)^*(\delta_\infty) = d^{n-1}\delta_\infty$ for any $n \geq 1$, which concludes the proof. \square

Question. — *Let \mathcal{E} be the set of points $(z_1, \dots, z_k) \in (\mathbb{P}^1)^k$ where $d^{-nk}(v_n)^*(\delta_{(z_1, \dots, z_k)})$ does not converge to $T_1 \wedge \dots \wedge T_k$.*

1. *Is \mathcal{E} empty for families of rational maps with generic empty exceptional set?*
2. *Is it reduced to $\bigcup_{j=1}^k (\mathbb{P}^1)^{j-1} \times \{\infty\} \times (\mathbb{P}^1)^{k-j}$ in any family of polynomials?*

References

- [BB1] Giovanni Bassanelli and François Berteloot. Bifurcation currents in holomorphic dynamics on \mathbb{P}^k . *J. Reine Angew. Math.*, 608:201–235, 2007.
- [BB2] Giovanni Bassanelli and François Berteloot. Lyapunov exponents, bifurcation currents and laminations in bifurcation loci. *Math. Ann.*, 345(1):1–23, 2009.
- [BB3] Giovanni Bassanelli and François Berteloot. Distribution of polynomials with cycles of a given multiplier. *Nagoya Math. J.*, 201:23–43, 2011.
- [BD] Jean-Yves Briend and Julien Duval. Exposants de Liapounoff et distribution des points périodiques d’un endomorphisme de \mathbf{CP}^k . *Acta Math.*, 182(2):143–157, 1999.
- [BG1] François Berteloot and Thomas Gauthier. On the geometry of bifurcation currents for quadratic rational maps. *Ergodic Theory and Dynamical Systems*, FirstView:1–11, 4 2015.
- [BG2] Xavier Buff and Thomas Gauthier. Quadratic polynomials, multipliers and equidistribution. *Proc. Amer. Math. Soc.*, 143(7):3011–3017, 2015.
- [BH] Bodil Branner and John H. Hubbard. The iteration of cubic polynomials. I. The global topology of parameter space. *Acta Math.*, 160(3-4):143–206, 1988.
- [BT] Eric Bedford and B. A. Taylor. The Dirichlet problem for a complex Monge-Ampere equation. *Bull. Amer. Math. Soc.*, 82(1):102–104, 1976.
- [C] E. M. Chirka. *Complex analytic sets*, volume 46 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1989. Translated from the Russian by R. A. M. Hoksbergen.
- [CG] Lennart Carleson and Theodore W. Gamelin. *Complex dynamics*. Universitext: Tracts in Mathematics. Springer-Verlag, New York, 1993.
- [Dem] J.-P. Demailly. Complex analytic and differential geometry, 2011. free accessible book (<http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>).

- [DeM1] Laura DeMarco. Dynamics of rational maps: a current on the bifurcation locus. *Math. Res. Lett.*, 8(1-2):57–66, 2001.
- [DeM2] Laura DeMarco. Dynamics of rational maps: Lyapunov exponents, bifurcations, and capacity. *Math. Ann.*, 326(1):43–73, 2003.
- [Du] Romain Dujardin. The supports of higher bifurcation currents. *Ann. Fac. Sci. Toulouse Math. (6)*, 22(3):445–464, 2013.
- [DF] Romain Dujardin and Charles Favre. Distribution of rational maps with a preperiodic critical point. *Amer. J. Math.*, 130(4):979–1032, 2008.
- [DH] A. Douady and J. H. Hubbard. *Étude dynamique des polynômes complexes. Partie II*, volume 85 of *Publications Mathématiques d’Orsay [Mathematical Publications of Orsay]*. Université de Paris-Sud, Département de Mathématiques, Orsay, 1985. With the collaboration of P. Lavaurs, Tan Lei and P. Sentenac.
- [DS1] Tien-Cuong Dinh and Nessim Sibony. Dynamics of regular birational maps in \mathbb{P}^k . *J. Funct. Anal.*, 222(1):202–216, 2005.
- [DS2] Tien-Cuong Dinh and Nessim Sibony. Super-potentials of positive closed currents, intersection theory and dynamics. *Acta Math.*, 203(1):1–82, 2009.
- [DS3] Tien-Cuong Dinh and Nessim Sibony. Dynamics in several complex variables: endomorphisms of projective spaces and polynomial-like mappings. In *Holomorphic dynamical systems*, volume 1998 of *Lecture Notes in Math.*, pages 165–294. Springer, Berlin, 2010.
- [DTV] Henry De Thélin and Gabriel Vigny. Entropy of meromorphic maps and dynamics of birational maps. *Mém. Soc. Math. Fr. (N.S.)*, 122:vi+98, 2010.
- [E] Adam Epstein. Transversality principles in holomorphic dynamics, 2009. preprint.
- [FG] Charles Favre and Thomas Gauthier. Distribution of postcritically finite polynomials, 2013. to appear in *Israël J. Math.*
- [FRL] C. Favre and J. Rivera-Letelier. Equidistribution quantitative des points de petite hauteur sur la droite projective. *Math. Ann.*, 335(2):311–361, 2006.
- [G1] Thomas Gauthier. Strong bifurcation loci of full Hausdorff dimension. *Ann. Sci. Éc. Norm. Supér. (4)*, 45(6):947–984, 2012.
- [G2] Thomas Gauthier. Equidistribution towards the bifurcation current I: Multipliers and degree d polynomials, 2013. preprint arXiv:1312.7554, submitted.
- [I] Patrick Ingram. A finiteness result for post-critically finite polynomials. *Int. Math. Res. Not. IMRN*, 3:524–543, 2012.
- [Le] Genadi Levin. On the theory of iterations of polynomial families in the complex plane. *J. Soviet Math.*, 52(6):3512–3522, 1990.
- [Lj] M. Ju. Ljubich. Entropy properties of rational endomorphisms of the Riemann sphere. *Ergodic Theory Dynam. Systems*, 3(3):351–385, 1983.
- [Ly] M. Yu. Lyubich. Some typical properties of the dynamics of rational mappings. *Uspekhi Mat. Nauk*, 38(5(233)):197–198, 1983.
- [M1] John Milnor. Geometry and dynamics of quadratic rational maps. *Experiment. Math.*, 2(1):37–83, 1993. With an appendix by the author and Lei Tan.
- [M2] John Milnor. Cubic polynomial maps with periodic critical orbit. I. In *Complex dynamics*, pages 333–411. A K Peters, Wellesley, MA, 2009.
- [MSS] R. Mañé, P. Sad, and D. Sullivan. On the dynamics of rational maps. *Ann. Sci. École Norm. Sup. (4)*, 16(2):193–217, 1983.
- [Oku1] Yūsuke Okuyama. Equidistribution of rational functions having a superattracting periodic point towards the activity current and the bifurcation current. *Conform. Geom. Dyn.*, 18:217–228, 2014.
- [Oku2] Yūsuke Okuyama. Quantitative approximations of the lyapunov exponent of a rational function over valued fields. *Mathematische Zeitschrift*, pages 1–16, 2015.

- [P] Feliks Przytycki. Lyapunov characteristic exponents are nonnegative. *Proc. Amer. Math. Soc.*, 119(1):309–317, 1993.
- [Si] Joseph H. Silverman. *The arithmetic of dynamical systems*, volume 241 of *Graduate Texts in Mathematics*. Springer, New York, 2007.
- [St] Norbert Steinmetz. *Rational iteration*, volume 16 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1993. Complex analytic dynamical systems.

THOMAS GAUTHIER & GABRIEL VIGNY, LAMFA, Université de Picardie Jules Verne, 33 rue Saint-Leu, 80039 AMIENS Cedex 1, FRANCE • *E-mail* : thomas.gauthier@u-picardie.fr, gabriel.vigny@u-picardie.fr