



HAL
open science

COMMUTATIVE RINGS WHOSE COTORSION MODULES ARE PURE-INJECTIVE

Francois Couchot

► **To cite this version:**

Francois Couchot. COMMUTATIVE RINGS WHOSE COTORSION MODULES ARE PURE-INJECTIVE. 2015. hal-01162032v1

HAL Id: hal-01162032

<https://hal.science/hal-01162032v1>

Preprint submitted on 9 Jun 2015 (v1), last revised 19 Mar 2016 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMMUTATIVE RINGS WHOSE COTORSION MODULES ARE PURE-INJECTIVE

FRANÇOIS COUCHOT

ABSTRACT. Let R be a ring (not necessarily commutative). A left R -module is said to be *cotorsion* if $\text{Ext}_R^1(G, M) = 0$ for any flat R -module G . It is well known that each pure-injective left R -module is cotorsion, but the converse does not hold: for instance, if R is left perfect but not left pure-semisimple then each left R -module is cotorsion but there exist non-pure-injective left modules. The aim of this paper is to describe the class \mathcal{C} of commutative rings R for which each cotorsion R -module is pure-injective. It is easy to see that \mathcal{C} contains the class of von Neumann regular rings and the one of pure-semisimple rings. We prove that \mathcal{C} is strictly contained in the class of locally pure-semisimple rings. We state that a commutative ring R belongs to \mathcal{C} if and only if R verifies one of the following conditions:

- (1) each Warfield cotorsion R -module is RD-injective;
- (2) R is coherent and each pure-essential extension of R -modules is essential;
- (3) R is coherent and each RD-essential extension of R -modules is essential;
- (4) any R -module M is pure-injective if and only if $\text{Ext}_R^1(R/A, M) = 0$ for each pure ideal A of R (Baer's criterion).

1. INTRODUCTION AND PRELIMINARIES

The aim of this study is to give a complete description of commutative rings for which each cotorsion module is pure-injective. In this first section we recall some definitions and some former results. Then, in section 2, we enunciate and show some partial results which are available even if the ring is not commutative. In particular, if R is a ring for which each cyclic flat left R -module is projective, then every cotorsion left R -module is pure-injective if and only if R is left pure-semisimple. Section 3 is devoted to the commutative case. We get our main result (Theorem 3.9) by using localizations and local rings. In the last section we show that a commutative ring R is locally perfect if and only if any R -module M for which $\text{Ext}(C, M) = 0$ for each cyclic flat module C is cotorsion, and we investigate the following question: give a characterization of rings R for which each flat-essential extension of R -modules is essential. When R is commutative and coherent a complete response is given. Throughout this paper other related questions are studied, where we use the following notions: Warfield cotorsion module, RD-injective module, RD-essential extension and so on...

Even in the commutative case some questions are open. For instance we do not know if there exist non-coherent commutative rings for which each pure-essential extension of modules is essential. Also, it should be interesting to study strongly perfect rings which are introduced in the last section.

2010 *Mathematics Subject Classification.* 13C11, 16D40, 16D80.

Key words and phrases. cotorsion module, pure-injective module, essential extension, perfect ring, pure-semisimple ring.

We shall assume that all rings are associative with identity and all modules are unitary. Given a ring R , any left module M is said to be **P-flat** (resp. **P-injective**) if $\text{Tor}_1^R(R/rR, M) = 0$ (resp. $\text{Ext}_R^1(R/Rr, M) = 0$) for each $r \in R$. We say that R is **left P-coherent** if each principal left ideal of R is finitely presented.

A left module M is **FP-injective** if $\text{Ext}_R^1(F, M) = 0$ for each finitely presented left module F .

Any left module M is called **cotorsion** (resp. **Warfield cotorsion**) if, for each flat (resp. P-flat) left module F , $\text{Ext}_R^1(F, M) = 0$.

A short exact sequence of left modules is **pure** (resp. **RD-pure**) if it remains exact when tensoring it by any right module (resp. module of the form R/rR , $r \in R$). A left module is **pure-injective** (resp. **RD-injective**) if it is injective relatively to each pure (resp. RD-pure) exact sequence of left modules.

R is said to be left **pure-semisimple** (resp. **RD-semisimple**) if each left R -module is pure-injective (resp. RD-injective). When R is commutative then R is pure-semisimple if and only if it is RD-semisimple if and only if it is an Artinian ring whose all ideals are principal ([6, Theorem 4.3]).

An R -module B is a **pure-essential extension** (resp. **RD-essential extension**) of a submodule A if A is a pure (resp. RD) submodule of B and, if for each submodule K of B , either $K \cap A \neq 0$ or $(A + K)/K$ is not a pure (resp. RD) submodule of B/K . We say that B is a **pure-injective hull** (resp. **RD-injective hull**) of A if B is pure-injective (resp. RD-injective) and a pure-essential (resp. RD-essential) extension of A .

Each R -module M has a pure-injective hull and an RD-injective hull ([8, Proposition 6]).

A left module B is a **flat extension** (resp. **P-flat extension**) of a submodule A if B/A is flat (resp. P-flat). Moreover, if there are no submodules S of B with $S \cap A = 0$ and B/S flat (resp. P-flat) extension of A , then B is a **flat essential extension** (resp. **P-flat essential extension**) of A . If B is cotorsion (resp. Warfield cotorsion) and a flat (resp. P-flat) essential extension of a submodule A then we say that B is a **cotorsion** (resp. **Warfield cotorsion**) **envelope** of A (by [10, Theorem 3.4.5] these definitions are equivalent to the usual ones).

Each left module M has a cotorsion (resp. Warfield cotorsion) envelope ([4, Theorem 6] and [10, Theorem 3.4.6]).

For each left module M we denote by $\mathcal{E}(M)$ its cotorsion envelope, $\mathcal{E}_W(M)$ its Warfield cotorsion envelope, $\text{PE}(M)$ its pure-injective hull and $\text{RDE}(M)$ its RD-injective hull.

Each pure(RD)-injective module is (Warfield) cotorsion, but [10, Example p.75] shows that the converse does not hold.

Theorem 1.1. [10, Theorem 3.5.1] *For any ring R the following are equivalent:*

- (1) *for any exact sequence of left modules $0 \rightarrow P' \rightarrow P \rightarrow P'' \rightarrow 0$ with P' and P'' pure-injective, P is also pure-injective;*
- (2) *for any left module M , $\text{PE}(M)/M$ is flat;*
- (3) *every cotorsion left module is pure-injective.*

Moreover if R is right coherent, then the above are equivalent to the following:

- (4) *for any exact sequence of left modules $0 \rightarrow P' \rightarrow P \rightarrow P'' \rightarrow 0$ with P' and P pure-injective, P'' is also pure-injective.*

Theorem 1.2. *For any ring R the following are equivalent:*

- (1) for any exact sequence of left modules $0 \rightarrow P' \rightarrow P \rightarrow P'' \rightarrow 0$ with P' and P'' RD-injective, P is also RD-injective;
- (2) for any left module M , $\text{RDE}(M)/M$ is P -flat;
- (3) every Warfield cotorsion left module is RD-injective.

Moreover if R is right P -coherent, then the above are equivalent to the following:

- (4) for any exact sequence of left modules $0 \rightarrow P' \rightarrow P \rightarrow P'' \rightarrow 0$ with P' and P RD-injective, P'' is also RD-injective.

The following proposition is well known. For convenience, a proof is given. We set 0_P the kernel of the natural map $R \rightarrow R_P$ where $P \in \text{Spec } R$.

Proposition 1.3. *Let R be a commutative ring. We assume that each prime ideal is maximal. Then:*

- (1) for any closed subset C of $\text{Spec } R$, $C = V(A)$ where $A = \bigcap_{P \in C} 0_P$ is a pure ideal;
- (2) for each maximal ideal P $R_P = R/0_P$;
- (3) each pure ideal of R is generated by idempotents.

Proof. (1). Let $C = V(B)$ where $B = \bigcap_{L \in C} L$. We put $A = \bigcap_{P \in C} 0_P$. Let $b \in B$ and $P \in C$. The image of b , by the natural map $R \rightarrow R_P$, belongs to the nilradical of R_P . It follows that there exist $0 \neq n_P \in \mathbb{N}$ and $s_P \in R \setminus P$ such that $s_P b^{n_P} = 0$. Hence, $\forall L \in D(s_P) \cap C$, $b^{n_P} \in 0_L$. A finite family $(D(s_{P_j}))_{1 \leq j \leq m}$ covers C . Let $n = \max\{n_{P_1}, \dots, n_{P_m}\}$. Then $b^n \in 0_L$, $\forall L \in C$, whence $b^n \in A$. We deduce that $C = V(A)$. Now, we have $A_P = 0$ if $P \in V(A)$ and $A_P = R_P$ if $P \in D(A)$. Hence A is a pure ideal.

(2) is a consequence of (1) by taking $C = \{P\}$.

(3). We know that $\text{Spec } R$ is homeomorphic to $\text{Spec } R/N$ where N is the nilradical of R . Since R/N is von Neumann regular its principal ideals are generated by idempotents. So, $\text{Spec } R$ has a base of clopen subsets (closed and open). Whence if A is a pure ideal then, for any $a \in A$ there exists an idempotent e_a such that $D(e_a) = D(a) \subseteq D(A)$. Clearly $D(A) = D(\sum_{a \in A} R e_a)$. Since $\sum_{a \in A} R e_a$ is a pure ideal then by (1) we conclude that $A = \sum_{a \in A} R e_a$. \square

2. WHEN COTORSION MODULES ARE PURE-INJECTIVE: GENERAL CASE

A left module M over a ring R is called **regular** if all its submodules are pure.

Proposition 2.1. *Let R be a ring, J its Jacobson radical. Let M be a left R -module satisfying the following property: for each submodule N of M and for each $r \in R$, $rM \cap N = rN$ (N is an RD-submodule of M). Then $R/\text{ann}(M)$ is von Neumann regular, M is regular, $JM = 0$ and, if in addition R is semilocal, M is semisimple.*

Proof. Let $R' = R/A$ where A is the annihilator of M . Then R' is isomorphic to a submodule of M^M . It is easy to check that $rN = N \cap rM^M$ for each $r \in R$ and any submodule N of M^M . Then R' satisfies this property too as left R -module. We deduce that R' is a von Neumann regular ring, and consequently M is regular. If $0 \neq x \in M$ then Rax is an RD-submodule of Rx for each $a \in R$. So, for each $a \in J$, there exists $b \in R$ such that $ax = abax$. It follows that $(1 - ab)ax = 0$, and from $a \in J$ we successively deduce that $(1 - ab)$ is a unit and $ax = 0$. \square

Proposition 2.2. *Let R be a ring. Assume there exists a family \mathfrak{E} of orthogonal central idempotents of R satisfying the following conditions:*

- (1) $R/R(1 - e)$ is a left pure-semisimple ring, for each $e \in \mathfrak{E}$;
- (2) R/A is a von Neumann regular ring where $A = \bigoplus_{e \in \mathfrak{E}} Re$.

Then:

- (1) each cotorsion left R -module is pure-injective;
- (2) each pure-essential extension of left R -modules is essential;
- (3) any left R -module M is pure-injective if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat left R -module C ;
- (4) for any left R -module M , $\text{PE}(M)/M$ is flat, FP-injective and regular.

Proof. (3) and (1). Let M be a left R -module satisfying $\text{Ext}_R^1(R/B, M) = 0$ for each pure left ideal B of R . Since A is a pure ideal then the following sequence is exact:

$$0 \rightarrow \text{Hom}_R(R/A, M) \rightarrow \text{Hom}_R(R, M) \rightarrow \text{Hom}_R(A, M) \rightarrow 0.$$

Let C be a left ideal of R/A . Since R/A is von Neumann regular, then C is a pure ideal and its inverse image B by the natural map $R \rightarrow R/A$ is a pure left ideal of R . From $\text{Ext}_{R/A}^1(R/B, \text{Hom}_R(R/A, M)) \cong \text{Ext}_R^1(R/B, M) = 0$ we deduce that $\text{Hom}_R(R/A, M)$ is injective over R/A and R . So, the above sequence splits. On the other hand $\text{Hom}_R(A, M) \cong \prod_{e \in \mathfrak{E}} eM$. Since $R/R(1 - e)$ is left pure-semisimple, it successively follows that eM is pure-injective for each $e \in \mathfrak{E}$, $\text{Hom}_R(A, M)$ is pure-injective and M too.

(2). Let M be a left R -module, $N = \text{Hom}_R(R/A, M)$, $E = E(N)$ and $L = \text{Hom}_R(A, M)$. As above L is pure-injective. So, $E \oplus L$ is pure injective. The inclusion map $N \rightarrow E$ extends to a homomorphism $f : M \rightarrow E$. Let $g : M \rightarrow L$ be the canonical map and L' its image. Then, it is easy to check that the homomorphism $\phi : M \rightarrow E \oplus L$ defined by $\phi(m) = (f(m), g(m))$ for each $m \in M$ is injective. Since R/A is von Neumann regular then E/N is flat. It is easy to see that $AL = AL'$. So, L/L' is also an R/A -module. It follows that $\text{coker}(\phi)$ is an R/A -module which is flat over R . Hence ϕ is a pure monomorphism. Let $(x, y) \in E \oplus L$. First assume that $y \neq 0$. There exists $e \in \mathfrak{E}$ such that $ey \neq 0$. So, there exists $z \in M$ such that $ey = g(z)$. It follows that $ey = g(ez)$ and $\phi(ez) = e(x, y) = (0, ey)$. If $y = 0$ then there exists $s \in R$ such that $0 \neq sx \in N$, whence $\phi(sx) = s(x, 0)$. Hence ϕ is an essential monomorphism.

(4). Since $\text{coker}(\phi)$ is a module over R/A which is a von Neumann regular ring and flat as right R -module then $\text{coker}(\phi)$ is flat, FP-injective and regular as R -module. \square

Proposition 2.3. *Let R be a ring. Assume there exists a family \mathfrak{E} of orthogonal central idempotents of R satisfying the following conditions:*

- (1) $R/R(1 - e)$ is a left RD-semisimple ring, for each $e \in \mathfrak{E}$;
- (2) R/A is a von Neumann regular ring where $A = \bigoplus_{e \in \mathfrak{E}} Re$.

Then:

- (1) each Warfield cotorsion left R -module is RD-injective;
- (2) each RD-essential extension of left R -modules is essential;
- (3) any left R -module M is RD-injective if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat left R -module C ;
- (4) for any left R -module M , $\text{RDE}(M)/M$ is flat, FP-injective and regular.

Proposition 2.4. *Let R be a ring. Assume that each cotorsion left R -module is pure-injective. Then, for each left module M , every submodule of $\mathcal{E}(M)/M$ is flat. Moreover, for each submodule C of $\mathcal{E}(M)/M$, there exists a pure submodule B of $\mathcal{E}(M)$ containing M such that $B/M \cong C$.*

Proof. In this case $\mathcal{E}(M)$ is also the pure-injective hull of M . By Theorem 1.1, $\mathcal{E}(M)/M$ is flat. Let C be a submodule of $\mathcal{E}(M)/M$, A its inverse image by the epimorphism $\mathcal{E}(M) \rightarrow \mathcal{E}(M)/M$. The inclusion map $M \rightarrow A \rightarrow \mathcal{E}(A)$ extends to a monomorphism $\phi : \mathcal{E}(M) \rightarrow \mathcal{E}(A)$. Let B be the inverse image of A by ϕ . Then A and B are isomorphic submodules of $\mathcal{E}(M)$ and B is a pure submodule of $\mathcal{E}(M)$. So, B/M is a pure submodule of $\mathcal{E}(M)/M$, and consequently it is flat. Since $\phi(x) = x$ for each $x \in M$, ϕ induces an isomorphism between B/M and C . We conclude that C is flat. \square

With a similar proof we get the following.

Proposition 2.5. *Let R be a ring. Assume that each Warfield cotorsion left R -module is RD -injective. Then, for each left module M , every submodule of $\mathcal{E}_W(M)/M$ is P -flat. Moreover, for each submodule C of $\mathcal{E}_W(M)/M$, there exists an RD -submodule B of $\mathcal{E}_W(M)$ containing M such that $B/M \cong C$.*

Corollary 2.6. *Let R be a ring. Assume that each left cotorsion R -module is pure-injective and each cyclic flat left module is projective. Then R is left pure-semisimple.*

Proof. Let M be a left R -module. By way of contradiction suppose that $\mathcal{E}(M)/M \neq 0$. Let C be a cyclic submodule of $\mathcal{E}(M)/M$. By Proposition 2.4 C is projective and there exists a pure submodule B of $\mathcal{E}(M)$ containing M and such that $B/M \cong C$. Then $B = M \oplus C'$ where C' is a submodule of A isomorphic to C . Then $M \cap C' = 0$ and $M \cong B/C'$ is a pure submodule of $\mathcal{E}(M)/C'$. This contradicts that $\mathcal{E}(M)$ is a pure-essential extension of M . Hence each left R -module is pure-injective. We conclude that R is left pure-semisimple. \square

Corollary 2.7. *Let R be a ring. Assume that each left Warfield cotorsion R -module is RD -injective and each cyclic flat left module is projective. Then R is left RD -semisimple.*

As in [7] a left R -module M is said to be **semi-compact** if every finitely solvable set of congruences $x \equiv x_\alpha \pmod{M[I_\alpha]}$ (where $\alpha \in \Lambda$, $x_\alpha \in M$ and I_α is a left ideal of R for each $\alpha \in \Lambda$) has a simultaneous solution in M .

Proposition 2.8. *Let R be a ring. Assume that each pure-essential extension of left R -modules is essential. Then each semi-compact left module is pure-injective.*

Proof. Let M be a semi-compact left R -module. By way of contradiction assume there exists $x \in \text{PE}(M) \setminus M$. Let $A = \{a \in R \mid ax \in M\}$. Then $A \neq 0$ since the extension $M \rightarrow \text{PE}(M)$ is essential. We consider the following system of equations: $aX = ax$, $a \in A$. Since M is a pure submodule then, for each finite subset B of A , there exists $x_B \in M$ such that $ax_B = ax$ for each $a \in B$. By [3, Proposition 1.2] the semi-compactness of M implies that there exists $y \in M$ such that $ax = ay$ for each $a \in A$. It follows that $R(x - y) \cap M = 0$ which contradicts that M is essential in $\text{PE}(M)$. \square

Proposition 2.9. *Let R be a ring. Assume that each pure-essential extension of left modules is essential. Then, for each FP-injective left R -module M , $\text{PE}(M)/M$ is regular.*

Proof. Let C be a submodule of $\text{PE}(M)/M$, A its inverse image by the epimorphism $\text{PE}(M) \rightarrow \text{PE}(M)/M$. The inclusion map $M \rightarrow A \rightarrow \text{PE}(A)$ is an essential extension. Hence $\text{PE}(A) \cong \text{PE}(M)$. Then $\text{PE}(A)$ is injective and A is FP-injective, whence A is a pure submodule of $\text{PE}(M)$ and C a pure submodule of $\text{PE}(M)/M$. \square

In the same way and by using Proposition 2.1 we get the following.

Proposition 2.10. *Let R be a ring. Assume that each RD-essential extension of left R -modules is essential. Then, for each P -injective left R -module M , $\text{RDE}(M)/M$ is regular.*

3. WHEN COTORSION MODULES ARE PURE-INJECTIVE: COMMUTATIVE CASE

Proposition 3.1. *Let R be a commutative ring. Assume that each pure(RD)-essential extension of R -modules is essential. Then for each multiplicative subset S of R , each pure(RD)-essential extension of $S^{-1}R$ -modules is essential.*

Proof. Let $A \rightarrow B$ be a pure-essential extension of $S^{-1}R$ -modules, and let C be an R -submodule of B such that $A \cap C = 0$ and A is a pure submodule of B/C . It is easy to check that $A \cap S^{-1}C = 0$ and A is a pure submodule of $B/S^{-1}C$. So, $S^{-1}C = 0$ and $C = 0$. Then $A \rightarrow B$ is a pure-essential extension of R -modules. Now it is easy to conclude. \square

Recall that a ring R is left **perfect** if each flat left R -module is projective.

Proposition 3.2. *Let R be a commutative ring. Assume that each semi-compact R -module is pure-injective. Then each prime ideal is maximal.*

Proof. Let L be a prime ideal of R , $R' = R/L$ and M a flat R' -module. Since R' is a domain, each flat R' -module is semi-compact over R' and over R too. It follows that each flat R' -module is pure-injective. There is a pure-exact sequence $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ where F is a free R' -module. So, K is flat and pure-injective over R' . We deduce that the above sequence splits and consequently M is projective. Hence R' is a perfect domain, whence R' is a field and L is maximal. \square

Theorem 3.3. *Let R be a commutative ring satisfying each pure-essential extension of R -modules is essential. Then R_P is pure-semisimple for any maximal ideal P .*

Proof. By Proposition 3.1 we may assume that R is local of maximal ideal P . Let $I = \text{E}_R(R/P)$, $M = I^{(\mathbb{N})}$, $E = \text{E}_R(M)$ and $S = E/M$. By Propositions 2.9 and 2.1 S is semisimple. Let $0 \neq a \in P$, $A = (0 : a)$ and $R_a = R/A$. By Propositions 2.8 and 3.2 P is the sole prime ideal. So, $A \neq 0$. For any R -module G we put $G' = \{g \in G \mid Ag = 0\}$. Then $I' = \text{E}_{R_a}(R/P) = aI$, $M' = I'^{(\mathbb{N})} = aM$, $E' = aE$ and E' is injective over R_a . Since $aS = 0$ then $M' = E'$. By [2, Theorem 25.3] R_a is Noetherian, and Artinian since P is the sole prime ideal. Let $(Ra_n)_{n \in \mathbb{N}}$ be a descending chain of proper ideals of R . We may assume that $a_0 \in P$. If we choose $a = a_0$, then Ra is an R_a -module. So, it is Artinian and consequently R satisfies the descending condition on principal ideals. We conclude that R is perfect by [9, 43.9]. It follows that $P^2 \neq P$. By way of contradiction suppose that P/P^2

is a vector space of dimension ≥ 2 over R/P . Then there is a Noetherian factor R' of R modulo a suitable ideal whose maximal ideal is generated by 2 elements. So, R' is not pure-semisimple. But, we successively get that each R' -module is semicomact (because R' is Noetherian), and pure-injective by Proposition 2.8. From this contradiction we deduce that P is principal and R pure-semisimple. \square

Theorem 3.4. *Let R be a commutative ring satisfying each RD-essential extension of R -modules is essential. Then R_P is pure-semisimple for any maximal ideal P .*

Proof. By Proposition 3.1 we may assume that R is local of maximal ideal P .

First assume that R is Noetherian. If $R' = R/P^2$, then each RD-essential extension of R' -modules is essential too. If P/P^2 is of dimension > 1 over R/P then, by [5, Lemma 2.4] applied to $A = P$, the RD-injective hull of R/P over R' is not an essential extension. Hence P is principal. Since $\bigcap_{n \in \mathbb{N}} P^n = 0$, then it is easy to show that $\{P^n \mid n \in \mathbb{N}\}$ is the set of principal ideals of R . Hence R is a chain ring. So, each RD-exact sequence is pure. Consequently R satisfies the assumption of Theorem 3.3, whence R is pure-semisimple.

Now, assume that R is not Noetherian. Let L be a prime ideal and $R'' = R/L$. Suppose that $L \neq P$. Then, from the first part of the proof R'' is not Noetherian. Let N be a FP-injective R'' -module which is not injective, $E = E_{R''}(N)$ and $T = E/N$. Let $a \in P \setminus L$. Then $E = aE$, whence $T = aT$. But, by Propositions 2.10 and 2.1 T is semisimple, whence $aT = 0$. From this contradiction we deduce that $L = P$. Now, we do as in the proof of Theorem 3.3 to show that R is perfect. So, P/P^2 is of infinite dimension over R/P . Whence there exists Noetherian factor rings of R which are not pure-semisimple. This contradicts the beginning of the proof. Hence R is pure-semisimple. \square

Proposition 3.5. *Let R be a ring, E a left R -module and U a pure submodule of E . Then the following conditions are equivalent:*

- (1) E/U is FP-injective if E is FP-injective;
- (2) E/U is FP-injective if E is an injective hull of U .

Proof. It is obvious that (1) \Rightarrow (2).

(2) \Rightarrow (1). First we assume that E is injective. Then E contains a submodule E' which is an injective hull of U . Since E/E' is injective and E'/U FP-injective, then E/U is FP-injective too. Now we assume that E is FP-injective. Let H be the injective hull of E . Then E/U is a pure submodule of H/U . We conclude that E/U is FP-injective. \square

Theorem 3.6. *Let R be a commutative ring. Assume that each cotorsion R -module is pure-injective. Then:*

- (1) for each maximal ideal P , R_P is pure-semisimple;
- (2) R is coherent.

Proof. (1). For any maximal ideal P , each cotorsion R_P -module is pure-injective over R_P . So, we conclude by Corollary 2.6

(2). We shall prove that E/U is FP-injective for any FP-injective module E and any pure submodule U of E . By Proposition 3.5 we may assume that E is the injective hull of U . So, $E \cong \mathcal{E}(U)$. By Proposition 2.4 E/U is flat. Then, for each maximal ideal P , $(E/U)_P$ is flat, hence free and injective since R_P is pure-semisimple. We conclude that E/U is FP-injective and R is coherent by [9, 35.9]. \square

In a same way we prove the following theorem.

Theorem 3.7. *Let R be a commutative ring. Assume that each Warfield cotorsion R -module is RD -injective. Then, for each maximal ideal P , R_P is pure-semisimple, and R is coherent.*

Proposition 3.8. *Let R be a commutative ring whose prime ideals are maximal. Let X be the set of all maximal ideals P such that $PR_P = 0$. We denote by A , the kernel of the naturel map $R \rightarrow \prod_{P \in X} R_P$. If R is P -coherent then, A is a pure submodule of R and $X = V(A)$.*

Proof. Since R/A is a subring of a product of fields, then R/A is reduced. Since each prime ideal is maximal we deduce that R/A is von Neumann regular. Thus R/A is a pure submodule of $\prod_{P \in X} R_P$ which is P -flat because R is P -coherent. It follows that A is a pure ideal. Since $A_P = 0$ for each $P \in X$ then $X \subseteq V(A)$. Let $P \in V(A)$. Then $A_P = 0$ since A is pure. It is obvious that $J \subseteq A$ where J is the Jacobson radical of R . Since J is also the nilradical of R , then $PR_P = JR_P = 0$. Hence $P \in X$. \square

Theorem 3.9. *Let R be a commutative ring. The following conditions are equivalent:*

- (1) *each cotorsion R -module is pure-injective;*
- (2) *each Warfield cotorsion R -module is RD -injective;*
- (3) *R is P -coherent and each pure-essential extension of R -modules is essential;*
- (4) *R is P -coherent and each RD -essential extension of R -modules is essential;*
- (5) *any R -module M is pure-injective if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat R -module C ;*
- (6) *any R -module M RD -injective if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat R -module C ;*
- (7) *there exists a family \mathfrak{E} of orthogonal irreducible idempotents of R satisfying the following conditions:*
 - (a) *$R/R(1 - e)$ is a pure-semisimple ring but not a field, for each $e \in \mathfrak{E}$;*
 - (b) *R/A is a von Neumann regular ring where $A = \bigoplus_{e \in \mathfrak{E}} Re$.*

Moreover, when these conditions hold, $\text{PE}(M)/M$ is flat, FP -injective and regular for each R -module M , where $\text{PE}(M)$ is the pure-injective hull of M .

Proof. It is obvious that (5) \Rightarrow (1) and (6) \Rightarrow (2). If R satisfies condition (1), (2), (3) or (4) then, by Theorems 3.6, 3.7, 3.3 or 3.4, R is arithmetical. It follows that (1) \Leftrightarrow (2) and (4) \Leftrightarrow (3).

(1) \Rightarrow (7). By Theorem 3.6, R is coherent and R_P is pure-semisimple for each maximal ideal P . Let A be the pure ideal of R defined in Proposition 3.8. By Proposition 1.3 A is generated by its idempotents. Let $e = e^2 \in A$. Then $R' = R/R(1 - e)$ satisfies (1). Since R' is locally self injective and coherent by Theorem 3.6 then each R' -module is flat if and only if it is FP -injective. Let M be an R' -module. By Proposition 2.4 each submodule of $\mathcal{E}(M)/M$ is flat, FP -injective and pure. So, $\mathcal{E}(M)/M$ is regular. Then, for each $P \in V(1 - e)$, $(\mathcal{E}(M)/M)_P$ is flat and semisimple by Proposition 2.4. Since $PR_P \neq 0$, it follows that $M_P = \mathcal{E}(M)_P$ for each $P \in V(1 - e)$, and $M = \mathcal{E}(M)$. Then R' is pure-semisimple. So, R' is a finite product of local rings. We deduce that e is a sum of orthogonal irreducible idempotents. So,

$$\mathfrak{E} = \{e_P \mid P \in D(A) \text{ and } D(e_P) = \{P\}\}.$$

(3) \Rightarrow (7). Since R is locally pure-semisimple by Theorem 3.3 and coherent we do as above to define the pure ideal A . Then, by using Proposition 2.9, we show that each FP-injective R' -module is injective. So, R' is Noetherian, and Artinian since each prime ideal is maximal. We end as above.

By Proposition 2.2 (7) \Rightarrow (5) and (3), and by Proposition 2.3 (7) \Rightarrow (6). \square

4. WHEN FLAT-ESSENTIAL EXTENSIONS ARE ESSENTIAL

Proposition 4.1. *Let R be a ring. Assume that each flat-essential extension of left R -modules is essential. Then, for each left module M , every submodule of $\mathcal{E}(M)/M$ is flat. Moreover, for each submodule C of $\mathcal{E}(M)/M$, there exists a pure submodule B of $\mathcal{E}(M)$ containing M such that $B/M \cong C$.*

Proof. Let C be a submodule of $\mathcal{E}(M)/M$, A its inverse image by the epimorphism $\mathcal{E}(M) \rightarrow \mathcal{E}(M)/M$ and $\mathcal{E}(A)$. The inclusion map $M \rightarrow A \rightarrow \mathcal{E}(A)$ extends to a monomorphism $\phi : \mathcal{E}(M) \rightarrow \mathcal{E}(A)$. Let B be the inverse image of A by ϕ . Then A and B are isomorphic submodules of $\mathcal{E}(M)$ and B is a pure submodule of $\mathcal{E}(M)$. So, B/M is a pure submodule of $\mathcal{E}(M)/M$, and consequently it is flat. Since $\phi(x) = x$ for each $x \in M$, ϕ induces an isomorphism between B/M and C . We conclude that C is flat. \square

In the same way we show the following.

Proposition 4.2. *Let R be a ring. Assume that each P -flat-essential extension of left R -modules is essential. Then, for each left module M , every submodule of $\mathcal{E}_W(M)/M$ is P -flat. Moreover, for each submodule C of $\mathcal{E}_W(M)/M$, there exists an RD -submodule B of $\mathcal{E}_W(M)$ containing M such that $B/M \cong C$.*

Corollary 4.3. *Let R be a ring. Assume that each flat-essential extension of left R -modules is essential and each cyclic flat left module is projective. Then R is left perfect.*

Proof. Let M be a left R -module and $\mathcal{E}(M)$ its cotorsion envelope. By way of contradiction suppose that $T = \mathcal{E}(M)/M \neq 0$. Let C be a cyclic submodule of T . By Proposition 4.1 C is projective and there exists a pure submodule B of $\mathcal{E}(M)$ containing M and such that $B/M \cong C$. So, if $C_1 = B/M$ then C_1 is a pure submodule of T and T/C_1 is flat. Moreover $B = M \oplus C'$ where C' is a submodule of A isomorphic to C . Thus $M \cap C' = 0$ and since $\mathcal{E}(M)/B \cong T/C_1$ then $\mathcal{E}(M)/C'$ is a flat extension of $M \cong B/C'$. This contradicts that $\mathcal{E}(M)$ is a flat-essential extension of M . Hence each left R -module is cotorsion. We conclude that R is left perfect. \square

We say that a ring R is left **strongly perfect** if each P -flat left R -module is projective. Clearly every left strongly perfect ring is perfect, but [1, Proposition 4.8] shows that there exist Artinian commutative rings which are not strongly perfect. And [3, Example 3.2] is strongly perfect by [1, Theorem 4.11] and it is non-Artinian if Λ is not finite.

Corollary 4.4. *Let R be a ring. Assume that each P -flat-essential extension of left R -modules is essential and each cyclic flat left module is projective. Then R is left strongly perfect.*

Proof. We show that each left R -module is Warfield cotorsion. It follows that each P -flat left module is projective. \square

Theorem 4.5. *Let R be a commutative ring. Assume that each flat-essential extension of R -modules is essential. Then R_P is perfect for each maximal ideal P .*

Proof. Let P be a maximal ideal. It is easy to check that any flat-essential of R_P -modules is essential. Since each flat cyclic R_P -module is free we apply Corollary 4.3 to conclude. \square

In a similar way we show the following theorem by using Corollary 4.4.

Theorem 4.6. *Let R be a commutative ring. Assume that each P -flat-essential extension of left R -modules is essential. Then R_P is strongly perfect for each maximal ideal P .*

Theorem 4.7. *Let R be a ring. Consider the following conditions:*

- (1) *there exists a family \mathfrak{E} of orthogonal central idempotents of R satisfying the following conditions:*
 - (a) *$R/R(1 - e)$ is left perfect for each $e \in \mathfrak{E}$;*
 - (b) *R/A is a von Neumann regular ring where $A = \bigoplus_{e \in \mathfrak{E}} Re$.*
- (2) *each flat-essential extension of left R -modules is essential;*
- (3) *any left R -module M is cotorsion if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat left R -module C .*

Then:

- (i) *(1) implies (2) and (3). Moreover, when this conditions holds, $\mathcal{E}(M)/M$ is flat, FP-injective and regular for each left R -module M .*
- (ii) *if R is commutative and P -coherent then (1) and (2) are equivalent, and in this case $R/(1 - e)R$ is Artinian for each $e \in \mathfrak{E}$.*

Proof. (i). We do as in the proof of Proposition 2.2.

(ii). Let A be the pure ideal of R defined in Proposition 3.8. We know that A is generated by its idempotents. Let $e = e^2 \in A$. Then $R' = R/R(1 - e)$ satisfies (2). Let M be an R' -module. By Proposition 4.1 each submodule of $\mathcal{E}(M)/M$ is flat. By way of contradiction suppose that $T = \mathcal{E}(M)/M \neq 0$. There exists a maximal ideal $P \in D(e)$ such that $T_P \neq 0$. Then each nonzero cyclic submodule of T_P is free. This implies that each principal ideal of R_P is free. But $PR_P \neq 0$ and all its elements are zerodivisor. Hence $T = 0$ and each R' -module is cotorsion. We conclude that R' is perfect. So, it is a finite product of local rings. Since R is coherent then R' is Artinian. Now we end as in the proof of (1) \Rightarrow (6) of Theorem 3.9. \square

Theorem 4.8. *Let R be a ring. Consider the following conditions:*

- (1) *there exists a family \mathfrak{E} of orthogonal central idempotents of R satisfying the following conditions:*
 - (a) *$R/R(1 - e)$ is left strongly perfect for each $e \in \mathfrak{E}$;*
 - (b) *R/A is a von Neumann regular ring where $A = \bigoplus_{e \in \mathfrak{E}} Re$.*
- (2) *each P -flat-essential extension of left R -modules is essential;*
- (3) *any left R -module M is Warfield cotorsion if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat left R -module C .*

Then:

- (i) *(1) implies (2) and (3). Moreover, when this conditions holds, $\mathcal{E}_W(M)/M$ is flat, FP-injective and regular for each left R -module M .*

- (ii) if R is commutative and coherent then (1) and (2) are equivalent, and in this case $R/(1-e)R$ is Artinian for each $e \in \mathfrak{E}$.

Theorem 4.9. *Let R be commutative ring. Then the following conditions are equivalent:*

- (1) R_P is perfect for each maximal ideal P ;
- (2) any R -module M is cotorsion if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat R -module C .

Proof. (2) \Rightarrow (1). Let P be a maximal ideal and M an R_P -module. If C is a nonzero cyclic flat R -module, then C_P is free over R_P . It follows that $\text{Ext}_R^1(C, M) \cong \text{Ext}_{R_P}^1(C_P, M) = 0$. So, M is cotorsion over R and R_P . Since each R_P -module is cotorsion then R_P is perfect.

(1) \Rightarrow (2). Let M be an R -module satisfying $\text{Ext}_R^1(C, M) = 0$ for any flat cyclic R -module C . Let F be a free R -module, K a pure submodule of F and $\alpha : K \rightarrow M$ a homomorphism. We must prove that α extends to F . We consider the family $\mathcal{F} = \{(N, \beta)\}$ where N is a pure submodule of F containing K and β an extension of α to N . We consider the following order on \mathcal{F} : $(N, \beta) \leq (L, \gamma)$ if and only if $N \subseteq L$ and $\gamma|_N = \beta$. It is easy to see that we can apply Zorn Lemma to \mathcal{F} . So, let (N, β) be a maximal element of \mathcal{F} . By way of contradiction suppose that $N \neq F$. Let $G = F/N$. There exists a maximal ideal P such that $G_P \neq 0$. Since R_P is perfect then G_P is free over R_P . Thus there exists $x \in F \setminus N$ such that its image y in G_P verifies $(0 :_{R_P} y) = 0$. It follows that $(N : x) = 0_P$ (see Proposition 1.3). Let $\delta : 0_P \rightarrow M$ be the homomorphism defined by $\delta(a) = \beta(ax)$ for any $a \in 0_P$. Then δ extends to R . Now, let $\phi : N + Rx \rightarrow M$ be the homomorphism defined by $\phi(n + rx) = \beta(n) + \delta(r)$ for any $n \in N$ and $r \in R$. It is easy to check that ϕ is well defined. Let $H = N + Rx/N$. Then $H \cong R_P$. So, H_P is a direct summand of G_P and if P' is another maximal ideal then $H_{P'} = 0$. We successively deduce that H is a pure submodule of G , $F/N + Rx$ is flat and $N + Rx$ is a pure submodule of F . This contradicts the maximality of (N, β) . Hence $N = F$ and M is cotorsion. \square

Corollary 4.10. *Let R be commutative ring. Then the following conditions are equivalent:*

- (1) R_P is strongly perfect for each maximal ideal P ;
- (2) any R -module M is Warfield cotorsion if and only if $\text{Ext}_R^1(C, M) = 0$ for each cyclic flat R -module C .

Proof. Let G be a P-flat R -module. For each maximal ideal P G_P is P-flat. Since R_P is strongly perfect then G_P is free. Hence G is flat. So, each cotorsion R -module is Warfield cotorsion. \square

ACKNOWLEDGEMENTS

This work was presented at the "second International Conference on Mathematics and Statistics (AUS-ICMS15)" held at American University of Sherjah, April 2-5, 2015. I thank again the organizers of this conference. I thank too the laboratory of mathematics Nicolas Oresme of the university of Caen Basse-Normandie which allowed me to participate to this conference.

REFERENCES

- [1] W. Al-Kawarit and F. Couchot. Comparison of some purities, flatnesses and injectivities. *Comm. Alg.*, 39(10):3879–3896, (2011).
- [2] F.W. Anderson and K.R. Fuller. *Rings and Categories of Modules*. Graduate Texts in Mathematics. Springer-Verlag, (1992).
- [3] M. Behboodi, F. Couchot, and S.H. Shojae. Σ -semi-compact rings and modules. *J.A.A.*, 13(8), (2014).
- [4] L. Bican, R. El Bashir, and E. Enochs. All modules have flat covers. *Bull. London Math. Soc.*, 33:385–390, (2001).
- [5] F. Couchot. RD-flatness and RD-injectivity. *Comm. Algebra*, 34:3675–3689, (2006).
- [6] P. Griffith. On the decomposition of modules and generalized left uniserial rings. *Math. Ann.*, 184:300–308, (1970).
- [7] E. Matlis. Injective modules over Prüfer rings. *Nagoya Math. J.*, 15:57–69, (1959).
- [8] R.B. Warfield. Purity and algebraic compactness for modules. *Pac. J. Math.*, 28(3):689–719, (1969).
- [9] R. Wisbauer. *Foundations of Module and Ring Theory*. (1991).
- [10] J. Xu. Flat covers of modules. volume 1634 of *Lecture Notes in Mathematics*. Springer, (1996).

UNIVERSITÉ DE CAEN BASSE-NORMANDIE, CNRS UMR 6139 LMNO, F-14032 CAEN, FRANCE
E-mail address: francois.couchot@unicaen.fr