



**HAL**  
open science

# ON THE LINKS-GOULD INVARIANT AND THE SQUARE OF THE ALEXANDER POLYNOMIAL

Ben-Michael Kohli

► **To cite this version:**

Ben-Michael Kohli. ON THE LINKS-GOULD INVARIANT AND THE SQUARE OF THE ALEXANDER POLYNOMIAL. 2015. hal-01165480v1

**HAL Id: hal-01165480**

**<https://hal.science/hal-01165480v1>**

Preprint submitted on 19 Jun 2015 (v1), last revised 29 Nov 2015 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE LINKS-GOULD INVARIANT AND THE SQUARE OF THE ALEXANDER POLYNOMIAL

BEN-MICHAEL KOHLI

ABSTRACT. This paper gives a connection between well chosen reductions of the Links-Gould invariants of oriented links and powers of the Alexander-Conway polynomial. We prove these formulas by showing the representations of the braid groups we derive the specialized Links-Gould polynomials from can be seen as exterior powers of copies of Burau representations.

## CONTENTS

Introduction	1
1. Definitions and main result	2
2. The reduced Links-Gould invariant expressed as a product	5
3. Proof of the main theorem	9
4. Generalizing the proof	11
5. Appendix : proof of proposition 2.4	16
References	19

## INTRODUCTION

The Links-Gould invariants of oriented links  $LG^{m,n}(L; t_0, t_1)$  are two variable polynomial quantum invariants. In [1], David De Wit, Atsushi Ishii and Jon Links proved the following equalities :

$$LG^{1,n}(L; t_0, e^{2i\pi/n}t_0^{-1}) = \Delta_L(t_0^n)$$

where  $\Delta_L(t)$  is the Alexander-Conway invariant of  $L$ . So the Links-Gould invariants contain some topological information. We reinforce that statement by proving the following identity, that the authors we just cited had already conjectured and proved for particular links :

$$LG^{n,1}(L; t_0, t_0^{-1}) = \Delta_L(t_0)^n$$

when  $n = 2, 3$ . There is no known set of complete skein relations for the square of the Alexander polynomial, so the ideas used in [1] cannot be transposed to our case easily. On the other hand, Ivan Marin and Emmanuel Wagner give a complete set of skein relations for  $LG^{2,1}$  in [10]. So evaluating them and testing whether the square of the Alexander polynomial satisfies these evaluated skein relations or not is a possible strategy. However, the cubic skein relation is barely practicable, and such an approach can not be generalized to  $n$  greater than 2.

Our strategy will be to use the robustness of the braid structure to encode links. We express the Alexander-Conway polynomial as a quantum trace as it is done in [11], appendix C. Then we prove the R-matrix representation of braid group  $B_n$  used to define reduced Links-Gould invariant  $LG^{2,1}$  (resp.  $LG^{3,1}$ ) is isomorphic to the exterior power of a direct sum of Burau representations. That way, the specialized Links-Gould invariants can be

---

*Key words and phrases.* Link, knot, Alexander-Conway polynomial, Links-Gould invariant, R-matrix.

written as products of terms, each of which can be identified with the Alexander polynomial of our link seen as a quantum trace.

Our result along with the one we cited at the beginning of this introduction can also be thought of as a counterpart to the well known result stating that the Jones polynomial and its square can both be recovered as evaluations of the two variable Kauffman polynomial. See [8], Proposition 16.6, p. 180.

Let us also mention the work of Nathan Geer and Bertrand Patereau-Mirand who extended the Links-Gould invariant to a multivariable link invariant in the same fashion the multivariable Alexander polynomial arises from its traditional counterpart [5]. We suspect that our results remain true in some sense in that multivariable context.

In section 1, we recall the definition of the Links-Gould invariant of oriented links, and an expression of the Alexander-Conway polynomial in terms of a partial trace. In section 2 we show that the specialized Links-Gould invariant  $LG^{2,1}$  can be written as a product by proving two representations of the braid group are isomorphic. We then identify in section 3 each part of the product with the Alexander-Conway invariant. Section 4 is dedicated to extending the proof to the next Links-Gould invariant  $LG^{3,1}$ .

## 1. DEFINITIONS AND MAIN RESULT

### 1.1. The Alexander-Conway polynomial.

**Definition 1.1.** (*Reduced and non-reduced Burau representations of a braid*)

Set  $\mathbb{K} := \mathbb{C}(t^{\pm\frac{1}{2}})$ . Let  $W_n = \langle f_1, \dots, f_n \rangle$  be a  $n$ -dimensional  $\mathbb{K}$ -vector space, and  $B_n$  be the braid group on  $n$  strands. We denote by  $\sigma_1, \dots, \sigma_{n-1}$  the standard Artin generators of the group. The non-reduced Burau representation  $\Psi_{W_n} : B_n \rightarrow GL(W_n)$  is given by :

$$\Psi_{W_n}(\sigma_i)(f_j) = \begin{cases} (1-t)f_i + t^{1/2}f_{i+1} & \text{if } j = i, \\ t^{1/2}f_i & \text{if } j = i+1, \\ f_j & \text{otherwise.} \end{cases}$$

Denote by  $\delta_n := t^{-(n-1)/2}f_1 + t^{-(n-2)/2}f_2 + \dots + t^{-1/2}f_{n-1} + f_n$ . One can verify that for any  $b \in B_n$ ,  $\Psi_{W_n}(b)(\delta_n) = \delta_n$ . Hence the reduced Burau representation  $\Psi_{\widehat{W_n}} : B_n \rightarrow GL(\widehat{W_n})$  is given by :

$$\Psi_{\widehat{W_n}}(b)(\bar{x}) = \overline{\Psi_{W_n}(b)(x)}$$

where  $\widehat{W_n} := W_n / \langle \delta_n \rangle$ .

Recall that the Alexander theorem states that any link can be obtained as the closure of a given braid. Moreover, the Markov theorem allows us to define link invariants through braids with closure the link. A possible definition of the classical Alexander link invariant uses that procedure.

**Definition 1.2.** (*Alexander polynomial of a link through the Burau representation*)

The Alexander polynomial of an oriented link  $L$  is defined as :

$$\Delta_L(t) \stackrel{\bullet}{=} \frac{1-t}{1-t^n} \det(I - \Psi_{\widehat{W_n}}(b))$$

where  $b$  is any braid in  $B_n$  with closure  $L$ , and the notation  $\stackrel{\bullet}{=}$  means equality up to multiplication by a unit of  $\mathbb{C}[t^{\pm 1}]$ .

However, it is not this definition of the Alexander polynomial that will be useful to us in the following. Next theorem gives another expression, that will be the one we will consider. In particular, this definition removes the ambiguity that relied in the multiplication by a unit.

**Definition 1.3.** Let  $V$  be a 2-dimensional  $\mathbb{K}$ -vector space, and  $(e_0, e_1)$  be a basis of  $V$ . We define a representation  $\Psi_{V^{\otimes n}} : B_n \rightarrow GL(V^{\otimes n})$  of  $B_n$  :

$$\Psi_{V^{\otimes n}}(\sigma_i) = id_V^{\otimes i-1} \otimes R_1 \otimes id_V^{\otimes n-i-1}$$

where

$$R_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & t^{1/2} & 0 \\ 0 & t^{1/2} & 1-t & 0 \\ 0 & 0 & 0 & -t \end{pmatrix} \in End(V \otimes V)$$

is an R-matrix, that is a solution of the Yang-Baxter equation.

**Theorem 1.4.** Let  $L$  be an oriented link and  $b \in B_n$  be any braid with closure  $L$ . We define

$$h = \begin{pmatrix} t^{1/2} & 0 \\ 0 & -t^{1/2} \end{pmatrix} \in End(V).$$

Then :

- 1)  $\exists c \in \mathbb{K}$  such that  $trace_{2,3,\dots,n}((id_V \otimes h^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b)) = c \cdot id_V$ ,
- 2)  $c$  is a link invariant and is equal to the Alexander polynomial of  $L$ ,  $\Delta_L(t)$ .

For a detailed proof, see [11], appendix C.

*Remark 1.5.* R-matrix  $R_1$  can be recovered from the universal R-matrix of ribbon Hopf algebra  $U_\zeta(sl_2)$  at root of unity  $\zeta = -1$  thanks to a one parameter family of irreducible representations of  $U_\zeta(sl_2)$  on  $V$ . For precise explanations, see [11], p.95-97, or [13]. It may also be derived from the quantized universal enveloping algebra of  $gl(1|1)$ , that is  $U_q(gl(1|1))$ . See [13] or [12] for details.

*Remark 1.6.* Identifying algebras  $End(V^{\otimes n})$  and  $End(V)^{\otimes n}$ , the partial trace operator verifies  $trace_{2,3,\dots,n}(f_1 \otimes \dots \otimes f_n) := trace(f_2)trace(f_3) \dots trace(f_n)f_1 \in End(V)$  for any  $f_1, \dots, f_n \in End(V)$ .

**Corollary 1.7.** With the same notations, this formula follows from theorem 1.4 :

$$\Delta_L(t) = \frac{1}{2} trace((id_V \otimes h^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b)).$$

*Proof.* Applying the trace operator on each side of the formula that defines constant  $c$ , we obtain :

$$2c = trace(trace_{2,3,\dots,n}((id_V \otimes h^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b))).$$

But :

$$\begin{aligned} trace(trace_{2,3,\dots,n}(f_1 \otimes \dots \otimes f_n)) &= trace(trace(f_2)trace(f_3) \dots trace(f_n)f_1) \\ &= trace(f_2)trace(f_3) \dots trace(f_n)trace(f_1) = trace(f_1 \otimes \dots \otimes f_n) \end{aligned}$$

Since the trace and the partial trace are linear maps, we can extend the equality to any  $f \in End(V^{\otimes n}) \simeq End(V)^{\otimes n}$ , which provides the result. CQFD

## 1.2. The Links-Gould invariant $LG^{2,1}$ of links.

**Definition 1.8.** Set  $\mathbb{L} := \mathbb{C}(t_0^{\pm \frac{1}{2}}, t_1^{\pm \frac{1}{2}})$ . Let  $W = \langle e_1, \dots, e_4 \rangle$  be a four-dimensional  $\mathbb{L}$ -vector space. The following linear map  $R$ , expressed in basis  $(e_1 \otimes e_1, e_1 \otimes e_2, e_1 \otimes e_3, e_1 \otimes e_4, e_2 \otimes e_1, e_2 \otimes e_2, e_2 \otimes e_3, \dots)$ , is an automorphism of  $W \otimes W$  and an R-matrix [2], p.186 :



In the next two sections, we prove case (2, 1). We express the conjecture in the set of variables we used to introduce  $LG^{2,1}$ . We want to prove :

$$LG^{2,1}(L; \tau, -1) = \Delta_L(\tau^2)^2.$$

Variables  $(\tau, q)$  and  $(t_0, t_1)$  are related by :  $t_1^{1/2} = \tau^{-1}q$ ,  $t_0^{1/2} = \tau$  (and therefore  $t_0^{1/2}t_1^{1/2} = q$ ). Since in our case  $q = -1$ , we obtain :

$$t_0^{1/2}t_1^{1/2} = -1, \tau^2 = t_0 = t_1^{-1}.$$

Thus, once it is formulated in a convenient way, our main result states :

**Theorem 1.12.** *For any oriented link  $L$ ,  $LG^{2,1}(L; t_0, t_0^{-1}) \stackrel{\bullet}{=} \Delta_L(t_0)^2$ .*

## 2. THE REDUCED LINKS-GOULD INVARIANT EXPRESSED AS A PRODUCT

We derive a representation of the braid group  $B_n$  from the Burau representation. We identify it with a specialization of the R-matrix representation given in subsection 1.2. Then we use this identification to express the specialized Links-Gould invariant as a product.

**2.1. A representation of  $B_n$  isomorphic to  $b_R^n(t_0, t_0^{-1})$ .** Denote by  $F$  the following Burau representation of  $B_n$  on vector space  $W_n = \langle f_1, \dots, f_n \rangle$  where we replace  $t_0$  by  $t_0^{-1}$  :

$$F(\sigma_i)(f_j) = \begin{cases} (1 - t_0^{-1})f_i + t_0^{-1/2}f_{i+1} & \text{if } j = i, \\ t_0^{-1/2}f_i & \text{if } j = i + 1, \\ f_j & \text{otherwise.} \end{cases}$$

In a similar way, let  $G$  be the representation of  $B_n$  on  $n$ -dimensional vector space  $W_n = \langle g_1, \dots, g_n \rangle$  given by :

$$G(\sigma_i)(g_j) = \begin{cases} -t_0^{1/2}g_{i+1} & \text{if } j = i, \\ -t_0^{1/2}g_i + (1 - t_0)g_{i+1} & \text{if } j = i + 1, \\ g_j & \text{otherwise.} \end{cases}$$

**Proposition 2.1.** *Representation  $G$  is isomorphic to the Burau representation of  $B_n$ .*

*Proof.* One can verify that for  $i = 1, 2, \dots, n - 1$  :  $J_n \circ \Psi_{W_n}(\sigma_i) = G(\sigma_i) \circ J_n$  where  $J_n$  can be defined inductively :  $J_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  and

$$J_n = \left( \begin{array}{ccc|c} & & & (-1)^{2t^{(n-2)/2}} \\ & & & \vdots \\ & & & (-1)^{n-1}t^{1/2} \\ & & & (-1)^nt^{0/2} \\ \hline & J_{n-1} & & 0 \\ (-1)^{n+1}t^{-(n-2)/2} & \dots & (-1)^{n+1}t^{-1/2} & (-1)^{n+1}t^{-0/2} \end{array} \right).$$

Moreover, evaluating the determinant of  $J_n$ , we deduce that  $J_n$  is an automorphism. Indeed,  $\det J_{n+1} = (-1)^{n+1}(t^{1/2} + t^{-1/2})\det J_n + \det J_{n-1}$ . So  $\det J_n \in \mathbb{Z}[t^{\pm 1/2}]$  is invertible in  $\mathbb{Q}(t^{\pm 1/2})$  since it has degree  $n - 2$  in both variables  $t^{1/2}$  and  $t^{-1/2}$ . CQFD

**Definition 2.2.** If  $F \oplus G$  is the representation of the  $n$ -strand braid group on  $W_n \oplus W_n$  built from  $F$  and  $G$ , we consider the exterior representation  $\Psi_n := \wedge(F \oplus G)$  on exterior algebra  $\wedge(W_n \oplus W_n)$ .

*Remark 2.3.* Note that  $b_R^n(t_0, t_0^{-1})$  and  $\Psi_n$  both are  $4^n = 2^{2n}$ -dimensional representations.

We are going to show these two representations are isomorphic. For that we study first the case where  $n = 2$ . Since  $t_1 = t_0^{-1}$ , we have a simpler R-matrix  $R$  :







2.2. **A convenient expression for  $LG^{2,1}$ .** Now we have built an exterior representation that is isomorphic to  $b_R^n(t_0, t_0^{-1})$ , we use it to write the reduction of the Links-Gould polynomial as the product of two quantities we will then identify.

Using proposition 2.4, we can write :

$$LG(L; t_0, t_0^{-1}) = \frac{1}{4} \text{trace}(\underbrace{I_n \circ (id_W \otimes \mu^{\otimes n-1}) \circ I_n^{-1}}_{\tilde{\mu}} \circ \Psi_n(b)).$$

We wish to explicit  $\tilde{\mu}$ .

**Lemma 2.5.** *Map  $\tilde{\mu}$  can be expressed on the reference basis of  $\bigwedge(W_n \oplus W_n)$  :*

$$\begin{aligned} \tilde{\mu}(f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m}) &= t_0^{-(n-1)} (-1)^{n-1} (-1)^{\#\{k \in \{2, \dots, n\} | f_k \text{ appears}\}} \\ &\quad (-1)^{\#\{k \in \{2, \dots, n\} | g_k \text{ appears}\}} f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m}. \end{aligned}$$

*Proof.* If  $f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m}$  is an element of the basis of  $\bigwedge(W_n \oplus W_n)$ , we denote by  $e_{l_1} \otimes \dots \otimes e_{l_n}$  it's image under  $I_n^{-1}$ . That way,  $I_n(e_{l_1} \otimes \dots \otimes e_{l_n}) = f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m}$ .

$$\begin{aligned} \tilde{\mu}(f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m}) &= I_n \circ (id_W \otimes \mu^{\otimes n-1})(e_{l_1} \otimes \dots \otimes e_{l_n}) \\ &= t_0^{-(n-1)} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 2\}} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 3\}} \\ &\quad I_n(e_{l_1} \otimes \dots \otimes e_{l_n}) \\ &= t_0^{-(n-1)} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 2\}} \\ &\quad (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 3\}} f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m} \end{aligned}$$

But :

$$\begin{aligned} &(-1)^{\#\{k \in \{2, \dots, n\} | l_k = 2\}} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 3\}} \\ &= (-1)^{n-1} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 1\}} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 4\}} \\ &= (-1)^{n-1} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 1\}} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 4\}} \left( (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 3\}} \right)^2 \\ &= (-1)^{n-1} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 3 \text{ or } l_k = 4\}} (-1)^{\#\{k \in \{2, \dots, n\} | l_k = 1 \text{ or } l_k = 3\}} \\ &= (-1)^{n-1} (-1)^{\#\{k \in \{2, \dots, n\} | f_k \text{ appears}\}} (-1)^{\#\{k \in \{2, \dots, n\} | g_k \text{ appears}\}} \end{aligned}$$

This provides the result.

CQFD

Given the expression for  $\tilde{\mu}$  we just obtained, and the special form of representation  $\Psi_n$  we have :

**Proposition 2.6.** *Invariant  $LG(L; t_0, t_0^{-1})$  can be written as a product, with each term depending only on one of the copies of the Burau representation.*

*Proof.* Recall  $LG(L; t_0, t_0^{-1}) = \frac{1}{4} \text{trace}((id_W \otimes \mu^{\otimes n-1}) \circ b_R^n(b))$ , where :

$$\mu = \begin{pmatrix} t_0^{-1} & \cdot & \cdot & \cdot \\ \cdot & -t_0^{-1} & \cdot & \cdot \\ \cdot & \cdot & -t_0^{-1} & \cdot \\ \cdot & \cdot & \cdot & t_0^{-1} \end{pmatrix}.$$

Using that we can write :

$$\begin{aligned} LG(L; t_0, t_0^{-1}) &= \frac{1}{4} \text{trace}(\tilde{\mu} \circ \Psi_n(b)) \\ &= \frac{1}{4} \sum_{\substack{1 \leq i_1 < \dots < i_p \leq n \\ 1 \leq j_1 < \dots < j_m \leq n}} (f_{i_1} \wedge \dots \wedge g_{j_m})^* (\tilde{\mu} \circ \Psi_n(b)(f_{i_1} \wedge \dots \wedge g_{j_m})) \end{aligned}$$

where  $(f_{i_1} \wedge \dots \wedge g_{j_m})^*$  indicates a vector of the dual basis of the reference basis. But given lemma 2.5,

$$\begin{aligned} (f_{i_1} \wedge \dots \wedge g_{j_m})^* (\tilde{\mu} \circ \Psi_n(b)(f_{i_1} \wedge \dots \wedge g_{j_m})) &= (-t_0)^{-(n-1)} (-1)^{\#\{k \in \{2, \dots, n\} | f_k \text{ appears}\}} \\ &\quad (-1)^{\#\{k \in \{2, \dots, n\} | g_k \text{ appears}\}} \\ &\quad (f_{i_1} \wedge \dots \wedge g_{j_m})^* (\Psi_n(b)(f_{i_1} \wedge \dots \wedge g_{j_m})) \end{aligned}$$

$$\begin{aligned} \text{Also, } (f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m})^* &\underbrace{(\Psi_n(b)(f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m}))}_{\wedge F(b)(f_{i_1} \wedge \dots \wedge f_{i_p}) \wedge \wedge G(b)(g_{j_1} \wedge \dots \wedge g_{j_m})} \\ &= (f_{i_1} \wedge \dots \wedge f_{i_p})^* (\wedge F(b)(f_{i_1} \wedge \dots \wedge f_{i_p})) (g_{j_1} \wedge \dots \wedge g_{j_m})^* (\wedge G(b)(g_{j_1} \wedge \dots \wedge g_{j_m})). \end{aligned}$$

That way we have the following expression for  $LG(L; t_0, t_0^{-1})$ :

$$\begin{aligned} \frac{1}{4} (-t_0)^{-(n-1)} \sum_{1 \leq i_1 < \dots < i_p \leq n} &\left( (-1)^{\#\{k \in \{2, \dots, n\} | f_k \text{ appears}\}} (f_{i_1} \wedge \dots \wedge f_{i_p})^* (\wedge F(b)(f_{i_1} \wedge \dots \wedge f_{i_p})) \right) \\ * \sum_{1 \leq j_1 < \dots < j_m \leq n} &\left( (-1)^{\#\{k \in \{2, \dots, n\} | g_k \text{ appears}\}} (g_{j_1} \wedge \dots \wedge g_{j_m})^* (\wedge G(b)(g_{j_1} \wedge \dots \wedge g_{j_m})) \right) \end{aligned}$$

CQFD

Now we wish to show that each of these two sums is equal to  $\Delta_L(t_0)$  up to multiplication by a unit of  $\mathbb{C}[t_0^{\pm 1}]$ , that is up to multiplication by  $\pm t_0^n$ ,  $n \in \mathbb{Z}$ .

### 3. PROOF OF THE MAIN THEOREM

A careful analysis of [11], appendix C, shows that we have a coefficient in front of the partial trace in the expression of the Alexander-Conway polynomial :

$$\Delta_L(t_0) \stackrel{\bullet}{=} t_0^{-(n-1)/2} \frac{1}{2} \text{trace}((id_V \otimes h^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b)).$$

So we can write more simply :

$$\Delta_L(t_0) \stackrel{\bullet}{=} \frac{1}{2} \text{trace}((id_V \otimes \tilde{h}^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b)), \text{ where } \tilde{h} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

**Proposition 3.1.** *Let  $J_n : V^{\otimes n} \longrightarrow \wedge W_n$ ,  $e_{i_1} \otimes \dots \otimes e_{i_n} \longmapsto \bigwedge_{k : i_k=1} f_k$ . Then  $J_n$  is a  $\mathbb{C}[B_n]$ -module automorphism :*

$$\bigwedge \Psi_{W_n}(b) \circ J_n = J_n \circ \Psi_{V^{\otimes n}}(b), \forall b \in B_n.$$

The proof is quite similar to the one we did in the previous section. It is detailed in [11], appendix C, where what we just called  $J_n$  is denoted by  $I_n$ , and is introduced by induction.

Applying  $J_n$ , we can express the Alexander polynomial differently :

$$\Delta_L(t_0) \stackrel{\bullet}{=} \frac{1}{2} \text{trace} \left( \underbrace{J_n \circ (id_V \otimes \tilde{h}^{\otimes n-1}) \circ J_n^{-1}}_{\mu_1} \circ \bigwedge \Psi_{W_n}(b) \right).$$

Where :

$$\begin{aligned} \mu_1(f_{i_1} \wedge \dots \wedge f_{i_p}) &= J_n \circ (id_V \otimes \tilde{h}^{\otimes n-1})(e_0 \otimes \dots \otimes e_0 \otimes \underbrace{e_1}_{i_1^{th} \text{ position}} \otimes e_0 \otimes \dots \otimes e_0 \otimes \underbrace{e_1}_{i_2^{th} \text{ position}} \otimes \dots) \\ &= (-1)^{\#\{k \in \{2, \dots, n\} | i_k = 1\}} f_{i_1} \wedge \dots \wedge f_{i_p} \\ &= (-1)^{\#\{k \in \{2, \dots, n\} | f_k \text{ appears}\}} f_{i_1} \wedge \dots \wedge f_{i_p}. \end{aligned}$$

Therefore,

$$\Delta_L(t_0) \stackrel{\bullet}{=} \frac{1}{2} \sum_{1 \leq i_1 < \dots < i_p \leq n} \left( (-1)^{\#\{k \in \{2, \dots, n\} | f_k \text{ appears}\}} (f_{i_1} \wedge \dots \wedge f_{i_p})^* \left( \bigwedge \Psi_{W_n}(b)(f_{i_1} \wedge \dots \wedge f_{i_p}) \right) \right).$$

But  $F$  and  $\Psi_{W_n}$  are identical once you change  $t_0$  into  $t_0^{-1}$ . That way we can identify the first factor of our product with  $\Delta_L(t_0^{-1})$ . But the Alexander polynomial is symmetric :  $\Delta_L(t_0) \stackrel{\bullet}{=} \Delta_L(t_0^{-1})$  [4]. So the only remaining problem is to identify the second sum with the Alexander invariant to be able to conclude. To do that we have to modify the representation of  $V^{\otimes n}$  we used up to now to define  $\Delta_L(t_0)$ , and especially R-matrix  $R_1$  we introduced at the beginning.

**Lemma 3.2.** *We can slightly modify R-matrix  $R_1$  so that the new representations  $\rho_{V^{\otimes n}}$  of the braid groups we obtain that way still verify :*

$$\Delta_L(t_0) \stackrel{\bullet}{=} \frac{1}{2} \text{trace}((id_V \otimes \tilde{h}^{\otimes n-1}) \circ \rho_{V^{\otimes n}}(b)).$$

*Proof.* For the moment, we can write :  $\Delta_L(t_0) \stackrel{\bullet}{=} \frac{1}{2} \text{trace}((id_V \otimes \tilde{h}^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b))$ , where

$$\Psi_{V^{\otimes n}} \text{ is the representation associated with R-matrix } R_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & t_0^{1/2} & 0 \\ 0 & t_0^{1/2} & 1 - t_0 & 0 \\ 0 & 0 & 0 & -t_0 \end{pmatrix}.$$

We can replace  $R_1$  by  $-t_0^{-1}R_2 = \begin{pmatrix} -t_0^{-1} & 0 & 0 & 0 \\ 0 & 0 & -t_0^{-1/2} & 0 \\ 0 & -t_0^{-1/2} & 1 - t_0^{-1} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$  in the definition of  $\Psi_{V^{\otimes n}}$ ,

and we will still have  $\Delta_L(t_0) \stackrel{\bullet}{=} \frac{1}{2} \text{trace}((id_V \otimes \tilde{h}^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b))$ . At last, we replace  $t_0$

by  $t_0^{-1}$  in  $R_2$  to obtain  $R_3 = \begin{pmatrix} -t_0 & 0 & 0 & 0 \\ 0 & 0 & -t_0^{1/2} & 0 \\ 0 & -t_0^{1/2} & 1 - t_0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ . We define the representation of

$B_n$  associated with  $R_3$  :

$$\rho_{V^{\otimes n}}(\sigma_i) = id_V^{\otimes i-1} \otimes R_3 \otimes id_V^{\otimes n-i-1}.$$

Since the Alexander polynomial is symmetric, we have the following expression for  $\Delta_L(t_0)$ , that will help us to conclude :

$$\Delta_L(t_0) \stackrel{\bullet}{=} \frac{1}{2} \text{trace}((id_V \otimes \tilde{h}^{\otimes n-1}) \circ \rho_{V^{\otimes n}}(b)).$$

Using the same strategy as previously, we wish to find  $K_n : V^{\otimes n} \longrightarrow \bigwedge W_n$  such that for any  $b \in B_n : \bigwedge G(b) \circ K_n = K_n \circ \rho_{V^{\otimes n}}(b)$ .

**Proposition 3.3.** We set  $K_1 = \begin{cases} V & \longrightarrow \bigwedge W_1 \\ e_1 & \longmapsto 1 \\ e_0 & \longmapsto g_1 \end{cases}$  and, for  $n \geq 2$ ,

$$K_n = \begin{cases} V^{\otimes n} & \longrightarrow \bigwedge W_n \\ e_{i_1} \otimes \dots \otimes e_{i_{n-1}} \otimes e_0 & \longmapsto K_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge g_n \\ e_{i_1} \otimes \dots \otimes e_{i_{n-1}} \otimes e_1 & \longmapsto K_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \end{cases} .$$

Then, for any  $b \in B_n : \bigwedge G(b) \circ K_n = K_n \circ \rho_{V^{\otimes n}}(b)$ .

*Proof.* We leave it to the reader to verify that a proof by induction resembling the one we did with  $I_n$  concludes. CQFD

That way,

$$\begin{aligned} \Delta_L(t_0) &\stackrel{\bullet}{=} \frac{1}{2} \text{trace}((id_V \otimes \tilde{h}^{\otimes n-1}) \circ \rho_{V^{\otimes n}}(b)) \\ &= \frac{1}{2} \text{trace}(\underbrace{K_n \circ (id_V \otimes \tilde{h}^{\otimes n-1}) \circ K_n^{-1}}_{\nu} \circ \bigwedge G(b)). \end{aligned}$$

We can set  $K_n(e_{i_1} \otimes \dots \otimes e_{i_n}) = g_{j_1} \wedge \dots \wedge g_{j_m}$ . That allows us to explicit the values of  $\nu$  on the natural basis of  $\bigwedge W_n$ .

$$\begin{aligned} \nu(g_{j_1} \wedge \dots \wedge g_{j_m}) &= K_n \circ id_V \otimes \tilde{h}^{\otimes n-1}(e_{i_1} \otimes \dots \otimes e_{i_n}) \\ &= (-1)^{\#\{k \in \{2, \dots, n\} | i_k = 1\}} K_n(e_{i_1} \otimes \dots \otimes e_{i_n}) \\ &= (-1)^{n-1} (-1)^{\#\{k \in \{2, \dots, n\} | i_k = 0\}} g_{j_1} \wedge \dots \wedge g_{j_m} \\ &= (-1)^{n-1} (-1)^{\#\{k \in \{2, \dots, n\} | g_k \text{ appears}\}} g_{j_1} \wedge \dots \wedge g_{j_m}. \end{aligned}$$

So :

$$\begin{aligned} \Delta_L(t_0) &= \frac{1}{2} \text{trace}(\nu \circ \bigwedge G(b)) \\ &\stackrel{\bullet}{=} \frac{1}{2} \sum_{1 \leq j_1 < \dots < j_m \leq n} \left( (-1)^{\#\{k \in \{2, \dots, n\} | g_k \text{ appears}\}} (g_{j_1} \wedge \dots \wedge g_{j_m})^* (\bigwedge G(b)(g_{j_1} \wedge \dots \wedge g_{j_m})) \right). \end{aligned}$$

And finally  $LG(L; t_0, t_0^{-1}) \stackrel{\bullet}{=} \Delta_L(t_0)^2$  for any link  $L$ .

#### 4. GENERALIZING THE PROOF

**4.1. Writing the conjecture in case  $(n, 1)$  and other considerations.** The completely general conjecture states, using variables  $(\tau, q)$  :

$$LG^{m,n}(L; \tau, e^{i\pi/n}) = \Delta_L(\tau^{2n})^m, \text{ for any link } L.$$

We can rewrite it using variables  $(t_0, t_1)$ . Indeed, since  $q = e^{i\pi/n}$ , the variables are related by  $t_1^{1/2} = \tau^{-1} e^{i\pi/n}$  and  $t_0^{1/2} = \tau$ . Therefore, the conjecture can be expressed the following way :

$$LG^{m,n}(L; t_0, e^{2i\pi/n} t_0^{-1}) = \Delta_L(t_0^n)^m, \text{ for any link } L.$$

We can now explore in which cases it seems reasonable to attempt to generalize the strategy we used to evaluate the reduction of  $LG^{2,1}$ . An obvious obstruction to that concerns the dimension of both representations we built and showed they were isomorphic. Let's

calculate the dimensions of the natural generalizations of these representations in case  $(m, n)$ .

The vector space corresponding to what we denoted  $W$  is the highest weight  $U_q(\mathfrak{gl}(m|n))$ -module used to define  $LG^{m,n}$ . It is  $2^{mm}$ -dimensional. So the representation of braid group  $B_p$  defined thanks to the corresponding R-matrix is  $2^{mp}$ -dimensional. On the other hand, the representation of  $B_p$  we want to define to produce  $\Delta_L(t_0^m)$  is :

$$\bigwedge \underbrace{(W_p \oplus \dots \oplus W_p)}_{m \text{ times}}$$

where each  $W_p$  is a  $\mathbb{C}[B_p]$ -module isomorphic to a version of  $\Psi_{W_p}$  where  $t_0$  is replaced by  $t_0^{\pm n}$ . Such a representation is  $2^{mp}$ -dimensional. These two representations can not be isomorphic if  $n > 1$ .

That is why a straightforward use of our method can only be applied to prove cases  $(m, 1)$ .

**4.2. Proof of case  $(3, 1)$ .** We give the essential steps to prove the result that interests us in the case  $(m, n) = (3, 1)$ . We follow the same ideas we used to study  $LG^{2,1}$ .

**Theorem 4.1.** *For any oriented link  $L$ ,  $LG^{3,1}(L; t_0, t_0^{-1}) \stackrel{\bullet}{=} \Delta_L(t_0)^3$ .*

For an explicit definition of  $LG^{3,1}$ , see [3], p.17. The author uses variables  $(\tau, q)$ , but denotes  $\tau = q^{-\alpha}$ . We will only use the reduced version of  $LG^{3,1}$ . It is obtained by setting  $q = -1$  and  $q^{-\alpha} = t_0^{1/2}$ .

*Remark 4.2.* Since we are going to set  $q = -1$  in the R-matrix of [3], we have to chose precisely what the roots that are written formally are. We have chosen :  $[\alpha + 1]^{1/2} = q^{-1/2}[\alpha]^{1/2}$  and  $[\alpha + 2]^{1/2} = -[\alpha]^{1/2}$ .

**Definition 4.3.** (*R-matrix  $S$* )

Set  $\mathbb{F} := \mathbb{C}(t_0^{\pm 1/2})$ . Let  $W = \langle e_1, \dots, e_8 \rangle$  be a 8-dimensional  $\mathbb{F}$ -vector space. We define  $S$  an automorphism of  $W \otimes W$  as the direct sum of the following automorphisms ( $S$  is globally multiplied by  $t_0^{-3/2}$  in comparison with the R-matrix explicited in [3]) :

$$\begin{pmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & -t_0^{-1} & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & -t_0^{-1} & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & -t_0^{-1} & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & t_0^{-2} & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & t_0^{-2} & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & t_0^{-2} & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & -t_0^{-3} \end{pmatrix}$$

in basis  $(e_1 \otimes e_1, e_2 \otimes e_2, e_3 \otimes e_3, e_4 \otimes e_4, e_5 \otimes e_5, e_6 \otimes e_6, e_7 \otimes e_7, e_8 \otimes e_8)$  ;

several copies of

$$\begin{pmatrix} 0 & t_0^{-1/2} \\ t_0^{-1/2} & 1 - t_0^{-1} \end{pmatrix}$$

in bases  $(e_1 \otimes e_2, e_2 \otimes e_1)$ ,  $(e_1 \otimes e_3, e_3 \otimes e_1)$  and  $(e_1 \otimes e_4, e_4 \otimes e_1)$  ;

several copies of

$$t_0^{-2} \begin{pmatrix} 0 & t_0^{-1/2} \\ t_0^{-1/2} & 1 - t_0^{-1} \end{pmatrix}$$

in bases  $(e_7 \otimes e_8, e_8 \otimes e_7)$ ,  $(e_6 \otimes e_8, e_8 \otimes e_6)$  and  $(e_5 \otimes e_8, e_8 \otimes e_5)$  ;

several copies of

$$-t_0^{-1} \begin{pmatrix} 0 & t_0^{-1/2} \\ t_0^{-1/2} & 1 - t_0^{-1} \end{pmatrix}$$

in bases  $(e_2 \otimes e_5, e_5 \otimes e_2)$ ,  $(e_3 \otimes e_5, e_5 \otimes e_3)$ ,  $(e_2 \otimes e_6, e_6 \otimes e_2)$ ,  $(e_4 \otimes e_6, e_6 \otimes e_4)$ ,  $(e_3 \otimes e_7, e_7 \otimes e_3)$  and  $(e_4 \otimes e_7, e_7 \otimes e_4)$  ;

several copies of

$$t_0^{-1} \begin{pmatrix} \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & 1 & t_0^{1/2} - t_0^{-1/2} \\ \cdot & 1 & \cdot & t_0^{1/2} - t_0^{-1/2} \\ 1 & t_0^{1/2} - t_0^{-1/2} & t_0^{1/2} - t_0^{-1/2} & (t_0^{1/2} - t_0^{-1/2})^2 \end{pmatrix}$$

in bases  $(e_1 \otimes e_5, e_2 \otimes e_3, e_3 \otimes e_2, e_5 \otimes e_1)$ ,  $(e_1 \otimes e_6, e_2 \otimes e_4, e_4 \otimes e_2, e_6 \otimes e_1)$  and  $(e_1 \otimes e_7, e_3 \otimes e_4, e_4 \otimes e_3, e_7 \otimes e_1)$  ;

several copies of

$$-t_0^{-2} \begin{pmatrix} \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & 1 & t_0^{1/2} - t_0^{-1/2} \\ \cdot & 1 & \cdot & t_0^{1/2} - t_0^{-1/2} \\ 1 & t_0^{1/2} - t_0^{-1/2} & t_0^{1/2} - t_0^{-1/2} & (t_0^{1/2} - t_0^{-1/2})^2 \end{pmatrix}$$

in bases  $(e_4 \otimes e_8, e_6 \otimes e_7, e_7 \otimes e_6, e_8 \otimes e_4)$ ,  $(e_3 \otimes e_8, e_5 \otimes e_7, e_7 \otimes e_5, e_8 \otimes e_3)$  and  $(e_2 \otimes e_8, e_5 \otimes e_6, e_6 \otimes e_5, e_8 \otimes e_2)$  ;

$$t_0^{-3/2} \begin{pmatrix} \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & t_0^{1/2} - t_0^{-1/2} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & t_0^{1/2} - t_0^{-1/2} \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & t_0^{1/2} - t_0^{-1/2} \\ \cdot & \cdot & \cdot & 1 & \cdot & t_0^{1/2} - t_0^{-1/2} & \cdot & t_0^{1/2} - t_0^{-1/2} & (t_0^{1/2} - t_0^{-1/2})^2 \\ \cdot & \cdot & 1 & \cdot & \cdot & t_0^{1/2} - t_0^{-1/2} & \cdot & t_0^{1/2} - t_0^{-1/2} & (t_0^{1/2} - t_0^{-1/2})^2 \\ \cdot & 1 & \cdot & \cdot & \cdot & t_0^{1/2} - t_0^{-1/2} & t_0^{1/2} - t_0^{-1/2} & \cdot & (t_0^{1/2} - t_0^{-1/2})^2 \\ 1 & t_0^{1/2} - t_0^{-1/2} & t_0^{1/2} - t_0^{-1/2} & t_0^{1/2} - t_0^{-1/2} & (t_0^{1/2} - t_0^{-1/2})^2 & (t_0^{1/2} - t_0^{-1/2})^2 & (t_0^{1/2} - t_0^{-1/2})^2 & (t_0^{1/2} - t_0^{-1/2})^2 & (t_0^{1/2} - t_0^{-1/2})^3 \end{pmatrix}$$

in basis  $(e_1 \otimes e_8, e_4 \otimes e_5, e_3 \otimes e_6, e_2 \otimes e_7, e_7 \otimes e_2, e_6 \otimes e_3, e_5 \otimes e_4, e_8 \otimes e_1)$ .

Then  $S$  is an R-matrix. So we can denote by  $b_S^n$  the representation of braid group  $B_n$  derived from  $S$ . It is given by the usual expression :

$$b_S^n(\sigma_i) = id_W^{\otimes i-1} \otimes S \otimes id_W^{\otimes n-i-1}, \quad i = 1, \dots, n-1.$$

**Definition 4.4.** *Reduced Links-Gould invariant  $LG^{3,1}$*

Let  $L$  be any oriented link, and  $b \in B_n$  be a braid with closure  $L$ . The reduced version of Links-Gould invariant  $LG^{3,1}$  is given by the following formula :

$$LG^{3,1}(L; t_0, t_0^{-1}) = \frac{1}{8} \text{trace}((id_W \otimes \mu^{\otimes n-1}) \circ b_S^n(b))$$

where

$$\mu = t_0^{3/2} \begin{pmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & -1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & -1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & -1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & -1 \end{pmatrix} \in \text{End}(W).$$

We set three  $n$ -dimensional vector spaces  $\langle f_1, \dots, f_n \rangle$ ,  $\langle g_1, \dots, g_n \rangle$  and  $\langle h_1, \dots, h_n \rangle$  that will be all referred to as  $W_n$ . On each of them, we define a representation isomorphic to the Burau representation :

$$F(\sigma_i)(f_j) = \begin{cases} t_0^{-1/2} f_{i+1} & \text{if } j = i, \\ t_0^{-1/2} f_i + (1 - t_0^{-1}) f_{i+1} & \text{if } j = i + 1, \\ f_j & \text{otherwise.} \end{cases}$$

We designate by  $G$  and  $H$  representations on  $\langle g_1, \dots, g_n \rangle$  and  $\langle h_1, \dots, h_n \rangle$  defined by the exact same formula. Then we set  $\Phi_n$  the representation of  $B_n$  on  $\bigwedge(W_n \oplus W_n \oplus W_n)$  given by :

$$\Phi_n := \bigwedge(F \oplus G \oplus H).$$

When  $n = 2$ , one can compute  $\Phi_2(\sigma_1)$  and notice that its matrix is equal to  $S$  in a well chosen basis. A precise look at this basis gave us the idea to define the following map by induction. Note that retrospectively one can recover this basis simply by computing the image by our map of the basis we used to express  $S$  when  $n = 2$ .

$$\text{Theorem 4.5. We set } I_1 = \begin{cases} V & \longrightarrow \bigwedge W_1 \\ e_1 & \longmapsto 1 \\ e_2 & \longmapsto f_1 \\ e_3 & \longmapsto g_1 \\ e_4 & \longmapsto h_1 \\ e_5 & \longmapsto f_1 \wedge g_1 \\ e_6 & \longmapsto f_1 \wedge h_1 \\ e_7 & \longmapsto g_1 \wedge h_1 \\ e_8 & \longmapsto f_1 \wedge g_1 \wedge h_1 \end{cases} \quad \text{and, for } n \geq 2,$$

$$I_n = \begin{cases} V^{\otimes n} & \longrightarrow \bigwedge W_n \\ e_1 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \\ e_2 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto \text{Reord}(I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \wedge f_n) \\ e_3 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto \text{Reord}(I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \wedge g_n) \\ e_4 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \wedge h_n \\ e_5 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto \text{Reord}(I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \wedge f_n \wedge g_n) \\ e_6 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto \text{Reord}(I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \wedge f_n \wedge h_n) \\ e_7 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto \text{Reord}(I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \wedge g_n \wedge h_n) \\ e_8 \otimes e_{i_{n-1}} \otimes \dots \otimes e_{i_1} & \longmapsto \text{Reord}(I_{n-1}(e_{i_{n-1}} \otimes \dots \otimes e_{i_1}) \wedge f_n \wedge g_n \wedge h_n) \end{cases}.$$

Then the following identity holds for  $n \geq 1$  and  $b \in B_n$  :

$$\Phi_n(b) \circ I_n = I_n \circ b_S^n(\hat{b}).$$

*Remark 4.6.* As in the previous sections, *Reord* refers to a reference basis of  $\bigwedge(W_n \oplus W_n \oplus W_n)$  that is :

$$(f_{i_1} \wedge \dots \wedge f_{i_p} \wedge g_{j_1} \wedge \dots \wedge g_{j_m} \wedge h_{k_1} \wedge \dots \wedge h_{k_q})_{1 \leq i_1 < \dots < i_p \leq n, 1 \leq j_1 < \dots < j_m \leq n, 1 \leq k_1 < \dots < k_q \leq n}$$

*Remark 4.7.* For  $b = \sigma_{i_1}^{\varepsilon_1} \dots \sigma_{i_p}^{\varepsilon_p} \in B_n$ , we define  $\hat{b} := \sigma_{n-i_1}^{\varepsilon_1} \dots \sigma_{n-i_p}^{\varepsilon_p}$ .  $\hat{b}$  is braid  $b$  "looked at from the other side". That way we have elementary properties :  $\text{closure}(b) = \text{closure}(\hat{b})$  ;  $\hat{\sigma}_k = \sigma_{n-k}$  ; for any  $\sigma, \tau \in B_n$  :  $\hat{\sigma\tau} = \hat{\sigma}\hat{\tau}$ .

We can use  $I_n$  to express  $LG^{3,1}$  differently.

$$\begin{aligned} LG^{3,1}(L; t_0, t_0^{-1}) &= \frac{1}{8} \text{trace}((id_W \otimes \mu^{\otimes n-1}) \circ b_S^n(b)) \\ &= \frac{1}{8} \text{trace}(\underbrace{I_n \circ (id_W \otimes \mu^{\otimes n-1}) \circ I_n^{-1}}_{\tilde{\mu}} \circ \Phi_n(\hat{b})). \end{aligned}$$

Denoting as we already did several times  $I_n(e_{i_n} \otimes \dots \otimes e_{i_1}) = f_{i_1} \wedge \dots \wedge h_{k_q}$ , we can compute  $\tilde{\mu}$  :

$$\begin{aligned} \tilde{\mu}(f_{i_1} \wedge \dots \wedge h_{k_q}) &= I_n \circ (id_W \otimes \mu^{\otimes n-1})(e_{i_n} \otimes \dots \otimes e_{i_1}) \\ &= t_0^{3(n-1)/2} (-1)^{\#\{k \in \{1, \dots, n-1\} | i_k \in \{2, 3, 4, 8\}\}} I_n(e_{i_n} \otimes \dots \otimes e_{i_1}) \\ &= t_0^{3(n-1)/2} (-1)^{\#\{k \in \{1, \dots, n-1\} | \text{an odd number of the following appear : } \{f_k, g_k, h_k\}\}} \\ &\quad f_{i_1} \wedge \dots \wedge h_{k_q} \\ &= t_0^{3(n-1)/2} (-1)^{\#\{k \in \{1, \dots, n-1\} | f_k \text{ appears}\}} (-1)^{\#\{k \in \{1, \dots, n-1\} | g_k \text{ appears}\}} \\ &\quad (-1)^{\#\{k \in \{1, \dots, n-1\} | h_k \text{ appears}\}} f_{i_1} \wedge \dots \wedge h_{k_q}. \end{aligned}$$

So

$$\begin{aligned} LG^{3,1}(L; t_0, t_0^{-1}) &= \frac{1}{8} \text{trace}(\tilde{\mu} \circ \Phi_n(\hat{b})) \\ &= \frac{1}{8} \sum_{\substack{1 \leq i_1 < \dots < i_p \leq n \\ 1 \leq j_1 < \dots < j_m \leq n \\ 1 \leq k_1 < \dots < k_q \leq n}} (f_{i_1} \wedge \dots \wedge h_{k_q})^* (\tilde{\mu} \circ \Phi_n(\hat{b})(f_{i_1} \wedge \dots \wedge h_{k_q})) \end{aligned}$$

But

$$\begin{aligned} &(f_{i_1} \wedge \dots \wedge h_{k_q})^* (\tilde{\mu} \circ \Phi_n(\hat{b})(f_{i_1} \wedge \dots \wedge h_{k_q})) \\ &= t_0^{3(n-1)/2} (-1)^{\#\{k \in \{1, \dots, n-1\} | f_k \text{ appears}\}} (-1)^{\#\{k \in \{1, \dots, n-1\} | g_k \text{ appears}\}} \\ &\quad (-1)^{\#\{k \in \{1, \dots, n-1\} | h_k \text{ appears}\}} (f_{i_1} \wedge \dots \wedge h_{k_q})^* (\Phi_n(\hat{b})(f_{i_1} \wedge \dots \wedge h_{k_q})) \\ &= t_0^{3(n-1)/2} (-1)^{\dots} (f_{i_1} \wedge \dots \wedge f_{i_p})^* (\bigwedge F(\hat{b})(f_{i_1} \wedge \dots \wedge f_{i_p})) \\ &\quad (g_{j_1} \wedge \dots \wedge g_{j_m})^* (\bigwedge G(\hat{b})(g_{j_1} \wedge \dots \wedge g_{j_m})) (h_{k_1} \wedge \dots \wedge h_{k_q})^* (\bigwedge H(\hat{b})(h_{k_1} \wedge \dots \wedge h_{k_q})) \end{aligned}$$

So finally

$$\begin{aligned} LG^{3,1}(L; t_0, t_0^{-1}) &= t_0^{3(n-1)/2} \left( \frac{1}{2} \sum_{1 \leq i_1 < \dots < i_p \leq n} (-1)^{\#\{k \in \{1, \dots, n-1\} | f_k \text{ appears}\}} \right. \\ &\quad \left. (f_{i_1} \wedge \dots \wedge f_{i_p})^* (\bigwedge F(\hat{b})(f_{i_1} \wedge \dots \wedge f_{i_p})) \right) * \dots \end{aligned}$$

The only thing that remains to be shown is that each of the three terms in the product is equal to  $\Delta_{\hat{b}}(t) = \Delta_b(t)$ . The proof is similar to the one we did for  $LG^{2,1}$ . The main



point is to find R-matrices associated to their representations on  $V^{\otimes n}$  such that

$$\Delta_L(t) = \frac{1}{2} \text{trace}((id_V \otimes h^{\otimes n-1}) \circ \Psi_{V^{\otimes n}}(b))$$

and that up to conjugation the trace is one of the three sums. We will not detail this argument.

4.3. **A remark around case  $(n, 1)$ .** To prove the identity

$$LG^{n,1}(L; t_0, t_0^{-1}) \stackrel{\bullet}{=} \Delta_L(t_0)^n$$

when  $n = 2, 3$ , we have used the crucial fact that we know an explicit formula for the R-matrix and the left handle (the maps we called  $\mu$ ) in these two cases. Solving the conjecture for any  $n$  using the same ideas therefore requires the R-matrix to be computed in all cases. In [3], the calculations are explicit up to  $n = 4$ . However, we believe it is possible, with a proper amount of sweat and will, to give a formula for any  $n$ . Indeed,  $U_q(gl(n|1))$  is a quantum super-algebra and one can find expressions for universal R-matrices in that context in [7] or [14]. More recently, M.D. Gould, P.S. Isaac and J.L. Werry wrote the representation that derives the R-matrix from it's universal counterpart in a practical basis [6]. This allows to project the universal R-matrix to find the corresponding map.

## 5. APPENDIX : PROOF OF PROPOSITION 2.4

Here we prove the result we stated in proposition 2.4. That is : for any  $b \in B_n$ ,

$$\Psi_n(b) \circ I_n = I_n \circ b_R^n(b).$$

*Proof.* We show the commutation by induction on  $n$ , the number of strands in the braid group we consider. Note that it has already been verified when  $n = 1, 2$ . Let us now suppose the equality holds for  $n - 1$ ,  $n \geq 3$ . We only need to prove the result for  $b = \sigma_k$ ,  $k = 1, \dots, n - 1$ .

For  $\sigma_k$ ,  $k \leq n - 2$  :

$$\begin{aligned} & I_n(b_R^n(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_n})) \\ &= I_n(e_{i_1} \otimes \dots \otimes R(e_{i_k} \otimes e_{i_{k+1}}) \otimes \dots \otimes e_{i_n}) \\ &= I_n(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \otimes e_{i_n}) \\ &= \begin{cases} I_{n-1}(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge g_n & \text{if } i_n = 1 \\ I_{n-1}(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) & \text{if } i_n = 2 \\ \text{Reord}(I_{n-1}(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n) & \text{if } i_n = 3 \\ \text{Reord}(I_{n-1}(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n) & \text{if } i_n = 4 \end{cases} \\ &= \begin{cases} \Psi_{n-1}(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge g_n & \text{if } i_n = 1 \\ \Psi_{n-1}(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) & \text{if } i_n = 2 \\ \text{Reord}(I_{n-1}(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n) & \text{if } i_n = 3 \\ \text{Reord}(I_{n-1}(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n) & \text{if } i_n = 4 \end{cases} \quad (\text{inductive hypothesis}) \end{aligned}$$

On the other hand :

$$\begin{aligned} & \Psi_n(\sigma_k)(I_n(e_{i_1} \otimes \dots \otimes e_{i_n})) \\ &= \begin{cases} \Psi_n(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge g_n) & \text{if } i_n = 1 \\ \Psi_n(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) & \text{if } i_n = 2 \\ \Psi_n(\sigma_k)(\text{Reord}(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge f_n \wedge g_n)) & \text{if } i_n = 3 \\ \Psi_n(\sigma_k)(\text{Reord}(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge f_n)) & \text{if } i_n = 4 \end{cases} \end{aligned}$$

$$= \begin{cases} \Psi_n(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge \Psi_n(\sigma_k)(g_n) & \text{if } i_n = 1 \\ \Psi_n(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) = \Psi_{n-1}(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) & \text{if } i_n = 2 \\ \Psi_n(\sigma_k)(\text{Reord}(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n) & \text{if } i_n = 3 \\ \Psi_n(\sigma_k)(\text{Reord}(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n) & \text{if } i_n = 4 \end{cases}$$

For  $i_n = 1$ , since  $\Psi_n(\sigma_k)(g_n) = g_n$ , we obtain the result. We also observe the equality holds when  $i_n = 2$ . We now study the two remaining cases.

Let  $\mu(e_{i_1} \otimes \dots \otimes e_{i_n})$  be the total of the number of  $e_1$  and the number of  $e_3$  in that elementary tensor. Given the expression of  $I_n$  and our reference basis of  $\bigwedge(W_n \oplus W_n)$ , it is obvious that :

$$\text{Reord}(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})} I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge f_n$$

and

$$\text{Reord}(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})} I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge f_n \wedge g_n.$$

Therefore :

$$\begin{aligned} & \Psi_n(\sigma_k)(\text{Reord}(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n) \\ &= (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})} \Psi_n(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n \\ &= (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})} \Psi_{n-1}(\sigma_k)(I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n \end{aligned}$$

Moreover, given the specific form of matrix  $R$ , every term that appears in  $b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})$  has the same total number of  $e_1$  and  $e_3$  as  $e_{i_1} \otimes \dots \otimes e_{i_{n-1}}$ . Hence :

$$\begin{aligned} & \text{Reord}(I_{n-1}(b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})) \wedge f_n \wedge g_n) \\ &= (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})} I_{n-1} \circ b_R^{n-1}(\sigma_k)(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge f_n \wedge g_n \\ &= (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-1}})} \Psi_{n-1}(\sigma_k) \circ I_{n-1}(e_{i_1} \otimes \dots \otimes e_{i_{n-1}}) \wedge f_n \wedge g_n \end{aligned}$$

Thus we obtain the identity in case  $i_n = 3$ . Similar calculations show it is also true when  $i_n = 4$ . The only remaining question is for the last generator of  $B_n$ .

For  $\sigma_{n-1}$  : We show that  $\Psi_n(\sigma_{n-1}) \circ I_n(e_{i_1} \otimes \dots \otimes e_{i_n}) = I_n \circ b_R^n(\sigma_{n-1})(e_{i_1} \otimes \dots \otimes e_{i_n})$  for each of the 16 possible ordered pairs  $(i_{n-1}, i_n)$  :

$$\begin{aligned} & \boxed{(1,1)} : \\ & \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_1 \otimes e_1)) \\ &= \Psi_n(\sigma_{n-1})(I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge g_{n-1} \wedge g_n) \\ &= I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge (-t_0^{1/2} g_n) \wedge (-t_0^{1/2} g_{n-1} + (1-t_0)g_n) \\ &= -t_0 I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge g_{n-1} \wedge g_n \end{aligned}$$

$$\begin{aligned} & I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_1 \otimes e_1) \\ &= I_n(e_{i_1} \otimes \dots \otimes e_{i_{n-2}} \otimes -t_0 e_1 \otimes e_1) \\ &= -t_0 I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge g_{n-1} \wedge g_n \end{aligned}$$

Now that we have explicitated one case, we give the results for the remaining ones.

$$\begin{aligned} & \boxed{(4,4)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_4 \otimes e_4)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_4 \otimes e_4) \\ &= -t_0^{-1} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge f_{n-1} \wedge f_n \end{aligned}$$

$$\boxed{(2,2)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_2 \otimes e_2)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_2 \otimes e_2) \\ = I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})$$

$$\boxed{(3,3)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_3 \otimes e_3)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_3 \otimes e_3) \\ = I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge f_{n-1} \wedge f_n \wedge g_{n-1} \wedge g_n$$

$$\boxed{(1,2)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_1 \otimes e_2)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_1 \otimes e_2) \\ = -t_0^{1/2} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge g_n$$

$$\boxed{(2,1)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_2 \otimes e_1)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_2 \otimes e_1) \\ = I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge (-t_0^{1/2} g_{n-1} + (1 - t_0) g_n)$$

$$\boxed{(1,3)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_1 \otimes e_3)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_1 \otimes e_3) \\ = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}} \otimes e_1)} t_0^{1/2} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge f_{n-1} \wedge g_{n-1} \wedge g_n$$

$$\boxed{(3,1)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_3 \otimes e_1)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_3 \otimes e_1) \\ = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge ((1 - t_0) f_{n-1} \wedge g_{n-1} \wedge g_n - t_0^{1/2} f_n \wedge g_{n-1} \wedge g_n)$$

$$\boxed{(3,4)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_3 \otimes e_4)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_3 \otimes e_4) \\ = t_0^{-1/2} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge f_{n-1} \wedge f_n \wedge g_n$$

$$\boxed{(4,3)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_4 \otimes e_3)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_4 \otimes e_3) \\ = I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge (t_0^{-1/2} f_{n-1} \wedge f_n \wedge g_{n-1} + (1 - t_0^{-1}) f_{n-1} \wedge f_n \wedge g_n)$$

$$\boxed{(2,4)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_2 \otimes e_4)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_2 \otimes e_4) \\ = t_0^{-1/2} (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge f_{n-1}$$

$$\boxed{(4,2)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_4 \otimes e_2)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_4 \otimes e_2) \\ = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge ((1 - t_0^{-1}) f_{n-1} + t_0^{-1/2} f_n)$$

$$\boxed{(1,4)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_1 \otimes e_4)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_1 \otimes e_4) \\ = -(-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge f_{n-1} \wedge g_n$$

$$\boxed{(2,3)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_2 \otimes e_3)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_2 \otimes e_3) \\ = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge (-f_{n-1} \wedge g_{n-1} - Y f_{n-1} \wedge g_n)$$

$$\boxed{(3,2)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_3 \otimes e_2)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_3 \otimes e_2) \\ = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge (-Y f_{n-1} \wedge g_n - f_n \wedge g_n)$$

$$\boxed{(4,1)} : \Psi_n(\sigma_{n-1})(I_n(e_{i_1} \otimes \dots \otimes e_4 \otimes e_1)) = I_n \circ (1 \otimes 1 \otimes \dots \otimes R)(e_{i_1} \otimes \dots \otimes e_4 \otimes e_1) \\ = (-1)^{\mu(e_{i_1} \otimes \dots \otimes e_{i_{n-2}})} I_{n-2}(e_{i_1} \otimes \dots \otimes e_{i_{n-2}}) \wedge (-Y f_{n-1} \wedge g_{n-1} - f_n \wedge g_{n-1} - Y^2 f_{n-1} \wedge g_n - \\ Y f_n \wedge g_n)$$

Which ends the proof.

CQFD

ACKNOWLEDGMENTS. I would like to thank my thesis advisor Emmanuel Wagner very warmly for his kind help and his sound advice, and for pointing me to the right direction in times of doubt.

#### REFERENCES

- [1] D. De Wit, A. Ishii, J. Links. *Infinitely many two-variable generalisations of the Alexander-Conway polynomial*. Algebraic and Geometric Topology, Volume 5, 405-418, 2005.
- [2] D. De Wit, L.H. Kauffman, J. Links. *On the Links-Gould invariant of links*. J. Knot Theory Ramifications. 8 (1999), no.2, 165-199. MR 2000j:57020
- [3] D. De Wit. *An infinite suite of Links-Gould invariants*. J. Knot Theory Ramifications. 10 (2001), no.1, 37-62. MR 2002e:57015
- [4] R.H. Fox, G. Torres. *Dual presentations of the group of a knot*. The Annals of Mathematics. Second Series, vol.59, no.2 (Mar., 1954), 211-218.
- [5] N. Geer, B. Patureau-Mirand. *Multivariable link invariants arising from  $sl(2|1)$  and the Alexander polynomial*. Journal of Pure and Applied Algebra, Volume 210, Issue 1, July 2007, 283-298.
- [6] M.D. Gould, P.S. Isaac, J.L. Werry. *Matrix elements for type 1 unitary irreducible representations of the Lie superalgebra  $gl(m|n)$* . Journal of Mathematical Physics, 55 011703: 011703.1-011703.32. doi:10.1063/1.4861706, 2014.
- [7] S.M. Khoroshkin, V.N. Tolstoy. *Universal R-matrix for Quantized (Super)Algebras*. Communications in Mathematical Physics. no.141 (1991), 599-617.
- [8] W.B.R. Lickorish. *An Introduction to Knot Theory*. Graduate Texts in Mathematics, Vol. 175. Springer, 1997.
- [9] J. Links, M.D. Gould. *Two variable link polynomials from quantum supergroups*. Letters in Mathematical Physics, 26(3):187-198, November 1992.
- [10] I. Marin, E. Wagner. *A cubic defining algebra for the Links-Gould polynomial*. Advances in Math., 248 (2013), 1332-1365.
- [11] T. Ohtsuki. *Quantum invariants. A study of knots, 3-manifolds, and their sets*. Number 29 in Series on Knots and Everything. World Scientific, 2002.
- [12] N. Reshetikhin, C. Stroppel, B. Webster. *Schur-Weyl-Type duality for quantized  $gl(1|1)$ , the Burau representation of braid groups, and invariants of tangled graphs*. I. Itenberg et al. (eds.), Perspectives in Analysis, Geometry, and Topology: On the Occasion of the 60th Birthday of Oleg Viro, Progress in Mathematics 296.
- [13] O. Viro. *Quantum relatives of the Alexander polynomial*. St. Petersburg Math. J., Vol. 18 (2007), no.3, 391-457.
- [14] H. Yamane. *Quantized Enveloping Algebras Associated with Simple Lie Superalgebras and Their Universal R-matrices*. Publ. RIMS, Kyoto University, vol.30 (1994), 15-87.

(Ben-Michael Kohli) IMB UMR5584, CNRS, UNIVERSITÉ BOURGOGNE FRANCHE-COMTÉ, F-21000 DIJON, FRANCE.

*E-mail address:* Ben-Michael.Kohli@u-bourgogne.fr